



2nd International Congress on

SUSTAINABLE AGRICULTURE

March 04-05, 2025 / Iğdır University, Türkiye



EDITORS

Prof. Dr. Mehmet Hakkı ALMA

Prof. Dr. Sefa ALTIKAT

ISBN: 979-8-89695-046-2

INTERNATIONAL CONGRESS ON SUSTAINABLE AGRICULTURE-II

March 04-05, 2025 / Iğdır University, Türkiye

20.03.2025

Liberty Publishing House

Water Street Corridor New York, NY 10038

www.libertyacademicbooks.com

+1 (314) 597-0372

ALL RIGHTS RESERVED NO PART OF THIS BOOK MAY BE REPRODUCED IN ANY FORM, BY PHOTOCOPYING OR BY ANY ELECTRONIC OR MECHANICAL MEANS, INCLUDING INFORMATION STORAGE OR RETRIEVAL SYSTEMS, WITHOUT PERMISSION IN WRITING FROM BOTH THE COPYRIGHT OWNER AND THE PUBLISHER OF THIS BOOK.

© Liberty Academic Publishers 2025

The digital PDF version of this title is available Open Access and distributed under the terms of the Creative Commons Attribution-Non-Commercial 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits adaptation, alteration, reproduction and distribution for noncommercial use, without further permission provided the original work is attributed. The derivative works do not need to be licensed on the same terms.

adopted by Mariam Rasulan

PROCEEDINGS BOOK

ISBN: 979-8-89695-046-2

CONGRESS ID

CONGRESS TITLE

INTERNATIONAL CONGRESS ON SUSTAINABLE AGRICULTURE-II

DATE AND PLACE

March 04-05, 2025 / Iğdır University, Türkiye

ORGANIZATION

IĞDIR UNIVERSITY
NAKHIVAN STATE UNIVERSITY
ARTVİN ÇORUH UNIVERSITY
IKSAD INSTITUTE

EDITORS

Prof. Dr. Mehmet Hakkı ALMA

Prof. Dr. Sefa ALTİKAT

COORDINATOR

Samet KUŞKIRAN

PARTICIPANTS COUNTRY (16 countries)

TÜRKİYE, MOROCCO, ALGERIA, AZERBAIJAN, FRANCE, TUNUSIA, PAKISTAN, SPAIN,
PHILIPPINES, INDIA, IRAQ, AUSTRALIA, CHINA, ALBANIA, VIETNAM, ROMANIA

Total Accepted Article: 173

Total Rejected Papers: 46

Accepted Article (Türkiye): 82

Accepted Article (Other Countries): 91

ISBN: 979-8-89695-046-2

2. INTERNATIONAL CONGRESS ON SUSTAINABLE AGRICULTURE

March 04-05, 2025 / Iğdır University, Türkiye



20.03.2025

REF: Akademik Teşvik

İlgili makama;

2. Uluslararası Sürdürülebilir Tarım Kongresi, 4-5 Mart 2025 tarihleri arasında Iğdır'da 16 farklı ülkenin (Türkiye 82 bildiri- Diğer ülkeler 91 bildiri) akademisyen/araştırmacılarının katılımıyla gerçekleşmiştir

Kongre 16 Ocak 2020 Akademik Teşvik Ödeneği Yönetmeliğine getirilen "Tebliğlerin sunulduğu yurt içinde veya yurt dışındaki etkinliğin uluslararası olarak nitelendirilebilmesi için Türkiye dışında en az beş farklı ülkeden sözlü tebliğ sunan konuşmacının katılım sağlaması ve tebliğlerin yarısından fazlasının Türkiye dışından katılımcılar tarafından sunulması esastır." değişikliğine uygun düzenlenmiştir.

Bilgilerinize arz edilir,

Saygılarımla

Prof. Dr. Sefa ALTİKAT
Iğdır University, Faculty of Agriculture Dean
CONGRESS PRESIDENT

T.C
IĞDIR ÜNİVERSİTESİ REKTÖRLÜĞÜ
Ziraat Fakültesi Dekanlığı



Sayı : E-26144459-000-168360
Konu : Kongre düzenleme kurulu

21.03.2025

İlgili Makama

4-5 Mart 2025 tarihleri arasında düzenlenen 2. Uluslararası Sürdürülebilir Tarım Kongresinde aşağıda unvan ve isimleri yazılı olan öğretim elemanları kongre düzenleme kurulunda görevlendirilmiştir.

Gereğini bilgilerinize arz/rica ederim

Düzenleme Kurulu

Prof. Dr. İbrahim DEMİRTAŞ - Iğdır University, Türkiye
Prof. Dr. Ahmet Zafer TEL - Iğdır University, Türkiye
Prof. Dr. Uğur ŞİMŞEK - Iğdır University, Türkiye
Prof. Dr. Celalettin GÖZÜAÇIK - Iğdır University, Türkiye
Assoc. Prof. Dr. Tuncay KAYA - Iğdır University, Türkiye
Assoc. Prof. Dr. Ramazan GÜRBÜZ - Iğdır University, Türkiye
Assoc. Prof. Dr. Elsever Esedov - Nakhcivan State University, Azerbaijan
Assoc. Prof. Dr. Behruz Memmedov - Nakhcivan State University, Azerbaijan
Assoc. Prof. Dr. Mahir Meherremov - Nakhcivan State University, Azerbaijan
Assoc. Prof. Dr. Hilal Qasimov - Nakhcivan State University, Azerbaijan
Assoc. Prof. Dr. Elji Tahirov - Nakhcivan State University, Azerbaijan
Assoc. Prof. Dr. Mirmahmud Seyidli - Nakhcivan State University, Azerbaijan
Assoc. Prof. Dr. İsmail ALASERHAT - Iğdır University, Türkiye
Assoc. Prof. Dr. Adnan AYDIN - Iğdır University, Türkiye
Assist. Prof. Dr. Barış EREN - Iğdır University, Türkiye

Prof. Dr. Sefa ALTİKAT
Dekan



ORGANIZING COMMITTEE

Prof. Dr. Mehmet Hakkı ALMA

Rector of the Iğdır University

CONGRESS HONORARY PRESIDENT

Assoc. Prof. Dr. Elbrus ISAYEV

Rector of the Nakhcivan State University

CONGRESS HONORARY PRESIDENT

Prof. Dr. Mustafa Sıtkı BİLGİN

Rector of the Artvin Coruh University

CONGRESS HONORARY PRESIDENT

Dr. Mustafa Latif EMEK

President of IKSAD Institute

CONGRESS HONORARY PRESIDENT

Prof. Dr. Sefa ALTIKAT

Iğdır University, Faculty of Agriculture Dean

CONGRESS PRESIDENT

MEMBERS OF THE ORGANIZING COMMITTEE

Prof. Dr. İbrahim DEMİRTAŞ
İğdır University, Türkiye

Prof. Dr. Ahmet Zafer TEL
İğdır University, Türkiye

Prof. Dr. Uğur ŞİMŞEK
İğdır University, Türkiye

Prof. Dr. Celalettin GÖZÜAÇIK
İğdır University, Türkiye

Assoc. Prof. Dr. Tuncay KAYA
İğdır University, Türkiye

Assoc. Prof. Dr. Ramazan GÜRBÜZ
İğdır University, Türkiye

Assoc. Prof. Dr. Elsever Esedov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Behruz Memmedov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Mahir Meherremov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Hilal Qasimov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. E|i Tahirov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Mirmahmud Seyidli
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. İsmail ALASERHAT
İğdır University, Türkiye

Assoc. Prof. Dr. Adnan AYDIN
İğdır University, Türkiye

Assist. Prof. Dr. Barış EREN
İğdır University, Türkiye

SCIENTIFIC AND ADVISORY BOARD

Prof. Dr. Bilal KESKİN
İğdır University, Türkiye

Prof. Dr. Melekşen AKIN
İğdır University, Türkiye

Prof. Dr. Bünyamin YILDIRIM
İğdır University, Türkiye

Prof. Dr. Şükrü İsmail İPEK
İğdır University, Türkiye

Prof. Dr. Köksal KARADAŞ
İğdır University, Türkiye

Prof. Dr. Süleyman TEMEL
İğdır University, Türkiye

Assoc. Prof. Dr. Elsever Esedov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Mahir Meherremov
Nakhcivan State University, Azerbaijan

Prof. Dr. Etibar Memmedov
Nakhcivan State University, Azerbaijan

Prof. Dr. Daşqın Qenberov
Nakhcivan State University, Azerbaijan

Prof. Dr. Talibov Tariyel
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Yunis Rustemli
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Hilal Qasimov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Eli Tahirov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Hacıyev Sahib
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Namiq Abbasov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Tofiq Eliyev
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. İsmayil Memmedov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Enzale Novruzova
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Akif Merdanlı
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Feride Seferova
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Arzu Memmedov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Loğman Bayramov
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. İbrahimov Enver
Nakhcivan State University, Azerbaijan

Assoc. Prof. Dr. Cabbar Necefov
Nakhcivan State University, Azerbaijan

Dr. Behruz Bayramov
Nakhcivan State University, Azerbaijan

Dr. Mürsel Seyidov
Nakhcivan State University, Azerbaijan

Dr. Mirvasif Seyidov
Nakhcivan State University, Azerbaijan

Dr. Afaq Eliyeva
Nakhcivan State University, Azerbaijan

Dr. Aynur İbrahimova
Nakhcivan State University, Azerbaijan

Dr. Raife Salmanova
Nakhcivan State University, Azerbaijan

Dr. Huseynova Ezize
Nakhcivan State University, Azerbaijan

PHOTO GALLERY



PHOTO GALLERY





PHOTO GALLERY






PHOTO GALLERY





PHOTO GALLERY

Workplace Meeting JMAILI Karima's screen

HALL-4 SESSION-1 DI... HALL-4 -S_1 Dr...

JMAILI Karima H4-Sarra Farjallah Amor HALL-4 SESSION-1 DI... HALL-4 -S_1 Dr. Mazhar

HIA FST LBEA

2nd INTERNATIONAL CONGRESS ON SUSTAINABLE AGRICULTURE
March 04-05, 2025 / Iğdır University, Türkiye

Impact of whey on the germination of tomatoes and wheat seeds :
Toward the valorization of dairy by-products for sustainable agriculture

Presented by: JMAILI Karima

Participants (5)

- Observer H4 (Co-host, me)
- JMAILI Karima
- H4-Sarra Farjallah Amor
- HALL-4 -S_1 Dr. Mazhar
- HALL-4 SESSION-1 DIAB Nacima

Mute all

Workplace Meeting - HALL 3

H-3: Pinar ADIGÜZEL H-3 Ezgi Günay SARGUT H1-Celalettin Gozuack HALL-3 / SESSION-1 / Ali ÇELİK

HALL-3, İlayda Gizem AKIN H3-Nagihan Kılıç Dilek DOĞAN H-3 Gönül USTAGEBEŞ

Reyyan Yergin Ö... HALL-3, SEDA K... HALL-3, Serkan... Özge SIZMAZ

Reyyan Yergin Ozkan HALL-3, SEDA KAÇAR HALL-3, Serkan BENLİ Özge SIZMAZ

Mehmet hakkı i... Mahdaoui Yazid

Mehmet hakkı iPhone'u Mahdaoui Yazid

Participants (15)

- Observer H3 (Co-host, me)
- H1-Celalettin Gözüağık
- HALL-3, İlayda Gizem AKIN
- Dilek DOĞAN
- H-3 Ezgi Günay SARGUT
- H-3 Gönül USTAGEBEŞ
- H-3, Pinar ADIGÜZEL
- H3-Nagihan Kılıç
- HALL-3 / SESSION-1 / Ali ÇELİK
- HALL-3, SEDA KAÇAR
- HALL-3, Serkan BENLİ
- Mahdaoui Yazid
- Mehmet hakkı iPhone'u

Mute all

11:43 AM

PHOTO GALLERY

Workplace

Participants (14)

Find a participant

- DD Dilek DOĞAN
- HA Hall-3, Aydın Akkaya
- HU Hall-3-Tahsin Uygun
- AA ALEV ALDEMİR
- HG H1-Celalettin Gözüağık
- A H-3 Aslı İşler Kaya
- HE H-3 Ezgi Demiral
- HG H3 Gönül USTA GEBEŞ
- HR H-3, RIDVAN TIK
- HM Hall-3, Muhammet ÖTER
- MÇ MESUT ÇOŞLU
- MY Mete Yiğit
- Y yağmur

Mute all

Workplace

Participants (6)

- Observer H3 (Co-host, me)
- H-3 Res. Asst. Alperay ALTİKAT
- Z H3 zeynep Asutay
- HR H-3, RIDVAN TIK
- HK H3-Tuncay Kaya
- HG Hall-3, Gizem TIRVAKI

Mute all

PHOTO GALLERY




PHOTO GALLERY



PHOTO GALLERY




PHOTO GALLERY




PHOTO GALLERY

Workplace

observer hall-2 Adnan Aydın H-2, Fatih KUMBASAR köksal karadağ tugba hanedan

EPİGENETİK

Günümüzde, epigenetik terimi DNA dizisindeki değişikliklerden kaynaklanmayan kalıtsal gen ifadesindeki varyasyonları inceleyen bilim dalı olarak tanımlanmaktadır.

Canlılar alemi içerisinde bitkilerin doğada var oldukları çevreyi değiştiremedikleri, içinde buldukları değişken ve çoğunlukla olumsuz iklim koşullarıyla başa çıkmak zorunda kaldıkları için sahip oldukları epigenom ve epigenetik mekanizmalar sayesinde gen aktivitesinde önemli değişiklikler yapabildikleri ve öngörülmeleyen ortamlarda başarılı bir şekilde hayatta kaldıkları ayrıca bu epigenetik özellikleri kalıtım yoluyla aktararak nesillerini devam ettirebildikleri düşünülmektedir.



Zoom Workplace

Участники (12)

Найти участника

- OH observe... (Соорганизатор, я)
- A Adnan Aydın
- AB Anil Baltacı-Kayseri Şeker
- M h-2 Mehmet Can
- HZ h-2 Zeki Acar
- HF H-2, Fatih KUMBASAR
- G H-2, Gülcan Kaymak Bayram
- HALL 2 Halit Burak Yüksekdağ
- HY HALL-2 Yusuf GÜMÜŞ
- I İlknur AYAN
- KK köksal karadağ
- tugba hanedan

Выключить звук для всех

Workplace

observer hall-2 Hall-2, Aysun Altikat H-2, Zekiye ALTUNTEKİN Hall-2/Boran KARATAŞ Hall-2, Soner AKCAN

Nanopestisitler, genellikle aşağıdaki yollarla etki gösterir:

- Kontrollü salınım mekanizması:** Nanopartiküller, aktif bileşenleri zaman içinde salarak pestisit etkinliğini uzatır ve sık tekrar gereksinimini azaltır.
- Hedefe yönelik etki:** Nanoteknoloji sayesinde nanopestisitler belirli organizmalara özel olarak tasarlanabilir ve zararlı olmayan böceklere veya bitkilere zarar vermeden uygulanabilir.
- Biyomoleküllerle etkileşim:** Nanopestisitlerin yüzeyleri modifiye edilerek belirli zararlı böceklerin sindirim sistemine veya solunum sistemine daha etkin şekilde etki edebilir.
- Yapısal kararlılık:** Nanopartiküller, geleneksel pestisitlere göre daha stabil olup, sıcaklık ve pH değişikliklerine karşı daha dayanıklıdır.
- Biyoaktif bileşen taşıyıcılar:** Nanopartiküller, biyolojik olarak aktif bileşenleri bitki veya zararlı dokularına taşıyabilir ve etkinliğini artırabilir.

Zoom Workplace

Участники (6)


- OH observe... (Соорганизатор, я)
- A Hall-2, Aysun Altikat
- HG H1-Celalettin Gözüağık
- HZ H-2, Zekiye ALTUNTEKİN
- S HALL-2, Soner AKCAN
- HK Hall-2/Boran KARATAŞ

Выключить звук для всех

Workplace

Observer Hall-4 Tawseef Ahmad... Hall4-BENBELKEM... Brikci

Observer Hall-4 Tawseef Ahmad Dar Hall4-BENBELKACEM... Dr. Ali Imran Brikci



Efficient Extraction and Antioxidant Profiling of Polyphenols from Cereal Industrial Waste for Sustainable Resource Utilization

Ali Imran^{1,2}
Postdoctoral Research Fellow/Associate Professor

¹School of Agriculture, Food and Ecosystem Sciences, Faculty of Science, The University of Melbourne, Parkville, Victoria, Australia.
²Department of Food Science, Faculty of Life Science, Government College University, Faisalabad, Pakistan

Zoom Workplace

Участники (5)

- OH Observ... (Соорганизатор, я)
- A Dr. Ali Imran
- TA Tawseef Ahmad Dar
- B Brikci
- HI Hall4-BENBELKACEM IMENE

Выключить звук для всех

2nd INTERNATIONAL CONGRESS ON SUSTAINABLE AGRICULTURE

March 04-05, 2025 / Iğdır University, Türkiye



CONGRESS PROGRAM

ONLINE



Meeting ID: 829 5595 6031
Passcode: 030405

<https://us02web.zoom.us/j/82955956031?pwd=8lPlbfR9tbJjn1146HunqS6L7KakUN.1>

Participant Countries (16)

Türkiye, Azerbaijan, Morocco, Algeria, France, Tunisia, Pakistan, Spain, Philippines,
India, Iraq, Australia, China, Albania, Vietnam, Romania

Önemli, Dikkatle Okuyunuz Lütfen

- Kongremizde Yazım Kurallarına uygun gönderilmiş ve bilim kurulundan geçen bildirimler için online (video konferans sistemi üzerinden) sunum imkanı sağlanmıştır.
- Online sunum yapabilmek için <https://zoom.us/join> sitesi üzerinden giriş yaparak "Meeting ID or Personal Link Name" yerine ID numarasını girerek oturuma katılabilirsiniz.
- Zoom uygulaması ücretsizdir ve hesap oluşturmaya gerek yoktur.
- Zoom uygulaması kaydolmadan kullanılabilir.
- Uygulama tablet, telefon ve PC'lerde çalışıyor.
- Her oturumdaki sunucular, sunum saatinden 5 dk öncesinde oturuma bağlanmış olmaları gerekmektedir.
- Tüm kongre katılımcıları canlı bağlanarak tüm oturumları dinleyebilir.
- Moderatör – oturumdaki sunum ve bilimsel tartışma (soru-cevap) kısmından sorumludur.

Dikkat Edilmesi Gerekenler - TEKNİK BİLGİLER

- Bilgisayarınızda mikrofon olduğuna ve çalıştığına emin olun.
- Zoom'da ekran paylaşma özelliğini kullanabilmelisiniz.
- Kabul edilen bildiri sahiplerinin mail adreslerine Zoom uygulamasında oluşturduğumuz oturuma ait ID numarası gönderilecektir.
- Katılım belgeleri kongre sonunda tarafınıza pdf olarak gönderilecektir.
- Kongre programında yer ve saat değişikliği gibi talepler dikkate alınmayacaktır.

Important, Please Read Carefully

- To be able to attend a meeting online, login via <https://zoom.us/join> site, enter ID "Meeting ID or Personal Link Name" and solidify the session.
- The Zoom application is free and no need to create an account.
- The Zoom application can be used without registration.
- The application works on tablets, phones and PCs.
- The participant must be connected to the session 5 minutes before the presentation time.
- All congress participants can connect live and listen to all sessions.
- Moderator is responsible for the presentation and scientific discussion (question-answer) section of the session.

Points to Take into Consideration - TECHNICAL INFORMATION

- Make sure your computer has a microphone and is working.
- You should be able to use screen sharing feature in Zoom.
- Attendance certificates will be sent to you as pdf at the end of the congress.
- Requests such as change of place and time will not be taken into consideration in the congress program.

Zoom'a giriş yapmadan önce lütfen örnekteki gibi salon numaranızı, adınızı ve soyadınızı belirtiniz
Before you login to Zoom please indicate your hall number, name and surname

exp. H-5, Radmila Janičić

-OPENING CEREMONY-

04.03.2025 / 13:10-13:50

Prof. Dr. Sefa ALTIKAT

Iğdır University, Faculty of Agriculture Dean
CONGRESS PRESIDENT

Dr. Mustafa Latif EMEK

President of IKSAD Institute
CONGRESS HONORARY PRESIDENT

Prof. Dr. Mustafa Sıtkı BİLGİN

Rector of the Artvin Coruh University
CONGRESS HONORARY PRESIDENT

Assoc. Prof. Dr. Elbrus ISAYEV

Rector of the Nakhcivan State University
CONGRESS HONORARY PRESIDENT

Prof. Dr. Mehmet Hakkı ALMA

Rector of the Iğdır University
CONGRESS HONORARY PRESIDENT



ONLINE PRESENTATIONS

04.03.2025 / HALL-1 / SESSION-1



ANKARA LOCAL TIME: 14⁰⁰ - 16⁰⁰

HEAD OF SESSION: **Prof. Dr. Sefa ALTİKAT**

AUTHORS	AFFILIATION	TOPIC TITLE
Prof. Dr. Sefa ALTİKAT	<i>Iğdır University TÜRKİYE</i>	CHANGE ON ALFALFA AGRICULTURE: 2023 AND 2025 COMPARATIVE ANALYSIS BASED ON SATELLITE DATA
Prof. Dr. Sefa ALTİKAT	<i>Iğdır University TÜRKİYE</i>	THE EFFECT OF ALFALFA SOWING ON SOIL ORGANIC MATTER CONTENT IN İĞDIR PROVINCE: A REMOTE SENSING BASED EVALUATION
Assist. Prof. Dr. Derya GÜLOĞLU	<i>Isparta University of Applied Sciences TÜRKİYE</i>	SOIL CONSERVATION MEASURES IN SUSTAINABLE AGRICULTURE
Assist. Prof. Dr. Haydar BALCI Lect. Dr. Muhsin YILDIZ Lect. Dr. Murat KARA	<i>Van Yüzüncü Yıl University TÜRKİYE</i>	EFFECTS OF PROLINE AND SALICYLIC ACID APPLICATION ON YIELD AND PLANT GROWTH IN GREEN ONION CULTIVATION
Assist. Prof. Dr. Haydar BALCI Lect. Dr. Murat KARA Lect. Dr. Muhsin YILDIZ	<i>Van Yüzüncü Yıl University TÜRKİYE</i>	EFFECTS OF LIQUID VERMICOMPOST, SALICYLIC ACID, AND PROLINE APPLICATIONS ON THE GROWTH OF GREEN ONION (ALLIUM CEPA L.)
Şeyma KAYA Feride MERCAN Assoc. Prof. Dr. Akide ÖZCAN Prof. Dr. Mehmet SÜTYEMEZ	<i>Kahramanmaraş Sütçü İmam University TÜRKİYE</i>	EFFECT OF HIGH TEMPERATURE ON WALNUT CULTIVATION
Agr. Eng. Abdullah İREÇ Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	DAMAGE SITUATION OF PEACH TRUNK BLOODBITE, PTEROCHLOROIDES PERSICAE (CHOLODKOVSKY) (HOMOPTERA APHIDIDAE) IN APRICOT AND PEACH ORCHARDS OF İĞDIR PROVINCE
Agr. Eng. Abdullah İREÇ Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	DAMAGE SITUATION OF MEDITERRANEAN FRUIT FLY, CERATITIS CAPITATA (WIEDEMANN) (DIPTERA: TEPHRITIDAE) IN PEACH GARDENS OF İĞDIR PROVINCE

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

04.03.2025 / HALL-2 / SESSION-1



ANKARA LOCAL TIME: 14⁰⁰ - 16⁰⁰

HEAD OF SESSION: **Prof. Dr. Ahmet Zafer TEL**

AUTHORS	AFFILIATION	TOPIC TITLE
Ergün ÖZUSLU Prof. Dr. Ahmet Zafer TEL	<i>Gaziantep Islam Science and Technology University TÜRKİYE İğdır University TÜRKİYE</i>	DETERMINATION OF EUNIS HABITAT TYPES OF SOME NATURAL PROTECTED AREAS IN ADANA (TÜRKİYE) PROVINCE
Lect. Musa KARADAĞ Dr. Yunus BAŞAR Prof. Dr. Mehmet Hakkı ALMA	<i>İğdır University TÜRKİYE</i>	PHYTOCHEMICAL CONTENT OF LAURUS NOBILIS LEAVES AND EXTRACT; POTENTIAL FOR USE IN VALUE-ADDED COSMETICS, HEALTH AND FOOD PRODUCTS
Dr. Murat TAK Prof. Dr. Ahmet Zafer TEL	<i>İğdır University TÜRKİYE</i>	RARE AND ENDEMIC PLANTS OF AKDAĞ (ADIYAMAN/MALATYA) FLORA
Assist. Prof. Dr. Emine ATALAY	<i>Selçuk University TÜRKİYE</i>	SEED COATING TECHNOLOGIES FROM A SUSTAINABLE AGRICULTURE PERSPECTIVE
Assist. Prof. Dr. Necibe KAYAK Prof. Dr. Önder TÜRKMEN	<i>Sakarya University of Applied Sciences TÜRKİYE Selçuk University TÜRKİYE</i>	USE OF MICROBIAL AGENTS AGAINST SOIL SALINITY STRESS IN VEGETABLES
Ayşe ALTINKAYA Melahat Özge ÖZEN Nur ÜLGER Büşra ÜLGER	<i>Multi Tohum Tar. Singing. Tic. Inc. Antalya, TÜRKİYE</i>	TOMATO YELLOW LEAF CURL VIRUS (TYLCV) AND CONTROL METHODS IN SOLANACEAE
Res. Assist. Tansu ALKAN Prof. Dr. Süleyman Savaş DURDURAN	<i>Niğde Ömer Halisdemir University TÜRKİYE Necmettin Erbakan University TÜRKİYE</i>	ANALYSING EXPERT OPINIONS ON AGRICULTURAL SUPPORTS IN RURAL AREAS IN KONYA CLOSED BASIN

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

04.03.2025 / HALL-3 / SESSION-1



ANKARA LOCAL TIME: 14⁰⁰ - 16⁰⁰

HEAD OF SESSION: **Prof. Dr. Ramazan GÜRBÜZ**

AUTHORS	AFFILIATION	TOPIC TITLE
Agr. Eng.Ayfer GÜNEY SARITAŞ Prof. Dr. Ramazan GÜRBÜZ	<i>Iğdır University TÜRKİYE</i>	INNOVATIVE BIOCONTROL STRATEGIES FOR SUSTAINABLE WEED MANAGEMENT
Agr. Eng.Ayfer GÜNEY SARITAŞ Prof. Dr. Ramazan GÜRBÜZ	<i>Iğdır University TÜRKİYE</i>	INTEGRATION OF COVER CROPS IN WEED SUPPRESSION AND SOIL FERTILITY
Assoc. Prof. Dr. Mehmet Zeki KOÇAK Prof. Dr. Bünyamin YILDIRIM	<i>Iğdır University TÜRKİYE</i>	USE OF FLAXSEED (<i>Linum usitatissimum</i> L.) WITH HIGH ADDED VALUE
Assoc. Prof. Dr. Mehmet Zeki KOÇAK Prof. Dr. Bünyamin YILDIRIM	<i>Iğdır University TÜRKİYE</i>	THE EFFECT OF ESSENTIAL OILS FROM MEDICINAL AND AROMATIC PLANTS ON PLANT GROWTH
Res. Assist. Dr. Muhammed İkbal ÇATAL Prof. Dr. Adil BAKOĞLU	<i>Recep Tayyip Erdoğan University TÜRKİYE</i>	EVALUATION OF THE FORAGE QUALITY AND NUTRITIONAL VALUE OF THE PASTURE AREA OF ULUTAŞ NEIGHBORHOOD OF İSPIR DISTRICT IN ERZURUM PROVINCE
Aslihan KULOGLU Assoc. Prof. Dr. Aysun ÇAVUŞOĞLU	<i>Kocaeli University TÜRKİYE</i>	TISSUE CULTURE STUDIES ON CYCLAMEN SPP.
Assist Prof. Dr. Ruşen AYDIN KARAAĞAÇ Ahmet Taner KALE	<i>Kastamonu University TÜRKİYE</i>	EVALUATION OF THE EFFECTS OF SHEEP MANURE, WORM COMPOSITION AND MELASSES APPLICATIONS ON BLACK ARONIA (<i>Aronia Melanocarpa</i>) YIELD.
Lect. Dr. Hasret ÖZTÜRK	<i>Iğdır University TÜRKİYE</i>	A REVIEW OF IGF-I GENE VARIABILITY AND ITS IMPACT ON SHEEP BREEDING IN TURKEY

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

04.03.2025 / HALL-4 / SESSION-1



ANKARA LOCAL TIME: 14⁰⁰ - 16⁰⁰

HEAD OF SESSION: **Sarra Farjallah**

AUTHORS	AFFILIATION	TOPIC TITLE
Karima JMAILI Khawla WADDI Prof. Dr. Bouchaib BAHLAOUAN Prof. Dr. Alla SILKINA Prof. Dr. Nadia BOUTALEB	<i>Hassan II University MOROCCO</i>	THE USE OF WHEY TO ENHANCE TOMATO AND WHEAT SEED GERMINATION: A PATHWAY TO VALORIZING DAIRY BY-PRODUCTS FOR SUSTAINABLE AGRICULTURE
Samira KAROU	<i>Department of Soil Science and Environment, Higher School of Agronomy ALGERIA</i>	UPDATE OF ALGERIA'S SOIL MAP AT A 1/500,000 SCALE: APPLICATION OF GOOGLE EARTH IMAGERY
Waffa ABDELMOUMENE Nabila BELYAGOUBI-BENHAMMOU Zineb KAID Asmaa DICH Larbi BELYAGOUBI	<i>Abou-Bekr Belkaïd University ALGERIA Djillali Liabes University ALGERIA</i>	COMPARATIVE ASSESSMENT OF ANTIOXIDANT ACTIVITY AND PHYTOCHEMICAL PROFILES OF AQUEOUS EXTRACTS FROM ATRIPLEX HALIMUS LEAVES AND STEMS
BELLIL Zahra MEYER Sylvain TILLOY Valentin BELLIL Dassine MAIRI Assia De CHAMPS Christophe BARRAUD Olivier TOUATI Abdelaziz	<i>Bejaia University ALGERIA Limoges University FRANCE Reims-Champagne-Ardenne University FRANCE</i>	OCCURRENCE AND CHARACTERIZATION OF ANTIBIOTIC-RESISTANT SALMONELLA ISOLATES FROM FRESH PRODUCE IN AGRICULTURAL ENVIRONMENTS
Sarra Farjallah	<i>Tunis El Manar University TUNUSIA</i>	DNA MOLECULAR MARKERS AND THEIR APPLICATIONS IN INVESTIGATING THE GENETIC VARIABILITY OF THE NEMATODE HETERAKIS GALLINARUM FROM TUNISIA
Sarra Farjallah	<i>Tunis El Manar University TUNUSIA</i>	EPIDEMIOLOGICAL SURVEY AND MOLECULAR CHARACTERIZATION OF MONOGENEAN PARASITE SPECIES INFESTING SPARUS AURATA FROM TUNISIA
Muhammad Mazhar Iqbal Muhammad Abdullah Tehmena Rashid Hafiz Qaiser Yasin Shafiq-ur-Rehmn	<i>Water Management Training Institute PAKISTAN Agriculture University PAKISTAN</i>	ANALYSIS OF HYDRO-METEOROLOGICAL TRENDS ACROSS MAJOR KOPPEN CLIMATE CLASSES USING GLDAS DATA AT A GLOBAL SCALE
Deghiche-Diab Nacima Saad Somia Belhamra Youcef Islem Ouamene Tarek Abdelmoneim Karoune Samira Djoudi Madjed Kechebar Mohamed Seif Allah Djazouli Zahr Eddine	<i>Scientific and Technical Research Center on Arid Regions (CRSTRA) ALGERIA Blida 1 University ALGERIA</i>	VALORISATION OF BIOPESTICIDE FROM DITTRICHIA VISCOSA (L.) GREUTER 1973 FROM ALGERIAN ARID REGION

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

04.03.2025 / HALL-5 / SESSION-1



ANKARA LOCAL TIME: 14⁰⁰ - 16⁰⁰

HEAD OF SESSION: Assoc. Prof. Dr. Nadira OUKALA & Assoc. Prof. Dr. Victoria PASTOR

AUTHORS	AFFILIATION	TOPIC TITLE
AOUFI Farida Dr. TAMERT Asma ABDI Yamina BARBARA Ferdaws	<i>Relizane University ALGERIA</i>	ANTIBACTERIAL AND ANTIFUNGAL ACTIVITY OF PLANT (MARRUBIUM VULGARE) FROM THE FAMILY OF LAMIACEAE IN WESTERN ALGERIA
Assoc. Prof. Dr. Nadira OUKALA ORERO BAYO Marta Assoc. Prof. Dr. Victoria PASTOR	<i>M'hamed Bougara University ALGERIA Jaume I University SPAIN</i>	ENDOPHYTIC BACTERIUM BACILLUS CERUS -MEDIATED SALT TOLERANCE IN WHEAT PLANTS
Nacéra TADJINE Prof. Dr. Saida MESSGO-MOUMENE	<i>Blida1 University ALGERIA</i>	IN VITRO EVALUATION OF THE ANTIFUNGAL POTENTIAL OF MYRTUS COMMUNIS L. AGAINST TOXIGENIC MOLDS OF HYDROPONIC BARLEY
HADJLOUNE Houria KACI Ahcène	<i>Agropastoralism Research Center (CRAPast), The "Agropastoral and Rural Economy" ALGERIA National Higher Agronomic School (ENSA), Department of Rural Economy ALGERIA</i>	THE INFLUENCE OF INSTITUTIONAL DYNAMICS AND GOVERNANCE STRUCTURES ON THE SUPPLY CHAIN OF THE MILK SECTOR IN ALGERIA: CASE OF THE DAIRY HODNA-LAIT (M'SILA)
Famela A. Foyo Bernadette I. Fillartos Cristine Angeli T. Mangampat Jane M. Balubar Klenton R. Torrevillas, MIHM	<i>Cavite State University PHILIPPINES</i>	INNOVATIVE BAKING: UTILIZING KAMIAS AND BANANA-ENRICHED FLOUR IN CAKE RECIPES INSPIRED BY LEMON CAKE
Muhammad Bilal Hussain Marwa Waheed Farhan Saeed Muhammad Afzaal Mariam Islam	<i>Government College University PAKISTAN Riphah International University PAKISTAN</i>	NUTRITIONAL AND PHARMACOLOGICAL PROPERTIES OF CANTALOUPE PEEL: AN OVERVIEW
Abdul Ahad Ansari Haris Hasan Khan	<i>Aligarh Muslim University INDIA</i>	ADVANCEMENTS IN REMOTE SENSING FOR CROP AND SOIL HEALTH MANAGEMENT IN SUSTAINABLE AGRICULTURE
N Anusuya V S Angulakshmi M Priyadharshini G Easterin Shejina	<i>P.S.G.R. Krishnammal College for Women INDIA</i>	GREEN MEDIATED SYNTHESIS, CHARACTERIZATION AND APPLICATION OF MANGANESE OXIDE NANOPARTICLES USING CITRUS LIMON PEEL EXTRACT

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

04.03.2025 / HALL-6 / SESSION-1



ANKARA LOCAL TIME: 14⁰⁰ - 16⁰⁰

HEAD OF SESSION: **Dr. Ihaddaden Akli**

AUTHORS	AFFILIATION	TOPIC TITLE
Dr. Ihaddaden Akli Prof. Dr. ABLA Safia Prof. Dr. Abdoun Fatiha	<i>Abdelhafid Ihaddaden University ALGERIA Algiers University ALGERIA University of Sciences and Technology Houari Boumediène ALGERIA</i>	PHYSIOMOMIC CHANGE OF VEGETATION AND DIVERSITY INDEX IN TWO CONSTRAINED ENVIRONMENT OF THE SEMI-ARID IN ALGERIA
Thinhinan KHEDIM	<i>University of Sciences and Technology Houari Boumediene ALGERIA</i>	THE CASE OF ROSY GARLIC (ALLIUM ROSEUM L.) POLYPLOID COMPLEX IN ALGERIA
MRABTI Imane GRIJJA Hassan BENZAHRHA Hayat AIT BADDOU Sanae BRHADDA Najiba AFECHTAL Mohamed ZIRI Rabea	<i>Ibn Toufail University MOROCCO National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra MOROCCO Ibn Toufail University MOROCCO</i>	PRECISE IDENTIFICATION OF THE COTTONY CUSHION SCALE, ICERYA PURCHASI IN MOROCCO THROUGH THE ADVANCED APPLICATION OF DNA BARCODING
Dr. Ahlem DEBIEB Prof. Dr. Leila KADIK Prof. Dr. Michel GODRON Dr. Zineb DJERRAD Dr. Nacere ADI	<i>University of Sciences and Technology Houari Boumediene ALGERIA University of Sciences and Technology Houari Boumediene ALGERIA Paris Diderot University FRANCE</i>	THE RESILIENT FOREST AFTER THE FIRE: ANALYSIS OF THE STRUCTURE AND DIVERSITY OF THE EL-HAMDANIA CORK OAK FOREST-ALGERIA
EL BAHJA Fatima ZIRI Rabea BRITAL Rania HAMRANI Meriem EL AMMARI Mohamed FAHAD Kaoutar	<i>Ibn Toufail University MOROCCO National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra MOROCCO</i>	STUDY OF CERATITIS CAPITATA WIEDEMANN (DIPTERA: TEPHRITIDAE) POPULATION DYNAMICS IN THE EL GHARB REGION OF MOROCCO FOR FUTURE PEST MANAGEMENT
Meriem RAFYA Prof. Dr. Naima ZEHAR Prof. Dr. Fatiha BENKHALTI	<i>Cadi Ayyad University MOROCCO</i>	THE NEGATIVE EFFECT OF SOIL AMENDMENTS WITH SOLID RESIDUES OF ROSMARINUS OFFICINALIS L. ON THE GROWTH OF SOLANUM LYCOPERSICUM MILL.
Kaouthar ELBIRGUI Houssam ASSIOUI Oussama RAFIK Nadia IOUNES Samir IBENMOUSSA	<i>Hassan II University MOROCCO</i>	INVESTIGATING THE ANTIOXIDANT POTENTIAL OF EBENUS PINNATA AITON: IN VITRO AND IN VIVO ASSESSMENTS OF HYDROETHANOLIC EXTRACTS
MRABTI Imane GRIJJA Hassan BRITAL Rania BRHADDA Najiba ZIRI Rabea AFECHTAL Mohamed	<i>Ibn Toufail University MOROCCO National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra MOROCCO</i>	IMPROVEMENT OF DNA EXTRACTION FOR THE PYRIFORM SCALE INSECT, PROTOPULVINARIA PYRIFORMIS, FOR MOLECULAR IDENTIFICATION OF CITRUS PESTS IN MOROCCO USING DNA BARCODING
Sarra Farjallah Khadija Doukali	<i>Tunis El Manar University TUNUSIA Carthage University TUNUSIA</i>	EPIDEMIOLOGICAL SURVEY AND MOLECULAR CHARACTERIZATION OF MONOGENEAN PARASITE SPECIES INFESTING SPARUS AURATA FROM TUNISIA

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-1 / SESSION-1



ANKARA LOCAL TIME: 10⁰⁰ - 12⁰⁰

HEAD OF SESSION: **Assoc. Prof. Dr. İsmail ALASERHAT**

AUTHORS	AFFILIATION	TOPIC TITLE
Eda ÖZÇALICI Assoc. Prof. Dr. İsmail ALASERHAT	<i>Agri Agricultural Quarantine Directorate TÜRKİYE İğdır University TÜRKİYE</i>	COTTON LEAFWORM, Spodoptera littoralis (Boisduval) (LEPIDOPTERA: NOCTUIDAE) AN IMPORTANT QUARANTINE PEST CAUSING HIGH LOSSES IN VEGETABLES
Hava ULUĞBEY Assoc. Prof. Dr. İsmail ALASERHAT	<i>Agri Agricultural Quarantine Directorate TÜRKİYE İğdır University TÜRKİYE</i>	ASIAN CITRUS PSYLLID, Diaphorina citri Kuwayama (HEMIPTERA: PSYLLIDAE), AN IMPORTANT QUARANTINE PEST CAUSING HIGH LOSSES IN CITRUS FRUIT
Assist. Prof. Dr. Berna BAŞ	<i>Gaziantep University TÜRKİYE</i>	SOILLESS PLANT CULTURE TECHNICAL METHODS
Assist. Prof. Dr. Berna BAŞ	<i>Gaziantep University TÜRKİYE</i>	IS VERTICAL FARMING SUSTAINABLE?
Agr. Eng. Hanife ÇEÇEN Assist. Prof. Dr. Murat KARACA	<i>Sector Tarım Agricultural, Konya TÜRKİYE Selçuk University TÜRKİYE</i>	WEED CONTROL IN CANNABIS
Mustafa ÇETİNER Prof. Dr. Ahmet Zafer TEL	<i>İğdır University TÜRKİYE</i>	FLORA AND ENDEMIC PLANTS OF ATATURK DAM BASIN (ŞANLIURFA/TURKEY)
İlknur İŞÇİ Res. Assist. Gülcan KAYMAK BAYRAM Zeki ACAR İlknur AYAN	<i>Ondokuz Mayıs University TÜRKİYE</i>	THE ROLE OF LOCAL GOVERNMENTS IN SUSTAINABLE AGRICULTURE: THE ISTANBUL CASE
Firas ALJANABI Assoc. Prof. Dr. Mert DEDEOĞLU	<i>Selçuk University TÜRKİYE</i>	PREDICTING LAND SUITABILITY FOR WHEAT CULTIVATION USING THE RANDOM FOREST MODEL
İlyas KILINÇER Dr. Özgür ÖZMEN Dr. Anıl Mehmet BALTACI Prof. Dr. Mehmet ARSLAN	<i>Kayseri Şeker R&D Center, TÜRKİYE Erciyes University TÜRKİYE</i>	DETERMINATION OF YIELD AND QUALITY PARAMETERS OF SUGAR BEET PARENTAL LINES UNDER LIMITED IRRIGATION

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-2 / SESSION-1



ANKARA LOCAL TIME: 10⁰⁰ - 12⁰⁰

HEAD OF SESSION: **Prof. Dr. Etibar MEMMEDOV**

AUTHORS	AFFILIATION	TOPIC TITLE
Assoc. Prof. Dr. Elsever Asadov Sevgi Kolaylı	<i>Nakhchivan State University</i> AZERBAIJAN <i>Karadeniz Technical University</i> TÜRKİYE	COMPARATIVE STUDY OF SOME SUGAR PROFILES OF HONEY AND FRUITS
Assoc. Prof. Dr. Mirmahmud SEYIDLİ Assoc. Prof. Dr. Jabbar NAJAFOV	<i>Nakhchivan State University</i> AZERBAIJAN	SPIDER MITE - TETRANYCHIDAE IN THE VINEYARDS OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC OF AZERBAIJAN AND MEASURES TO FIGHT AGAINST IT
Akim AKHUNDOV Assoc. Prof. Dr. Sahib HACIYEV Assoc. Prof. Dr. Elnara SEYİDOVA	<i>Nahçivan State University</i> AZERBAIJAN	ECOLOGICAL ASSESSMENT OF THE SOILS OF ABRAGUNUS DEPRESSION
Logman BAYRAMOV	<i>Nahcivan State University</i> AZERBAIJAN	RESEARCH AND GENE FUND COLLECTION OF STONE FRUIT PLANTS CULTIVATED IN THE TERRITORY OF NAKHCHIVAN AUTONOMOUS REPUBLIC
Prof. Dr. Etibar MEMMEDOV	<i>Nahcivan State University</i> AZERBAIJAN	THE ANTHELMINTIC EFFECT OF PREPARATIVE FORMS OF TANACERUM VULGARE L. ON GASTROINTESTINAL NEMATODES IN SHEEP
Novruzlu Aytan Ali Gurbanov	<i>Nahcivan State University</i> AZERBAIJAN	THE ECOLOGICAL RESULTS OF IMPACT OF THE ARPACHAY RESERVOIR IRRIGATION SYSTEMS TO THE ENVIRONMENT
Assoc. Prof. Dr. Ali TAHİROV Assoc. Prof. Dr. Yunis RUSTAMLİ	<i>Nahcivan State University</i> AZERBAIJAN	IMPACT OF GLOBAL CLIMATE CHANGE ON BEEKEEPING
Assoc. Prof. Dr. Yunis RUSTAMLİ Assoc. Prof. Dr. Ali TAHİROV	<i>Nahcivan State University</i> AZERBAIJAN	IN THE REPUBLIC OF AZERBAIJAN LEGAL BASIS OF SWITCHING TO ORGANIC BEEKEEPING
Ayaz Hajiyev	<i>Academy of Public Administration under the Presidency</i> AZERBAIJAN	THE TERM "GREEN ECONOMY" IN SUSTAINABLE DEVELOPMENT

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-3 / SESSION-1



ANKARA LOCAL TIME: 10⁰⁰ - 12⁰⁰

HEAD OF SESSION: **Prof. Dr. Celalettin GÖZÜAÇIK**

AUTHORS	AFFILIATION	TOPIC TITLE
Gönül USTA GEBEŞ Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	BIOLOGICAL CONTROL OF STORED PRODUCT DAMAGE
Ezgi Günay SARGUT Assoc. Prof. Dr. Reyvan Yergin ÖZKAN	<i>Van Yüzüncü Yıl University TÜRKİYE</i>	HOMEOPATHY IN PLANT PROTECTION
Ali ÇELİK Assoc. Prof. Dr. Furkan DİNÇER	<i>Kahramanmaraş Sütçü İmam University TÜRKİYE</i>	SUSTAINABILITY OF SALEP PRODUCTION THROUGH AGRIVOLTAIC APPLICATIONS: THE KALE TARIMGES MODEL
Ali ÇELİK Assoc. Prof. Dr. Furkan DİNÇER	<i>Kahramanmaraş Sütçü İmam University TÜRKİYE</i>	INTEGRATION OF RENEWABLE ENERGY AND AGRICULTURE WITH AGRIVOLTAICS: THE CASE OF KAHRAMANMARAŞ BULANIK TARIMGES
Lect. Nagihan KILIÇ Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	THE IMPACT OF CLIMATE CHANGE ON INTEGRATED PEST MANAGEMENT (IPM) IN TURKEY
Lect. Nagihan KILIÇ Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	PESTS OF GREENHOUSE VEGETABLE GROWING IN IĞDIR PROVINCE
Pınar ADIGÜZEL Assoc. Prof. Dr. Şenay KARABIYIK Prof. Dr. İlknur SOLMAZ	<i>Çukurova University TÜRKİYE</i>	APPLICATIONS TO ENHANCE POLLEN QUALITY IN WATERMELONS WITH DIFFERENT CHROMOSOME STRUCTURES
İlayda Gizem AKIN Prof. Dr. Özge SIZMAZ	<i>Ankara University TÜRKİYE</i>	USE OF THE ASTAXANTHIN IN TILAPIA (OREOCHROMIS NILOTICUS)

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-4 / SESSION-1



ANKARA LOCAL TIME: 10⁰⁰ - 12⁰⁰

HEAD OF SESSION: **Dr. K. SELVAM**

AUTHORS	AFFILIATION	TOPIC TITLE
VIGNESH K Dr. K. SELVAM ARSHA G	<i>Palar Agricultural College INDIA</i>	EFFECT OF NEW GENERATION FUNGICIDES AGAINST <i>Pyricularia oryzae</i> CAUSES BLAST IN RICE (<i>Oryza sativa</i> L.)
BRITAL Rania Abail Zhor MRABTI Imane EL bahja Fatima IBRIZ Mohammed	<i>Ibn Toufail University MOROCCO</i> <i>Regional Center for Agricultural Research of Kenitra MOROCCO</i> <i>Ibn Toufail University MOROCCO</i>	SURVEY ON FERTILIZATION PRACTICES AND NUTRIENT MANAGEMENT IN VALENCIA LATE CITRUS ORCHARDS: A CASE STUDY FROM THE GHARB REGION (NORTHEASTERN), MOROCCO
M.M Akibu M.S. Na-Allah A.Muhammad I. J. Yusuf	<i>Kebbi State University of Science and Technology NIGERIA</i>	YIELD AND STORABILITY OF ONION ECOTYPES AS INFLUENCED BY IRRIGATION INTERVAL IN SUDAN SAVANNA ZONE OF NIGERIA
ZAGGOUMI Hasna BOUDA Said NEG Imane MARGHALI Sonia TRIFI Neila HADDIOUI Abdelmajid	<i>Sultan Moulay Slimane University MOROCCO</i> <i>El Manar University TUNUSIA</i>	THE GENETIC CHARACTERIZATION OF THE MOROCCAN CAROB TREE (<i>CERATONIA SILIQUA</i> L.) USING ISSR MARKERS
Muhammad Muzammil Nazir Munaza Yasmeen Iqra Farzeen Sana Inam Asma Ashraf	<i>Government College University PAKISTAN</i>	THE STUDY OF ANTIDIARRHEAL POTENTIAL OF <i>HELICTERES ISORA</i> AND <i>PERSICARIA VIVIPARA</i> METHANOLIC EXTRACTS IN SWISS ALBINO MICE
Said Babou Miloud Chakit Abdelhalem Mesfioui Youssef Sqalli Houssaini	<i>Ibn Tofail University MOROCCO</i>	ANTILIPIDEMIC AND NEPHRO-HEPATOPROTECTIVE ACTIVITIES OF <i>ROSMARINUS OFFICINALIS</i> ETHANOLIC EXTRACT IN WISTAR RATS
Said Babou Miloud Chakit Radia El Gui Abdelhalem Mesfioui Youssef Sqalli Houssaini	<i>Ibn Tofail University MOROCCO</i>	TOXICITY ASSESSMENT OF ETHANOLIC EXTRACT OF <i>ROSMARINUS OFFICINALIS</i> LEAVES IN FEMALE WISTAR RATS
Fjorela MEMA Selma MYSLIHAKA Peçi NAQELLARI Nikoleta KALLAJXHIU Silvana TURKU Anxhela DAUTI	<i>Aleksandër Xhuvani University of Elbasan ALBANIA</i>	GEOBOTANICAL DATA ON FLORA AND VEGETATION IN SOME AREAS OF ELBASAN

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-5 / SESSION-1



ANKARA LOCAL TIME: 10⁰⁰ - 12⁰⁰

HEAD OF SESSION: **Dr. Radhia ARAB**

AUTHORS	AFFILIATION	TOPIC TITLE
Kaouthar ELBIRGUI Houssam ASSIOUI Oussama RAFIK Ali ZOURIF Houda ELALAMI Hanane ENNAJI Nadia IOUNES Samir IBENMOUSSA	<i>Laboratory of Chemistry and Biochemistry, Nutrition, Environment and Health MOROCCO Hassan II-University MOROCCO</i>	INVESTIGATING THE ANTIOXIDANT POTENTIAL OF EBENUS PINNATA AITON: IN VITRO AND IN VIVO ASSESSMENTS OF HYDROETHANOLIC EXTRACTS
Muslim Yusuf Sulaiman Sani Yusuf Abdulaziz Hassan Maidabino	<i>Umaru Musa Yar'adua University NIGERIA</i>	PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL ACTIVITY ON THE ETHANOLIC EXTRACT OF GUIERA SENEGALENSIS LEAVES
A. Attahiru A.U Birnin-Yauri M. Mahdi-Adamu A. Yahaya	<i>Kebbi State University of Science and Technology NIGERIA</i>	PHYTOCHEMICAL CONSTITUENTS OF MORINGA OLEIFERA ROOTS EXTRACT
M. Mahdi-Adamu F. Halidu A. Attahiru A. Yahaya R. Adamu	<i>Kebbi State University of Science and Technology NIGERIA Usmanu Danfodiyo University NIGERIA</i>	NUTRITIONAL AND PHYTOCHEMICAL ANALYSIS OF OREGANO (ORIGANUM VULGARE) LEAVES
IOANA POROSNICU Madalina DAVIDESCU Bianca-Maria MADESCU	<i>Lasi University of Life Sciences ROMANIA</i>	MIGRATION, CLIMATE CHANGE AND SUSTAINABLE AGRICULTURE: CHALLENGES AND OPPORTUNITIES FOR ECONOMIC DEVELOPMENT
Soumaya Atouife Jamila Elbiyad Halima Belaoufi Abdellah El Habazi Rekia Belahsen	<i>Chouaib Doukkali University MOROCCO</i>	UNDERSTANDING MOROCCAN POPULATION'S PERCEPTION AND AWARENESS OF ORGANIC FOOD
Hana ZAKKOURMI Assoc. Prof. Dr. Yazid MAHDAOUI Assoc. Prof. Dr. Houria HADJ-ARAB	<i>University of Sciences and Technology Houari Boumediene (USTHB) ALGERIA Algiers 1 Benyoucef Benkhedda University ALGERIA</i>	MORPHOLOGICAL DIVERSITY AND STYLAR POLYMORPHISM IN NARCISSUS SPECIES (AMARYLLIDACEAE) FROM NORTHERN ALGERIA
Dr. Radhia ARAB Nour-el-Houda KHELIFI Aicha AOUINA	<i>Mouhamed Boudiaf University ALGERIA</i>	REPELLENT ACTIVITY OF ESSENTIAL OILS OF TOW SPICES OF ALGERIAN KITCHEN AGAINST STORED CERALS BEETLE

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-1 / SESSION-2



ANKARA LOCAL TIME: 12³⁰ - 14³⁰

HEAD OF SESSION: **Assist. Prof. Dr. Barış EREN**

AUTHORS	AFFILIATION	TOPIC TITLE
Assist. Prof. Dr. Barış EREN İbrahim YILDIZHAN	<i>İğdır University TÜRKİYE</i>	EPIGENETIC INHERITANCE OF ENVIRONMENTAL CHANGES IN PLANTS: MECHANISMS AND IMPLICATIONS
Hasine KÜÇÜKYILDIRIM Abdulkadir BÜYÜK	<i>Harran University TÜRKİYE</i> <i>İğdır University TÜRKİYE</i>	WASTE MANAGEMENT IN AGRICULTURE: RECYCLING OF ORGANIC WASTE AND COMPOSTING TECHNIQUES
Assoc. Prof. Dr. Asuman ÇAPAR	<i>Sivas Cumhuriyet University TÜRKİYE</i>	SUSTAINABLE AGRICULTURE AS A REGULATORY AND SUPPORTIVE ACTIVITY OF PUBLIC ADMINISTRATION
Lect. Afşin AYHAN Prof. Dr. Banu ÇİÇEK KURDOĞLU	<i>Kastamonu University TÜRKİYE</i>	MULTI-CRITERIA DECISION-MAKING METHODS (MCDM) FOR SUSTAINABLE ENVIRONMENTAL MANAGEMENT: OPTIMIZED DECISION SUPPORT SYSTEMS WITH RSTUDIO
Lect. Rüştü ÇALLI Assist. Prof. Dr. Emirhan ÖZDEMİR Lect. Mitat Can YILDIZ Assoc. Prof. Dr. Alihsan ŞEKERTEKİN	<i>İğdır University TÜRKİYE</i>	DETECTION AND ANALYSIS OF BURNED AREAS WITH SENTINEL-2 SATELLITE IMAGERY AND FIRE SEVERITY MAP: 2025 LOS ANGELES FIRE
Lect. Mitat Can YILDIZ Assist. Prof. Dr. Emirhan ÖZDEMİR Lect. Rüştü ÇALLI Assoc. Prof. Dr. Alihsan ŞEKERTEKİN	<i>İğdır University TÜRKİYE</i>	BIBLIOMETRIC ANALYSIS OF THE USE OF SATELLITE-BASED REMOTE SENSING TECHNOLOGIES IN CROP MAPPING (2018-2025)
Assist. Prof. Dr. Orhan GÜNGÖR	<i>Burdur Mehmet Akif Ersoy University TÜRKİYE</i>	EFFECTS OF BATTERY TECHNOLOGIES ON AGRICULTURAL MACHINERY: EFFICIENCY, SUSTAINABILITY AND FUTURE PERSPECTIVES
Osman İNAN Assist. Prof. Dr. Sema KONYALI	<i>Independent Researcher İstanbul TÜRKİYE</i> <i>Tekirdağ Namık Kemal University TÜRKİYE</i>	DEVELOPMENT OF FOOD FROM PAST TO PRESENT AND FOOD SECURITY

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-2 / SESSION-2



ANKARA LOCAL TIME: 12³⁰ - 14³⁰

HEAD OF SESSION: **Assoc. Prof. Dr. Adnan AYDIN**

AUTHORS	AFFILIATION	TOPIC TITLE
Gülcan TAŞ Assoc. Prof. Dr. Adnan AYDIN	<i>Iğdır University TÜRKİYE</i>	DETECTION OF DNA MARKERS IN MEDICINAL SAGE (<i>Salvia officinalis</i> L.) FOR USE IN EPIGENETIC STUDIES
Assoc. Prof. Dr. Adnan AYDIN	<i>Iğdır University TÜRKİYE</i>	MOLECULAR MARKER TECHNIQUES FOR THE DETECTION OF DNA CYTOSINE METHYLATION
Fatih KUMBASAR Mehmet CAN Res. Assist. Gülcan KAYMAK BAYRAM İlknur AYAN Zeki ACAR	<i>Ondokuz Mayıs University TÜRKİYE</i>	THE IMPORTANCE OF RANGELANDS AS CARBON SINKS
Halit Burak YÜKSEKDAĞ Hayrullah ÖZBEK Mustafa Can SARIOĞLU Mustafa ARSLAN Mustafa BÖLÜK	<i>Kayseri Sugar Factory TÜRKİYE</i>	SUSTAINABLE AGRICULTURE AND SUGAR BEET: AN ANALYSIS OF CARBON FOOTPRINT, WATER USAGE, AND EFFICIENCY
Hayrullah ÖZBEK Halit Burak YUKSEKDAĞ Oguzhan ULU Anil Mehmet BALTACI	<i>Kayseri Sugar Factory TÜRKİYE</i>	EFFECT OF BIOLOGICAL AND ORGANIC FERTILIZERS ON YIELD AND QUALITY IN ORGANIC SUGAR BEET FARMING
Prof. Dr. Köksal KARADAŞ Assist. Prof. Dr. Osman Doğan BULUT	<i>Iğdır University TÜRKİYE</i>	SİİRT PISTACHIO PRODUCERS' DEMOGRAPHIC CHARACTERISTICS AND PRODUCTION PROBLEMS: THE CASE OF SİİRT PROVINCE
Assist. Prof. Dr. Osman Doğan BULUT Prof. Dr. Köksal KARADAŞ	<i>Iğdır University TÜRKİYE</i>	EXAMINATION OF FARMERS' KNOWLEDGE AND AWARENESS ON THE ENVIRONMENTAL EFFECTS OF PESTICIDE USE: THE CASE OF İĞDIR PROVINCE
Dr. Aysun UYSAL Prof. Dr. Şener KURT Tuğba HANEDAN	<i>Hatay Mustafa Kemal University TÜRKİYE</i>	ABIOTIC STRESS FACTORS AND THEIR EFFECTS IN STONE FRUIT ORCHARDS OF AMIK PLAIN
Dr. Aysun UYSAL Res. Assist. Yusuf GÜMÜŞ Prof. Dr. Soner SOYLU Assoc. Prof. Dr. Merve OĞUZ Prof. Dr. Emine Mine SOYLU	<i>Hatay Mustafa Kemal University TÜRKİYE</i>	DETERMINATION OF IN VITRO ANTIFUNGAL ACTIVITY OF WOOD VINEGAR AGAINST ROSALLINIA ROOT ROT DISEASE AGENT IN FIG TREES

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-3 / SESSION-2



ANKARA LOCAL TIME: 12³⁰ - 14³⁰

HEAD OF SESSION: **Prof. Dr. Aydın AKKAYA**

AUTHORS	AFFILIATION	TOPIC TITLE
Prof. Dr. Aydın AKKAYA	<i>Muş Alparslan University TÜRKİYE</i>	REASONS FOR WHITE SPIKE FORMATION IN WHEAT AND DIAGNOSIS
Prof. Dr. Aydın AKKAYA	<i>Muş Alparslan University TÜRKİYE</i>	WHEAT BASED INTERCROPPING
Res. Assist. Dilek DOĞAN Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	HARMFUL INSECTS OF THE HETEROPTERA (HEMIPTERA) SUBORDER SEEN ON FRUIT TREES IN İĞDIR PROVINCE
Assist. Prof. Dr. Tahsin UYGUN Assist. Prof. Dr. Mesut ÇOŞLU Assist. Prof. Dr. Mete YİĞİT Assoc. Prof. Dr. Iker ÜNAL	<i>Burdur Mehmet Akif Ersoy University TÜRKİYE</i>	DETERMINATION OF ORANGE FRUIT RIPENESS LEVELS USING DEEP LEARNING-BASED INSTANCE SEGMENTATION METHOD
Muhammet ÖTER Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	PESTICIDE USE IN İĞDIR PROVINCE
Res. Assist. Dr. Ezgi DEMİRAL	<i>Doğuş University TÜRKİYE</i>	A CRITICAL ANALYSIS OF TURKEY'S AGRICULTURAL POLICIES IN THE CONTEXT OF COMBATING CLIMATE CHANGE AND SUSTAINABLE AGRICULTURE
Alev ALDEMİR Prof. Dr. İsmail KIYAK Faruk AYDIN Ahmet KIRIK	<i>Marmara University TÜRKİYE Filizhane Biotechnology R&D Agricultural Applications Tic.Ltd. Şti, İstanbul TÜRKİYE</i>	PORTABLE MINI-SCALE VERTICAL FARMING SYSTEM DESIGN: GROWING MICROSPOUT IN SOIL-LESS MEDIA
Assist. Prof. Dr. Aslı İŞLER KAYA	<i>Turkish German University TÜRKİYE</i>	SUSTAINABLE AGRICULTURE FOR OILSEED CROPS
Yağmur TÜRKMEN Prof. Dr. Filiz ERTUNÇ	<i>Ordu University TÜRKİYE Ankara University TÜRKİYE</i>	INCIDENCE OF GRAPEVINE PINOT GRIS VIRUS IN WESTERN BLACK SEA REGION, TURKEY AND PHYLOGENETIC ANALYSIS BASED ON COAT PROTEIN AND MOVEMENT PROTEIN GENE

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-4 / SESSION-2



ANKARA LOCAL TIME: 12³⁰ - 14³⁰

HEAD OF SESSION: **Dr. Sundhararajan R.**

AUTHORS	AFFILIATION	TOPIC TITLE
LAZREG Halima SEBAA Hanane Siham CELIK Muhammed Suleyman Yildiz Ilyas Ramazan ERENLER ABDEDDAIM Katia Khadidja DJABEUR Abd Erezzak	<i>University of Sciences and Technology Oran Mohammed Boudiaf ALGERIA</i> <i>Iğdır University TÜRKİYE</i> <i>University of Sciences and Technology Oran Mohammed Boudiaf ALGERIA</i>	PHENOLIC COMPOUNDS OF OPUNTIA STRICTA CLADODES
Soukaina El Bourachdi Abdelhay El Amri Ali Raza Ayub Fatima Moussaoui Amal Lahkimi	<i>Sidi Mohamed Ben Abdellah University MOROCCO</i> <i>Ibn Tofaïl University MOROCCO</i> <i>Beijing Institute of Technology CHINA</i> <i>Sidi Mohamed Ben Abdellah University MOROCCO</i>	DEVELOPMENT OF A NOVEL LOW- COST ADSORBENT CHITOSAN@EDTA@CELLULOSE COMPOSITE TO EFFECTIVELY REMOVE METHYL ORANGE DYE FROM WASTEWATER: EXPERIMENTAL AND THEORETICAL INVESTIGATION
Hafiza Zara Saeed Muhammad Wajid Musrat Shaheen	<i>Government College University PAKISTAN</i>	EFFICACY OF ESSENTIAL ELEMENTS AGAINST SUCKING INSECT PESTS OF COTTON UNDER DIFFERENT CONDITION
Dr. Sundhararajan R. Mrs. Jothilakshmi. R. Mr. S.G.Raman	<i>Mohammed Sathak AJ College INDIA</i>	GREEN INNOVATIONS IN MEDICINAL AND AROMATIC PLANT RESEARCH
Farrah Deebe Muhammad Huzaifah Khalid Anas Sarwar Qureshi Muhammad Adil	<i>Agriculture University PAKISTAN</i> <i>Riphah International University PAKISTAN</i>	THERAPEUTIC STUDY OF CENTRATHERUM ANTHELMINTICUM FOR THE CONTROL OF SUBCLINICAL MASTITIS IN BUBALINE
Sadiq, M.S Singh, I.P Ahmad, M.M Sani, B.S Nazifi, I.K	<i>Department of Agricultural Economics and Agribusiness NIGERIA</i> <i>Federal Inland Revenue Services NIGERIA</i>	SEED-TO-SHELF VENTURES: BUILDING A YOUTH-LED AGRIBUSINESS MODEL FOR FARM-TO-MARKET TRANSPARENCY
Oyewole Mojisola Fauziyah	<i>Ibadan University NIGERIA</i>	ACCEPTABILITY OF ORANGE FLESHED SWEET POTATO CONFECTIONARIES AMONG FACULTY OF AGRICULTURE STUDENTS, UNIVERSITY OF IBADAN. NIGERIA
GUNASEKAR G	<i>Tamil Nadu Agricultural University INDIA</i>	EVOLUTION, BEHAVIOR AND ECOLOGY OF FEATHER MITE

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-5 / SESSION-2



ANKARA LOCAL TIME: 12³⁰ - 14³⁰

HEAD OF SESSION: **Dr. Le Dinh Trong & Assoc. Prof. Dr. Pham Duc Thuan**

AUTHORS	AFFILIATION	TOPIC TITLE
Fedwa BEGHDAI El-Hadj DRICHE	<i>Hassiba Benbouali University of Chlef ALGERIA</i>	PHYSIOLOGICAL AND BIOCHEMICAL CHARACTERIZATION OF SOME STRAINS OF ACTENOBACTERIA ISOLATED FROM AN ALGERIAN SAHARAN SOIL WITH A POTENTIAL FOR INHIBITION OF CERTAIN FUNGI AND PHYTOPATHOGENIC BACTERIA
Zineb El Hamri M. Karaoui M. Alami M. Assouag	<i>Moulay Ismail University MOROCCO</i>	THE IMPACT OF ADDING WALNUT SHELL POWDERS ON THE MORPHOLOGY, THERMAL PROPERTIES, AND MECHANICAL PERFORMANCE OF POLY(LACTIC ACID) (PLA)
Dhivya C. Arunkumar R.	<i>Tamil Nadu Agricultural University INDIA</i>	TECHNOLOGY TRANSFER AND ITS ROLE IN ENHANCING GLOBAL AGRICULTURAL PRODUCTIVITY
Oluwaseun Adekoya Adelaja Babafemi Raphael Babaniyi Olayinka Abigail Ajongbolo	<i>Federal University of Technology NIGERIA National Biotechnology Research and Development Agency NIGERIA</i>	BIOSORPTION OF METHYL RED, THYMOL BLUE AND THEIR DYE MIXTURES USING MODIFIED ELEPHANT GRASS, <i>Pennisetum purpureum</i>
Youssef El Hilali Alaoui Said Bouda Hajar Mhammedi Adil Essarioui	<i>Sultan Moulay Slimane University MOROCCO National Institute of Agricultural Research MOROCCO</i>	FUSARIUM ANTAGONISM IN SOIL AND RESOURCE COMPETITION DRIVE THE SUPPRESSION OF BAYOUD DISEASE IN MOROCCAN DATE PALM PLANTATIONS
Dr. Le Dinh Trong Assoc. Prof. Dr. Pham Duc Thuan	<i>Dong Thap University VIETNAM Can Tho University VIETNAM</i>	THE IMPACT OF THE U.S. DEVELOPMENT AID ON THE AGRICULTURE OF SOUTH VIETNAM (1954-1975)
Amira DEBAGHI Prof. Dr. Saida MESSGO-MOUMENE	<i>Saad Dahleb Blida 1 University ALGERIA</i>	VALORIZATION OF AGRICULTURAL RESIDUES FROM TOMATO PLANTS GROWN WITH ENDEMIC BACTERIAL ISOLATES
Dona Mary Eldhose Sejian V	<i>College of Veterinary and Animal Science INDIA Rajiv Gandhi Institute of Veterinary Education and Research INDIA</i>	PLANT SECONDARY METABOLITES TO REDUCE ENTERIC METHANE EMISSION IN DAIRY CATTLE
Nassima BAHA Sabrina BEHAIRI Roumaissa MOULKAF Massakib BEKKAYE	<i>University of Sciences and Technology Houari Boumedienne ALGERIA</i>	APPLICATION OF NEW BIOTECHNOLOGICAL PRACTICES FOR SUSTAINABLE AGRICULTURE: BIOFERTILIZATION

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-1 / SESSION-3



ANKARA LOCAL TIME: 15⁰⁰ - 17⁰⁰

HEAD OF SESSION: **Assoc. Prof. Dr. Hilal GASİMOV**

AUTHORS	AFFILIATION	TOPIC TITLE
Ağalarova Səma	<i>Nahcivan State University AZERBAIJAN</i>	CLINICAL USE OF HORMONES
Cavadxan QASIMOV Əşraf ƏLƏKBƏROV	<i>Nahcivan State University AZERBAIJAN</i>	THE IMPORTANCE OF AGRICULTURE IN THE SUSTAINABLE ECONOMIC DEVELOPMENT OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC
Assoc. Prof. Dr. Hilal GASİMOV Assoc. Prof. Dr. Anvar İBRAHİMOV Dr. Mursel SEYİDOV Assoc. Prof. Dr. Namig ABBASOV	<i>Nahcivan State University AZERBAIJAN</i>	USEFUL PROPERTIES AND USES OF SPECIES BELONGING TO THE GENERA BARBERRY (BERBERIS L.) AND IRGA (AMELANCHER MEDIK) COMMON IN THE AREA HAVUSH VILLAGE
Dr. Mursel SEYİDOV Assoc. Prof. Dr. Hilal GASİMOV Assoc. Prof. Dr. Namig ABBASOV Assoc. Prof. Dr. Anvar İBRAHİMOV Fidan HUSEYNOVA	<i>Nahcivan State University AZERBAIJAN</i>	NATURAL RESOURCES OF FRUIT PLANTS BELONGING TO THE GENERA APPLE (MALUS MILL.), PEAR (PYRUS L.) AND CHERRY (PRUNUS L.) FOUND IN THE HAYFIELD AND FOREST EDGE AREAS
Assoc. Prof. Dr. Namig ABBASOV Assoc. Prof. Dr. Hilal GASİMOV Dr. Mursel SEYİDOV Assoc. Prof. Dr. Bahruz MAMMEDOV Semennaz ABBASLI	<i>Nahcivan State University AZERBAIJAN</i>	SPECIES OF THE GENUS LONICERA L. IN THE FLORA OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC, THEIR PHYTOCHEMICAL PROPERTIES AND PROSPECTS OF USE (AZERBAIJAN)
Dashgin GANBAROV Safura BABAYEVA	<i>Nahcivan State University AZERBAIJAN</i>	USAGE DIRECTIONS OF SPECIES BELONGING TO THE GENUS ALCHEMILLA L. IN THE FLORA OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC
Leyla İBRAHİMOVA	<i>Nahcivan State University AZERBAIJAN</i>	AGRO-PRODUCTION CHARACTERISTICS OF SOILS IN THE NAKHCHIVAN AUTONOMOUS REPUBLIC (ON THE EXAMPLE OF SHAHBUZ AND ORDUBAD DISTRICTS)
Aydan MAHARRAMLİ Assoc. Prof. Dr. Mahir MAHARRAMOV	<i>Nahcivan State University AZERBAIJAN</i>	THE ROLE OF BEES OF THE GENUS HYLAEUS FABRICIUS, 1793 IN POLLINATION OF CORIANDER AND FENNEL PLANTS IN NAKHCHIVAN AUTONOMOUS REPUBLIC

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-2 / SESSION-3



ANKARA LOCAL TIME: 15⁰⁰ - 17⁰⁰

HEAD OF SESSION: **Assoc. Prof. Dr. Aysun ALTIKAT**

AUTHORS	AFFILIATION	TOPIC TITLE
Assoc. Prof. Dr. Aysun ALTIKAT	<i>Iğdır University TÜRKİYE</i>	NANOPESTICIDE USAGE IN SUSTAINABLE AGRICULTURE: AN ENVIRONMENTALLY FRIENDLY ALTERNATIVE?
Assoc. Prof. Dr. Aysun ALTIKAT	<i>Iğdır University TÜRKİYE</i>	CLIMATE THREATS ON AGRICULTURAL PRODUCTION: EXTREME WEATHER EVENTS AND WATER MANAGEMENT
Prof. Dr. İsmail GÜVENÇ Arzu AKBAŞ Soner AKCAN	<i>Kahramanmaraş Sütçü İmam University TÜRKİYE</i>	AGRICULTURE IN KAHRAMANMARAŞ
Agr. Eng. Zekiye ALTUNTEKİN Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	FRUIT SPECIES DAMAGED BY SANJOSE SCULBI (QUADRASPIDIOTUS PERNICIOSUS COMST.) (HEMIPTERA: DIASPIDIDAE) SEEN IN ORCHARDS IN İĞDIR PROVINCE AND THEIR DAMAGE STATUS
Agr. Eng. Zekiye ALTUNTEKİN Prof. Dr. Celalettin GÖZÜAÇIK	<i>Iğdır University TÜRKİYE</i>	DAMAGE SITUATION OF PEACH SHOOT MOTH, ANARSIA LINEATELLA (ZELLER) (LEPIDOPTERA: GELECHIIDAE) IN PEACH GARDENS OF İĞDIR PROVINCE
Assist. Prof. Dr. Boran KARATAŞ	<i>Van Yüzüncü Yıl University TÜRKİYE</i>	IMPACTS OF TEMPERATURE FLUCTUATIONS ON AQUACULTURE AND MITIGATION STRATEGIES
Nihal Ceren ALICI Prof. Dr. Mehmet Ali KIRPIK	<i>Kafkas University TÜRKİYE</i>	EFFECTS OF VERMICOMPOST APPLICATIONS ON YIELD PARAMETERS OF KAVILCA WHEAT

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-3 / SESSION-3



ANKARA LOCAL TIME: 15⁰⁰ - 17⁰⁰

HEAD OF SESSION: **Assoc. Prof. Dr. Tuncay KAYA**

AUTHORS	AFFILIATION	TOPIC TITLE
Res. Assist. Rıdvan TİK Assoc. Prof. Dr. Tuncay KAYA	<i>Iğdır University TÜRKİYE</i>	LANDSCAPE ARCHITECTURE AND ENERGY IN SMART CITIES
Res. Assist. Rıdvan TİK Assoc. Prof. Dr. Tuncay KAYA	<i>Iğdır University TÜRKİYE</i>	THE EFFECTS OF GREEN ROOF SYSTEMS ON ENERGY EFFICIENCY
Assoc. Prof. Dr. Emine NAKİLCİOĞLU Selin ALİHANOĞLU Gizem TİRYAKİ	<i>Ege University TÜRKİYE Aegean Agricultural Research Institute, Beekeeping Research Center İzmir TÜRKİYE</i>	INVESTIGATION OF POLYPHENOL CONTENTS AND ANTIOXIDANT CAPACITIES OF SOME MONOFLORAL POLLENS FROM THE AEGEAN REGION
Assoc. Prof. Dr. Emine NAKİLCİOĞLU Gizem TİRYAKİ	<i>Ege University TÜRKİYE</i>	SUPERFOOD OF THE FUTURE: SEAWEED
Res. Assist. Alperay ALTIKAT Prof. Dr. Mehmet Hakkı ALMA	<i>Iğdır University TÜRKİYE</i>	APPLICATIONS OF BIOPOLYMERS AND BIOPLASTICS IN SMART AGRICULTURE: SENSORS, PACKAGING, AND COATING MATERIALS
Res. Assist. Alperay ALTIKAT Prof. Dr. Mehmet Hakkı ALMA	<i>Iğdır University TÜRKİYE</i>	BIOPLASTIC USE IN AGRICULTURE: EFFECTS ON SOIL HEALTH AND PLANT DEVELOPMENT
Res. Assist. Alperay ALTIKAT Prof. Dr. Mehmet Hakkı ALMA	<i>Iğdır University TÜRKİYE</i>	BIOPOLYMER PRODUCTION FROM AGRICULTURAL WASTE: CIRCULAR ECONOMY AND SUSTAINABILITY
Lect. Zeynep ASUTAY Prof. Dr. Hakan İNCİ	<i>Bitlis Eren University TÜRKİYE Bingöl University TÜRKİYE</i>	BIOTECHNOLOGICAL CONTROL METHODS AGAINST VARROA DESTRUCTOR IN HONEY BEES (<i>Apis mellifera</i>)
Lect. Zeynep ASUTAY Prof. Dr. Hakan İNCİ	<i>Bitlis Eren University TÜRKİYE Bingöl University TÜRKİYE</i>	THE ROLE OF BEES IN THE POLLINATION ECOSYSTEM: A CRITICAL ACTOR IN BIODIVERSITY CONSERVATION

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-4 / SESSION-3



ANKARA LOCAL TIME: 15⁰⁰ - 17⁰⁰

HEAD OF SESSION: **Prof. Dr. Saida MESSGO-MOUMENE**

AUTHORS	AFFILIATION	TOPIC TITLE
Imene BENBELKACEM Prof. Dr. Aicha Bouhafoun Dr. Rawaa JAMALADDEEN Assoc. Prof. Dr. Bruno COUDOUR Res. Eng. Dr. Laurent LEMEE	<i>University of Science and Technology Mohamed Boudiaf (USTO-MB) ALGERIA</i> <i>University of Science and Technology Mohamed Boudiaf (USTO-MB) ALGERIA</i> <i>Poitiers University FRANCE</i>	BIO OIL FROM CORK: PYROLYSIS-GC-MS STUDY
Ali Imran Hafiz Suleria Safura Kousar Muhammad Usama Tahir	<i>Government College University PAKISTAN</i> <i>Melbourne University AUSTRALIA</i>	BIOACTIVE COMPOUNDS IN OILSEED WASTE: ANTIOXIDANT PROPERTIES AND LC-ESI-QTOF-MS/MS-BASED PROFILING
Ahmed Osmane Khadija Zidan Moustapha Belmouden	<i>Ibn Zohr University MOROCCO</i> <i>Cadi Ayyad University MOROCCO</i>	POTENTIAL FUNCTIONAL IMPLICATIONS OF PEARL MILLET IN HEALTH AND DISEASE, IRRIGATED BY TRICKLING FILTER-TREATED WASTEWATER IN SEMI-ARID CLIMATES IN NORTH AFRICA
Yousra DEBBAH Prof. Mohamed BENCHERCHALI Prof. Massgo-Saida MOUMEN	<i>Blida 1 University ALGERIA</i>	EVALUATING THE ROLE OF ENDEMIC BENEFICIAL MICROORGANISMS STRAINS IN BOOSTING THE NUTRITIONAL QUALITY OF <i>Medicago sativa</i> L.
Tawseef Ahmad Dar Assist. Prof. Dr. Yash Paul Assoc. Prof. Dr. Mohammad Ahsan Chishti	<i>Central University INDIA</i> <i>National Institute of Technology Srinagar INDIA</i>	PRECISION AGRICULTURE: "EFFICIENT DEEP LEARNING FOR REAL-TIME AGRICULTURAL IOT SYSTEMS"
BRIKCI Fatiha BOUHAFSOUN Aicha BOKHARI Hassiba DJABEUR Abderrezak	<i>University of Science and Technology of Oran Mohamed Boudiaf ALGERIA</i>	BIOMETRIC ANALYSIS OF THE FIBERS OF A LOCAL ALGERIAN PLANT SPECIES AND THEIR VALORIZATION
Amira DEBAGHI Prof. Dr. Saida MESSGO-MOUMENE	<i>Saad Dahleb Blida 1 University ALGERIA</i>	INFLUENCE OF BACTERIAL BIOSTIMULANTS ON THE BIOCHEMICAL COMPOSITION OF <i>SOLANUM TUBEROSUM</i> L
Moufida ABABOU Halima SAIAH	<i>Hassiba Ben Bouali University ALGERIA</i>	ANTIOXIDANT ACTIVITY, BIOACTIVE MOLECULES AND TOXICITY STUDY OF AN ALIMENTARY BYPRODUCT HYDROETHANOLIC EXTRACT

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.



ONLINE PRESENTATIONS

05.03.2025 / HALL-5 / SESSION-3



ANKARA LOCAL TIME: 15⁰⁰ - 17⁰⁰

HEAD OF SESSION: **Prof. Dr. Lynda LAMOUDI**

AUTHORS	AFFILIATION	TOPIC TITLE
Marwa Waheed Muhammad Bilal Hussain Fatima Waheed Nida Kanwal	<i>Riphah International University</i> PAKISTAN <i>Government College University</i> PAKISTAN <i>Agriculture University</i> PAKISTAN	TRANSFORMING FOOD SAFETY AND SUSTAINABILITY: THE ROLE OF MODERN FOOD PROCESSING TECHNOLOGIES
V. S. Angulakshmi N. Anusuya R Nithya S Kalaiselvan	<i>PSGR Krishnammal College for Women</i> INDIA	GREEN SYNTHESIS OF COPPER OXIDE NANOPARTICLES USING CITRUS SINENSIS PEEL EXTRACT AND EXPLORATION OF THEIR ANTIBACTERIAL ACTIVITIES
Rabar Mohammed Hussein	<i>Noble Technical Institute</i> IRAQ	GROWTH DYNAMICS OF MARINE MICROALGAE IN BATCH CULTURES
Mohamed LAHRAIRI Karima JMAILI Bouchaib BAHLAOUAN Hlima BESSI Nadia BOUTALEB	<i>Hassan II University</i> MOROCCO	COMPARATIVE ANALYSIS OF FERMENTED ALMOND MILK USING MOROCCAN ALMOND VARIETIES: A STUDY ON MICROBIOLOGICAL AND NUTRITIONAL PROPERTIES
AKRETCHÉ Soraya BOUSBIA Nabil LAMOUDI Lynda KERBOUCHE Lamia DJILI Manel KACI Ismahan	<i>University of Science and Technology Houari Boumediene (U.S.T.H.B.)</i> ALGERIA	STUDY OF THE QUALITY OF A DAIRY PRODUCT ENRICHED WITH POLLEN
AKRETCHÉ Soraya LAMOUDI Lynda BOUSBIA Nabil KERBOUCHE Lamia DAHMANI Chaima	<i>University of Science and Technology Houari Boumediene (U.S.T.H.B.)</i> ALGERIA	QUALITY CONTROL OF ALGERIAN OLIVE OIL
Prof. Dr. Lynda LAMOUDI Prof. Dr. Soraya AKRETCHÉ Assoc. Prof. Dr. Nabil BOUSBIA Assoc. Prof. Dr. Nadjet GUEMRAS Prof. Dr. Kamel DAOUD	<i>University of Science and Technology Houari Boumediene (U.S.T.H.B.)</i> ALGERIA	EVALUATION OF THE INSECTICIDAL ACTIVITY OF ESSENTIAL OILS FROM LAVANDULA STOECHAS AND ORIGANUM FLORIBUNDUM FROM NORTHERN ALGERIA
Farzana Iftikhar Faisal Iftikhar	<i>Women University</i> PAKISTAN <i>Bahauddin Zakariya University Multan</i> PAKISTAN	EVALUATION OF GENETIC DIVERSITY AND GENOME FINGERPRINTING OF BITTER GOURD GENOTYPES (MOMORDICA CHARANTIA) BY MOLECULAR MARKER

All participants must join the conference 10 minutes before the session time. Every presentation should last no longer than 10-12 minutes. Kindly keep your cameras on till the end of the session.

CONTENTS

AUTHOR	TITLE	No
Muhammad Mazhar Iqbal Muhammad Abdullah Tehmena Rashid Hafiz Qaiser Yasin Shafiq-ur-Rehmn	ANALYSIS OF HYDRO-METEOROLOGICAL TRENDS ACROSS MAJOR KOPPEN CLIMATE CLASSES USING GLDAS DATA AT A GLOBAL SCALE	1
Kaouthar ELBIRGUI Houssam ASSIOUI Oussama RAFIK Nadia IOUNES Samir IBENMOUSSA	INVESTIGATING THE ANTIOXIDANT POTENTIAL OF EBENUS PINNATA AITON: IN VITRO AND IN VIVO ASSESSMENTS OF HYDROETHANOLIC EXTRACTS	2
GUNASEKAR G	EVOLUTION, BEHAVIOR AND ECOLOGY OF FEATHER MITE	4
EL BAHJA Fatima ZIRI Rabea BRITAL Rania HAMRANI Meriem EL AMMARI Mohamed FAHAD Kaoutar	STUDY OF CERATITIS CAPITATA WIEDEMANN (DIPTERA: TEPHRITIDAE) POPULATION DYNAMICS IN THE EL GHARB REGION OF MOROCCO FOR FUTURE PEST MANAGEMENT	5
Sundhararajan R Jothilakshmi.R S.G.Raman	GREEN INNOVATIONS IN MEDICINAL AND AROMATIC PLANT RESEARCH	7
N Anusuya V S Angulakshmi M Priyadharshini G Easterin Shejina	GREEN MEDIATED SYNTHESIS, CHARACTERIZATION AND APPLICATION OF MANGANESE OXIDE NANOPARTICLES USING CITRUS LIMON PEEL EXTRACT	8
Le Dinh Trong Pham Duc Thuan	THE IMPACT OF THE U.S. DEVELOPMENT AID ON THE AGRICULTURE OF SOUTH VIETNAM (1954-1975)	9
IOANA POROSNICU Madalina DAVIDESCU Bianca-Maria MADESCU	MIGRATION, CLIMATE CHANGE AND SUSTAINABLE AGRICULTURE: CHALLENGES AND OPPORTUNITIES FOR ECONOMIC DEVELOPMENT	17
A. Attahiru A.U Birnin-Yauri M. Mahdi-Adamu A. Yahaya	PHYTOCHEMICAL CONSTITUENTS OF MORINGA OLEIFERA ROOTS EXTRACT	18
Muslim Yusuf Sulaiman Sani Yusuf Abdulaziz Hassan Maidabino	PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL ACTIVITY ON THE ETHANOLIC EXTRACT OF GUIERA SENEGALENSIS LEAVES	19
Marwa Waheed Muhammad Bilal Hussain Fatima Waheed Nida Kanwal	TRANSFORMING FOOD SAFETY AND SUSTAINABILITY: THE ROLE OF MODERN FOOD PROCESSING TECHNOLOGIES	20
M. Mahdi-Adamu F. Halidu A. Attahiru A. Yahaya R. Adamu	NUTRITIONAL AND PHYTOCHEMICAL ANALYSIS OF OREGANO (ORIGANUM VULGARE) LEAVES	21

Oyewole Mojisola Fauziyah	ACCEPTABILITY OF ORANGE FLESHED SWEET POTATO CONFECTIONARIES AMONG FACULTY OF AGRICULTURE STUDENTS, UNIVERSITY OF IBADAN, NIGERIA	22
BELLIL Zahra MEYER Sylvain BELLIL Dassine MAIRI Assia BARRAUD Olivier TOUATI Abdelaziz	OCCURRENCE AND CHARACTERIZATION OF ANTIBIOTIC-RESISTANT SALMONELLA ISOLATES FROM FRESH PRODUCE IN AGRICULTURAL ENVIRONMENTS	23
M.M Akibu M.S. Na-Allah A.Muhammad I. J. Yusuf	YIELD AND STORABILITY OF ONION ECOTYPES AS INFLUENCED BY IRRIGATION INTERVAL IN SUDAN SAVANNA ZONE OF NIGERIA	25
ZAGGOUMI Hasna BOUDA Said NEG Imane MARGHALI Sonia TRIFI Neila HADDIOUI Abdelmajid	THE GENETIC CHARACTERIZATION OF THE MOROCCAN CAROB TREE (CERATONIA SILIQUA L.) USING ISSR MARKERS	27
Hafiza Zara Saeed Muhammad Wajid Musrat Shaheen	EFFICACY OF ESSENTIAL ELEMENTS AGAINST SUCKING INSECT PESTS OF COTTON UNDER DIFFERENT CONDITION	29
Karima JMAILI Khawla WADDI Bouchaib BAHLAOUAN Alla SILKINA Nadia BOUTALEB	THE USE OF WHEY TO ENHANCE TOMATO AND WHEAT SEED GERMINATION: A PATHWAY TO VALORIZING DAIRY BY-PRODUCTS FOR SUSTAINABLE AGRICULTURE	30
Mohamed LAHRAIRI Karima JMAILI Bouchaib BAHLAOUAN Hlima BESSI Nadia BOUTALEB	COMPARATIVE ANALYSIS OF FERMENTED ALMOND MILK USING MOROCCAN ALMOND VARIETIES: A STUDY ON MICROBIOLOGICAL AND NUTRITIONAL PROPERTIES	32
Imene BENBELKACEM Aicha Bouhafoun Rawaa JAMALADDEEN Bruno COUDOUR Laurent LEMEE	BIO OIL FROM CORK : PYROLYSIS-GC-MS STUDY	34
BRIKCI Fatiha BOUHAFSOUN Aicha BOKHARI Hassiba DJABEUR Abderrezak	BIOMETRIC ANALYSIS OF THE FIBERS OF A LOCAL ALGERIAN PLANT SPECIES AND THEIR VALORIZATION	36
Oluwaseun Adekoya Adelaja Babafemi Raphael Babaniyi Olayinka Abigail Ajongbolo	BIOSORPTION OF METHYL RED, THYMOL BLUE AND THEIR DYE MIXTURES USING MODIFIED ELEPHANT GRASS, Pennisetum purpureum	37
Meriem RAFYA Naima ZEHAR Fatiha BENKHALTI	THE NEGATIVE EFFECT OF SOIL AMENDMENTS WITH SOLID RESIDUES OF ROSMARINUS OFFICINALIS L. ON THE GROWTH OF SOLANUM LYCOPERSICUM MILL	38
Ahlem DEBIEB Leila KADIK Michel GODRON Zineb DJERRAD Nacere ADI	THE RESILIENT FOREST AFTER THE FIRE: ANALYSIS OF THE STRUCTURE AND DIVERSITY OF THE EL-HAMDANIA CORK OAK FOREST- ALGERIA	39

Yousra DEBBAH Mohamed BENCHERCHALI Massgo-Saida MOUMEN	EVALUATING THE ROLE OF ENDEMIC BENEFICIAL MICROORGANISMS STRAINS IN BOOSTING THE NUTRITIONAL QUALITY OF <i>Medicago sativa</i> L.	41
Samira KAROU	UPDATE OF ALGERIA'S SOIL MAP AT A 1/500,000 SCALE: APPLICATION OF GOOGLE EARTH IMAGERY	42
Lynda LAMOUDI Soraya AKRETICHE Nabil BOUSBIA Nadjet GUEMRAS Kamel DAOUD	EVALUATION OF THE INSECTICIDAL ACTIVITY OF ESSENTIAL OILS FROM <i>LAVANDULA STOECHAS</i> AND <i>ORIGANUM FLORIBUNDUM</i> FROM NORTHERN ALGERIA	43
Nadira OUKALA ORERO BAYO Marta Victoria PASTOR	ENDOPHYTIC BACTERIUM <i>BACILLUS CEREUS</i> - MEDIATED SALT TOLERANCE IN WHEAT PLANTS	45
Thinhinan KHEDIM	THE CASE OF ROSY GARLIC (<i>ALLIUM ROSEUM</i> L.) POLYPLOID COMPLEX IN ALGERIA	47
HADJLOUNE Houria KACI Ahcène	THE INFLUENCE OF INSTITUTIONAL DYNAMICS AND GOVERNANCE STRUCTURES ON THE SUPPLY CHAIN OF THE MILK SECTOR IN ALGERIA: CASE OF THE DAIRY HODNA-LAIT (M'SILA)	48
Moufida ABABOU Halima SAIHAH	ANTIOXIDANT ACTIVITY, BIOACTIVE MOLECULES AND TOXICITY STUDY OF AN ALIMENTARY BYPRODUCT HYDROETHANOLIC EXTRACT	49
Dona Mary Eldhose Sejian V	PLANT SECONDARY METABOLITES TO REDUCE ENTERIC METHANE EMISSION IN DAIRY CATTLE	50
Sarra Farjallah	DNA MOLECULAR MARKERS AND THEIR APPLICATIONS IN INVESTIGATING THE GENETIC VARIABILITY OF THE NEMATODE <i>HETERAKIS GALLINARUM</i> FROM TUNISIA	51
Fedwa BEGHDADI El-Hadj DRICHE	PHYSIOLOGICAL AND BIOCHEMICAL CHARACTERIZATION OF SOME STRAINS OF ACTENOBACTERIA ISOLATED FROM AN ALGERIAN SAHARAN SOIL WITH A POTENTIAL FOR INHIBITION OF CERTAIN FUNGI AND PHYTOPATHOGENIC BACTERIA	53
AOUFI Farida TAMERT Asma ABDI Yamina BARBARA Ferdaws	ANTIBACTERIAL AND ANTIFUNGAL ACTIVITY OF PLANT (<i>MARRUBIUM VULGARE</i>) FROM THE FAMILY OF LAMIACEAE IN WESTERN ALGERIA	54
V. S. Angulakshmi N. Anusuya R Nithya S Kalaiselvan	GREEN SYNTHESIS OF COPPER OXIDE NANOPARTICLES USING CITRUS SINENSIS PEEL EXTRACT AND EXPLORATION OF THEIR ANTIBACTERIAL ACTIVITIES	55
Said Babou Miloud Chakit Radia El Gui Abdelhalem Mesfioui Youssef Sqalli-Houssaini	TOXICITY ASSESSMENT OF ETHANOLIC EXTRACT OF <i>ROSMARINUS OFFICINALIS</i> LEAVES IN FEMALE WISTAR RATS	56
VIGNESH K K. SELVAM ARSHA G	EFFECT OF NEW GENERATION FUNGICIDES AGAINST <i>Pyricularia oryzae</i> CAUSES BLAST IN RICE (<i>Oryza sativa</i> L.)	57
Ihaddaden Akli ABLA Safia Abdoun Fatiha	PHYSIONOMIC CHANGE OF VEGETATION AND DIVERSITY INDEX IN TWO CONSTRAINED ENVIRONMENT OF THE SEMI-ARID IN ALGERIA	58

Soumaya Atouife Jamila Elbiyad Halima Belaoufi Abdellah El Habazi Rekia Belahsen	UNDERSTANDING MOROCCAN POPULATION'S PERCEPTION AND AWARENESS OF ORGANIC FOOD	59
Said Babou Miloud Chakit Abdelhalem Mesfioui Youssef Sqalli Houssaini	ANTILIPIDEMIC AND NEPHRO-HEPATOPROTECTIVE ACTIVITIES OF ROSMARINUS OFFICINALIS ETHANOLIC EXTRACT IN WISTAR RATS	60
Youssef El Hilali Alaoui Said Bouda Hajar Mhammedi Adil Essarioui	FUSARIUM ANTAGONISM IN SOIL AND RESOURCE COMPETITION DRIVE THE SUPPRESSION OF BAYOUD DISEASE IN MOROCCAN DATE PALM PLANTATIONS	61
Ali Imran Hafiz Suleria Safura Kousar Muhammad Usama Tahir	BIOACTIVE COMPOUNDS IN OILSEED WASTE: ANTIOXIDANT PROPERTIES AND LC-ESI-QTOF- MS/MS-BASED PROFILING	62
Muhammad Muzammil Nazir Munaza Yasmeen Iqra Farzeen Sana Inam Asma Ashraf	THE STUDY OF ANTIDIARRHEAL POTENTIAL OF HELICTERES ISORA AND PERSICARIA VIVIPARA METHANOLIC EXTRACTS IN SWISS ALBINO MICE	63
Dhivya C Arunkumar R	TECHNOLOGY TRANSFER AND ITS ROLE IN ENHANCING GLOBAL AGRICULTURAL PRODUCTIVITY	65
Nassima BAHA Sabrina BEHAIRI Roumaissa MOULKAF Massakib BEKKAYE	APPLICATION OF NEW BIOTECHNOLOGICAL PRACTICES FOR SUSTAINABLE AGRICULTURE: BIOFERTILIZATION	66
Farzana Iftikhar Faisal Iftikhar	EVALUATION OF GENETIC DIVERSITY AND GENOME FINGERPRINTING OF BITTER GOURD GENOTYPES(MOMORDICA CHARANTIA) BY MOLECULAR MARKER	67
LAZREG Halima SEBAA Hanane Siham CELIK Muhammed Suleyman ERENLER Ramazan ABDEDDAIM Katia Khadidja DJABEUR Abd Erezzak	PHENOLIC COMPOUNDS OF OPUNTIA STRICTA CLADODES	68
Soukaina El Bourachdi Abdelhay El Amri Ali Raza Ayub Fatima Moussaoui Amal Lahkimi	DEVELOPMENT OF A NOVEL LOW-COST ADSORBENT CHITOSAN@EDTA@CELLULOSE COMPOSITE TO EFFECTIVELY REMOVE METHYL ORANGE DYE FROM WASTEWATER: EXPERIMENTAL AND THEORETICAL INVESTIGATION	69
Nacéra TADJINE Saida MESSGO-MOUMENE	IN VITRO EVALUATION OF THE ANTIFUNGAL POTENTIAL OF MYRTUS COMMUNIS L. AGAINST TOXIGENIC MOLDS OF HYDROPONIC BARLEY	70
Abdul Ahad Ansari Haris Hasan Khan	ADVANCEMENTS IN REMOTE SENSING FOR CROP AND SOIL HEALTH MANAGEMENT IN SUSTAINABLE AGRICULTURE	71
Zineb El Hamri M. Karaoui M. Alami M. Assouag	THE IMPACT OF ADDING WALNUT SHELL POWDERS ON THE MORPHOLOGY, THERMAL PROPERTIES, AND MECHANICAL PERFORMANCE OF POLY(LACTIC ACID) (PLA)	72

Fjorela MEMA Selma MYSLIHAKA Peçi NAQELLARI Nikoleta KALLAJXHIU Silvana TURKU Anxhela DAUTI	GEOBOTANICAL DATA ON FLORA AND VEGETATION IN SOME AREAS OF ELBASAN	73
AKRETCHE Soraya LAMOUDI Lynda BOUSBIA Nabil KERBOUCHE Lamia DAHMANI Chaima	QUALITY CONTROL OF ALGERIAN OLIVE OIL	74
AKRETCHE Soraya BOUSBIA Nabil LAMOUDI Lynda KERBOUCHE Lamia DJILI Manel KACI Ismahan	STUDY OF THE QUALITY OF A DAIRY PRODUCT ENRICHED WITH POLLEN	76
BRITAL Rania Abail Zhor MRABTI Imane EL bahja Fatima IBRIZ Mohammed	SURVEY ON FERTILIZATION PRACTICES AND NUTRIENT MANAGEMENT IN VALENCIA LATE CITRUS ORCHARDS: A CASE STUDY FROM THE GHARB REGION (NORTHEASTERN), MOROCCO	78
MRABTI Imane GRIJJA Hassan BRITAL Rania BRHADDA Najiba ZIRI Rabea AFECHTAL Mohamed	IMPROVEMENT OF DNA EXTRACTION FOR THE PYRIFORM SCALE INSECT, PROTOPULVINARIA PYRIFORMIS, FOR MOLECULAR IDENTIFICATION OF CITRUS PESTS IN MOROCCO USING DNA BARCODING	80
MRABTI Imane GRIJJA Hassan BENZAHERA Hayat AIT BADDOU Sanae BRHADDA Najiba AFECHTAL Mohamed ZIRI Rabea	PRECISE IDENTIFICATION OF THE COTTONY CUSHION SCALE, ICERYA PURCHASI IN MOROCCO THROUGH THE ADVANCED APPLICATION OF DNA BARCODING	82
Hana ZAKKOUMI Yazid MAHDAOUI Houria HADJ-ARAB	MORPHOLOGICAL DIVERSITY AND STYLAR POLYMORPHISM IN NARCISSUS SPECIES (AMARYLLIDACEAE) FROM NORTHERN ALGERIA	84
Famela A. Foyo Bernadette I. Fillartos Cristine Angeli T. Mangampat Jane M. Balubar Klenton R. Torrevillas, MIHM	INNOVATIVE BAKING: UTILIZING KAMIAS AND BANANA-ENRICHED FLOUR IN CAKE RECIPES INSPIRED BY LEMON CAKE	94
Sadiq, M.S Singh, I.P Ahmad, M.M Sani, B.S Nazifi, I.K	SEED-TO-SHELF VENTURES: BUILDING A YOUTH- LED AGRIBUSINESS MODEL FOR FARM-TO-MARKET TRANSPARENCY	116
Farrah Deebe Muhammad Huzaifah Khalid Anas Sarwar Qureshi Muhammad Adil	THERAPEUTIC STUDY OF CENTRATHERUM ANTHELMINTICUM FOR THE CONTROL OF SUBCLINICAL MASTITIS IN BUBALINE	136
Muhammad Bilal Hussain Marwa Waheed Farhan Saeed Muhammad Afzaal Mariam Islam	NUTRITIONAL AND PHARMACOLOGICAL PROPERTIES OF CANTALOUPE PEEL: AN OVERVIEW	137

Rabar Mohammed Hussein	GROWTH DYNAMICS OF MARINE MICROALGAE IN BATCH CULTURES	138
Ahmed Osmane Khadija Zidan Moustapha Belmouden	POTENTIAL FUNCTIONAL IMPLICATIONS OF PEARL MILLET IN HEALTH AND DISEASE , IRRIGATED BY TRICKLING FILTER-TREATED WASTEWATER IN SEMI-ARID CLIMATES IN NORTH AFRICA	146
Amira DEBAGHI Saida MESSGO-MOUMENE	INFLUENCE OF BACTERIAL BIOSTIMULANTS ON THE BIOCHEMICAL COMPOSITION OF SOLANUM TUBEROSUM L	147
Sefa ALTIKAT	THE EFFECT OF ALFALFA SOWING ON SOIL ORGANIC MATTER CONTENT IN İĞDIR PROVINCE: A REMOTE SENSING BASED EVALUATION	148
Sefa ALTIKAT	CHANGE ON ALFALFA AGRICULTURE: 2023 AND 2025 COMPARATIVE ANALYSIS BASED ON SATELLITE DATA	157
Pınar ADIGÜZEL Şenay KARABIYIK İlknur SOLMAZ	APPLICATIONS TO ENHANCE POLLEN QUALITY IN WATERMELONS WITH DIFFERENT CHROMOSOME STRUCTURES	173
Afşin AYHAN Banu ÇİÇEK KURDOĞLU	MULTI-CRITERIA DECISION-MAKING METHODS (MCDM) FOR SUSTAINABLE ENVIRONMENTAL MANAGEMENT: OPTIMIZED DECISION SUPPORT SYSTEMS WITH RSTUDIO	186
Abdullah İREÇ Celalettin GÖZÜAÇIK	DAMAGE SITUATION OF MEDITERRANEAN FRUIT FLY, CERATITIS CAPITATA (WIEDEMANN) (DIPTERA: TEPHRITIDAE) IN PEACH GARDENS OF İĞDIR PROVINCE	204
Rıdvan TİK Tuncay KAYA	LANDSCAPE ARCHITECTURE AND ENERGY IN SMART CITIES	211
Haydar BALCI Murat KARA Muhsin YILDIZ	EFFECTS OF LIQUID VERMICOMPOST, SALICYLIC ACID, AND PROLINE APPLICATIONS ON THE GROWTH OF GREEN ONION (ALLIUM CEPA L.)	223
Alperay ALTIKAT Mehmet Hakkı ALMA	APPLICATIONS OF BIOPOLYMERS AND BIOPLASTICS IN SMART AGRICULTURE: SENSORS, PACKAGING, AND COATING MATERIALS	225
Aysun UYSAL Yusuf GÜMÜŞ Soner SOYLU Merve OĞUZ Emine Mine SOYLU	DETERMINATION OF IN VITRO ANTIFUNGAL ACTIVITY OF WOOD VINEGAR AGAINST ROSALLINIA ROOT ROT DISEASE AGENT IN FIG TREES	238
Haydar BALCI Muhsin YILDIZ Murat KARA	EFFECTS OF PROLINE AND SALICYLIC ACID APPLICATION ON YIELD AND PLANT GROWTH IN GREEN ONION CULTIVATION	245
Ezgi DEMİRAL	A CRITICAL ANALYSIS OF TURKEY'S AGRICULTURAL POLICIES IN THE CONTEXT OF COMBATING CLIMATE CHANGE AND SUSTAINABLE AGRICULTURE	247
Alperay ALTIKAT Mehmet Hakkı ALMA	BIOPLASTIC USE IN AGRICULTURE: EFFECTS ON SOIL HEALTH AND PLANT DEVELOPMENT	255
Aysun UYSAL Şener KURT Tuğba HANEDAN	ABIOTIC STRESS FACTORS AND THEIR EFFECTS IN STONE FRUIT ORCHARDS OF AMIK PLAIN	265
Alperay ALTIKAT Mehmet Hakkı ALMA	BIOPOLYMER PRODUCTION FROM AGRICULTURAL WASTE: CIRCULAR ECONOMY AND SUSTAINABILITY	272

Akim AKHUNDOV Sahib HACIYEV Elnara SEYİDOVA	ECOLOGICAL ASSESSMENT OF THE SOILS OF ABRAGUNUS DEPRESSION	284
Aydan MAHARRAMLI Mahir MAHARRAMOV	THE ROLE OF BEES OF THE GENUS HYLAEUS FABRICIUS, 1793 IN POLLINATION OF CORIANDER AND FENNEL PLANTS IN NAKHCHIVAN AUTONOMOUS REPUBLIC	291
Ali TAHİROV Yunis RUSTAMLI	IMPACT OF GLOBAL CLIMATE CHANGE ON BEEKEEPING	293
Novruzlu Aytən Əli Qurbanov	THE ECOLOGICAL RESULTS OF IMPACT OF THE ARPACHAY RESERVOIR IRRIGATION SYSTEMS TO THE ENVIRONMENT	295
Mustafa ÇETİNER Ahmet Zafer TEL	FLORA AND ENDEMIC PLANTS OF ATATURK DAM BASIN (ŞANLIURFA/TURKEY)	305
Waffa ABDELMOUMENE Nabila BELYAGOUBI- BENHAMMOU Zineb KAID Larbi BELYAGOUBI	COMPARATIVE ASSESSMENT OF ANTIOXIDANT ACTIVITY AND PHYTOCHEMICAL PROFILES OF AQUEOUS EXTRACTS FROM ATRIPLEX HALIMUS LEAVES AND STEMS	306
Orhan GÜNGÖR	EFFECTS OF BATTERY TECHNOLOGIES ON AGRICULTURAL MACHINERY: EFFICIENCY, SUSTAINABILITY AND FUTURE PERSPECTIVES	308
Aydın AKKAYA	REASONS FOR WHITE SPIKE FORMATION IN WHEAT AND DIAGNOSIS	315
Ali ÇELİK Furkan DİNÇER	SUSTAINABILITY OF SALEP PRODUCTION THROUGH AGRIVOLTAIC APPLICATIONS: THE KALE TARIMGES MODEL	327
Ali ÇELİK Furkan DİNÇER	INTEGRATION OF RENEWABLE ENERGY AND AGRICULTURE WITH AGRIVOLTAICS: THE CASE OF KAHRAMANMARAŞ BULANIK TARIMGES	332
Boran KARATAŞ	IMPACTS OF TEMPERATURE FLUCTUATIONS ON AQUACULTURE AND MITIGATION STRATEGIES	339
Aydın AKKAYA	WHEAT BASED INTERCROPPING	341
Necibe KAYAK Önder TÜRKMEN	USE OF MICROBIAL AGENTS AGAINST SOIL SALINITY STRESS IN VEGETABLES	352
Ergün ÖZUSLU Ahmet Zafer TEL	DETERMINATION OF EUNIS HABITAT TYPES OF SOME NATURAL PROTECTED AREAS IN ADANA (TÜRKİYE) PROVINCE	362
Berna BAŞ	IS VERTICAL FARMING SUSTAINABLE?	381
Dilek DOĞAN Celalettin GÖZÜAÇIK	HARMFUL INSECTS OF THE HETEROPTERA (HEMIPTERA) SUBORDER SEEN ON FRUIT TREES IN IĞDIR PROVINCE	391
Etibar MEMMEDOV	THE ANTHELMINTIC EFFECT OF PREPARATIVE FORMS OF TANACERUM VULGARE L. ON GASTROINTESTINAL NEMATODES IN SHEEP	403
Şeyma KAYA Feride MERCAN Akide ÖZCAN Mehmet SÜTYEMEZ	EFFECT OF HIGH TEMPERATURE ON WALNUT CULTIVATION	405
Halit Burak YÜKSEKDAĞ Hayrullah ÖZBEK Mustafa Can SARIOĞLU Mustafa ARSLAN Mustafa BÖLÜK	SUSTAINABLE AGRICULTURE AND SUGAR BEET: AN ANALYSIS OF CARBON FOOTPRINT, WATER USAGE, AND EFFICIENCY	407

Tahsin UYGUN Mesut OŞLU Mete YİĞİT İlker ÜNAL	DETERMINATION OF ORANGE FRUIT RIPENESS LEVELS USING DEEP LEARNING-BASED INSTANCE SEGMENTATION METHOD	409
İlknur İŞÇİ Gülcan KAYMAK BAYRAM Zeki ACAR İlknur AYAN	THE ROLE OF LOCAL GOVERNMENTS IN SUSTAINABLE AGRICULTURE: THE ISTANBUL CASE	425
İlyas KILINÇER Özgür ÖZMEN Anil Mehmet BALTACI Mehmet ARSLAN	DETERMINATION OF YIELD AND QUALITY PARAMETERS OF SUGAR BEET PARENTAL LINES UNDER LIMITED IRRIGATION	435
Yunus RUSTAMLI Ali TAHIROV	IN THE REPUBLIC OF AZERBAIJAN LEGAL BASIS OF SWITCHING TO ORGANIC BEEKEEPING	437
Muhammed İkbal ÇATAL Adil BAKOĞLU	EVALUATION OF THE FORAGE QUALITY AND NUTRITIONAL VALUE OF THE PASTURE AREA OF ULUTAŞ NEIGHBORHOOD OF İSPİR DISTRICT IN ERZURUM PROVINCE	439
Ayfer GÜNEY SARITAŞ Ramazan GÜRBÜZ	INNOVATIVE BIOCONTROL STRATEGIES FOR SUSTAINABLE WEED MANAGEMENT	449
Ayfer GÜNEY SARITAŞ Ramazan GÜRBÜZ	INTEGRATION OF COVER CROPS IN WEED SUPPRESSION AND SOIL FERTILITY	459
İsmail GÜVENÇ Arzu AKBAŞ Soner AKCAN	AGRICULTURE IN KAHRAMANMARAŞ	466
İlayda Gizem AKIN Ozge SIZMAZ	USE OF THE ASTAXANTHIN IN TILAPIA (OREOCHROMIS NILOTICUS)	468
Hayrullah OZBEK Halit Burak YUKSEKDAĞ Oguzhan ULU Anil Mehmet BALTACI	EFFECT OF BIOLOGICAL AND ORGANIC FERTILIZERS ON YIELD AND QUALITY IN ORGANIC SUGAR BEET FARMING	476
Elsever Asadov Sevgi Kolaylı	COMPARATIVE STUDY OF SOME SUGAR PROFILES OF HONEY AND FRUITS	478
Emine ATALAY	SEED COATING TECHNOLOGIES FROM A SUSTAINABLE AGRICULTURE PERSPECTIVE	482
Adnan AYDIN	MOLECULAR MARKER TECHNIQUES FOR THE DETECTION OF DNA CYTOSINE METHYLATION	495
Fatih KUMBASAR Mehmet CAN Gülcan KAYMAK BAYRAM İlknur AYAN Zeki ACAR	THE IMPORTANCE OF RANGELANDS AS CARBON SINKS	502
Musa KARADAĞ Yunus BAŞAR Mehmet Hakkı ALMA	PHYTOCHEMICAL CONTENT OF LAURUS NOBILIS LEAVES AND EXTRACT; POTENTIAL FOR USE IN VALUE-ADDED COSMETICS, HEALTH AND FOOD PRODUCTS	515
Gönül USTA GEBEŞ Celalettin GÖZÜAÇIK	BIOLOGICAL CONTROL OF STORED PRODUCT DAMAGE	522
Ruhşen AYDIN KARAAĞAÇ Ahmet Taner KALE	EVALUATION OF THE EFFECTS OF SHEEP MANURE, WORM COMPOSITION AND MELASSES APPLICATIONS ON BLACK ARONIA (Aronia Melanocarpa) YIELD	530

Rüştü ÇALLI Emirhan ÖZDEMİR Mitat Can YILDIZ Aliihsan ŞEKERTEKİN	DETECTION AND ANALYSIS OF BURNED AREAS WITH SENTINEL-2 SATELLITE IMAGERY AND FIRE SEVERITY MAP: 2025 LOS ANGELES FIRE	539
Mitat Can YILDIZ Emirhan ÖZDEMİR Rüştü ÇALLI Aliihsan ŞEKERTEKİN	BIBLIOMETRIC ANALYSIS OF THE USE OF SATELLITE-BASED REMOTE SENSING TECHNOLOGIES IN CROP MAPPING (2018-2025)	549
Aslihan KULOGLU Aysun CAVUSOGLU	TISSUE CULTURE STUDIES ON CYCLAMEN SPP.	563
Leyla IBRAHIMOVA	AGRO-PRODUCTION CHARACTERISTICS OF SOILS IN THE NAKHCHIVAN AUTONOMOUS REPUBLIC (ON THE EXAMPLE OF SHAHBUZ AND ORDUBAD DISTRICTS)	569
Muhammet ÖTER Celalettin GÖZÜAÇIK	PESTICIDE USE IN İĞDIR PROVINCE	574
Ağalarova Səma	CLINICAL USE OF HORMONES	584
Nagihan KILIÇ Celalettin GÖZÜAÇIK	IMPACT OF CLIMATE CHANGE ON INTEGRATED PEST MANAGEMENT (IPM) IN TURKEY	587
Ayşe Altinkaya Melahat Özge Özen Nur Ülger Büşra Ülger	TOMATO YELLOW LEAF CURL VIRUS (TYLCV) AND CONTROL METHODS IN SOLANACEAE	595
Osman Doğan BULUT Köksal KARADAŞ	EXAMINATION OF FARMERS' KNOWLEDGE AND AWARENESS ON THE ENVIRONMENTAL EFFECTS OF PESTICIDE USE: THE CASE OF İĞDIR PROVINCE	626
Emine NAKİLCİOĞLU Selin ALİHANOĞLU Gizem TİRYAKİ	INVESTIGATION OF POLYPHENOL CONTENTS AND ANTIOXIDANT CAPACITIES OF SOME MONOFLOREAL POLLENS FROM THE AEGEAN REGION	634
Cavadxan QASIMOV Əşrəf ƏLƏKBƏROV	THE IMPORTANCE OF AGRICULTURE IN THE SUSTAINABLE ECONOMIC DEVELOPMENT OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC	636
Aslı İŞLER KAYA	SUSTAINABLE AGRICULTURE FOR OILSEED CROPS	638
Loğman BAYRAMOV	RESEARCH AND GENE FUND COLLECTION OF STONE FRUIT PLANTS CULTIVATED IN THE TERRITORY OF NAKHCHIVAN AUTONOMOUS REPUBLIC	640
Dashgin GANBAROV Safura BABAYEVA	USAGE DIRECTIONS OF SPECIES BELONGING TO THE GENUS ALCHEMILLA L. IN THE FLORA OF THE NAKHCHIVAN AUTONOMOUS REPUBUL	644
Nihal Ceren ALICI Mehmet Ali KIRPIK	EFFECTS OF VERMICOMPOST APPLICATIONS ON YIELD PARAMETERS OF KAVILCA WHEAT	645
Zeynep ASUTAY Hakan İNCİ	BIOTECHNOLOGICAL CONTROL METHODS AGAINST VARROA DESTRUCTOR IN HONEY BEES (<i>Apis mellifera</i>)	647
Yagmur TÜRKMEN Filiz ERTUNÇ	INCIDENCE OF GRAPEVINE PINOT GRIS VIRUS IN WESTERN BLACK SEA REGION, TURKEY AND PHYLOGENETIC ANALYSIS BASED ON COAT PROTEIN AND MOVEMENT PROTEIN GENE	660
Firas ALJANABI Mert DEDEOĞLU	PREDICTING LAND SUITABILITY FOR WHEAT CULTIVATION USING THE RANDOM FOREST MODEL	662
Derya GÜLOĞLU	SOIL CONSERVATION MEASURES IN SUSTAINABLE AGRICULTURE	673

Gülcan TAŞ Adnan AYDIN	DETECTION OF DNA MARKERS IN MEDICINAL SAGE (<i>Salvia officinalis</i> L.) FOR USE IN EPIGENETIC STUDIES	675
Zekiye ALTUNTEKİN Celalettin GÖZÜAÇIK	FRUIT SPECIES DAMAGED BY SAN-JOSE SCULBI (<i>QUADRASPIDIOTUS PERNICIOSUS</i> COMST.) (HEMIPTERA: DIASPIDIDAE) SEEN IN ORCHARDS IN İĞDIR PROVINCE AND THEIR DAMAGE STATUS	681
Zekiye ALTUNTEKİN Celalettin GÖZÜAÇIK	DAMAGE SITUATION OF PEACH SHOOT MOTH, ANARSIA LINEATELLA (ZELLER) (LEPIDOPTERA: GELECHIIDAE) IN PEACH GARDENS OF İĞDIR PROVINCE	683
Abdullah İREÇ Celalettin GÖZÜAÇIK	DAMAGE SITUATION OF PEACH TRUNK BLOODBITE, PTEROCHLOROIDES PERSICAE (CHOLODKOVSKY) (HOMOPTERA APHIDIDAE) IN APRICOT AND PEACH ORCHARDS OF İĞDIR PROVINCE	685
Hilal GASİMOV Anvar İBRAHİMOV Mursel SEYİDOV Namig ABBASOV	USEFUL PROPERTIES AND USES OF SPECIES BELONGING TO THE GENERA BARBERRY (<i>BERBERIS</i> L.) AND İRGA (<i>AMELANCHER MEDİK</i>) COMMON IN THE AREA HAVUSH VILLAGE	692
Mursel SEYİDOV Hilal GASİMOV Namig ABBASOV Anvar İBRAHİMOV Fidan HUSEYNOVA	NATURAL RESOURCES OF FRUIT PLANTS BELONGING TO THE GENERA APPLE (<i>MALUS</i> MILL.), PEAR (<i>PYRUS</i> L.) AND CHERRY (<i>PRUNUS</i> L.) FOUND IN THE HAYFIELD AND FOREST EDGE AREAS	697
Namig ABBASOV Hilal GASİMOV Mursel SEYİDOV Bahruz MAMMEDOV Semennaz ABBASLI	SPECIES OF THE GENUS <i>LONICERA</i> L. IN THE FLORA OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC, THEIR PHYTOCHEMICAL PROPERTIES AND PROSPECTS OF USE (AZERBAIJAN)	705
Mirmahmud Seyidli Jabbar Najafov	SPIDER MITE - TETRANYCHIDAE IN THE VINEYARDS OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC OF AZERBAIJAN AND MEASURES TO FIGHT AGAINST IT	710
Hava ULUĞBEY İsmail ALASERHAT	ASIAN CITRUS PSYLLID, <i>Diaphorina citri</i> Kuwayama (HEMIPTERA: PSYLLIDAE), AN IMPORTANT QUARANTINE PEST CAUSING HIGH LOSSES IN CITRUS FRUIT	716
İbrahim YILDIZHAN Baris EREN	EPIGENETIC INHERITANCE OF ENVIRONMENTAL CHANGES IN PLANTS: MECHANISMS AND IMPLICATIONS	724
Hasret ÖZTÜRK	REVIEW OF IGF-IGENE VARIABILITY AND ITS IMPACT ON SHEEP BREEDING IN TURKEY	731
Asuman ÇAPAR	SUSTAINABLE AGRICULTURE AS A REGULATORY AND SUPPORTIVE ACTIVITY OF PUBLIC ADMINISTRATION	740
Mehmet Zeki KOÇAK Bünyamin YILDIRIM	USE OF FLAXSEED (<i>Linum usitatissimum</i> L.) WITH HIGH ADDED VALUE	749
Alev ALDEMİR İsmail KIYAK Faruk AYDIN Ahmet KIRIK	PORTABLE MINI-SCALE VERTICAL FARMING SYSTEM DESIGN: GROWING MICROSPROUT IN SOIL- LESS MEDIA	755
Mehmet Zeki KOÇAK Bünyamin YILDIRIM	THE EFFECT OF ESSENTIAL OILS FROM MEDICINAL AND AROMATIC PLANTS ON PLANT GROWTH	774

Eda ÖZÇALICI İsmail ALASERHAT	COTTON LEAFWORM, <i>Spodoptera littoralis</i> (Boisduval) (LEPIDOPTERA: NOCTUIDAE) AN IMPORTANT QUARANTINE PEST CAUSING HIGH LOSSES IN VEGETABLES	780
Hasine KÜÇÜKYILDIRIM Abdulkadir BÜYÜK	WASTE MANAGEMENT IN AGRICULTURE: RECYCLING OF ORGANIC WASTE AND COMPOSTING TECHNIQUES	787
Osman İNAN Sema KONYALI	DEVELOPMENT OF FOOD FROM PAST TO PRESENT AND FOOD SECURITY	796
Tansu ALKAN Süleyman Savaş DURDURAN	ANALYSING EXPERT OPINIONS ON AGRICULTURAL SUPPORTS IN RURAL AREAS IN KONYA CLOSED BASIN	812
Aysun ALTIKAT	CLIMATE THREATS ON AGRICULTURAL PRODUCTION: EXTREME WEATHER EVENTS AND WATER MANAGEMENT	820
Ayaz Hajiyev	THE TERM "GREEN ECONOMY" IN SUSTAINABLE DEVELOPMENT	833
Rıdvan TİK Tuncay KAYA	THE EFFECTS OF GREEN ROOF SYSTEMS ON ENERGY EFFICIENCY	841
Ezgi Günay SARGUT Reyyan Yergin ÖZKAN	HOMEOPATHY IN PLANT PROTECTION	852
Köksal KARADAŞ Osman Doğan BULUT	SIİRT PISTACHIO PRODUCERS' DEMOGRAPHIC CHARACTERISTICS AND PRODUCTION PROBLEMS: THE CASE OF SIİRT PROVINCE	854
Berna BAŞ	SOILLESS PLANT CULTURE TECHNICAL METHODS	863
Nagihan KILIÇ Celalettin GÖZÜAÇIK	PESTS OF GREENHOUSE VEGETABLE GROWING IN IĞDIR PROVINCE	870
Emine NAKİLCİOĞLU Gizem TİRYAKİ	SUPERFOOD OF THE FUTURE: SEAWEED	890
Murat KARACA Hanife ÇEÇEN	WEED CONTROL IN CANNABIS	892
Deghiche-Diab Nacima Saad Somia Belhamra Youcef Islem Ouamene Tarek Abdelmoneim Karoune Samira Djoudi Madjed Kechebar Mohamed Seif Allah Djazouli Zahr Eddine	VALORISATION OF BIOPESTICIDE FROM DITTRICHIA VISCOSA (L.) GREUTER 1973 FROM ALGERIAN ARID REGION	911

**ANALYSIS OF HYDRO-METEOROLOGICAL TRENDS ACROSS MAJOR
KOPPEN CLIMATE CLASSES USING GLDAS DATA AT A GLOBAL SCALE**

Muhammad Mazhar Iqbal

Water Management Training Institute, Department of Agriculture (On Farm Water Management), Government of Punjab, Lahore 54000, Pakistan

Muhammad Abdullah

University of Agriculture, Faisalabad, Punjab Pakistan.

Tehmena Rashid

Agriculture Mechanization Research Institute (AMRI), Agriculture Department, Government of Punjab, Multan, Pakistan

Hafiz Qaiser Yasin

Directorate General of Agriculture Punjab, Department of Agriculture (On Farm Water Management), Government of the Punjab, Lahore 54000, Pakistan

Shafiq-ur-Rehmn

Water Management Training Institute, Department of Agriculture (On Farm Water Management), Government of Punjab, Lahore 54000, Pakistan

Abstract

This study investigates the spatial and temporal trends of key hydrometeorological variables—evapotranspiration, soil moisture, precipitation, temperature, and runoff—across global land areas, categorized into five major Köppen climate classes. Monthly data from the Global Land Data Assimilation System (GLDAS) were analyzed to evaluate long-term trends using the Mann-Kendall test and Sen's slope estimator. Results reveal significant spatial and temporal variability in trends, driven by regional climatic characteristics.

In tropical climates, evapotranspiration and precipitation exhibit increasing trends, whereas soil moisture and runoff show moderate declines, reflecting amplified water scarcity. Dry climates demonstrate significant warming trends with marked decreases in soil moisture and runoff, highlighting vulnerability to desertification. Continental climates experience notable increases in temperature, coupled with declining runoff and mixed trends in soil moisture and precipitation. Temperate climates exhibit pronounced increases in evapotranspiration and temperature, with precipitation trends varying regionally. Polar climates indicate substantial warming trends, accompanied by significant reductions in snow-driven runoff.

Spatially, warming trends dominate across all climate classes, with dry and polar regions showing the steepest increases. Variability in precipitation trends across regions suggests complex interactions between climate drivers. The findings underscore the need for region-specific water resource management strategies to mitigate climate change impacts on hydro-meteorological processes.

Keywords: GLDAS, Köppen climate classes, Mann-Kendall, Sen's slope, evapotranspiration, soil moisture, precipitation, surface runoff

INVESTIGATING THE ANTIOXIDANT POTENTIAL OF *EBENUS PINNATA AITON*: *IN VITRO* AND *IN VIVO* ASSESSMENTS OF HYDROETHANOLIC EXTRACTS

Kaouthar ELBIRGUI

Laboratory of Chemistry, Biochemistry, Nutrition, and Environment, Faculty of Medicine and Pharmacy, University Hassan II, Casablanca, Morocco

ORCID ID: <https://orcid.org/0000-0002-0891-6924>

Houssam ASSIOUI

Laboratory of Immunology and biodiversity -Faculty of Sciences Ain Chock, University Hassan II, 8 Route d'El Jadida, B.P 5366 Maarif 20100, Casablanca. Morocco.²

ORCID ID: <https://orcid.org/0009-0006-2951-9043>

Oussama RAFIK

Laboratory of Chemistry, Biochemistry, Nutrition, and Environment, Faculty of Medicine and Pharmacy, University Hassan II, Casablanca, Morocco

ORCID ID: <https://orcid.org/0000-0001-5931-1792>

Nadia IOUNES

⁴Environment, Faculty of Sciences Ben M'Sick, University Hassan II. Avenue Cdt Driss El Harti, B.P.7955 Sidi Othmane. Casablanca. Morocco

⁴ORCID ID: <https://orcid.org/0000-0002-4207-8610>

Samir IBENMOUSSA

Laboratory of Chemistry, Biochemistry, Nutrition, and Environment, Faculty of Medicine and Pharmacy, University Hassan II, Casablanca, Morocco

ORCID ID: <https://orcid.org/0009-0008-2346-9289>

Abstract

Introduction and Purpose: *Ebenus pinnata Aiton*, a perennial herb native to North Africa and the Iberian Peninsula, is commonly known as Sofat Ibagra. This plant has a rich history of traditional use in folk medicine for treating various ailments, including inflammation, wounds, and digestive disorders. Despite its traditional applications, scientific research on the pharmacological potential of *Ebenus pinnata Aiton* remains limited.

This study aims to elucidate the phytochemical composition of *Ebenus pinnata Aiton* and investigate its potential biological activities, specifically focusing on antioxidant properties.

Materials and Methods: Antioxidant activity was assessed both *in vitro* and *in vivo*. *In vitro*, antioxidant capacity was evaluated using the ABTS (2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid)) assay, which measures the ability of the sample to scavenge free radicals, the

ferric reducing power (FRP) assay, which measures the electron-donating capacity of the sample, and the thiobarbituric acid reactive substances (TBARS) assay, which measures lipid peroxidation products. *In vivo*, antioxidant activity was assessed by measuring lipid peroxidation (LPO) levels in serum, an indicator of oxidative stress in the body.

Results: Leaf extracts were administered to mice subjected to H₂O₂ oxidative stressor, following the treatment period, liver tissue was collected, and malondialdehyde (MDA) levels, a marker of lipid peroxidation, were quantified using the TBARS assay. Results demonstrated a significant decrease in MDA levels in the mice receiving the *Ebenus pinnata* leaf extract compared to the control group receiving the oxidative stressor alone.

Conclusion : This finding suggests that *Ebenus pinnata* leaf extracts possess *in vivo* antioxidant activity, capable of mitigating lipid peroxidation induced by H₂O₂ in mice.

Key Words: *Ebenus pinnata* Aiton ; *Phytochemical screening* ; *Antioxidant activity*

EVOLUTION, BEHAVIOR AND ECOLOGY OF FEATHER MITE

GUNASEKAR G

PG Scholar Department of Agricultural Entomology,

Tamil Nadu Agricultural University

ORCID: 0009-0007-2566-8348

Abstract

Birds host many lineages of symbiotic mites, but the greatest diversity is shown by the three superfamilies of astigmatan feather mites: Analgoidea, Pterolichoidea, and Freyanoidea. Members of this diphyletic grouping have colonized all parts of the avian integument from their ancestral nidicolous habitat. Whereas some clearly feed on feather pith or skin, acting as parasites, other feather mites are paraphages and consume feather oils without causing structural damage. Sexual dimorphism in feather mites is often extreme, and little is known of the function of many elaborate male structures. Abundance and location of vane-dwelling mites is affected by season, temperature, light, humidity, and host body condition. Because transmission between hosts usually depends on host body contact, it is unsurprising that feather mite phylogeny often parallels host phylogeny; however, recent cladistic analyses have also found evidence of host-jumping and “missing the boat” in several mite lineages.

Keywords: Birds, diversity, habitat, mite, feather

**STUDY OF *CERATITIS CAPITATA* WIEDEMANN (DIPTERA:
TEPHRITIDAE) POPULATION DYNAMICS IN THE EL GHARB REGION OF
MOROCCO FOR FUTURE PEST MANAGEMENT**

EL BAHJA Fatima

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0007-0503-6327>

ZIRI Rabea

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-5223-8235>

BRITAL Rania

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0002-9971-3790>

HAMRANI Meriem

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco

EL AMMARI Mohamed

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0009-9822-1306>

FAHAD Kaoutar

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco

Abstract

The Mediterranean fruit fly, *Ceratitidis capitata* Wiedemann (Diptera: Tephritidae), is a major pest that causes significant damage to fruit crops in the El Gharb region. Understanding its population dynamics is essential for developing effective pest management strategies. This study aims to investigate the life cycle, and environmental factors influencing *C. capitata* both in laboratory conditions and through field monitoring outside the crops. Laboratory rearing was conducted to track the development stages, survival rates, sex ratio. Additionally, an evolutionary monitoring was carried out outside of crop areas to observe population fluctuations in natural environments. The data obtained from both controlled and field settings allowed us to assess the effects of environmental variables such as temperature, humidity, on the population dynamics of *C. capitata*. Our findings offer insights into the biology and ecology of

this pest and provide a basis for developing integrated pest management strategies tailored to the specific conditions of the El Gharb region, ultimately supporting sustainable agricultural practices.

Keywords: *Ceratitis capitata*, population dynamics, Mediterranean fruit fly, pest management, El Gharb region, Morocco

GREEN INNOVATIONS IN MEDICINAL AND AROMATIC PLANT RESEARCH

Dr.Sundhararajan R

Principal and Professor, Mohammed Sathak AJ College of Pharmacy

Mrs.Jothilakshmi.R

Associate Professor, Mohammed Sathak AJ College of Pharmacy

Mr.S.G.Raman

Associate Professor, Mohammed Sathak AJ College of Pharmacy

Abstract

The increasing global demand for natural products, coupled with the need for sustainable practices, has led to significant advancements in the research and development of green technologies within medicinal and aromatic plants. Green innovations focus on environmentally friendly methodologies for cultivating, extracting, and processing plant-based bioactive compounds. These technologies aim to reduce the environmental footprint of plant-derived products while enhancing the efficiency and efficacy of the bioactive compounds present in these plants. Key strategies include the use of organic farming techniques, biotechnological innovations, and solvent-free extraction methods, all of which minimize ecological damage and improve sustainability.

Bioactive profiling of medicinal and aromatic plants plays a crucial role in identifying the specific compounds responsible for therapeutic effects, including antioxidants, anti-inflammatory agents, and antimicrobial substances. Recent advancements in analytical techniques, such as chromatography and mass spectrometry, enable precise identification of these bioactive molecules, allowing for the development of targeted therapeutic applications. Additionally, green technologies such as micropropagation and bioreactor-based plant production systems offer promising alternatives to traditional cultivation methods, promoting resource efficiency and reducing waste.

This evolving area of research not only supports the sustainable use of medicinal plants but also contributes to the global push for greener and more sustainable healthcare solutions. The integration of green technologies with bioactive profiling is expected to revolutionize the production of plant-based medicines, making them more accessible, environmentally friendly, and effective in treating a wide range of health conditions.

Keywords: Green technologies, medicinal plants, aromatic plants, bioactive profiling, sustainable practices, natural products, plant-based medicines, eco-friendly extraction, biotechnology, sustainable agriculture.

GREEN MEDIATED SYNTHESIS, CHARACTERIZATION AND APPLICATION OF MANGANESE OXIDE NANOPARTICLES USING *CITRUS LIMON* PEEL EXTRACT

N Anusuya

Assistant Professor, Department of Chemistry, P.S.G.R. Krishnammal College for Women, Peelamedu, Coimbatore – 641004, Tamil Nadu, India.

PG Student, PSGR Krishnammal College for Women, Coimbatore, TN, India

V S Angulakshmi

Assistant Professor, Department of Chemistry, P.S.G.R. Krishnammal College for Women, Peelamedu, Coimbatore – 641004, Tamil Nadu, India.

PG Student, PSGR Krishnammal College for Women, Coimbatore, TN, India

M Priyadharshini

Assistant Professor, Department of Chemistry, P.S.G.R. Krishnammal College for Women, Peelamedu, Coimbatore – 641004, Tamil Nadu, India.

PG Student, PSGR Krishnammal College for Women, Coimbatore, TN, India

G Easterin Shejina

Assistant Professor, Department of Chemistry, P.S.G.R. Krishnammal College for Women, Peelamedu, Coimbatore – 641004, Tamil Nadu, India.

PG Student, PSGR Krishnammal College for Women, Coimbatore, TN, India

Abstract

The growing demand for sustainable and eco-friendly methods in nanomaterial synthesis has prompted the use of plant-based extracts as potential reducing and stabilizing agents. This research focuses on the green synthesis of Manganese oxide (MnO) nanoparticles using Citrus limon peel extract, a readily accessible and cost-effective source of bioactive compounds. The antioxidants and phytochemicals present in lemon peel facilitate the reduction of metal ions into nanoparticles while preventing agglomeration. The synthesized (MnO)nanoparticles were characterized using various techniques including IR, UV-Vis spectroscopy, XRD and SEM. UV-Vis analysis revealed a distinct absorption peak at 370 nm, confirming the presence of MnO nanoparticles. X-ray diffraction (XRD) confirmed their crystalline structure, with an average particle size ranging from 32 to 35 nm. The antibacterial activity of the MnO nanoparticles was tested against *Escherichia coli* and *Staphylococcus aureus* using the agar well diffusion method. The green synthesis approach offers numerous advantages, including environmental friendliness, simplicity, and the elimination of harmful chemicals.

Key words: Green synthesis, Manganese oxide nanoparticle, biosynthesis

THE IMPACT OF THE U.S. DEVELOPMENT AID ON THE AGRICULTURE OF SOUTH VIETNAM (1954-1975)

Dr. Le Dinh Trong

School of Education, DongThap University

ORCID: 0000-0002-1388-1818

Assoc. Prof. Dr. Pham Duc Thuan

School of Education, CanTho University

Abstract

After the Geneva Agreement on Indochina in 1954, France left Indochina and Vietnam. The United States quickly replaced it and became the largest aid provider to the Republic of Vietnam government. US aid in the form of “development aid” played an important role in the economic development of South Vietnam and greatly influenced the US war policy in Vietnam. On the positive side, that aid had a strong impact on the transformation of rural areas in South Vietnam. Modern capitalist economic methods were increasingly developed in rural areas in South Vietnam. However, in the context of US aid policy in Vietnam, it was completely linked to the war with the terrible destruction caused by bombs and toxic chemicals that the US army caused to agriculture and the lives of farmers in South Vietnam. In international relations in Southeast Asia during this period, the US aid policy caused geopolitical complications, contributed to the increasingly fierce war in Vietnam and had a great impact on security and politics in the region.

This research paper aims to clarify the US aid policy for the Republic of Vietnam government in agricultural policy. Research from national archival sources, combined with historical factual methods, will contribute to clarifying important historical issues of the relationship between the Republic of Vietnam and the United States from 1954 to 1975.

Keywords: development aid, agriculture, South Vietnam, Republic of Vietnam, US, Vietnam War

Introduction

Before 1975, the United States government acted as both the “midwife” and the “lifeline” that nurtured the emergence and existence of the so-called Republic of Vietnam (RVN) in Southern Vietnam. This “lifeline” was none other than “US aid”. Without American intervention and the influence of its aid, Vietnam’s history might not have endured more than two additional decades of suffering and devastation. However, this study does not examine US aid in general, nor does it focus on its military assistance to the RVN. Instead, the primary concern is the most economically relevant form of aid - specifically, the assistance that had an impact on the agricultural economy of the RVN.

Results and discuss

Types of US Economic Aid to the Republic of Vietnam (RVN)

The most economically significant forms of US aid to the RVN can be broadly categorized into two types:

- **Commercial Import Programme (CIP):** Implemented through two mechanisms - *Commercial Aid & Commodity Sales*

Commercial Aid: The United States provided foreign currency to the RVN to finance the import of goods. This was the largest form of economic aid, primarily focused on consumer products, which contributed significantly to inflationary pressures.

Commodity Sales: Governed by Title I of Public Law 480 (PL-480), this was the second-largest economic aid program after Commercial Aid. It involved surplus US agricultural products, which temporarily alleviated food shortages. However, these products were often not suited to Vietnamese tastes, and some of them created competition that negatively affected Southern Vietnam's agricultural production.

- **Development Aid:** Consisting of **Technical Assistance, Project Aid, and Development Credit.**

Technical Assistance: In principle, this form of aid focused on human resources, knowledge, and methodologies. It was the smallest component of US aid, primarily targeting agriculture, healthcare, education, and social sectors, making it beneficial for long-term development.

Project Aid: This was direct US government assistance, implemented through "agreements" or "action plans." It generally fell into three categories: Counterinsurgency Programs, Infrastructure Warfare Programs, and Socioeconomic Development Programs. Despite being limited in quantity and value, its impacts were often mixed - some projects contributed to economic development, while others had unintended consequences.

Development Credit: A form of long-term capital lending with low interest rates, beneficial to both parties, though it was provided in very limited amounts.

From this classification, it is evident that aid specifically aimed at agricultural development in the RVN constituted only a small portion of the second category - "*Development Aid*" - characterized by low quantity and limited value.

Volume of Aid

An overview of the proportion of the most economically significant aid within the total US assistance to the RVN from 1955 to 1975 can be observed in the following summary table:

Fiscal year	AID ^a	PL 480	Piaster subsidy	Total economic aid	Military assistance	Total assistance
1955	320,2	2,2		322,4		322,4
1956	195,7	14,3		210,0	176,5	386,5
1957	259,4	22,8		282,2	119,8	402,0
1958	179,4	9,6		189,0	79,3	268,3
1959	200,9	6,5		207,4	52,4	259,8
1960	170,6	11,3		181,8	72,7	254,5
1961	140,5	11,5		152,0	71,0	223,0
1962	124,1	31,9		156,0	237,2	393,2
1963	143,3	52,6		195,9	275,9	471,8
1964	165,7	59,1	5,8	230,6	190,9	421,5
1965	225,0	49,9	15,4	290,3	318,6	608,9
1966	593,5	143,0	57,4	793,9	686,2	1.480,1
1967	494,4	73,7	98,5	666,6	662,5	1.329,1
1968	398,2	138,5	114,4	651,1	1.243,4	1.894,5
1969	314,2	99,4	146,9	560,5	1.534,0	2.094,5
1970	365,9	110,8	178,7	655,4	1.577,3	2.232,7
1971	387,7	188,0	202,3	778,0	1.945,6	2.723,6
1972	386,8	67,8	133,1	587,7	2.602,6	3.190,3
1973	313,3	188,3	29,5	531,2	3.349,4	3.880,6
1974	384,3	269,9	3,2	657,4	941,9	1.599,3
1975	191,3	49,6		240,9	625,1	866,0
Total	5.954,4	1.600,7	985,2	8.540,3	16.762,3	25.302,6

Table 1: US economic and military assistance to South Vietnam by fiscal year, 1955-1975 (million dollars) [3, p.200]

According to the table, the total volume of US aid to the RVN from 1954 to 1975 was enormous, exceeding \$25 billion. Military aid accounted for more than two-thirds of this total. Although economic aid was also substantial, amounting to over \$8.5 billion, the majority of this amount did not directly contribute to agricultural development in the RVN.

Aid specifically aimed at agricultural development represented only a very small proportion. This can be further illustrated through the following statistical table on project aid categories:

Class	Project Aid	Project-type aid	PL 480-Title II	Total
Relief & rehabilitation	87,7	128,7	330,6	547,0
Political development	284,5			284,5
Public Safety & public adm	159,0			
Pacification	125,5			
Economic development	516,0	58,9		574,9
Agriculture & natural resources	85,5	58,9		
Industrial & mineral development	103,9			
Labor	6,0			
Transportation	320,6			
Social development	267,4			267,4
Health & sanitation	209,9			
Education	57,5			
Technical support	231,5			231,5
Other	192,8	197,1		389,9
Discrepancy ^a	-25,3			-25,3
Total	1.554,6	384,7	330,6	2.269,9

Table 2: Noncommercial aid to South Vietnam by class of project, cumulative to end of period (million dollars) [3, p.208]

According to the statistics, over a span of more than 20 years, the most directly relevant form of US aid to the RVN's agriculture was project aid. However, agricultural-related projects accounted for only about 5% of the total project aid value, amounting to less than \$150 million - equivalent to approximately 1.7% of total economic aid and just 0.6% of total US aid to the RVN.

US assistance related to agriculture was also reflected in loan-based aid. However, a distinctive characteristic of the RVN was that US loans were significantly smaller than grants, as the United States prioritised its global strategic interests in Vietnam over financial returns. "From 1958 to 1969, the United States lent Vietnam \$90 million" [2, p.17]. Over the course of 20 years, total US credit to the RVN was estimated at nearly \$200 million, though only a small portion of these loans was directly allocated to agriculture or used by borrowers for agricultural purposes. For instance, US funding partially contributed to the establishment of the National Agricultural Credit Bureau under Ngô Đình Diệm in 1957 and the Agricultural Development Bank under Nguyễn Văn Thiệu in 1967. In 1970, the US also provided a loan to Nguyễn Văn Thiệu's regime to implement the "Land-to-the-Tiller" law, etc.

Beyond financial assistance, US agricultural aid to the RVN also included technical experts accompanying various projects. Specifically, during the period 1966-1973, the total number of US technical advisors was as follows:

	Number of US technicians	AID-employed	Participating agency	Contractor technicians	Number of participants
1966	1.014	781	97	136	344
1967	2.074	1.591	219	264	1.160
1968	2.780	1.908	371	501	643
1969	2.864	1.771	306	787	691
1970	2.177	1.480	255	442	974
1971	1.649	1.248	108	293	1.039
1972	1.109	912	67	130	1.223
1973	882	579	81	222	1.030

Table 3: Number of technicians supported by AID project funds, 1966-1973 [3, p.210]

However, it is important to note that the total figures in the table do not exclusively represent agricultural experts, as they were involved in various projects, with agricultural initiatives constituting only a small fraction. Additionally, the salaries of these technical advisors were included in the total funding for each project, further reducing the actual financial benefits that Southern Vietnam's agriculture received from US aid.

Beyond the United States, the RVN also received aid from various allied nations and international organisations. A 1974 study conducted in the RVN reported: "According to the Ministry of Foreign Affairs, from 1964 to 1970, Vietnam received \$145,835,000 from 45 countries" [2, p.13]⁵. In the following years (1970-1974), these nations continued to provide aid, amounting to several hundred million dollars. For instance, annual aid figures were as follows:

- 1970: \$21.80 million
- 1971: \$18.93 million
- 1972: \$61.60 million
- 1973: \$151.40 million

The aid and loan distribution for the RVN in 1974 is outlined as follows:

Nation	Aid	Loan	Total
Australia	9		9
Canada	3		3
Taiwan	2	2	4
Denmark	2	3	5
France	8	12	20
West Germany	8	4	12
Japan	24	35	59
South Korea	3		3
Netherlands		3	3
Other countries	2	1	3
United Nations	5		5
Asian Development Bank		4	4
Other Multilateral Aid	2		2
Total	68	64	132

Table 4: Aid and Loan Assistance to the RVN from Various Countries and International Organisations in 1974 (Excluding the US) [5, p.190]

However, not all aid from non-US sources was directed toward agricultural development in the RVN, as these contributions spanned various sectors. Notable agricultural-

related aid came from countries participating in the “**Colombo Plan**” (United Kingdom, Australia, Japan, etc.) and members of the “**Free Bloc**”: **France**: Sent numerous agricultural and irrigation experts, professors, and scholars; provided scholarships for overseas study; established laboratories for the Agricultural and Forestry College; and funded experimental cotton plantation projects, etc. **West Germany (FRG)**: Supplied agriculture and forestry experts, imported German machinery, and granted scholarships, etc. **United Kingdom**: Dispatched experts to research and develop tea factories, provided fishing equipment, supplied agricultural school facilities, and awarded scholarships, etc. **Republic of China (Taiwan)**: Supplied tractors, grain storage facilities, farming techniques, production models, and agriculture-related scholarships, etc. Other contributing nations included **Japan, Israel, Philippines, South Korea**, and others. International organisations also played a role, such as: *Asian Cultural Agency (since 1956)*; *Asian Productivity Organization (since 1965)*; *United Nations*, etc.

The Impact of US “Development Aid” on the Agriculture of the RVN

From the historical reality of the 1954-1975 period, it is evident that US development aid influenced various aspects of the RVN’s agriculture, including land reform, agricultural credit, the introduction of new rice and crop varieties, cultivation techniques,...and the agricultural market. For instance, during Ngô Đình Diệm’s administration, the US played a significant role in supporting land resettlement projects, the establishment of “prosperous hamlets” and “strategic hamlets.” Under Nguyễn Văn Thiệu, the US helped introduce IR rice varieties, provided funds for land purchases to implement the Land-to-the-Tiller Law (NCCR), financed and trained personnel for the establishment of the Agricultural Bank and Rural Bank, and supported programs such as the New Life Hamlets, Rejuvenated Hamlets, and the Rural Pacification & Reconstruction Program, etc.

How should the impact of US aid on RVN’s agriculture be assessed? Was it significant or minimal? Beneficial or detrimental? Sincere or merely strategic?

From the categorisation of US aid, the portion truly beneficial to agricultural development was only a small fraction of what was classified as “development aid,” particularly under “project-based aid.” However, as previously noted, the total value of agricultural and natural resource development projects accounted for less than 5.5% of all project-based aid over two decades. Furthermore, “project-based aid” was often divided into three main programs: “Counterinsurgency,” “Infrastructure Support,” and “Socioeconomic Development.” Historical evidence suggests that not all projects under these categories actually contributed to agricultural progress. For example, the Strategic Hamlet Program not only failed to foster agricultural development but also caused severe social disruptions and ultimately collapsed.

Additionally, an analysis of the “National Budget” distribution, based on foreign aid allocations to RVN ministries, further highlights the lack of investment in agriculture. Even during the peak years of the Land-to-the-Tiller Law’s implementation, the Ministry of Agriculture consistently received one of the lowest budget allocations:

Ministry/Year	1970	1971
Office of the President	3.031.000	160.000
Ministry of Public Reconciliation	1.170.000	915.000
Ministry of the Interior	177.000	210.000
Ministry of Education	171.000	-
Ministry of Health	27.000	-
Ministry of Agriculture	147.000	226.000
Ministry of Public Works	-	1.206.000
Ministry of Finance	-	4.800.000

Bảng 5: Allocation of Foreign Aid Budget for Selected Ministries of the Republic of Vietnam (RVN) in 1970-1971 [4, p.49]

Thus, the proportion and value of aid beneficial to the development of agriculture in the Republic of Vietnam (RVN) within the total economic aid in particular, and total US aid in general, were extremely minimal, not to mention the losses and reductions through various means. Placed in the context of South Vietnam, where agriculture was the dominant economic sector, and considering the damages of war that the US simultaneously brought, the amount of US aid for the agricultural development of the RVN was even more insignificant.

Nevertheless, it must be acknowledged that certain aid under the category of “development aid” from the United States, other countries, and international organizations for the RVN in the agricultural sector yielded some results, initially introducing certain innovations to the agriculture of South Vietnam during that period. Thanks to US aid, as well as support from other countries and organizations, South Vietnam’s agricultural economy underwent some transformations compared to before. For instance, in terms of science and technology, South Vietnam’s agriculture was introduced to new high-yield crop varieties; modern farming techniques with the initial elements of mechanization; the use of chemical fertilizers was significantly intensified compared to agriculture in the North during the same period; and improvements in the preservation and processing of agricultural products were initiated. US aid also contributed to the early effectiveness of irrigation projects on certain areas of land, the implementation of technical demonstration projects, support for agricultural-forestry-animal husbandry schools, farmer training, scholarships for specialists to study abroad, and the deployment of foreign agricultural experts to South Vietnam, among others. These efforts helped a portion of farmers and the agricultural sector in South Vietnam gain greater exposure to scientific and technical advancements, etc. In general, although US development aid, along with aid from other countries and international organizations, was extremely limited, it still had certain effects, causing agriculture and farmers in South Vietnam to experience some changes, including the introduction of new elements. The agriculture of South Vietnam during the 1954-1975 period was, on one hand, heavily influenced by war, fluctuating with the rhythm of the conflict, but on the other hand, it also began to show signs of becoming a market-oriented agricultural economy.

So, what were the limitations of US development aid to South Vietnamese agriculture at the time?

First and foremost, the total value of the aid was far too small, insufficient for a “nation” whose economic foundation and the majority of its population depended on agriculture.

Secondly, not all agricultural and rural development aid projects were successful in “winning the hearts and minds of South Vietnamese farmers,” as the US and the RVN had hoped. On the contrary, many projects had counterproductive effects, with economic factors serving merely as a façade for military and political objectives pursued by the US and the RVN government. These objectives often ran counter to the aspirations and traditions of the people, disrupting rural life in South Vietnam and negatively impacting agriculture.

Thirdly, at the same time, military aid and commercialized aid also had severe consequences, harming agricultural production in South Vietnam. Bombs, defoliants, and surplus agricultural products provided as aid effectively nullified the already minimal positive effects of US development aid for South Vietnamese agriculture. Moreover, rigid mechanisms in procedures, aid reception, imports and exports, etc., also imposed numerous negative effects, further restricting agricultural production in the RVN.

To add an objective perspective on the limitations of US aid to South Vietnamese agriculture specifically, as well as the overall impact of US aid on the RVN, we cite an assessment from a contemporary study conducted under the RVN regime: “After 18 years of receiving US aid, we still see that Vietnam’s industry remains underdeveloped, **agriculture is in decline (emphasis added by the author)**, the budget is in deficit, foreign trade is in imbalance, and inflation is rising” [1, p. 55].

Conclusion

Finally, did the US government at that time truly intend to provide aid to help the Republic of Vietnam (RVN) develop its agriculture? Based on the considerations above - especially when taking into account the political objectives of the US government in its intervention in Vietnam and the intentions behind its aid to the Saigon administration - it can be affirmed that there was no genuine intention of assisting the RVN in developing its agriculture. The meager aid provided to South Vietnamese agriculture amidst the devastation of war was merely an attempt to win over the rural population of South Vietnam at a minimal cost in the battle for hearts and minds against the communists, with the ultimate goal of pacifying the countryside. The so-called development aid, economic aid, and, alongside it, the massive military aid from the US all served a single political purpose: to pacify and control South Vietnam, turning the RVN into the frontline in the war against communism and implementing US security and political strategies. At the same time, this aid transformed the RVN into a market for American capitalist goods. In essence, what was termed “development aid” from the US for South Vietnamese agriculture, at best, can only be considered a form of “stabilization aid” aimed at achieving the aforementioned political objectives.

Examining the various aspects of US (and allied) aid to the RVN in Vietnam’s history remains relevant even in the present day.

REFERENCES

- Dacy, D. C. (1986). *Foreign aid, war, and economic development: South Vietnam, 1955-1975*. Cambridge University Press.
- Gawthorpe, A. J. (2022). Modernization, agricultural economics, and US policy towards land reform in South Vietnam. *The International History Review*, 44(2), 282-299.
- Logan, W. J. (1971). How Deep is the Green Revolution in South Vietnam?: The Story of the Agricultural Turn-Around in South Vietnam. *Asian Survey*, 321-330.
- Lawrence, M. A. (2010). *The Vietnam War: A concise international history*. Oxford University Press.
- Wiest, A. (2003). *The Vietnam War 1956-1975*. Routledge.

**MIGRATION, CLIMATE CHANGE AND SUSTAINABLE AGRICULTURE:
CHALLENGES AND OPPORTUNITIES FOR ECONOMIC DEVELOPMENT**

IOANA POROSNICU

Ion Ionescu de la Brad” Iasi University of Life Sciences / Research and Development Station
for Cattle Breeding Dancu, Romania

ORCID ID: 0009-0006-2687-6694

Madalina DAVIDESCU

Ion Ionescu de la Brad” Iasi University of Life Sciences, Romania

ORCID ID:0000-0002-8611-9787

Bianca-Maria MADESCU

Ion Ionescu de la Brad” Iasi University of Life Sciences, Romania

ORCID ID: 0000-0003-0810-5088

Abstract

Global migration is increasingly influenced by climate change, which has a significant impact on the agricultural and livestock sectors. Rural regions, which depend on these sectors for their livelihoods, are among the most vulnerable to these changes. Extreme weather events, such as droughts, floods, and temperature fluctuations, directly affect agricultural production and animal health. The development of mycotoxins, generated by fungi that infest crops and animal feed, also represents an additional threat to global food security. These conditions caused by climate change contribute to the forced migration of populations from affected areas, who are seeking more stable and secure economic conditions. This work aims to highlight the interdependence between forced migration, climate change, and sustainable economic development, highlighting both the challenges and opportunities generated by these processes. Forced migration can be seen as both a challenge and an opportunity for sustainable economic development. Although migration can lead to forced labor loss in rural areas, migrants who integrate into the economies of host countries can bring with them advanced skills and technologies in agriculture and animal husbandry. They can thus contribute to improving sustainable agricultural techniques and optimizing natural resource management, generating significant economic benefits. For economic development to be sustainable, economic and agricultural policies must address both the effects of migration and the challenges of climate change in an integrated manner. Investment in agricultural research, the promotion of environmentally friendly technologies, and the adoption of sustainable economic strategies are essential for building resilient economies capable of responding to these complex challenges.

Keywords: agriculture, climate change, food security, migration, sustainable development.

PHYTOCHEMICAL CONSTITUENTS OF MORINGA OLEIFERA ROOTS EXTRACT

A. Attahiru

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State
University of Science and Technology, Aliero. Nigeria

A.U Birnin-Yauri

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State
University of Science and Technology, Aliero. Nigeria

M. Mahdi-Adamu

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State
University of Science and Technology, Aliero. Nigeria

A. Yahaya

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State
University of Science and Technology, Aliero. Nigeria

Abstract

Moringa oleifera, also known as the drumstick tree or horseradish tree, is a fast-growing, tropical deciduous tree native to India. It is known for its many uses, including food, medicine, and cosmetics. *Moringa* has many pharmacological properties, including anti-inflammatory, anti-diabetic, anti-cancer, and anti-bacterial. The leaves are used to treat malnutrition and increase breast milk production. *Moringa* oil can restore liver enzymes and reduce oxidative stress. All parts of the *moringa* tree are edible, including the leaves, flowers, seeds, roots, and pods. The leaves are rich in protein, minerals, and antioxidants. The seeds are a natural coagulant and contain up to 40% oil, which is used in cosmetics and as a food supplement. This research work aimed to investigate the qualitative and quantitative analysis of *Moringa oleifera* roots. The plant sample was collected in Unguwar Galadima, "Aliero." Cold maceration was the method used for extraction of plant roots, with methanol as solvent for the extraction. The standard methods were used in carrying out the qualitative and quantitative phytochemical analysis. The result of qualitative phytochemical screening showed the presence of glycosides, phenols, flavonoids, steroids, anthraquinones, alkaloids, tannins, saponins, and terpenoids. The result of quantitative investigation showed the content of glycosides (6.33 ± 0.58 %), phenols (7.50 ± 0.50 %), flavonoids (7.33 ± 0.58 %), steroids (3.67 ± 0.58 %), anthraquinones (8.0 ± 0.00 %), alkaloids (13.50 ± 0.50 %), tannins (11.30 ± 0.29 %), saponins (9.39 ± 1.17 %), and terpenoids (6.50 ± 0.05 %). The presence of biochemical compounds in the roots extracts is an indication that this plant might contain several compounds that are used in the treatment of different diseases caused by microorganisms. The presence of this phytochemicals in the roots extract can be as a guide for further studies regarding this plant.

Keywords: *Moringa oleifera*, Roots, Qualitative phytochemical, Quantitative phytochemical, Methanol.

PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL ACTIVITY ON THE ETHANOLIC EXTRACT OF *GUIERA SENEGALENSIS* LEAVES

Muslim Yusuf

Department of Pure and Industrial Chemistry, Faculty of Natural and Applied Science, Umaru Musa Yar'adua University, P. M. B 2218, Katsina, Nigeria.

ORCID: 0009-0009-2777-3429

Sulaiman Sani Yusuf

Department of Pure and Industrial Chemistry, Faculty of Natural and Applied Science, Umaru Musa Yar'adua University, P. M. B 2218, Katsina, Nigeria.

Abdulaziz Hassan Maidabino

Department of Pure and Industrial Chemistry, Faculty of Natural and Applied Science, Umaru Musa Yar'adua University, P. M. B 2218, Katsina, Nigeria.

Abstract

The leaves of *Guiera senegalensis* are account to possess medicinal properties according to Nigerian mythology. Medicinal properties of the plant may be attributed to the quality of their bioactive constituents. This study investigated the phytochemical constituents and antibacterial activity of the leaves extract of *G. senegalensis*. Powdered mixture of the leaves of *Guiera senegalensis* was subjected to maceration (extraction) using ethanol (90%, v/v). After extraction, the crude extract was obtained, which is oily and sticky in texture and brownish green in color. The phytochemical screening of the extract was carried out using standard methods. The result for phytochemical screening test showed the presence of phenolic, tannins, saponins, terpenoids, steroids, alkaloids and absence of anthraquinone, and also partially presence of flavonoids which all accounted to its medicinal properties. The antibacterial activity test was evaluated qualitatively through agar disc diffusion method against *Staphylococcus aureus* and *Escherichia coli* bacterial isolates. The results showed that *E. coli* is highly susceptible to *G. senegalensis* leaves extract at high concentration of 250mg/ml with highest zone of inhibition 16mm. The extract showed moderate activity against *Staphylococcus aureus* at high concentration of 250mg/ml with a zone of inhibition 14mm. However, the result showed that the plant *G. senegalensis* can be a promising source of naturally occurring antibiotics and hence justify the claim made by traditional medicine and practitioners.

Keywords: Phytochemical, antibacterial, ethanolic extract and leaves

TRANSFORMING FOOD SAFETY AND SUSTAINABILITY: THE ROLE OF MODERN FOOD PROCESSING TECHNOLOGIES

Marwa Waheed

Department of Food Science and Technology, Riphah International University, Faisalabad, Pakistan

Muhammad Bilal Hussain

Department of Food Sciences, Government College University Faisalabad, Pakistan

Fatima Waheed

National Institute of Food Science and Technology, University of Agriculture Faisalabad, Pakistan

Nida Kanwal

Department of Food Science and Technology, Riphah International University, Faisalabad, Pakistan

Abstract

Food processing technologies play a significant role in guaranteeing the quality, sustainability and safety of global food supply chain. This research explores the recent innovations in these food processing technologies and their effect on product quality and safety. Several preservation techniques including pulsed electric field (PEF), high-pressure processing (HPP) and hurdle technology had been shown to boost microbial safety which further preserve the sensory parameters and nutrition. Advanced thermal techniques including ohmic heating, radio frequency heating and microwave heating decrease energy consumption, processing time and maintain the quality of product. Non-thermal techniques for example cold plasma, ultraviolet radiation and ultrasound prolong the shelf life deprived of compromising the sensory profile and nutritional value. Innovative packaging methods contribute freshness, reduce food waste and prolong the shelf life. This current research highlights the sustainable practices including optimized water usage, energy-efficient equipment, eco-friendly packaging materials and waste reduction. Implementation of the above mentioned advanced techniques can reduce waste, improve consumer confidence and create a worthwhile food supply chain. This research serves as an appreciated resource for researchers and industry professionals looking for implementing advanced food processing techniques.

Keywords: Innovative technologies, High pressure processing, Hurdle technology, Ohmic heating, Cold plasma technology, Ultrasound

NUTRITIONAL AND PHYTOCHEMICAL ANALYSIS OF OREGANO (*ORIGANUM VULGARE*) LEAVES**M. Mahdi-Adamu**

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State University of Science and Technology, Aliero. Nigeria

F. Halidu

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State University of Science and Technology, Aliero. Nigeria

A. Attahiru

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State University of Science and Technology, Aliero. Nigeria

A. Yahaya

Department of Pure and Industrial Chemistry, Faculty of Physical Sciences, Kebbi State University of Science and Technology, Aliero. Nigeria

R. Adamu

Department of Pure and Environmental Chemistry, Faculty of Physical Sciences, Usmanu Danfodiyo University Sokoto, Nigeria.

Abstract

Oregano (*Origanum vulgare*) is a perennial herb that belongs to the mint family, *Lamiaceae* and native to the Mediterranean region and some parts of Western Asia, it is widely used for its strong aroma and distinct, slightly bitter flavor. Oregano is an essential culinary herb in many global cuisines, particularly in Mediterranean, Mexican, and Italian dishes. It is also known for its medicinal properties, which have been recognized since ancient times. Oregano has established itself in folkloric medicine as relaxant, antibacterial and can boost the immune system. Studies suggests that it is an antioxidant that can help prevent cancer. The aim of this study was to determine the nutritional composition and phytochemical constituents of *Origanum vulgare* leaves. The Standard methods were used to in carrying out this research. The result of proximate analysis showed the content of carbohydrate, Crude protein, Crude lipids (fats and oil), Crude fibre, moisture and ash respectively (18.23 ± 0.78 %, 13.44 ± 0.28 %, 17.00 ± 0.70 %, 17.50 ± 2.00 %, 32.83 ± 1.76 % and 1.00 ± 0.50 %). The result of phytochemical constituent indicates the presence of tannins, alkaloids, flavonoids, steriod, terpinoids and saponins. The phytochemical screening identified the key bioactive chemical compounds, flavonoids, Alkaloids, terpenoids, tannins, saponins and steroids which contribute to oregano's antioxidant and antimicrobial properties and other pharmacological activities. The proximate analysis revealed a balanced composition, highlighting essential nutrients such as protein, lipids, and carbohydrates, alongside a low moisture content, indicating their potential for preservation and use in various applications.

Keywords: Proximate, Phytochemical, Oregano Leaves, Terpenes.

**ACCEPTABILITY OF ORANGE FLESHED SWEET POTATO
CONFECTIONARIES AMONG FACULTY OF AGRICULTURE STUDENTS,
UNIVERSITY OF IBADAN. NIGERIA**

Oyewole Mojisola Fauziyah

Department of Agricultural Extension and Rural Development, University of Ibadan,
Nigeria

ORCID : <https://orcid.org/0000-0003-07333-4348>

Abstract

The study examined the acceptability of orange fleshed sweet potato confectionaries among agricultural students at university of Ibadan. A simple random sampling technique was used to select 233 respondents for the study and data collected through questionnaire were analyzed using both descriptive and inferential statistics. Result shows that (74.1%) of the respondents had mean age of 23 ± 4 years, female (52.3%), Christians (80.8%), Yorubas (83.9%), they also receive mean monthly allowance of $25,217 \pm 15,000$, Sweet potato chips (91.2%) were more available to the respondents in the study area. The majority (51.3%) of the respondents had high nutritional knowledge about orange-fleshed sweet potatoes. The shelf life of preservation of confectionaries with highest means of 1.24, ranked 1st as the most severe constraint faced by the respondents in the study area. More than half of the respondents (55.0%) had high level of acceptance of the varieties of orange-fleshed sweet potato confectionaries in the study area. A significant relationship exists between respondent's level of nutritional knowledge about orange fleshed sweet potato ($r=0.259$, $p=0.015$) and the respondents level of acceptability of orange-fleshed sweet potatoes confectionaries in the study area. A significant difference ($F=1.166$, $p=0.346$) does not exist in the acceptability of different formulations of orange-fleshed sweet potatoes confectionaries in the study area. The study concluded that majority of the respondents had high acceptance of the varieties of orange-fleshed sweet potato confectionaries. It is therefore recommended that nutritional educationists should sensitize people with nutritional values of OFSP, to enhance the acceptability of OFSP confectionaries among students and promote the consumption of OFSP confectionaries as nutritious and delicious snacks.

Keywords: Acceptability, varieties, orange-fleshed, sweet potato, confectionaries.

**OCCURRENCE AND CHARACTERIZATION OF ANTIBIOTIC-RESISTANT
SALMONELLA ISOLATES FROM FRESH PRODUCE IN AGRICULTURAL
ENVIRONMENTS**

BELLIL Zahra

Laboratoire d'Ecologie Microbienne, FSNV, Université de Bejaia, Algérie

MEYER Sylvain

Université de Limoges, INSERM, CHU Limoges, RESINFIT, U1092, Limoges, France

TILLOY Valentin

CNR Herpesvirus, UF 9481 Bioinformatique, CHU Limoges, France

BELLIL Dassine

Laboratoire d'Ecologie Microbienne, FSNV, Université de Bejaia, Algérie

MAIRI Assia

Laboratoire d'Ecologie Microbienne, FSNV, Université de Bejaia, Algérie

De CHAMPS Christophe

INSERM UMR-S 1250 P3Cell, SFR CAP-Santé, Université de Reims-Champagne-Ardenne, Reims, France. Laboratoire de Bactériologie-Virologie-Hygiène Hospitalière-Parasitologie-Mycologie, CHU Reims, Hôpital Robert Debré, Reims, France.

BARRAUD Olivier

Université de Limoges, INSERM, CHU Limoges, RESINFIT, U1092, Limoges, France

TOUATI Abdelaziz^{1*}

Laboratoire d'Ecologie Microbienne, FSNV, Université de Bejaia, Algérie

Abstract

Introduction and Purpose: *Salmonella* contamination of fresh produce, including watermelon, poses a significant public health concern. This study aimed to investigate the presence and characteristics of *Salmonella* isolates in watermelon, soil and water samples obtained from agricultural sources.

Materials and Methods: A total of 1450 watermelon samples were collected from different farms in Bejaia and subjected to microbiological analysis. *Salmonella* spp. were isolated using standard culture methods and confirmed by molecular techniques. The isolates were further characterized for antimicrobial susceptibility, serotype determination, and genetic relatedness using whole genome sequencing.

Results: The results revealed the presence of *Salmonella* in 19% of the total samples, indicating a potential risk of foodborne illness associated with watermelon consumption. Antimicrobial susceptibility testing demonstrated varying resistance patterns among the isolates. Serotyping revealed diverse *Salmonella* serotypes, suggesting multiple contamination sources. Molecular

analysis indicated genetic diversity among the isolates, with some strains showing close relatedness to clinical isolates.

Discussion and Conclusion: These findings underscore the importance of implementing stringent food safety measures throughout the vegetables production and distribution chain to mitigate the risk of *Salmonella* contamination and subsequent foodborne outbreaks.

Keywords: *Salmonella*, antimicrobial resistance, watermelon, agricultural environments, genotypic characterization, food safety

YIELD AND STORABILITY OF ONION ECOTYPES AS INFLUENCED BY IRRIGATION INTERVAL IN SUDAN SAVANNA ZONE OF NIGERIA

M.M Akibu

Department of Crop Science, Kebbi State University of Science and Technology, Aliero
Nigeria

M.S. Na-Allah

Department of Crop Science, Kebbi State University of Science and Technology, Aliero
Nigeria

A.Muhammad

Department of Crop Science, Kebbi State University of Science and Technology, Aliero
Nigeria

I. J. Yusuf

Department of Crop Science, Kebbi State University of Science and Technology, Aliero
Nigeria

Abstract

Field trials were conducted simultaneously during the 2023/2024 dry seasons at the Kebbi State University of Science and Technology, Aliero University orchard located at Aliero and The second experiment were conducted at Usmanu Danfodio University Sokoto Teaching and research Fadama farm. The area is located at Sokoto State, both locations are in Sudan Savannah agro-ecological zone of Nigeria. The aim was to study the effects of irrigation interval on yield and the storability of twelve onion ecotypes (*Allium cepa* L.). Treatments consisted of three irrigation intervals (3, 6 and 9 days), and twelve onion ecotypes (Ex-Dikwa, Ex-Kaffe, Ex-Bama, Ex-Alieru, Ex-Gothege, Ex-Gada, Ex-Wurno, Ex-GHM, Ex-Gaya, Ex-Borno, Ex-Lahodu, and Ex-Goronyo). Factorial combinations of irrigation intervals were allocated to the main-plots, while ecotypes were assigned to the sub-plots in a randomized completely block design, replicated three times. The size of each sub-plot was 1x2m (4m²). Result revealed that plant height, number of leaves, above ground biomass (AGBM), total biomass (TBM), fresh bulb weight (FBW), cured bulb diameter (CBD) and bulb diameter (BD) were higher with 3 to 6 days irrigation interval, while weight loose and weight loose in (%) were higher in 3 days irrigation interval. Higher percentage of rotted bulbs and sprouting were recorded by frequent irrigation of 3 days interval. Plant height, number of leaves especially at early growth stages, and number of leaves throughout the crop cycle were higher at 3 days irrigation, ecotypes Ex-Kaffe and Ex-Dikwa has the highest plant height and leave number at 4, 8 and 12 WAT in the both locations, also total biomass (TBM), above ground biomass (AGBM), fresh bulb weight (FBW) recorded the highest weight value followed by Ex-Dikwa. Similarly, 3 days irrigated ecotypes has the highest weight value followed by 6 days irrigation interval while 9 days irrigation ecotypes recorded the least weight value. Ex-Borno, Ex-Dikwa and Ex-Kaffe

recorded the highest weight value after undergoing curing period in the two locations, while ExGHM, Ex-Lohodu has the least weight value. Likewise in sokoto Ex-Dikwa and Ex-Kaffe has the highest weight value. In terms of bulb diameter (cm) Ex-Dikwa recorded the highest value in Aliero while the least was recorded in Ex-Goronyo, similarly in Sokoto Ex-Dikwa, Ex-Kaffe and Ex-Borno has the highest weight values after curing while Ex-Aliero, Ex-Gothege, Ex-Gada, Ex-GHM, Ex-Lohodu and Ex-Goronyo followed. In terms of weight loose, Ex-Borno loses weight drastically for the whole of the storage period followed by Ex-Gaya and Ex-Goronyo. Equally, Ex-Wurno and Ex-Kaffe recorded less weight loose during the storage period. As well as high number of rotten, sprouting were recorded in 3 days irrigation interval. Based on the result of this study, 3 days irrigation scheduling proved the best for more number of leaves plant height and fresh bulb yield in the study area, ecotypes Ex-Dikwa and Ex-Kaffe could be adopted due to its potential for higher yielding, while 9 days irrigation interval is the best for post-harvest storage and Ex-Goronyo were found superior.

Key words: Onion, Ecotypes, Irrigation, Yield, Storability

**THE GENETIC CHARACTERIZATION OF THE MOROCCAN CAROB TREE
(*CERATONIA SILIQUA* L.) USING ISSR MARKERS**

ZAGGOUMI Hasna

Laboratory of Agro-industrial and Medical Biotechnologies, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, Béni-Mellal, Morocco
Tunisian-Moroccan Joint Laboratory: Plant/Pathogen/Environment Interactions of Phylogenetic Resources of Agronomic and Medicinal Interest: Responses to Biotic and Abiotic Stresses

BOUDA Said

Laboratory of Agro-industrial and Medical Biotechnologies, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, Béni-Mellal, Morocco
Tunisian-Moroccan Joint Laboratory: Plant/Pathogen/Environment Interactions of Phylogenetic Resources of Agronomic and Medicinal Interest: Responses to Biotic and Abiotic Stresses

NEG Imane

Laboratory of Agro-industrial and Medical Biotechnologies, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, Béni-Mellal, Morocco
Tunisian-Moroccan Joint Laboratory: Plant/Pathogen/Environment Interactions of Phylogenetic Resources of Agronomic and Medicinal Interest: Responses to Biotic and Abiotic Stresses

MARGHALI Sonia

Laboratory of microorganisms and actives Biomolecules, Faculty of Sciences of Tunis, University Tunis El Manar, Tunisia
Tunisian-Moroccan Joint Laboratory: Plant/Pathogen/Environment Interactions of Phylogenetic Resources of Agronomic and Medicinal Interest: Responses to Biotic and Abiotic Stresses

TRIFI Neila

Laboratory of microorganisms and actives Biomolecules, Faculty of Sciences of Tunis, University Tunis El Manar, Tunisia
Tunisian-Moroccan Joint Laboratory: Plant/Pathogen/Environment Interactions of Phylogenetic Resources of Agronomic and Medicinal Interest: Responses to Biotic and Abiotic Stresses

HADDIOUI Abdelmajid

Laboratory of Agro-industrial and Medical Biotechnologies, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, Béni-Mellal, Morocco
Tunisian-Moroccan Joint Laboratory: Plant/Pathogen/Environment Interactions of Phylogenetic Resources of Agronomic and Medicinal Interest: Responses to Biotic and Abiotic Stresses

Abstract

The carob tree (*Ceratonia siliqua* L.) is a species with significant socio-economic and ecological importance. It possesses cytotoxic, antimicrobial, antioxidant, and anti-

inflammatory properties, and is used in various sectors, including pharmaceuticals, cosmetics, biotechnology, textiles, food sciences, and nutrition. It is frequently found in Morocco in different bioclimatic zones, ranging from humid to desert, with a warm and temperate climate. To establish a conservation strategy for the carob tree in Morocco, the first step towards its rational exploitation is to analyze the genetic diversity of the local genetic material. In this context, 12 populations from different regions of Morocco were analyzed using 18 ISSR molecular markers. The use of appropriate statistical analysis techniques demonstrated significant genetic diversity both within (74.20%) and between (25.80%) carob populations. The mean values of PIC and R_p is 0.47 and 3.33, respectively. In addition, hierarchical AMOVA revealed very little genetic differentiation between groups of mountain range, bioclimatic and altitudinal conditions (FCT=1.4%, FCT=5% and FCT=0.1%, respectively). The results showed that the ISSR markers used are informative and effective in detecting polymorphism and characterizing the populations of this species. Furthermore, there was very little genetic differentiation between the groups of geographic, bioclimatic, and altitudinal conditions.

The data obtained will be useful for defining conservation strategies and breeding programs for this species.

Keywords: *Ceratonia siliqua*, population, polymorphism, ISSR markers, Morocco.

EFFICACY OF ESSENTIAL ELEMENTS AGAINST SUCKING INSECT PESTS OF COTTON UNDER DIFFERENT CONDITION

Hafiza Zara Saeed

Department of Botany, Government College University Faisalabad, 38000 Pakistan

Muhammad Wajid

Department of Botany, Government College University Faisalabad, 38000 Pakistan

Musrat Shaheen

Department of Botany, Government College University Faisalabad, 38000 Pakistan

Abstract

Cotton production faced a major challenge from sucking insect pests, which inflicted substantial economic damage by diminishing both yield and fiber quality. The effectiveness of various essential elements in controlling primary sucking pests under natural and semi-natural condition. These elements, were found to boost plant resistance by reinforcing cell walls, enhancing metabolic functions, and triggering defense mechanisms, thereby mitigating pest infestations and their associated damage. The foliar applications revealed that silicon and potassium notably repelled sucking pests by increasing leaf tissue toughness, while zinc and magnesium enhanced overall plant health, indirectly reducing pest populations. The potential of essential elements in enhancing cotton crop resilience against sucking pests and promoting sustainable agriculture.

**THE USE OF WHEY TO ENHANCE TOMATO AND WHEAT SEED
GERMINATION: A PATHWAY TO VALORIZING DAIRY BY-PRODUCTS FOR
SUSTAINABLE AGRICULTURE**

Karima JMAILI

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0009-0009-3153-0825>

Khawla WADDI

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0000-0001-9820-2265>

Bouchaib BAHLAOUAN

Higher Institutes of the Nursing Professions and Techniques of Health ISPITS Casablanca
22500, Morocco

ORCID ID: <https://orcid.org/0000-0002-0976-1274>

Alla SILKINA

Algal Research Group, Bioscience Department, College of Science, Swansea University,
Singleton Park, Swansea, SA2 8PP, UK

ORCID ID: <https://orcid.org/0000-0002-1804-8083>

Nadia BOUTALEB

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0000-0003-1628-4311>

Abstract

Introduction and Purpose: The management of whey, a major by-product of the dairy industry, presents significant environmental and economic challenges due to its high organic charge. Reusing whey meets the growing demand for sustainable agricultural practices and offers a valuable solution by making use of its nutrient-rich composition. This study explores the potential of whey as a biostimulant to improve seed germination and early growth in

economically significant crops, specifically tomato and wheat, to assess whey's impact on germination rates, root length, and biomass production, and to determine the optimal conditions for its application.

Materials and Methods: Synthetic whey was prepared by heating to 80°C and adding 2% citric acid for coagulation and filtration. Anaerobic digestion was conducted in a mesophilic digester at 35–40°C, producing a nutrient-rich digestate (nitrogen, phosphorus, potassium, calcium). The digestate was diluted into four treatment levels (0%, 25%, 50%, 75%) and applied to plants grown in controlled conditions. Parameters such as germination rate, plant height, and root and shoot biomass were measured. The data were statistically analysed to assess the observed effects and identify potential inhibitory impacts related to digestate concentrations.

Results: Tomato plants treated with whey exhibited a notable increase in vegetative biomass, particularly in shoot growth, indicating enhanced vigour. Wheat seeds demonstrated improved germination rates under whey treatment. However, higher whey concentrations showed inhibitory effects, likely due to its high organic and mineral content, emphasising the need for optimal dilution.

Discussion and Conclusion: This study confirms whey's potential as a natural biostimulant for improving crop performance while addressing dairy waste management challenges. Optimised whey formulations and concentrations could improve crop productivity and valorise dairy by-products, reducing environmental impact. Future research should refine application protocols and expanding its use across diverse cropping systems.

Key Words: Whey, Sustainable agriculture, Biostimulant, Germination, Tomato, Wheat.

COMPARATIVE ANALYSIS OF FERMENTED ALMOND MILK USING MOROCCAN ALMOND VARIETIES: A STUDY ON MICROBIOLOGICAL AND NUTRITIONAL PROPERTIES

Mohamed LAHRAIRI

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0009-0006-9408-9190>

Karima JMAILI

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0009-0009-3153-0825>

Bouchaib BAHLAOUAN

Higher Institutes of the Nursing Professions and Techniques of Health ISPITS Casablanca
22500, Morocco

ORCID ID: <https://orcid.org/0000-0002-0976-1274>

Hlima BESSI

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0000-0001-5068-7814>

Nadia BOUTALEB

Hassan II University of Casablanca, Faculty of sciences and techniques Mohammedia,
Laboratory of Biochemistry, Environment and Agri-Food, URAC36, 20650 Morocco

ORCID ID: <https://orcid.org/0000-0003-1628-4311>

Abstract

Introduction and Purpose: The growing demand for plant-based dairy alternatives has led to increased interest in fermented nut-based beverages. Almond milk, particularly when fermented, offers a promising alternative to traditional dairy products due to its potential probiotic benefits and rich nutritional profile. This study aims to evaluate the microbiological and nutritional properties of fermented almond milk produced from three Moroccan almond varieties and compare them with kefir. The research focuses on assessing fermentation kinetics, probiotic viability, antioxidant activity, and key physicochemical characteristics to determine the influence of almond variety on the final product's quality.

Materials and Methods: Fermented almond milk was prepared using three Moroccan almond varieties and probiotic cultures. Fermentation progress was monitored through pH, titratable acidity, and microbial viability. Nutritional analysis included protein content, lipid composition, total phenolic content, and antioxidant activity. A comparative evaluation with kefir was conducted to assess differences in fermentation efficiency and nutritional properties.

Results: The results indicate that almond variety significantly influences fermentation kinetics, probiotic retention, and nutritional composition. Certain varieties promoted higher microbial viability and exhibited enhanced antioxidant properties. The comparison with kefir revealed distinct physicochemical characteristics, with fermented almond milk showing variations in acidity, microbial activity, and nutrient retention. Differences in lipid and protein profiles suggest that specific almond varieties may be more suitable for fermentation-based functional beverages.

Discussion and Conclusion: Fermented almond milk presents a viable probiotic-rich dairy alternative, with its properties influenced by almond variety. Optimizing fermentation conditions can enhance its functional benefits. Future studies should explore sensory attributes and large-scale production feasibility to support the development of sustainable, plant-based probiotic beverages.

Key Words: Moroccan almond, Probiotics, Kefir, Fermentation, Functional beverage.

BIO OIL FROM CORK : PYROLYSIS-GC-MS STUDY

Imene BENBELKACEM

University of Science and Technology Mohamed Boudiaf (USTO-MB), Laboratory of Plant and Microbial Production and Valorization, Oran, Algeria

ORCID ID: <https://orcid.org/0000-0002-2198-7671>

Aicha Bouhafsoun

University of Science and Technology Mohamed Boudiaf (USTO-MB), Laboratory of Plant and Microbial Production and Valorization, Oran, Algeria

ORCID ID: <https://orcid.org/0000-0001-8522-8925>

Rawaa JAMALADDEEN

University of Poitiers, P' Institute, Fluids-Thermal-Combustion CNRS-UPR3346, ENSMA, poitiers, France

ORCID ID: <https://orcid.org/0000-0001-8029-9142>

Bruno COUDOUR

University of Poitiers, P' Institute, Fluids-Thermal-Combustion CNRS-UPR3346, ENSMA, poitiers, France

Laurent LEMEE

University of Poitiers, Institute of Chemistry of Environments and Materials (IC2MP), CNRS UMR 7285, poitiers, France

ORCID ID: <https://orcid.org/0000-0001-9520-3289>

Abstract

The use of fossil resources is one of the main causes of global warming, making it urgent to find alternative solutions. Biomass has attracted the attention of many researchers due to the economic and environmental benefits of its use.

Cork oak belongs to the Fagaceae family and is known for its production of cork, a highly valued material in various industries. However, the use of this material generates a significant amount of residues. The aim of this study was to characterize the products derived from cork to assess its potential for bio-oil production.

Cork was pyrolyzed using the double-shot method, meaning it was subjected to two successive temperatures (350°C and 550°C) on the same sample. The pyrolysis products were separated by gas chromatography and identified by mass spectrometry.

The first shot at 350°C mainly produced phenolic compounds and triterpenes, with a high content of friedelan-3-one. The second shot at 550°C generated a greater amount of pyrolysate, primarily composed of aliphatic hydrocarbons resulting from suberin degradation.

Cork is an excellent feedstock for bio-oil production due to its richness in aliphatic hydrocarbons and triterpenes. Moreover, these compounds are widely used in various industries, including the pharmaceutical and chemical sectors.

Keywords: cork; bio-oil; Py-GC/MS; suberin

BIOMETRIC ANALYSIS OF THE FIBERS OF A LOCAL ALGERIAN PLANT SPECIES AND THEIR VALORIZATION

BRIKCI Fatiha

Laboratory of Production, Plant and Microbial Valorization (LP2VM)
<https://orcid.org/my-orcid?orcid=0009-0005-5309-7314>

BOUHAFSOUN Aicha

Department of Biotechnology, Faculty of Natural and Life Sciences,
<https://orcid.org/0000-0001-8522-8925>

BOKHARI Hassiba

University of Science and Technology of Oran Mohamed Boudiaf
<https://orcid.org/0009-0004-2833-1247>

DJABEUR Abderrezak

BP, 1505, El-M'naouer, Oran 31000, Algeria
<https://orcid.org/0000-0003-2599-9757>

ABSTRACT

Introduction and Purpose: Studying the biometric characteristics of the stems of plant species has been of great interest to researchers in the wood and paper industry. The use of plant fibers has been widespread in the fields of composites, buildings, insulation, plastics and automobiles. The present study aimed to investigate the biometric characteristics of the stem fibers of local Algerian plant species. The dimensions of these “fibers” vary from one plant to another. Natural fibers, especially plant fibers, are attracting the attention of researchers more than ever because plant fibers are economical, available and environmentally friendly. The applications of plant fiber composites are increasing day by day in various fields.

Materials and Methods: The samples of this study are the leaves and stems from a medicinal plant, The fibers were extracted with glacial acetic acid and hydrogen peroxide and placed in the oven for 48 hours, then measured using a micrometer which allowed us to observe different types of morphologies and variable sizes, 30 fibers were studied and photographed after this step a statistical study is established to calculate the average.

Results:The results show that this species has fibers of different sizes ranging from (800 mm to 1920 mm) for the leaves and (from 280 mm to 910 mm) and which allowed us to classify these fibers in relation to other fibers studied in monocot species such as (fibers having a different size and shape).

Discussion and Conclusion:Many natural and local species have various beneficial interests on health and the environment, our research is established with the aim of protecting our environment and our sea planet.

Key Words: Medicinal plant, fibers, micrometer.

BIOSORPTION OF METHYL RED, THYMOL BLUE AND THEIR DYE MIXTURES USING MODIFIED ELEPHANT GRASS, *Pennisetum purpureum*

Oluwaseun Adekoya Adelaja

Department of Chemistry, School of physical sciences, Federal University of Technology Akure, Nigeria.

Babafemi Raphael Babaniyi

Bioresources Development Centre, National Biotechnology Research and Development Agency, Nigeria

Olayinka Abigail Ajongbolo

Bioresources Development Centre, National Biotechnology Research and Development Agency, Nigeria

ABSTRACT

The existence of dye effluents in surface water poses significant challenges to environmental sustainability and public health. The technology of biosorption, which employs natural plant materials and agricultural by-products, provides a sustainable methodology for the elimination of colorants from aqueous environments. This investigation assesses the efficacy of *Pennisetum purpureum* (commonly known as elephant grass) in the adsorption of methyl red and thymol blue dyes. The characterization of the adsorbent was conducted through Fourier Transform Infrared Spectroscopy (FT-IR), UV-Visible spectrophotometry, and adsorption isotherm analyses utilizing the Langmuir and Freundlich models. Batch adsorption experiments were performed to evaluate the effects of pH (ranging from 4 to 9), initial dye concentration (10-50 mg/L), contact duration (30-180 minutes), and adsorbent mass (0.05-0.3 g) on the removal efficiency of dyes. FT-IR analysis indicated that both untreated and potassium persulfate-treated *P. purpureum* exhibited functional groups such as carbonyl, carboxyl, hydroxyl, and amine groups that have the capacity to bind with positively charged ions. The optimal parameters for the removal of dyes were determined to be a pH of 4, an adsorbent dosage of 0.1 g, a contact time of 30 minutes, an initial concentration of 10 mg/L, and a temperature maintained at 30°C. The findings underscore the considerable potential of this biosorbent in the removal of methyl red and thymol blue dyes from aqueous solutions.

Keywords: Adsorbent; Biosorption; Biosorbent; FT-IR; cadmium ions

**THE NEGATIVE EFFECT OF SOIL AMENDMENTS WITH SOLID RESIDUES OF
ROSMARINUS OFFICINALIS L. ON THE GROWTH OF *SOLANUM
LYCOPERSICUM* MILL**

Meriem RAFYA

Sustainable Development and Health Research Laboratory, Cadi Ayyad University,
Marrakech, 40000, Morocco

Naima ZEHAR

Agrobiotechnology and Bioengineering Center, CNRST-labeled Research Unit (AgroBiotech-
URL-CNRST-05 Center), Faculty of Science and Technology, Cadi Ayyad University,
Marrakech, 40000, Morocco

Fatiha BENKHALTI

Sustainable Development and Health Research Laboratory, Cadi Ayyad University,
Marrakech, 40000, Morocco

Abstract

Introduction and Purpose This study explores the environmental implications of waste produced from the hydrodistillation process, a common method for extracting essential oils from medicinal and aromatic plants.

Materials and Methods: The focus is on the effects of the solid residue (SR) of *Rosmarinus officinalis* L., a byproduct of this process, on soil properties and the growth of *Solanum lycopersicum*, or tomato plants.

Results: The addition of SR to soil was found to alter key soil properties. It slightly decreased the soil's pH, making it more acidic, while increasing both the electrical conductivity, indicating a higher concentration of dissolved salts, and the organic matter content. However, these changes did not prove beneficial for all types of plants. In the case of tomato plants, their growth abruptly ceased following transplantation into soils amended with 2% and 4% SR. This suggests that the plants experienced abiotic stress, a form of non-living stressors such as changes in soil composition.

Discussion and Conclusion: The findings underscore the need for careful management of residues resulting from hydrodistillation. Therefore, strategies need to be developed to mitigate the potential adverse effects of SR on the environment and plant growth. This study contributes to our understanding of sustainable waste management in the context of essential oil extraction.

Keywords: *Rosmarinus officinalis*, solid residue, hydrodistillation, plant growth, negative impact.

THE RESILIENT FOREST AFTER THE FIRE: ANALYSIS OF THE STRUCTURE AND DIVERSITY OF THE EL-HAMDANIA CORK OAK FOREST- ALGERIA

Ahlem DEBIEB

University of Sciences and Technology Houari Boumediene, Faculty of Biological Sciences,
Laboratory of Ecology and Environment, Algiers, Algeria

ORCID ID: <https://orcid.org/0000-0001-5212-8151>

Leila KADIK

University of Sciences and Technology Houari Boumediene, Faculty of Biological Sciences,
Laboratory of Ecology and Environment, Algiers, Algeria

Michel GODRON

University of Paris, 7. Grands Moulins street, Laboratory of Ladyss, Paris, France

Zineb DJERRAD

University of Sciences and Technology Houari Boumediene, Faculty of Biological Sciences,
Laboratory of Ecology and Environment, Algiers, Algeria

Nacere ADI

University of Sciences and Technology Houari Boumediene, Faculty of Biological Sciences,
Laboratory of Ecology and Environment, Algiers, Algeria

Abstract

Introduction and Purpose: Fire is a disturbance that impacts many terrestrial ecosystems, especially forests, by altering the floristic composition of plant communities. Some species may disappear, while others colonize the burned area. This study aims to assess the diversity and structure of the El-Hamdania cork oak forest, located in the Chr ea National Park in the Blidean Atlas, after a fire.

Materials and Methods: To do so, we conducted a systematic sampling of four vegetation transects, using a synchronic approach, in both conserved and burned areas, characterized by a sub-humid Mediterranean climate at altitudes between 480 m and 700 m. The vegetation analysis focused on the horizontal structure, qualitatively establishing biological spectra and calculating biodiversity indices. This highlighted fire indicator species. We then analyzed the vertical structure, grouping species into distinct strata.

Results and Discussion: The floristic inventory revealed 173 species belonging to 52 botanical families, with a dominance of Asteraceae, which are general indicators of anthropization. The predominant presence of herbaceous therophytes suggests environmental degradation. However, a high biodiversity of heliophytic species was observed after the fire, representing the early stages of the evolutionary dynamics of the cork oak forest during the post-fire succession, likes *Ampelodesmos mauritanicus*, *Calycotome spinosa*, *Cistus salvifolius*, *Genista tricuspidata* and *Erica arborea*. Vertical stratification revealed a low density of the tree layer,

primarily composed of fire-resistant cork oaks "passive pyrophytes", whose cork enables the development of a dense herbaceous layer. This reflects the disturbance within the ecosystem.

Conclusion: The results of this work, provide valuable insight into the structure of the cork oak forest after the fire. We can conclude that we are dealing with a resilient cork oak forest, adapted to fire. These findings pave the way for informed decisions in restoration and conservation programs for this important ecosystem, which remains exposed to environmental constraints.

Key Words: Forest fires, cork oak forest (*Quercus suber*), horizontal and vertical vegetation structure, biodiversity indices, heliophytic species.

EVALUATING THE ROLE OF ENDEMIC BENEFICIAL MICROORGANISMS STRAINS IN BOOSTING THE NUTRITIONAL QUALITY OF *Medicago sativa* L.

Yousra DEBBAH

Laboratory of Research on Medicinal and Aromatic Plants, Science Nature and Life Faculty,
University of Blida 1, Blida, Algeria

Mohamed BENCHERHALI

Laboratory of Research on Medicinal and Aromatic Plants, Science Nature and Life Faculty,
University of Blida 1, Blida, Algeria

Massgo-Saida MOUMEN

Laboratory of Research on Medicinal and Aromatic Plants, Science Nature and Life Faculty,
University of Blida 1, Blida, Algeria

Abstract

Medicago sativa L., commonly known as alfalfa, is a highly valued perennial forage crop widely cultivated for its exceptional nutritional profile. As a key species in sustainable livestock production, its quality directly influences the nutritional value of livestock products. This study evaluated the effect of specific beneficial bacteria strains on the nutritional quality of *Medicago sativa* L. The study was conducted under open-field conditions using a randomized complete block design with five replicates. *Medicago sativa* L. plants were inoculated with specific strains of plant growth-promoting bacteria (PGPB) through seed treatment while a non-inoculated control group was maintained. Forage samples were collected at the vegetative stage and analyzed. DM was determined by drying at 105°C until constant weight, while ash content was measured by incineration at 500°C. CP was quantified using the Kjeldahl method, and CF was analyzed using the Weende method. The results revealed that these treatments significantly enhanced the nutritional value of *Medicago sativa* L. compared to the control. Dry matter content increased by 0.46% and 2.07% in treated groups. Crude protein content rose from 25.12% in the control to 27.09% and 26.18% in the experimental groups, while Ash content increased from 13.10 to 14.71% and 13.93%. In contrast, crude fiber content decreased from 23.87% in the control to 20.68% and 22.88% in the treated group. These findings demonstrate the ability of treatments to boost dry matter, protein, and mineral levels while reducing fiber content, making the forage more suitable for livestock nutrition. This study highlights the potential of PGPB as a sustainable alternative to chemical inputs for improving the quality of forage crops. By enhancing the nutritional profile of *Medicago sativa* L., PGPB applications can contribute to the production of high-quality livestock products. Future research should examine its long-term benefits for livestock health and productivity.

Key Words: *Medicago sativa* L.; beneficial microorganisms; Nutritive value; Plant Growth-Promoting bacteria; Forage crop.

UPDATE OF ALGERIA'S SOIL MAP AT A 1/500,000 SCALE: APPLICATION OF GOOGLE EARTH IMAGERY

Samira KAROU

Department of Soil Science and Environment, Higher School of Agronomy, Mostaganem
(ESAM), Algeria)

ORCID ID: <https://orcid.org/0009-0001-5212-1993>

Abstract

Soil mapping aims to define, classify, and group various soil types into cartographic units. According to the FAO (1990), "A map is the simplest tool to provide an accurate representation of the geographical distribution of various objects, including soils." At the global scale, few studies have focused on soil mapping at very small scales. In Algeria, the soil map at a 1/500,000 scale (Durand, 1954) remains a key reference. However, this document, created using methods now considered outdated, requires revision and updating to enhance its relevance and accuracy.

The objective of this study is to assess the feasibility of updating the soil map of Algeria (Durand, 1954). This update was carried out using Google Earth Pro (GEP), which allowed for the identification of several anomalies and the implementation of necessary modifications through the advanced features of this tool.

The results reveal the presence of inconsistencies in the original map, and the corrections made have significantly improved the geographic accuracy and classification of the soils. This update provides a more accurate and current representation of Algerian soils, better suited to the needs of environmental analysis and modern cartographic applications.

Key Words: Mapping; Soil Map; Update; Google Earth Pr

EVALUATION OF THE INSECTICIDAL ACTIVITY OF ESSENTIAL OILS FROM LAVANDULA STOECHAS AND ORIGANUM FLORIBUNDUM FROM NORTHERN ALGERIA

Lynda LAMOUDI

Laboratory of Transfer Phenomena. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

ORCID ID: <https://orcid.org/0000-0002-9138-7321>

Soraya AKRETCHE

Laboratory of Industrial Process Engineering Sciences. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria

ORCID ID: <https://orcid.org/0000-0003-1937-4267>

Nabil BOUSBIA

Laboratory of Industrial Process Engineering Sciences. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria

ORCID ID: <https://orcid.org/0009-0007-4755-9062>

Nadjet GUEMRAS

Laboratory of Transfer Phenomena. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

ORCID ID: <https://orcid.org/0009-0002-6584-7726>

Kamel DAOUD

Laboratory of Transfer Phenomena. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

ORCID ID: : <https://orcid.org/0000-0000-0000-0000>

Abstract

Introduction and Purpose : This study evaluates the insecticidal activity of essential oils extracted from two Mediterranean plants, *Lavandula stoechas* and *Origanum floribundum*, from northern Algeria, against *Sitophilus oryzae*, a cereal pest. The main objective is to analyze the effect of these essential oils at different concentrations on the mortality of adult *S. oryzae*.

Materials and Methods : The tests were conducted under controlled breeding conditions, with five concentrations of each essential oil (5, 10, 15, 20, and 25 μl /1 ml of acetone) used for the experimentation. The results were measured in terms of insect mortality after prolonged exposure to these concentrations.

Results : The results show that the essential oil of *O. floribundum* has a significant insecticidal effect, particularly at low concentrations (5 μl), with mortalities reaching 54%. Furthermore, insect mortality reached 100% between the 3rd and 6th days of exposure. In comparison, the essential oil of *L. stoechas* shows low efficacy at 5 μl , but the effect becomes significant from 10 μl , also reaching 100% mortality after 6 days at a concentration of 25 μl .

Discussion and Conclusion : The results demonstrate that these essential oils possess insecticidal and repellent properties, with their efficacy increasing proportionally with concentration. The essential oil of *Origanum floribundum* proved to be more potent compared to that of *Lavandula stoechas*, making it more effective in controlling *S. oryzae* at lower concentrations. These findings suggest that these essential oils could serve as natural alternatives for controlling *Sitophilus oryzae* in cereal pest management. Their use would provide a more ecological and sustainable approach compared to chemical insecticides. In conclusion, the essential oil of *O. floribundum* appears particularly promising for future applications in pest control, while *L. stoechas* could also play a role, although higher concentrations would be required to achieve similar efficacy.

Key Words: Insecticidal activity, Essential oils, *Sitophilus oryzae*, *Lavandula stoechas*, *Origanum floribundum*.

ENDOPHYTIC BACTERIUM *BACILLUS CEREUS* -MEDIATED SALT TOLERANCE IN WHEAT PLANTS

Nadira OUKALA

Laboratory of Valorization and Conservation of Biological Resources, University of M'hamed Bougara, Boumerdes, Algeria.

ORCID ID : <https://orcid.org/0000-0003-1317-7211>

ORERO BAYO Marta

Laboratory of Metabolic Integration and Cell Signalling, Plant Physiology Section, University of Jaume I, Spain

Victoria PASTOR

Laboratory of Metabolic Integration and Cell Signalling, Plant Physiology Section, University of Jaume I, Spain

ORCID ID : <https://orcid.org/0000-0003-3750-4184>

Abstract

Introduction and Purpose: Soil salinity has emerged as a serious issue for global food security. The current study investigated the plant growth promotion of bacterial strain *B.cereus* B2 under salinity stress conditions.

Methods: The present study explored the plant growth-promoting effects, as well as the biochemical and molecular characteristics of the endophytic bacterial strain *Bacillus cereus* B2 under salinity stress (0, 300, 600, and 900 mM).

Results: *B.cereus* B2 showed tolerance of up to 900mM NaCl when subjected to salinity stress. The bio-surfactant-producing genes *sfp* and *urfAA* were amplified in this strain, indicating its potential to withstand abiotic stresses. Interestingly, seeds inoculation with *B.cereus* B2 enhanced significantly plant growth even under salinity stress. *B.cereus* B2 showed distinctive salinity tolerance and plant growth-promoting traits including Abscisic acid (ABA), indole-3-acetic acid (IAA), ACC deaminase, siderophore, and exopolysaccharides (EPS) production and phosphate solubilization under salinity stress. Elevated levels of antioxidant enzymes and osmoprotectants such as free amino acids, glycine betaine, and proline were noticed in inoculated plants. *B.cereus* B2 significantly reduced oxidative stress markers (H₂O₂) under salinity stress in inoculated plants.

Discussion and Conclusion: The current study has demonstrated that *B. cereus* B2 may promote plant growth by alleviating salt stress, offering a sustainable and eco-friendly strategy to tackle salinity challenges in agriculture.

Keywords: salt stress, halo-tolerant PGPR, *Bacillus cereus*, plant-microbe interactions

PRECISION AGRICULTURE: "EFFICIENT DEEP LEARNING FOR REAL-TIME AGRICULTURAL IOT SYSTEMS"

Tawseef Ahmad Dar

Central University of Kashmir Ganderbal, Department of Information Technology, J & K,
India

Dr Yash Paul

Central University of Kashmir Ganderbal, Department of Information Technology, J & K,
India

Dr Mohammad Ahsan Chishti

National Institute of Technology Srinagar, Department of Computer Science and Engineering,
J & K, India

Abstract

The agriculture industry plays a crucial role in global food production and economic stability. With the rising global population and increasing food demand, traditional farming methods are becoming inadequate. Recent advancements in Artificial Intelligence (AI) and the Internet of Things (IoT) have introduced automation-driven agriculture, enhancing productivity, efficiency, and resource optimization. These newly adopted methods decrease the scarcity of labour, work efficiently, and fulfil the demand for food. Main theme of this paper is to review the use of Artificial Intelligence and the Internet of Things in different major activities of agriculture. These activities include soil preparation, irrigation, weeding, monitoring, harvesting and spraying with the help of sensors, cloud computing, machine learning assisted databases and other technologies implanted in drones and robots. However, the deployment of deep learning models in real-time agricultural IoT systems remains a challenge due to the resource constraints of edge devices. This paper also presents a comprehensive review of deep neural network models and its compression to enable efficient AI-driven data analytics in smart farming applications. These techniques help in reducing energy consumption, and enhancing real-time decision-making for agricultural tasks. Furthermore, we discuss the integration of lightweight AI models with IoT devices, including drones and edge sensors, to facilitate scalable and cost-effective solutions for precision agriculture.

Discussion and Conclusion: Maintenance of soil fertility, reducing the usage of pesticide, water, herbicides, optimum workforce utilization and finally boosting of productivity and quality is done by these technologies. Hence giving rise to sustainable agriculture with minimal damage to ecology and environment. The utilization of numerous automatic ways of soil preparation, water sensing systems, monitoring, harvesting and spraying drones assisted by machine learning has greater positive impact on sustainable agriculture than traditional practices.

Key words: Artificial Intelligence, Deep Learning, Drones, Precision Agriculture.

THE CASE OF ROSY GARLIC (*ALLIUM ROSEUM L.*) POLYPLOID COMPLEX IN ALGERIA**Thinhinan KHEDIM**

University of Sciences and Technology Houari Boumediene, Departement of Biology and Physiology of Organismes, BPO Laboratory. Algiers, Algeria.

ORCID ID: <https://orcid.org/0000-0002-8904-9438>**Abstract**

Introduction and Purpose: The rosy garlic (*A. roseum L. sens lato*) represents a traditional medicinal herb. All parts of the plant were used in herbal medicine for sulfur compounds involved in antiproliferative and antimicrobial properties. This species complex native from the Mediterranean region includes several subspecies and varieties taxonomically difficult to circumscribe, especially in North Africa due to high polymorphism of local populations. At last, 12 infraspecific taxa were described following ancient floras including the rare endemic *A. roseum* subsp. *odoratissimum*. Unfortunately, unreasonable ethnobotanical collection, overgrazing, climate changes and strong anthropization of localities within its range have a strong impact on the occurrence and survival of its natural populations.

Materials and Methods: This context led us to carry out an ecological and systematic survey in order to understand the extent of this impact and to identify conservation priorities. Plant material from 50 populations sampled in contrasted bioclimatic locations was subjected to multivariate analyses based on morphometric measurements. Chromosome counting was established for each population. Comparative molecular study covered both of Algerian and related Mediterranean populations. Phylogenetic analysis based on nuclear ribosomal ITS1-5.8S-ITS2 DNA sequences, was performed.

Results: Results explain substantial variability of flower segments correlated with diverse bulb structures suggesting occurrence of different breeding strategies. Morphological analyzes revealed two main groups structured in a north-south bioclimatic gradient, suggesting adaptation to arid environments. The phylogenetic tree topology provided a common evolutionary history of the north African populations, which form a single well-supported clade, despite the cytogenetic diversity highlighted with two chromosome numbers $2n=16$ and $2n=32$ as well as structurally variable karyotypes.

Discussion and Conclusion: This work showed the diversity of Algerian material but highlights the scarcity of natural populations. It is therefore essential to integrate our systematic evolutionary data into future conservation programs.

Key Words: Diversity, Ecological survey, Phylogenetics, Rosy garlic, Algeria.

**THE INFLUENCE OF INSTITUTIONAL DYNAMICS AND GOVERNANCE
STRUCTURES ON THE SUPPLY CHAIN OF THE MILK SECTOR IN ALGERIA:
CASE OF THE DAIRY HODNA-LAIT (M'SILA)**

HADJLOUNE Houria

Agropastoralism Research Center (CRAPast), The “Agropastoral and Rural Economy”
Division, Djelfa, Algeria.

ORCID ID: <https://orcid.org/0000-0002-7003-9133>

KACI Ahcène

National Higher Agronomic School (ENSA), Department of Rural Economy, Algiers,
Algeria

ORCID ID: <https://orcid.org/0009-0006-6499-3015>

Abstract

The agri-food sector is the one around which the challenges of food security for the population crystallize and the milk sector is no exception. Moreover, Algeria is the 2nd importer of milk and derivatives. The milk sector is defined through 4 links (production, collection, processing-marketing and consumption) supervised and regulated by the public authorities MADR, ONIL, DSA...etc. The downstream part of the sector is the most dynamic link thanks to the consumer price subsidy policy and the growth of approved dairies (227 dairies according to ONIL) such as the case of Hodna-Lait. However, milk production in Algeria does not allow self-sufficiency (70% of availability is imported). This situation is generally due to the fact that the upstream part of the sector is the weakest link. In this context, the objective of this study is to evaluate the strategies adopted by the dairy industries, institutions and governance structures in this sector and demonstrate whether they will be able to initiate and develop a real dynamic of integration of the milk sector. The analysis consists of 2 stages: the 1st consists of an external diagnosis of the environment according to the Pestel approach in order to measure its degree of influence on the operation of the Hodna-Lait dairy. The 2nd consists of an internal diagnosis of the dairy according to the SWOT approach, in order to characterize and evaluate the performance of its supply strategy in its immediate environment. The main results show almost total control of the dairy supply chain by the State and dependence on the external market (imports), the Hodna-Lait dairy has 3 sources of supply: ONIL (the State), imports and collection from its own fleet of breeders in M'Sila (464 breeders spread over the wilaya of M'Sila). Despite the limits of the generalization of the results, this exploratory research provides useful information to public authorities to identify gaps in supply chains, develop strategies and indicate action levers to design and implement policies to promote a sustainable diet in Algeria in milk and dairy products.

Key Words: Milk supply chain; Hodna-Lait Dairy; ONIL; Breeders; Collectors.

**ANTIOXIDANT ACTIVITY, BIOACTIVE MOLECULES AND TOXICITY STUDY
OF AN ALIMENTARY BYPRODUCT HYDROETHANOLIC EXTRACT**

Moufida ABABOU

Hassiba ben bouali University, Faculty of nature and life sciences, Department of nutrition,
Chlef , Algeria

ORCID ID: <https://orcid.org/0000-0000-0000-0000>

Halima SAIAH

Hassiba ben bouali University, Faculty of nature and life sciences, Department of nutrition,
Chlef , Algeria

ORCID ID: <https://orcid.org/0000-0000-0000-0000>

Abstract

Introduction and Purpose: Nowadays, economic actors as well as consumers are increasingly interested in reducing waste in the agri-food industry. One of the solutions chosen is the study of by-products in terms of nutritional composition and bioactive compounds. The objective of our study is to evaluate the effects of bioactive molecules from by-products of fruit processing on cognitive dysfunctions induced in mice.

Materials and Methods: Evaluation of antioxidant activity(dpph/ ABTS assays).Dosage of bioactive molecules (polyphenols, flavonoids) Study of in vivo toxicity on balb/c mice , every groupe (6 mice) had a different concentration of extract comparing with the blank.

RESULTS: Presence of bioactive molecules.expressing the results on ug / mg of extract according to the standards curves (galic acid and quercetin) Presence of antioxidant capacity. Absence of signs of toxicity. No death no skin or eye lesions.

CONCLUSION: The remaining work is to prove the effectiveness of the extract on the nervous system.

Key Words: bioactive molecules, ethanolic extract, byproduct , antioxidant activity, toxicity.

PLANT SECONDARY METABOLITES TO REDUCE ENTERIC METHANE EMISSION IN DAIRY CATTLE

Dona Mary Eldhose

College of Veterinary and Animal Science, Pookode, Wayanad, India

Sejian V

Rajiv Gandhi Institute of Veterinary Education and Research, Kurumbapet, Puducherry, India

Abstract

Methane emissions from livestock species, including cows, sheep, and goats, pose a significant environmental threat, accounting for the largest share of greenhouse gas emissions from animal agriculture. The rumen microflora plays a vital role in regulating methane production, and researchers have been exploring innovative strategies to mitigate these emissions. The economic viability of using concentrated feeds as a strategy to mitigate methane emissions from livestock is limited in several parts of the world due to the escalating costs of cereal grains and their competing uses in human food systems. To mitigate these emissions, researchers have investigated the potential of plant secondary metabolites, such as polyphenols, terpenoids, and alkaloids. These naturally occurring compounds have been shown to modulate rumen microbial communities, reducing methane production. Certain plant species, including those containing condensed tannins like *Leucaena leucocephala*, have demonstrated notable methane-mitigating effects. Additionally, saponins, essential oils, and flavonoids have shown potential in reducing enteric methane emissions by up to 26% in ruminant livestock. Certain secondary metabolites have been found to alter ruminal fermentation patterns, increasing the production of propionic acid and redirecting hydrogen away from methane production, thereby reducing methanogenesis. Therefore, incorporating these plant compounds into livestock feed could provide a cost-effective and sustainable solution for mitigating methane emissions from ruminant agriculture. Plant secondary metabolites, which serve as natural defense mechanisms, are emerging as a promising and cost-effective feeding strategy for reducing greenhouse gas emissions from livestock. By incorporating these compounds into livestock feed, farmers can promote sustainable livestock production while minimizing environmental impact.

Keywords: Methane, Tannins, Saponins, Essential oils, Secondary metabolite

**DNA MOLECULAR MARKERS AND THEIR APPLICATIONS IN
INVESTIGATING THE GENETIC VARIABILITY OF THE NEMATODE
HETERAKIS GALLINARUM FROM TUNISIA**

Sarra Farjallah

Laboratory of Ecology, Biology and Physiology of Aquatic organisms LR18ES41, Faculty of
Sciences of Tunis, University of Tunis El Manar, 2092, Tunis, Tunisia

Abstract

The nematode *Heterakis gallinarum* is widely distributed across various hosts and regions and is frequently reported in studies on avian helminths. It is one of the most commonly detected nematodes in the gastrointestinal systems of galliform birds. Additionally, *H. gallinarum* acts as a vector for the protozoan parasite *Histomonas meleagridis*, the causative agent of histomoniasis. This disease induces severe lesions in the gut and liver, significantly increasing host mortality and causing substantial economic losses in the poultry industry. Despite its importance, molecular research on *H. gallinarum* populations in Tunisia remains limited. This study aims to investigate the genetic variability of *H. gallinarum* samples collected from poultry across various locations in Tunisia using nuclear and mitochondrial molecular markers (ITS and COI). Analyses of haplotype and nucleotide diversity reveals low values, suggesting that the caecal nematode exhibits very limited genetic diversity. Haplotype network and phylogenetic tree analyses showed no significant geographical structuring of *H. gallinarum* populations within Tunisia. Furthermore, mismatch distribution and neutrality tests indicated that at least one population expansion event has occurred in the history of *H. gallinarum*. Both molecular markers showed an absence of genetic differentiation ($F_{ST} = -0.02/F_{ST} = -0.001$, for ITS and COI sequences, respectively) among populations, indicating considerable gene flow facilitated by the movement of infected birds. This finding highlights the critical role of poultry trade in promoting gene flow within *H. gallinarum* populations.

Keywords: *Heterakis gallinarum*//ITS/COI/ genetic variability /gene flow/ poultry trade/Tunisia.

EPIDEMIOLOGICAL SURVEY AND MOLECULAR CHARACTERIZATION OF MONOGENEAN PARASITE SPECIES INFESTING *SPARUS AURATA* FROM TUNISIA

Sarra Farjallah

Laboratory of Ecology, Biology and Physiology of Aquatic organisms LR18ES41, Faculty of Sciences of Tunis, University of Tunis El Manar, 2092, Tunis, Tunisia

Khadija Doukali

Laboratory of Ecology, Biology and Physiology of Aquatic organisms LR18ES41, Higher Institute of Marine Sciences, University of Carthage, Errimel B.P.15., 7080 Bizerte, Tunisia

Abstract

The expansion of marine aquaculture has raised significant environmental concerns, impacting wild fish communities and increasing the prevalence of pathogens, with potential for pathogen transmission between wild and farmed fish. This study investigates the occurrence and molecular identification of gill monogenean parasites in gilthead seabream (*Sparus aurata* Linnaeus, 1758) from multiple locations in Tunisia. Unlike previous studies that primarily relied on either morphological or genetic identification, this study integrates both approaches. A total of 97 fish were examined, resulting in 451 monogenean specimens recovered from their gills. Morphological analysis initially classified the monogeneans into two species from distinct subclasses: *Lamellodiscus echeneis* (Wagener, 1857) of the Monopisthocotylea and *Sparicotyle chrysophrii* (Van Beneden & Hesse, 1863) Mamaev, 1984, of the Polyopisthocotylea. The study assessed infection rates of *L. echeneis* and *S. chrysophrii* in both wild and farmed populations of gilthead seabream. *L. echeneis* exhibited a prevalence of 92.3% in wild seabream and 91.1% in cage-farmed seabream, while *S. chrysophrii* showed a prevalence of 45% in wild seabream and 45.61% in farmed seabream. Molecular characterization was performed using nuclear ITS rDNA markers and a partial fragment of the mitochondrial cytochrome oxidase subunit I (COI) gene for *L. echeneis*, as well as the large subunit 28S and COI gene for *S. chrysophrii*. Phylogenetic analyses (NJ, ML, and BI) placed all *L. echeneis* samples from various locations into a single clade. Similarly, sequences of *S. chrysophrii* from different sites were highly conserved, clustering into a distinct clade with strong bootstrap and posterior probability support. These findings confirm the widespread presence of these monogenean species across the studied regions. This study provides valuable epidemiological data on *L. echeneis* and *S. chrysophrii* in gilthead seabream from Tunisia, offering precise molecular characterization using nuclear and mitochondrial markers. These results lay the groundwork for future research on monogenean parasites in Tunisian waters and other economically important aquaculture species.

Keywords: *Sparus aurata* /Infection rates/Molecular characterization/*L. echeneis*/*S. chrysophrii*.

PHYSIOLOGICAL AND BIOCHEMICAL CHARACTERIZATION OF SOME STRAINS OF ACTENOBACTERIA ISOLATED FROM AN ALGERIAN SAHARAN SOIL WITH A POTENTIAL FOR INHIBITION OF CERTAIN FUNGI AND PHYTHOPATHOGENIC BACTERIA

Fedwa BEGHDAI

Laboratory of Molecular Biology, Genomics and Bioinformatics (LBMGB), Faculty of Natural Sciences and

El-Hadj DRICHE

Life (SNV), Hassiba Benbouali University of Chlef, Hay Salem, 02000 Chlef, Algeria.

Abstract

Algerian Saharan soil is an ecological product widely explored as a niche for microorganisms that produce biologically active natural compounds. Among the microorganisms of biotechnological interest, actinomycetes or actinobacteria.

The importance of actinobacteria has been highlighted in various fields: industrial, medical and veterinary, as well as in the field of agriculture and agri-food. Actinobacteria, especially those with a mycelial structure, are known for the production of antibiotics, especially the genus *Streptomyces*.

Other bioactive molecules are also secreted by actinobacteria such as, enzymes, vitamins, immunosuppressants, insecticides and herbicides, and *Streptomyces* are the best candidates for the production of biologically active secondary metabolites. Indeed, this bacterial genus is responsible for about 70% of the antibiotic molecules used in medicine and 60% of the antifungals used in agriculture.

The different physiological tests consist of inoculating the isolates on ISP2 medium in the presence of different NaCl concentrations, at different pH and temperature according to the method of Athalye et al, (1985). Different moderate concentrations of NaCl (3, 5, 7%) are added. For pH, the ISP2 medium is adjusted to pH values (5, 9). Growth is visually assessed after 8 days of incubation at 30°C.

And to determine the biochemical characteristics of our isolates from the tests carried out such as: use of carbohydrate compounds, assimilation of amino acids, degradation of gelatin, production of melanoid pigments...

The results of the study of the enzymatic activities of actinomycetes showed that the genera *Streptomyces*, *Micromonospora*, *Nocardia* and *Streptosporangium* possess significant enzymatic activity (**Okpukpara and George-okafor, 2016**).

Keywords: Actinobacteria, characterization, Bioactive molecules, Inhibition.

ANTIBACTERIAL AND ANTIFUNGAL ACTIVITY OF PLANT (MARRUBIUM VULGARE) FROM THE FAMILY OF LAMIACEAE IN WESTERN ALGERIA

AOUFI Farida

1Laboratoire d'environnement et développement durable, Faculty of Science and Technology,
Relizane University

TAMERT Asma

1Laboratoire d'environnement et développement durable, Faculty of Science and Technology,
Relizane University

ABDI Yamina

1Laboratoire d'environnement et développement durable, Faculty of Science and Technology,
Relizane University

BARBARA Ferdaws

1Laboratoire d'environnement et développement durable, Faculty of Science and Technology,
Relizane University

Abstract

Antibacterial activity was evaluated for three essential oils (*M.v*) against five Bacterial strains: *E.coli*, *s.aureus*, *Pseudomonas aeruginosa*, *Candida albican*, and *Aspergillus niger*. The study also evaluated the antibacterial and antifungal activity of essential oils: Essential oil L showed promising antibacterial and antifungal activity. It inhibited the growth of *E. coli*, *S. aureus*, and *Candida albicans* with inhibition zones ranging from 12mm to 50mm. However, it had no effect on *Aspergillus niger* and *Pseudomonas aeruginosa*. This suggests potential for essential oil L as an antimicrobial agent, but Further research is needed. Essential oil S exhibits effective antibacterial activity against *E. coli* and *Staphylococcus aureus*, with increasing inhibition zones at higher concentrations (100% to 50%). The antibacterial activity of essential oil (M.V) is négative, indicating that it has no effect on the growth of (*E.coli*), (*s.aureus*), or *Aspergillus niger*. However, it does exhibit a Slight positive activity against *seudomonas aeruginosa* at concentrations of 50 mg/ml and 100 mg/ml, with inhibition zones of 9mm and 13mm.

Key words: (Marrubium vulgare), antibacterial, *Candida albicans*, *Aspergillus niger*, *Pseudomonas aeruginosa*, (*E. coli*), (*S. aureus*)

GREEN SYNTHESIS OF COPPER OXIDE NANOPARTICLES USING CITRUS SINENSIS PEEL EXTRACT AND EXPLORATION OF THEIR ANTIBACTERIAL ACTIVITIES

V. S. Angulakshmi

Assistant Professor, Department of Chemistry, PSGR Krishnammal College for Women,
Coimbatore, TN, India

N. Anusuya

Assistant Professor, Department of Chemistry, PSGR Krishnammal College for Women,
Coimbatore, TN, India

R Nithya

PG Student, PSGR Krishnammal College for Women, Coimbatore, TN, India Associate
Professor, Department of Chemistry, Rathinam Technical Campus, Tamilnadu, India

S Kalaiselvan

Assistant Professor, Department of Chemistry, PSGR Krishnammal College for Women,
Coimbatore, TN, India

Abstract

Nanotechnology represents cutting-edge technology due to its potential to revolutionize various industries and address complex challenges through manipulation at the nanoscale. Nanomaterials can enhance microbial activity and biodegradation rates by serving as carriers for microbial cells, enzymes or nutrients, protecting them from environmental stressors and improving their accessibility to pollutants. Nanoparticles synthesized from plant extracts, often referred to as green nanoparticles, have gained noteworthy consideration due to their eco-friendly and sustainable nature, as well as their potential applications in various fields including medicine, agriculture, and environmental remediation. This paper focused on the synthesis of Copper oxide nanoparticle using *Citrus sinensis* peel extract and investigation of their antimicrobial activities. The synthesized nanoparticles were characterized to determine their size, shape, composition and stability using techniques such as UV-Vis spectroscopy, scanning electron microscopy (SEM), X-ray diffraction (XRD), and Fourier-transform infrared spectroscopy (FTIR).

Keywords: Copper oxide nanoparticle, *Citrus sinensis* peel, Green synthesis

TOXICITY ASSESSMENT OF ETHANOLIC EXTRACT OF *ROSMARINUS OFFICINALIS* LEAVES IN FEMALE WISTAR RATS

Said Babou

Biology and Health laboratory, Faculty of sciences, Ibn Tofail University, Kenitra, Morocco.

Miloud Chakit

Biology and Health laboratory, Faculty of sciences, Ibn Tofail University, Kenitra, Morocco.

Radia El Gui

Biology and Health laboratory, Faculty of sciences, Ibn Tofail University, Kenitra, Morocco.

Abdelhalem Mesfioui

Biology and Health laboratory, Faculty of sciences, Ibn Tofail University, Kenitra, Morocco.

Youssef Sqalli-Houssaini

Biology and Health laboratory, Faculty of sciences, Ibn Tofail University, Kenitra, Morocco.

Abstract

Rosmarinus officinalis (Rosemary), an aromatic plant rich in phenolic antioxidant compounds, is widely used in traditional therapy by Moroccan population. The objective of the current study was to assess the acute and subacute toxicity of ethanolic extract of *Rosmarinus officinalis* (EERO) in Wistar rats.

30 rats were divided into 5 groups (control, 1000, 2500, 3500, and 5000 mg/kg dose). Similarly, in the subacute toxicity study, 30 rats were divided into 5 groups (control, 200, 300, 500 and 1000mg/kg dose). The sub-acute toxicity study aimed to identify the NOAEL dose, by following the same signs of toxicity, hematological parameters, biochemical parameters, markers of oxidative stress, and histological analyses were evaluated.

No significant differences were observed in acute and sub-acute toxicity parameters (body weight, food, water), nor acute toxicity was any mortality observed, suggesting that the LD50 is greater than 5000 mg/kg. The sub-acute toxicity study showed no significant differences in hematological, biochemical, and histological parameters, while the extract increased CAT levels and decreased NO in the liver and kidney.

RO extract is non-toxic up to the doses tested and exerts antioxidant activity, manifested by a nitric oxide levels decrease and antioxidant enzyme catalase levels increase in in liver and kidneys.

Keywords: *Rosmarinus officinalis*, hematological analysis, acute and subacute toxicity, catalase (CAT), nitric oxide (NO)

**EFFECT OF NEW GENERATION FUNGICIDES AGAINST *Pyricularia oryzae*
CAUSES BLAST IN RICE (*Oryza sativa* L.)**

VIGNESH K

Assistant Professor, Palar Agricultural College, Ambur
ORCID: 0000-0003-4484-3862

Dr. K. SELVAM

Assistant Professor, Palar Agricultural College, Ambur
ORCID: 0000-0002-0060-2314

ARSHA G

Assistant Professor, Palar Agricultural College, Ambur
ORCID: 0000-0009-1694-9716

Abstract

Rice blast, caused by *Pyricularia oryzae*, is one of the most destructive fungal diseases affecting *Oryza sativa* L., leading to severe yield losses globally. The emergence of resistant fungal strains necessitates the development and evaluation of new-generation fungicides for effective disease management. This study assessed the efficacy of novel fungicides, including strobilurins, triazoles, and SDHI (succinate dehydrogenase inhibitors), against *P. oryzae* under laboratory and field conditions. Results indicated that fungicides such as azoxystrobin, tebuconazole, and fluxapyroxad exhibited strong inhibitory effects on spore germination, mycelial growth, and lesion development. Field applications of these fungicides significantly reduced disease incidence and improved plant health without causing adverse environmental effects. The findings suggest that new-generation fungicides provide a promising and sustainable approach for managing rice blast while minimizing the risks of resistance development and environmental toxicity.

Key Words: Rice, Blast, *Pyricularia oryzae*, Fungicide.

PHYSIONOMIC CHANGE OF VEGETATION AND DIVERSITY INDEX IN TWO CONSTRAINED ENVIRONMENT OF THE SEMI-ARID IN ALGERIA

Ihaddaden Akli

High National School of Nanoscience and Nanotechnology. University Abdelhafid Ihaddaden

ABLA Safia

Département SNV, Faculty of sciences University of Algiers

Abdoun Fatiha

University of Sciences and Technology Houari Boumediène, *USTHB, FSB, LEVE*.

Abstract

Introduction and Purpose: We have analyzed the influence of biodiversity and the environment factors on species diversity of the pre-forested and pre-steppic Green Oak formations, along a short gradient of rainfall in two semi-arid constrained regions. We looked at the changes induced on vegetation i) what are the changes on vegetation cover and vegetation physiognomy ii) and what are the changes on the specific richness and specific diversity.

Methods: We conducted a floristic sampling in two semi-arid regions, following two approaches: The ecological approach, which defines ecological diversity as a qualitative component of diversity, where species are grouped based on the similarity of their functional traits. The specific approach (specific diversity), which is a quantitative approach to diversity, involving the calculation of alpha diversity using the Shannon-Weaver index.

Results: However, the physiognomic changes on vegetation at Theniet El Had and Aflou show that the more environmental constraint is important, the more the changes on diversity of dominant species are important

Quantitative diversity analysis shows that low specific richness does not necessarily have a negative effect on diversity, so a region with relatively more rainfall is not synonymous with better diversity than a region with lower precipitations, it is the heterogeneity of environment that gives a higher diversity.

Key Words: Biodiversity, Vegetation cover, Alpha diversity, Species richness, constrained environments.

UNDERSTANDING MOROCCAN POPULATION'S PERCEPTION AND AWARENESS OF ORGANIC FOOD

Soumaya Atouife

Training and Research Unit on “Nutrition & Food Science”, LABS, Faculty of Sciences,
Chouaïb Doukkali University, El Jadida, Morocco

Jamila Elbiyad

Training and Research Unit on “Nutrition & Food Science”, LABS, Faculty of Sciences,
Chouaïb Doukkali University, El Jadida, Morocco

Halima Belaoufi

Training and Research Unit on “Nutrition & Food Science”, LABS, Faculty of Sciences,
Chouaïb Doukkali University, El Jadida, Morocco

Abdellah El Habazi

Training and Research Unit on “Nutrition & Food Science”, LABS, Faculty of Sciences,
Chouaïb Doukkali University, El Jadida, Morocco

Rekia Belahsen

Training and Research Unit on “Nutrition & Food Science”, LABS, Faculty of Sciences,
Chouaïb Doukkali University, El Jadida, Morocco

Abstract

Over recent decades, significant changes in food production and consumption patterns have reshaped the global food system, with notable impacts on both the environment and human health. These shifts have been primarily driven by the need to increase food production to address food security challenges and the demands of a growing global population. This increase in production has influenced dietary habits by improving food availability and enhancing food security. However, these changes have also led to serious environmental issues, including the depletion of natural resources, deforestation, biodiversity loss, pollution, and climate change. Furthermore, unsustainable food consumption patterns exacerbate these challenges, both in terms of quantity and quality.

In this context, organic agriculture has emerged as a promising alternative to promote sustainable food consumption. This study aims to evaluate Moroccans' perceptions of organic agriculture through an online questionnaire completed by 423 participants. The findings indicate that 81.7% of respondents consume organic foods, primarily for their perceived health and environmental benefits, while 85.5% cited high costs as the main barrier to consumption. Additionally, the study revealed significant confusion between organic and local products among 66.3% of participants.

In conclusion, the Moroccan population generally perceives organic food as superior to conventionally grown products in terms of quality and taste. However, the consumption of organic products remains limited. Despite the relatively underdeveloped state of organic agriculture in Morocco, the findings suggest that it holds promise as a pathway toward sustainable food consumption.

Key words: Organic food; local food; consumption; perception.

**ANTILIPIDEMIC AND NEPHRO-HEPATOPROTECTIVE ACTIVITIES OF
ROSMARINUS OFFICINALIS ETHANOLIC EXTRACT IN WISTAR RATS**

Said Babou

Biology and Health Laboratory, Faculty of Sciences, Ibn Tofail University, Kenitra, Morocco.

Miloud Chakit

Biology and Health Laboratory, Faculty of Sciences, Ibn Tofail University, Kenitra, Morocco.

Abdelhalem Mesfioui

Biology and Health Laboratory, Faculty of Sciences, Ibn Tofail University, Kenitra, Morocco.

Youssef Sqalli Houssaini

Biology and Health Laboratory, Faculty of Sciences, Ibn Tofail University, Kenitra, Morocco.

Abstract

Rosemary, *Rosmarinus officinalis*, from the vernacular name Azir, is medicinal plant used by Moroccan population, it contains chemical compounds with great nutritional value and multiple phytotherapeutic virtues. The study aims to assess the effect of ethanolic extract of *Rosmarinus officinalis* on lipidic status and blood parameters in rats.

Male rats of Wistar strain were divided into 3 groups: control groups and two groups receiving the *R. officinalis* extract, one fed a standard diet and the other receiving a fat-enriched diet. The ethanolic extract of the *Rosmarinus officinalis* leaves was administered orally at a rate of 300mg/Kg for the group 2 and 1000mg/Kg for the group3. mg/kg of rat weight for 4 weeks. Serum parameters were measured using a spectrophotometer including: alanine aminotransferase (ALAT) and aspartate aminotransferase (ASAT) at 340nm, creatinine (Cr), cholesterol, glucose, total cholesterol (CHO), triglycerides (TG) at 500nm and urea at 600nm.

The results show that EERO administration for 28 days decreases the levels of ALT and AST urea and glucose ($P < 0.05$), indicating a nephron-hepatoprotective activities, and without effect on cholesterol levels in treated rats with 300 mg/Kg and 100mg/Kg in comparison with control rats however, the triglyceride level was decreased in rats treated by EERO indicating an antilipidemic propriety of rosemary.

These findings show that EERO has a benefic effect on lipidemic, hepatic and nephrotic parameters.

Keywords: *Rosmarinus officinalis*, blood, liver, kidney, biochemical parameter, Rats.

**FUSARIUM ANTAGONISM IN SOIL AND RESOURCE COMPETITION DRIVE
THE SUPPRESSION OF BAYOUD DISEASE IN MOROCCAN DATE PALM
PLANTATIONS**

Youssef El Hilali Alaoui

Laboratory of Agro-Industrial and Medical Biotechnology, Faculty of Sciences and Technics,
Sultan Moulay Slimane University, B.P. 523, Beni Mellal, Morocco

Said Bouda

Laboratory of Agro-Industrial and Medical Biotechnology, Faculty of Sciences and Technics,
Sultan Moulay Slimane University, B.P. 523, Beni Mellal, Morocco

Hajar Mhammedi

Laboratory of Agro-Industrial and Medical Biotechnology, Faculty of Sciences and Technics,
Sultan Moulay Slimane University, B.P. 523, Beni Mellal, Morocco

Adil Essarioui

Oasis System Research Unit, Regional Center of Agricultural Research of Errachidia,
National Institute of Agricultural Research, PO. Box 415, Rabat 10090, Morocco

Abstract

Bayoud disease, caused by the soilborne fungus *Fusarium oxysporum* f. sp. *albedinis*, threatens date palm cultivation in Morocco. Certain soils exhibit a natural ability to suppress this disease, although the mechanisms remain unclear. This study investigated the functional traits of *Fusarium* isolates and the composition of bacterial and fungal communities in two conducive soils and one suppressive soil to better understand Bayoud suppression. Thirty saprophytic *Fusarium* isolates were randomly selected from each soil sample. The in vitro inhibitory activity of each isolate against the Bayoud pathogen was evaluated, and their growth on 95 carbon substrates was assessed using Biolog plates. ITS1 and 16S ribosomal genes were sequenced to analyze fungal and bacterial diversity and composition in each soil sample. The results revealed that *Fusarium* isolates from one conducive soil exhibited strong antagonistic activity but slower growth, indicating trade-offs between inhibition and growth. In contrast, isolates from the second conducive soil showed growth efficiency similar to the pathogen but less frequent and weaker antagonistic behavior. Isolates from the suppressive soil demonstrated intermediate antagonism and the highest growth efficiency. Microbial diversity was greater in suppressive soils than in conducive soils, with enrichment in microbial taxa such as *Fusarium*, *Aspergillus*, and Actinomycetes, known for producing antimicrobial compounds. These findings suggest that microbial antagonism and resource competition work together in suppressive soils to limit the spread of Bayoud disease in Moroccan date palm plantations.

Keywords: Soilborne fusarium, date palm, Moroccan oases, nutritional competition, *Fusarium oxysporum* f. sp. *albedinis* (*Foa*), Pair niche overlap.

BIOACTIVE COMPOUNDS IN OILSEED WASTE: ANTIOXIDANT PROPERTIES AND LC-ESI-QTOF-MS/MS-BASED PROFILING

Ali Imran

Department of Food Science, Faculty of Life Science, Government College University,
Faisalabad, Pakistan,

School of Agriculture, Food and Ecosystem Sciences, Faculty of Science, The University of
Melbourne, Parkville, Victoria, Australia

Hafiz Suleria

School of Agriculture, Food and Ecosystem Sciences, Faculty of Science, The University of
Melbourne, Parkville, Victoria, Australia

Safura Kousar

Department of Food Science, Faculty of Life Science, Government College University,
Faisalabad, Pakistan,

Muhammad Usama Tahir

Department of Food Science, Faculty of Life Science, Government College University,
Faisalabad, Pakistan,

Abstract

Oilseed waste, a significant byproduct of agro-industrial processing, represents a promising yet underutilized source of bioactive compounds. This study investigates the antioxidant properties and phenolic content of various oilseed waste types, complemented by detailed compound profiling using LC-ESI-QTOF-MS/MS. Antioxidant activities were assessed through established spectrophotometric methods, while the Folin-Ciocalteu assay was employed to quantify total phenolic content. Advanced LC-ESI-QTOF-MS/MS analysis enabled precise identification of key bioactive compounds, including flavonoids, phenolic acids, and other phytochemicals. The results revealed considerable variability in antioxidant capacity and phenolic composition among different oilseed waste samples, reflecting their diverse chemical profiles. This research highlights the potential of oilseed waste as a valuable resource for bioactive compounds, opening new avenues for applications in functional foods, nutraceuticals, and pharmaceuticals within a sustainable circular economy framework.

Keywords: waste valorization, Bioactive characterization, Antioxidants Potential

THE STUDY OF ANTIDIARRHEAL POTENTIAL OF HELICTERES ISORA AND PERSICARIA VIVIPARA METHANOLIC EXTRACTS IN SWISS ALBINO MICE

Muhammad Muzammil Nazir

Department of Zoology, Government College University, Faisalabad 38000, Pakistan

Munaza Yasmeen

Department of Zoology, Government College University, Faisalabad 38000, Pakistan

Iqra Farzeen

Department of Zoology, Government College University, Faisalabad 38000, Pakistan

Sana Inam

Department of Pharmaceutics, Government College University, Faisalabad 38000, Pakistan

Asma Ashraf

Department of Zoology, Government College University, Faisalabad 38000, Pakistan

Abstract

Diarrhea has been a major cause of death, especially among children in developing countries. Medicinal plants are the major sources of traditional treatment of disease due to abundant presence of plants and vast side effects of synthetic drugs. The increasing antibiotic resistance exhibited by microorganisms has led to the phytochemical screening of medicinal plants for antidiarrheal activity. The roots of *Periscaria vivipara* and fruit of *Helicteres isora* are used in folk medicine. The present study was conducted to analyse phytochemicals and in vivo antidiarrheal activity of *H. isora* and *P. vivipara* was evaluated. The extraction of plant parts was carried out by using Soxhlet apparatus and methanol as solvent. For evaluation of phytochemicals GC-MS analysis was carried out. For determining flavonoid and phenolic compound TFC and TPC were utilized respectively. For evaluating antibacterial activity well diffusion method was employed. Fresh roots of *P. vivipara* and fruits of *H. isora* were dried, powdered (250 g), and macerated in 500 ml methanol. Mice (32–36 g, 50:50 sex ratio) were acclimatized for a week and divided into five groups each of six mice: control, standard (1 mg/kg loperamide), and three treatment groups (100, 200, and 400 mg/kg extracts). The control group received 1 ml/kg saline. Statistical analysis was conducted using GraphPad Prism, with significance set at $p < 0.05$. Castor oil-induced diarrhea and gastro-intestinal motility tests were employed to assess the antidiarrheal efficacy of methanolic extracts of *P. vivipara* and *H. isora*. GC-MS analysis of *P. vivipara* root methanolic extract revealed presence of Oleic Acid, 6-Octadecenoic acid, 9-Octadecenoic acid, methyl ester, (E)- and Toluene abundantly. Whereas, Oleic Acid, 9,12-Octadecadienoic acid (Z, Z)-, methyl ester, Oleic Acid, cis-13-Octadecenoic acid, methyl ester and Methyl stearate phytochemicals were abundantly present in methanolic fruit extract of *H. isora*. Total phenolic and total flavonoid contents at different concentrations (25, 50, 75 and 100 mg/ml) of methanolic extract of *H. isora* were 90.09 ± 0.33 , 91.17 ± 0.66 , 91.39 ± 1.00 , 92.76 ± 0.33 mg GAE/g and 22.76 ± 0.66 , 98 ± 0.33 , 23.39 ± 0.83 and 24.12 ± 0.66 mg QE/g respectively. Moreover, total phenolic and total flavonoid contents at different

concentrations (25, 50, 75 and 100 mg/ml) of methanolic extract of *P. vivipara* were 120.23±0.33, 121.53±0.66, 123.39±1.03, 124.76±1.33 mg GAE/g and 31.76±0.33, 32.14±0.66, 33.26±1.00, 34.68±1.35 mg QE/g respectively. MEPV showed inhibition zones of 12±0.66 mm and 11±1.0 mm, while MEHI showed 5±0.33 mm and 14±1.33 mm against *E. coli* and *S. aureus* respectively, compared to ciprofloxacin's 27±1.33 mm and 28±2.0 mm. Results of antidiarrheal evaluation revealed that MEHI and MEPV indicated varying degrees of diarrhea inhibition, with percentages of 11.32%, 26.38%, 52.83% and 21.01%, 34.2%, 42.16% found at serial doses respectively, compared to loperamide. Both plant extracts showed significant delay in onset of diarrhea, suppression in weight and number of diarrheal stools in a dose dependent manner. MEHI and MEPV exhibited significant percentage inhibition in gastrointestinal transit as 33.94%, 36.54%, 41.94% and 18.44%, 27.35%, 40.36% respectively. Moreover, dose dependent reduction in peristaltic index was observed for MEPV and MEHI. From the outcomes of this study, it is inferred that methanolic extract from roots of *P. vivipara* and fruit of *H. isora* exhibited substantial antidiarrheal efficacy.

Keywords: Antidiarrheal, Mice, *Helicteres isora*, *Periscaria vivipara*, GC-MS, Castor oil, Gastrointestinal motility.

TECHNOLOGY TRANSFER AND ITS ROLE IN ENHANCING GLOBAL AGRICULTURAL PRODUCTIVITY

Dhivya C

Research Scholar (Agricultural Extension Education), Department of Agricultural Extension and Rural Sociology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India

ORCID: 0009-0004-8984-1812

Arunkumar R

Research Scholar (Agricultural Extension Education), Department of Agricultural Extension and Rural Sociology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India

ORCID: 0000-0001-5580-5170

Abstract

Global agricultural productivity has witnessed significant transformations due to the rapid transfer of technology across borders. Technology transfer in agriculture encompasses the dissemination of innovations such as precision farming, biotechnology, artificial intelligence, and climate-smart practices, which play a crucial role in improving efficiency, sustainability, and food security. In an era of globalization, international collaborations, public-private partnerships, and research institutions have facilitated the exchange of knowledge, helping to bridge the technological divide between developed and developing nations. This paper explores the mechanisms of agricultural technology transfer, including intellectual property rights, foreign direct investment, and knowledge-sharing platforms. It analyzes the role of multinational corporations, government policies, and international organizations in fostering technology dissemination. Additionally, it highlights case studies of successful adoption of agricultural technologies in various regions, demonstrating their impact on productivity, resource optimization, and climate resilience. Despite its benefits, technology transfer in agriculture faces challenges such as affordability, lack of infrastructure, resistance to change, and disparities in access. The paper discusses strategies to overcome these barriers, emphasizing the need for inclusive policies, localized adaptation of technologies, and capacity-building initiatives for farmers. By fostering effective technology transfer, global agriculture can address critical issues such as food security, climate change adaptation, and sustainable resource management. Strengthening international cooperation and promoting equitable access to agricultural innovations will be key to ensuring that productivity gains benefit farmers worldwide, contributing to a resilient and sustainable global food system.

Keywords: Technology Transfer, Agriculture, Productivity and Sustainable food system

APPLICATION OF NEW BIOTECHNOLOGICAL PRACTICES FOR SUSTAINABLE AGRICULTURE: BIOFERTILIZATION

Nassima BAHA

University of Sciences and Technology Houari Boumedienne, Algiers, Algeria,
<https://orcid.org/0000-0001-7548-2743>

Sabrina BEHAIRI

University of Sciences and Technology Houari Boumedienne, Algiers, Algeria

Roumaissa MOULKAF

University of Sciences and Technology Houari Boumedienne, Algiers, Algeria

Massakib BEKKAYE

University of Sciences and Technology Houari Boumedienne, Algiers, Algeria

Abstract

Introduction and Purpose: In arid and semi-arid regions, salinization is more critical as precipitation decreases and temperatures increase. It is also caused by agricultural practices, including poor water management, intensive irrigation, and exposure to seawater. Salinity has many effects on plant growth by restricting the flow of water to plants (osmotic stress), or by directly damaging plant cells through the accumulation of specific ions, nutritional imbalance, or a combination of these factors. Sustainable agriculture, which now incorporates biotechnological practices, has become a key concept. It involves the use of beneficial microorganisms, such as plant growth-promoting rhizobacteria (PGPR), which improve plant growth by various means.

Materials and Methods: In the same context, this study was carried out to investigate the effects of bio-fertilization by two halotolerant PGPR on the salt tolerance of lucerne (*Medicago sativa L.*) at germination and growth stages. At vegetative growth, photosynthetic pigments, mineral and relative water (RWC), proline, total soluble sugars and soluble proteins contents were monitored.

Results: In absence of salt stress the PGPR seems to have little or no effect on germination when inoculated singly. In fact, only a mixture of these PGPR showed a positive effect on germination. Inversely, all inoculations positively affect growth parameters. In opposite, under salt stress, all single inoculation and co-inoculation alleviate the adverse effects caused by salt stress on lucerne germination. Regarding to inoculated plants, salinity affected negatively growth parameters which remained better than without bacterial inoculation.

Discussion and Conclusion: These results indicate that all inoculations and co-inoculation with both PGPRs and especially at the proportion 1/2 could mitigate the negative effects of salinity on alfalfa. Suggesting therefore that this bio-fertilization could be used as a novel biotechnology that increases plant productivity, constitutes a promising alternative to chemical fertilizers and also represents an effective way to restore and make good use of soils affected by salinity.

Keywords: Biofertilization .Salt stress. Lucerne. Plant growth. PGPR

EVALUATION OF GENETIC DIVERSITY AND GENOME FINGERPRINTING OF BITTER GOURD GENOTYPES(*MOMORDICA CHARANTIA*) BY MOLECULAR MARKER

Farzana Iftikhar

Department of Zoology, The Women University Multan.

Faisal Iftikhar

Department of Agronomy, Faculty of agricultural sciences and technology, Bahauddin Zakariya University Multan

Abstract

Random Amplified Polymorphic DNA (RAPD) was used to evaluate such relationship among three genotypes of bitter gourd. Amplification of the genomic DNA from each of the three genotypes of bitter gourd using all the 10 decamer primers revealed a variety of RAPD patterns. Out of these 10 primers, five showed stable pattern with a set of major DNA fragments and were the most effective in detection of genetic polymorphism in the studied populations of Bitter gourd. The 10 decamer oligonucleotides on an average generated 244 fragments, of which 85 (34.8%) were polymorphic. A range of 15 to 47 bands was amplified from RAPD primer used in the study. The degree of polymorphism among primers was different. The approximate size of the amplified fragments ranged from 250 to 1000 bp. Coefficients were utilized to generate dendrogram by using an unweighted pair group method with arithmetic average (UPGMA) tree. According to the genetic similarity dendrogram constructed for all 244 traits for three genotypes detected using 10 primers, all studied Bitter gourd from the three different varieties of Bitter gourd can be subdivided into two clusters, of which cluster one comprises of two varieties of Bitter gourd (CBT-36 & PKBT-1), and cluster two comprises of one variety (BG-7017). The mean genetic distance between varieties of Bitter gourd within cluster 1 is 0.3232 (32 %), within cluster 2 is 0.4419 (44 %). The genetic distance between clusters 1 and 2 was near about 0.1187 (11.8%).

Key words: Amplified Polymorphic DNA, Bitter gourd, genotype, primer ,DNA fragments.

PHENOLIC COMPOUNDS OF OPUNTIA STRICTA CLADODES

LAZREG Halima

Laboratoire de Productions, Valorisation Végétale et Microbienne.Oran.Algeria

SEBAA Hanane Siham

Laboratoire de Productions, Valorisation Végétale et Microbienne.Oran.Algeria
Biotechnology Departement. Faculty of Sciences ; life and Nature, University of Sciences and
Technology Oran Mohammed Boudiaf USTO-MB.31000.Oran,Algeria.

CELIK Muhammed Suleyman

Alum Center Iğdir University ;Turkey

ERENLER Ramazan

Alum Center Iğdir University ;Turkey

ABDEDDAIM Katia Khadidja

Laboratoire de Productions, Valorisation Végétale et Microbienne.Oran.Algeria
Biotechnology Departement. Faculty of Sciences ; life and Nature, University of Sciences and
Technology Oran Mohammed Boudiaf USTO-MB.31000.Oran,Algeria.

DJABEUR Abd Erezzak

Laboratoire de Productions, Valorisation Végétale et Microbienne.Oran.Algeria
Biotechnology Departement. Faculty of Sciences ; life and Nature, University of Sciences and
Technology Oran Mohammed Boudiaf USTO-MB.31000.Oran,Algeria.

Abstract

Introduction: *Opuntia stricta* is a species of large cactus that is endemic to the subtropical and tropical coastal areas native of the Americas. Its fruits have a purplish color with a small green cladodes. It remains under-exploited in Algeria. This plant prefers high temperate and dry condition areas and can tolerate cold temperatures. This plant is known for its medicinal properties due to its phytochemical compounds. It contains bioactive components with beneficial health effects. The aim of this study was to extract total phenolic compounds with water and identified via LC-MS/MS.

Methodology: the cladodes of *Opuntia stricta* were recollected, cleaned and cut into small pieces, then oven-dried at 45°C. The extraction process was the maceration in pure water for 24 hours then it was filtered and dried via a rotavap and finely identified the different phenolic compounds via LC-MS/MS.

The results of the extract obtained showed that the phenolic compound composition is as follows (Gentisic acid ; Syringic acid ; Vanillic acid ; Vanillin ; trans-ferulic acid) ; with the respective content (10.24 ; 0.499 ; 6.63 ; 0.895 ; 2.23 µg/mg of extract).

Conclusion: *O.stricta* cladodes contain different important molecules such as phenolic acids and flavonoids which can be used as sources of bioactive molecules for pharmacological purposes and cosmetic products in order to their effect such as antioxidant activities and other biological activities.

Keywords: *Opuntia stricta* ; phenolic compounds ; bioactive molecules.

**DEVELOPMENT OF A NOVEL LOW-COST ADSORBENT
CHITOSAN@EDTA@CELLULOSE COMPOSITE TO EFFECTIVELY REMOVE
METHYL ORANGE DYE FROM WASTEWATER: EXPERIMENTAL AND
THEORETICAL INVESTIGATION**

Soukaina El Bourachdi

Laboratory of Engineering, Electrochemistry, Modelling and Environment, Faculty of Sciences Dhar El Mehraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco

Abdelhay El Amri

Laboratory of Advanced Materials and Process Engineering (LAMPE), Faculty of Sciences, Ibn Tofail University, B.P. 133, 14000 Kenitra, Morocco

Ali Raza Ayub

Key Laboratory of Clusters Science of Ministry of Education, School of Chemistry and Chemical Engineering, Beijing Institute of Technology, Beijing, 100081, P. R. China

Fatima Moussaoui

Laboratory of Engineering, Electrochemistry, Modelling and Environment, Faculty of Sciences Dhar El Mehraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco

Amal Lahkimi

Laboratory of Engineering, Electrochemistry, Modelling and Environment, Faculty of Sciences Dhar El Mehraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco

Abstract

Methyl Orange, a toxic and persistent azo dye, poses significant environmental challenges in aquatic ecosystems. This study investigates the efficiency of a novel Chitosan@EDTA@Cellulose composite, synthesized by linking shrimp-derived chitosan and cactus-derived cellulose using EDTA as a linking agent. Comprehensive characterization techniques, including Fourier-transform infrared spectroscopy, scanning electron microscopy, X-ray diffraction, and Brunauer-Emmett-Teller surface area analysis, were employed. Under optimal conditions (pH 5, 50 mg/L dye concentration, 55 min, 0.1 g adsorbent), the composite achieved a maximum adsorption capacity of 55.87 mg/g, significantly outperforming chitosan (7.29 mg/g) and cellulose (5.69 mg/g). Adsorption followed the pseudo-second-order kinetic model and the Langmuir isotherm model, with thermodynamic analysis confirming a spontaneous and endothermic process. Competitive adsorption tests demonstrated more than 90% removal efficiency despite the presence of interfering ions, attributed to the chelating properties of EDTA and the synergistic effect of the composite structure. Reusability tests showed a slight efficiency decline from 97.8% to 81.86% after four cycles. Box-Behnken Design optimization identified adsorbent mass, pH, and dye concentration as key factors in removal efficiency. Density functional theory analysis clarified the functional group interactions driving adsorption. These findings underscore the composite's potential as an effective and eco-friendly adsorbent for Methyl Orange removal.

Keywords: Adsorption, Box-Behnken Design, Cellulose, Chitosan, Density Functional Theory, Response Surface Methodology.

IN VITRO EVALUATION OF THE ANTIFUNGAL POTENTIAL OF *MYRTUS COMMUNIS L.* AGAINST TOXIGENIC MOLDS OF HYDROPONIC BARLEY

Nacéra TADJINE

Aromatic and Medicinal Plants Research Laboratory, Department of Biotechnology, Faculty of Natural and Life Sciences, University of Blida1

ORCID: <https://orcid.org/0000-0002-5209-6468>

Saida MESSGO-MOUMENE

Aromatic and Medicinal Plants Research Laboratory, Department of Biotechnology, Faculty of Natural and Life Sciences, University of Blida1

Abstract

Introduction and Purpose: This study aimed to focus on the evaluation of the antifungal activity of essential oil prepared from *Myrtus communis L.* leaves harvested from the mountains of the Medea region in the north of Algeria.

Materials and Methods: The extracts were prepared by maceration of leaves in sterile distilled water, methanol, chloroform and ethyl acetate. In vitro antifungal activity of the extracts was evaluated at concentrations of 5, 2.5, 1.25 and 0.625 mg mL⁻¹ by the direct contact method on the mycelial growth of toxigenic molds of hydroponic barley namely *Fusarium aveaceanum*, *Fusarium culmorum*, *Alternaria sp.* and *Aspergillus niger* ethyl acetate and Chloroform extracts exhibited pronounced antifungal potential. The inhibition potential varied proportionally with the concentrations.

Discussion and Conclusion: The fungal growth inhibition potentials varied significantly from 15 – 96%. Thus *Myrtus. communis* ethyl acetate extract could be used in the food industry for better phytoprotection of barley hydroponic culture and as a natural conservative of its grains in places of storage.

Keywords: Antifungal activity, *Hordeum vulgare*, Molds, Polyphenolic extracts, *Myrtus communis L.*

**ADVANCEMENTS IN REMOTE SENSING FOR CROP AND SOIL HEALTH
MANAGEMENT IN SUSTAINABLE AGRICULTURE**

Abdul Ahad Ansari

Interdisciplinary Department of Remote Sensing and GIS Applications, Aligarh Muslim
University, Aligarh, U.P., India PIN-202002

Haris Hasan Khan

Interdisciplinary Department of Remote Sensing and GIS Applications, Aligarh Muslim
University, Aligarh, U.P., India PIN-202002

Abstract

Agricultural sustainability is critical to addressing global food demands while ensuring the conservation of natural resources. Monitoring crop health and soil quality in real time is essential for maximizing agricultural productivity and minimizing environmental degradation. The role of remote sensing technologies in crop monitoring and soil health assessment is to enhance soil nutrition and manage environmental impacts by leveraging multispectral and hyperspectral satellite imagery and GIS-based spatial analysis. The study examines essential factors influencing crop growth, such as soil moisture, nutrient content, and early indicators of crop stress, across various agricultural landscapes.

A key focus is employing the Normalized Difference Vegetation Index (NDVI) and Soil-Adjusted Vegetation Index (SAVI) as reliable metrics to assess crop vigour and detect early signs of soil degradation. By integrating these indices with temporal and spatial data, the study provides insights into dynamic crop-soil interactions, allowing for targeted interventions in nutrient management and irrigation practices. Precision agriculture methods informed by these insights, such as site-specific fertilization and efficient water resource allocation, can significantly reduce input usage while optimizing crop yields.

The potential of remote sensing is to support sustainable agricultural practices by offering a high-resolution, cost-effective means of monitoring crop conditions and soil health. This approach helps enhance farm management strategies and aligns with global sustainability goals, addressing the urgent need for food security and environmental resilience.

Keywords: Remote Sensing, Crop Monitoring, Soil Health, Sustainable Agriculture, GIS, NDVI, Environmental Resilience

THE IMPACT OF ADDING WALNUT SHELL POWDERS ON THE MORPHOLOGY, THERMAL PROPERTIES, AND MECHANICAL PERFORMANCE OF POLY(LACTIC ACID) (PLA)

Zineb El Hamri

Team of Innovative Materials and Mechanical Manufacturing Processes, ENSAM, University
Moulay Ismail, B.P. 15290, Al Mansour, Meknes, Morocco

M. Karaoui

Team of Innovative Materials and Mechanical Manufacturing Processes, ENSAM, University
Moulay Ismail, B.P. 15290, Al Mansour, Meknes, Morocco

M. Alami

Team of Innovative Materials and Mechanical Manufacturing Processes, ENSAM, University
Moulay Ismail, B.P. 15290, Al Mansour, Meknes, Morocco

M. Assouag

Team of Innovative Materials and Mechanical Manufacturing Processes, ENSAM, University
Moulay Ismail, B.P. 15290, Al Mansour, Meknes, Morocco

Abstract

Poly(lactic acid) (PLA) is an environmentally friendly material; however, its widespread application is limited by its brittleness, slow crystallization rate, and low heat distortion temperature. To address these shortcomings, this study incorporated walnut shell (WS) powders into PLA. The influence of WS powders on the morphology, as well as the thermal and mechanical properties of PLA, was thoroughly examined. Characterization techniques such as differential scanning calorimetry (DSC), infrared (IR) spectroscopy, polarizing optical microscopy (POM), and various mechanical property tests were employed. The findings revealed that WS powders significantly affected the morphology and improved the thermal and mechanical behavior of PLA. The tensile strength, impact strength, and elongation at break of the PLA/WS composites initially increased with rising WS powder content, peaking at 0.5 wt%, where maximum values of 51.2 MPa, 23.3 MPa, and 19.0% were observed, respectively. Beyond this concentration, these properties declined. Compared to pure PLA, the composites exhibited reduced spherulite grain sizes and the formation of irregular polygonal structures during crystallization. Additionally, the melting, cold crystallization, and glass-transition temperatures of the PLA/WS composites were lower than those of unmodified PLA.

Keywords: Poly(lactic acid), walnut shells, mechanical properties, composites, thermal properties.

References

- [1]: Raquez JM, Habibi Y, Murariu M, et al. Polylactide (PLA)-based nanocomposites. *Prog Polym Sci* 2013; 38: 1504–1542..
- [2]: Cong DV, Hoang T, Giang NV, et al. A novel enzymatic biodegradable route for PLA/EVA blends under agricultural soil of Vietnam. *Mater Sci Eng C* 2012; 32: 558–563.
- [3]: Jonoobi M, Harun J, Mathew AP, et al. Mechanical properties of cellulose nanofiber (CNF) reinforced polylactic acid (PLA) prepared by twin screw extrusion. *Compos Sci Technol* 2010; 70: 1742–1747.

GEOBOTANICAL DATA ON FLORA AND VEGETATION IN SOME AREAS OF ELBASAN

Fjorela MEMA

Department of Biology, Faculty of Natural Sciences, University of Elbasan “Aleksandër Xhuvani”, Elbasan, Albania

Selma MYSLIHAKA

Department of Biology, Faculty of Natural Sciences, University of Elbasan “Aleksandër Xhuvani”, Elbasan, Albania

Peçi NAQELLARI

Department of Biology, Faculty of Natural Sciences, University of Elbasan “Aleksandër Xhuvani”, Elbasan, Albania

Nikoleta KALLAJXHIU

Department of Biology, Faculty of Natural Sciences, University of Elbasan “Aleksandër Xhuvani”, Elbasan, Albania

Silvana TURKU

Department of Biology, Faculty of Natural Sciences, University of Elbasan “Aleksandër Xhuvani”, Elbasan, Albania

Anxhela DAUTI

Department of Biology, Faculty of Natural Sciences, University of Elbasan “Aleksandër Xhuvani”, Elbasan, Albania

Abstract

The Elbasan region, located in central Albania, is characterized by a rich diversity of flora and vegetation, shaped by the interaction of Mediterranean and continental climatic conditions, as well as a varied geological and topographical landscape. However, human activities such as natural resource exploitation and urban development can impact the area’s natural habitats and biodiversity. It is essential to implement measures to minimize these negative effects and maintain the ecological balance of the region.

This study aims to present geobotanical data on the flora and vegetation of selected areas in Elbasan. The research focuses on key strategic zones, including Byshek, Qafë-Kërrabë, Ullishtë, Krastë, Xibrakë-Lugina e Shkumbini, and Elbasan, where field expeditions were conducted. Based on the collected data, floristic spectra were constructed to analyze life forms, floristic elements, and the percentage distribution of plant families in the region.

The findings indicate that the most widespread biological forms are Hemicryptophytes, Phanerophytes, and Therophytes. Regarding floristic elements, the dominant types include Eurymediterranean, Eury-Southeast, Central European, and Eurasian species. The most abundant plant families are *Fabaceae*, *Poaceae*, *Asteraceae*, and *Labiatae*. Additionally, a significant number of medicinal plants were identified, primarily from the *Asteraceae*, *Labiatae*, *Rosaceae*, and *Fabaceae* families.

Keywords: biological forms, floristic diversity, flora, medicinal plants,

QUALITY CONTROL OF ALGERIAN OLIVE OIL

AKRETCHE Soraya

Laboratory of Industrial Process Engineering Sciences (LSGPI). Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

LAMOUDI Lynda

Transfer Phenomena Laboratory. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

BOUSBIA Nabil

Laboratory of Industrial Process Engineering Sciences (LSGPI). Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

KERBOUCHE Lamia

Laboratory of Industrial Process Engineering Sciences (LSGPI). Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

DAHMANI Chaima

Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria

Abstract

Olive oil is a good source of dietary fat in the Mediterranean diet. It is obtained from the fruit of the olive tree (*Olea europaea* L) by mechanical or other means under specific thermal conditions (Criado et al., 2004).

In Algeria, the olive tree has approximately 32 million trees (Bensemmane, 2009) spread over an area of approximately 328,884 hectares (FAOSTAT, 2013), or 34.09% of the national agricultural orchard. Algerian olive growing is mainly found in the northern part of the country, where most orchards (80%) are located in mountainous areas.

The quality of olive oil depends on several factors such as maturity, extraction method, soil, climate, variety and also storage conditions. The evaluation of the quality of olive oil involves a series of important physicochemical parameters including acidity, peroxide index, saponification index are part of (Gharbi et al., 2015). The quality of olive oil is also characterized by its particular composition in saturated and unsaturated fatty acids, as well as its minor compounds which belong to the unsaponifiable fraction (tocopherols, chlorophylls, etc.) (Mezghache et al., 2010).

Olive oil is called green gold because of the beneficial and nutritional effects it presents such as preventing aging, strengthening the immune system, and preventing cardiovascular diseases (Ghedira, 2008).

This work was carried out with the aim of carrying out a study on the physicochemical characteristics of olive oil from the Bouira region, as well as on its fatty acid profile by gas chromatography.

The results obtained from the analyzes carried out show that the values of refractive index, saponification index and specific extinction at 232 nm comply with the limits set by the Codex Alimentarius (2017) for extra virgin olive oil.

It should be noted that the sample studied presented a K270 coefficient and a relative density slightly higher than the limits established by the Codex Alimentarius (2017). In addition, an acidity of around 0.64 and a peroxide index of 14 which are an excellent indicator of good quality of olive oil.

The results of the present study also show that the fatty acid composition complies with the standards; Oleic acid is the dominant fatty acid in the oil, with a proportion greater than 66% followed by palmitic and linoleic acid.

At the end of this study, we note that the olive oil analyzed has a high content of oleic acid as well as a richness in short-chain fatty acids, which allows it to be classified among extra virgin olive oils.

Key words: Olive oil, quality, characterization, physicochemical indices, fatty acid composition.

STUDY OF THE QUALITY OF A DAIRY PRODUCT ENRICHED WITH POLLEN

AKRETCHE Soraya

Laboratory of Industrial Process Engineering Sciences (LSGPI). Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

BOUSBIA Nabil

Laboratory of Industrial Process Engineering Sciences (LSGPI). Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

LAMOUDI Lynda

Transfer Phenomena Laboratory. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

KERBOUCHE Lamia

Laboratory of Industrial Process Engineering Sciences (LSGPI). Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

DJILI Manel

Transfer Phenomena Laboratory. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

KACI Ismahan

Transfer Phenomena Laboratory. Faculty of Mechanical Engineering and Process Engineering, University of Science and Technology Houari Boumediene (U.S.T.H.B.), P.O. Box 32 El Alia, 16111 Bab Ezzouar, Algeria.

Abstract

Milk and dairy products occupy an essential place in the daily diet by their taste, nature, diversity and particularly their great nutritional value or even their richness in proteins, calcium, vitamins, trace elements and minerals, in particular yogurt which is one of the fastest growing segments in the dairy sector and cheese is no longer the leading product.

It is often consumed as a fresh dessert, light, pleasant to the eye, to the taste and a real alternative to milk in lactose intolerant individuals, therefore it is suitable for all age groups.

The products of the hive (honey, pollen, royal jelly...) are also appreciated in terms of food, medicine and industry, their consumption is gradually increasing thanks to several beneficial effects that they confer on the health environment.

Among these products; bee pollen, it is first of all a natural product and a remarkable food in api therapy. These are microscopic seeds contained in the anthers of the stamens. Its colors vary from pale yellow to black, passing through all shades of brown and red. It is considered to be a source of antioxidants, such as polyphenols, flavonoids and carotenoids. It is rich in protein, lipid, mineral salts. A good generator of carbohydrates, mainly fructose and sucrose, and essential amino acids that the body cannot synthesize.

In this regard, a valuation of the pollen grains incorporated in the yogurt, as well as a physicochemical, microbiological and sensory characterization of the yoghurt produced were carried out.

At the end of this work, the aim of which is to study the effect of incorporating pollen into stirred yoghurt, given its high nutritional value, it has been observed that:

- For pollen: has a water content of 4.41%, a pH of 4.73, which can be as a factor that slows down the growth and development of some microorganisms, an ash content of 2.6% , and in a degree of brix of 5.55%, which makes it a natural material of interesting varied composition. On the other hand, the microbiological analyzes (search for yeasts and molds, salmonella, Staphylococcus aureus, enumeration of total and fecal coliforms, total germs), reflected the compliance of the pollen used with the required standards, which provides information on its good hygienic quality.

- The formulated yogurt has new organoleptic characteristics: a flavor and a texture appreciated with a nutritional advantage of pollen because it is rich in amino acids, vitamins, minerals, carotenoids and flavonoids. The yogurt is characterized by an acceptable acidity (88 ° D), a pH evaluated at 4.56 and a total dry extract of 25.73%. In addition, it is healthy and has a satisfactory microbiological quality.

- The sensory analysis of the three samples revealed that the tasting panel appreciated the pollen ground yogurt the most, these results clearly reveal the performance of the organoleptic characteristics developed due to the addition of pollen within the product finished.

Key words: formulation, pollen, sensory analyzes, physicochemical analyzes, microbiological analysis

SURVEY ON FERTILIZATION PRACTICES AND NUTRIENT MANAGEMENT IN VALENCIA LATE CITRUS ORCHARDS: A CASE STUDY FROM THE GHARB REGION (NORTHEASTERN), MOROCCO

BRITAL Rania

National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University Ibn Toufail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0002-9971-3790>

Abail Zhor

National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0001-7819-9400>

MRABTI Imane

National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University Ibn Toufail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-6580-4291>

EL bahja Fatima

National Institute for Agricultural Research, Regional Center for Agricultural Research of Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University Ibn Toufail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0007-0503-6327>

IBRIZ Mohammed

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University Ibn Toufail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-9024-912X>

Abstract

The cultivation of *Valencia Late* citrus represents a major economic challenge in many producing regions. However, fertilization management remains largely empirical, leading to high costs and significant environmental impacts. This study aims to analyze current fertilization practices among farmers to identify key gaps and propose optimization strategies. A survey was conducted among 20 producers in the Gharb region, located in northeastern Morocco, covering the provinces of Kenitra, Sidi Slimane, and Sidi Kacem. The study examined the types of fertilizers used, application frequencies, and methods for assessing orchard nutritional needs. The results reveal significant discrepancies between observed practices and agronomic recommendations, highlighting the need for improved fertilization management. Furthermore, only 25% of the farms regularly perform soil analyses, contributing to chronic nutrient imbalances. Implementing soil and plant tissue analysis-based approaches could enhance nutrient application efficiency, mitigate nutrient imbalances, and reduce

environmental impacts. These findings underscore the importance of strengthening farmer training and support to promote a more rational and sustainable fertilization strategy for *Valencia Late* orchards.

Keywords: *Valencia Late, fertilization, sustainability, citrus nutrition, soil analysis.*

IMPROVEMENT OF DNA EXTRACTION FOR THE PYRIFORM SCALE INSECT, *PROTOPULVINARIA PYRIFORMIS*, FOR MOLECULAR IDENTIFICATION OF CITRUS PESTS IN MOROCCO USING DNA BARCODING

MRABTI Imane

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-658-4291>

GRIJJA Hassan

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0002-3905-8173>

BRITAL Rania

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0002-9971-3790>

BRHADDA Najiba

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0003-1994-0310>

ZIRI Rabea

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-5223-8235>

AFECHTAL Mohamed

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra,
Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0003-4409-1226>

Abstract

DNA extraction from scale insects is particularly challenging due to their small size and waxy coating. In this regard, a study was conducted between the summer and autumn of 2024 in the north-west of Morocco to assess the effectiveness of ten DNA extraction methods on scale insects. The target species of this study, *Protopulvinaria pyriformis*, is a major pest of citrus crops in Morocco. The main objective of this research was to identify extraction methods that are quick, cost-effective, and environmentally friendly.

The results showed that fresh samples of *P. pyriformis* provided the best DNA yields. Among the methods tested, the use of NaCl (5M) produced the highest yield, while the Kac and SDS

methods yielded moderate results, all while being cost-effective and environmentally sustainable.

The results of this study provide valuable recommendations for selecting the optimal DNA extraction method for *P. pyriformis*. These suggestions can help improve the identification and management of this pest, thereby optimizing pest control and monitoring strategies in agricultural settings.

Keywords: *Protopulvinaria pyriformis*, DNA extraction, citrus, cost-effectiveness, Morocco.

PRECISE IDENTIFICATION OF THE COTTONY CUSHION SCALE, *ICERYA PURCHASI* IN MOROCCO THROUGH THE ADVANCED APPLICATION OF DNA BARCODING

MRABTI Imane

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-658-4291>

GRIJJA Hassan

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0002-3905-8173>

BENZAHERA Hayat

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0003-9708-5468>

AIT BADDOU Sanae

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra, Kenitra, Morocco
Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0009-0002-1128-9316>

BRHADDA Najiba

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0003-1994-0310>

AFECHTAL Mohamed

National Institute for Agricultural Research, Regional Center for Agricultural Research of
Kenitra,
Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0003-4409-1226>

ZIRI Rabea

Faculty of Sciences, Laboratory of Plant, Animal, and Agro-Industry Productions, University
Ibn Tofail, Kenitra, Morocco
ORCID ID: <https://orcid.org/0000-0002-5223-8235>

Abstract

DNA extraction from scale insects is particularly challenging due to their small size and waxy coating. In this regard, a study was conducted between the summer and autumn of 2024 in the north-west of Morocco to assess the effectiveness of ten DNA extraction methods on scale

insects. The target species of this study, *Icerya purchasi*, is a major pest of citrus crops in Morocco. The main objective of this research was to identify extraction methods that are quick, cost-effective, and environmentally friendly.

The results showed that fresh samples of *I. purchasi* provided the best DNA yields. Among the methods tested, the use of NaCl (5M) produced the highest yield, while the Kac and SDS methods yielded moderate results, all while being cost-effective and environmentally sustainable.

The results of this study provide valuable recommendations for selecting the optimal DNA extraction method for *I. purchasi*. These suggestions can help improve the identification and management of this pest, thereby optimizing pest control and monitoring strategies in agricultural settings.

Keywords: *Icerya purchasi*, DNA extraction, citrus, cost-effectiveness, Morocco.

MORPHOLOGICAL DIVERSITY AND STYLAR POLYMORPHISM IN *NARCISSUS* SPECIES (AMARYLLIDACEAE) FROM NORTHERN ALGERIA**Ph.D. Hana ZAKKOUMI**

University of Sciences and Technology Houari Boumediene (USTHB), Faculty of Biological Sciences

Yazid MAHDAOUI

University of Algiers 1 Benyoucef Benkhedda, Faculty of Sciences

Houria HADJ-ARAB

University of Sciences and Technology Houari Boumediene (USTHB), Faculty of Biological Sciences

Abstract

The *Narcissus* genus, belonging to the Amaryllidaceae family, is known for its horticultural diversity, shaped by frequent hybridization and polyploidization events. These evolutionary processes have made of *Narcissus* an important model for studying evolutionary patterns, but have also complicated its taxonomy due to the extensive morphological and sexual variation observed within the genus. This study investigated the morphological diversity of 48 populations from five *Narcissus* species in northern Algeria, by analyzing 45 vegetative and floral traits through biometric analyses. The results showed that *N. serotinus* and *N. elegans* exhibit stylar monomorphism, while *N. papyraceus* and *N. tazetta* display stylar dimorphism, featuring two sexual morphs: L-morph (long style, short stamens) and S-morph (short style, long stamens). The results also showed that *N. cantabricus* is characterized by a homomorphic long-styled floral type. On the other hand, morphometric analysis grouped the five species into three distinct phenotypic clusters, revealing strong morphological affinities and significant polymorphism. These findings suggest close genetic relationships between *N. tazetta* and *N. papyraceus*, as well as between *N. elegans* and *N. serotinus*. The study highlights a strong correlation between floral traits and morphological variation within the *Narcissus* genus, which offers valuable insights into the evolutionary adaptations driving floral morphology and enhances our understanding of the relationships among species in this genus.

Keywords: *Narcissus*, morphological variation, interspecific diversity, stylar polymorphism.

INTRODUCTION

The genus *Narcissus*, belonging to the Amaryllidaceae family, is known for its remarkable morphological diversity, driven by frequent hybridization and polyploidization events (Fernandes, 1968). These evolutionary processes have made of *Narcissus* a valuable model for studying evolutionary patterns, the same processes that have also complicated its taxonomy due to the extensive morphological variation observed within the genus. With approximately 100 species (Breiterová et al., 2020), *Narcissus* is the most diverse genus in its clade, and its taxonomy remains controversial. It is subdivided into ten sections, grouped into two major lineages corresponding to the subgenera *Narcissus* and *Hermione* (Blanchard, 1990; Fernandes, 1968; Marques, Aguilar, Martins-Louçao, Moharrek, & Feliner, 2017). This diversity is closely linked to its unique reproductive system, characterized by heterostyly, a form of sexual

polymorphism involving two or three distinct floral morphs with reciprocal positioning of the reproductive organs (Spencer C.H. Barrett & Harder, 2005).

Heterostyly (a key feature of *Narcissus*) and other style polymorphisms, previously known to be present in 28 phylogenetically diverse angiosperm families, including Linaceae, Oxalidaceae, and Amaryllidaceae (S. C.H. Barrett & Shore, 2008); have recently been expanded to include a total of 247 genera, belonging to 34 families (Simón-Porcar et al., 2024). Styler polymorphism, a transitional stage towards heterostyly, has been widely used as a model system to study the origin and maintenance of variability within populations (Spencer C.H. Barrett, Jesson, & Baker, 2000; Webb & Lloyd, 1986). Various systems of styler polymorphism, such as styler dimorphism, distyly, and tristily, have evolved independently from ancestral styler monomorphism in numerous genera, including *Turnera*, *Fagopyrum*, *Linum*, and *Primula* (Kappel, Huu, & Lenhard, 2017), but particularly in *Narcissus* (Graham & Barrett, 2004). Heterostylous populations often exhibit reciprocal herkogamy, a phenomenon where reproductive organs are reciprocally arranged (Webb & Lloyd, 1986). In *Narcissus*, styler dimorphism is common, differing from distyly by less pronounced reciprocity in stamen positioning (Charlesworth, 1992; Darwin, 1897) and the absence of pollen size variation between morphs (Baker, Thompson, & Barrett, 2000).

Narcissus species are distinguished by their morphological diversity, particularly in the positioning of reproductive organs and perianth shape (Dhiman, Kumar, Parkash, Gautam, & Singh, 2019). Traits such as the size, shape, and color of the corona are key to the classification of *Narcissus* cultivars (Diaz Lifante & Andres Camacho, 2007; Ma et al., 2023). This diversity makes *Narcissus* an unparalleled model for studying floral morphology and the evolution of styler polymorphism, offering valuable insights into the mechanisms driving such variability (Spencer C.H. Barrett & Harder, 2005; Graham & Barrett, 2004). Notably, *Narcissus* is the only taxon that exhibits all known types of styler variations, with an impressive range of floral morphology (Spencer C.H. Barrett & Harder, 2005). This unique combination of traits emphasizes the importance of *Narcissus* in understanding the evolutionary dynamics of floral diversity and reproductive systems. In this present study, we aimed at the investigation of the morphological and sexual diversity in populations of the *Narcissus* species present in Algeria.

METHODOLOGY

This study investigated the morphological diversity of five *Narcissus* species in northern Algeria. A comprehensive biometric analysis was conducted to evaluate phenotypic variation across these populations.

The research focused on *Narcissus* species present in the Algerian flora, with preliminary taxonomic identifications guided by key historical floras of the region, including works by Battandier & Trabut (1895), Desfontaines (1789), Maire (1952) and Quézel & Santa (1962).

Study of morphological variability and intra- and interspecific relationships

The first phase of the study focused on evaluating intra- and interspecific variability by examining a comprehensive set of morphological traits. A total of 44 qualitative and quantitative traits were measured, including 10 vegetative, 6 inflorescence, and 28 floral traits (Table 1). Measurements were conducted on an average of 30 individuals per population (41 populations), with a total of 1172 individuals, ensuring that all sampled flowers were at the same developmental stage.

To analyze the data, multivariate statistical methods were employed using R software v. 3.4, with the packages missMDA, factoextra, and FactoMineR. Principal Component Analysis (PCA) was performed based on the correlation matrix between variables.

Table 1. Measured characteristics of studied organs in *Narcissus* species

Organ / Characteristic	Abbreviation
Vegetative traits	
Number of bulbs	Nb
Length of the bulb (mm)	Lb
Width of the bulb (mm)	lb
Number of sheath layers	Ng
Length of the sheath (mm)	Lghf
Width of the sheath (mm)	lghf
Length of the floral stem (from sheath to base of spathe) (mm)	Lhf
Width of the floral stem (mm)	lhf
Length of the outermost leaf blade (mm)	Lf
Width of the leaf blade at its widest point (mm)	lf
Inflorescence	
Spathe color: whitish to pale brown, brown to dark brown	Csp
Spathe: scarious, non-scarious	Tsp
Total length of the spathe (mm)	Lsp
Length of the spathe enclosing the inflorescence (mm)	Lesp
Width of the spathe at its midpoint (mm)	lsp
Number of flowers	Nfl
Flower	
Tepal color: white, light yellow, dark yellow	Cf
Corona color: white, yellow, orange, or saffron	Cp
Shape of the floral tube: narrow, cylindrical, conical	Fth
Shape of the corona: circular, circular to triangular, triangular	Crp
Corona margin: toothed, entire, truncated, sinuate, retuse, crenate	Lbp
Length from the base of the ovary to the tip of the outer tepal (mm)	LTf
Length of the floral pedicel (from insertion point to the base of the ovary) (mm)	Lpd
Length of the ovary (mm)	Lo
Width of the ovary (mm)	lo
Length of the floral tube (from the top of the ovary to the insertion point of the tepals) (mm)	Lth
Width of the floral tube (mm)	lth
Height of the corona (mm)	Hp
Diameter of the corona (mm)	Dp
Length of the outer tepal (mm)	Lte
Width of the outer tepal (mm)	lte
Length of the mucro of the outer tepal (mm)	Lbte
Length of the inner tepal (mm)	Lti
Width of the inner tepal (mm)	lti
Length of the external stamen (including anther and filament) (mm)	LES
Length of the external stamen anther (mm)	Las
Height of the external stamen filament (mm)	Hfs
Distance from the insertion point of the external stamen filament to the base of the floral tube (mm)	Hpfs
Length of the internal stamen (mm)	LEI
Length of the internal stamen anther (mm)	Lai
Height of the internal stamen filament (mm)	Hfi
Distance from the insertion point of the internal stamen filament to the base of the floral tube (mm)	Hpfi
Length of the style (including stigma) (mm)	Ls
Total number of ovules in the ovary	No

Morph determination

Morph determination was conducted for all the 48 populations (3758 individuals), with over 100 individuals analyzed per population, except for small populations. The stigma position relative to the two stamen whorls was determined through direct visual examination of longitudinal flower sections.

RESULTS AND DISCUSSION

Study of morphological variability and interspecific relationships

The results identified two primary sets of correlated traits: vegetative traits were predominantly associated with axis 1, while floral traits were primarily linked to axis 2 (Figure 1). Two distinct groups of individuals emerged from the analysis: *N. cantabricus* was positioned in the positive region of axis 2, whereas all other species were distributed along axis 1.




Figure 1. Principal component analysis (PCA) of the five studied taxa within the genus *Narcissus*.

(a) Projection of the 44 morphological variables onto the correlation circle. (b) Distribution of 1172 individuals (representing 41 populations) defined by PCA axes 1 and 2.

N. cantabricus is characterized by a uniquely wide obconical floral tube, a large corona, long stamens, and a prominently exerted long style. Its presence in Algeria was historically debated until it was synonymized with *N. bulbocodium* by Quézel & Santa.

Partial analysis of groups *N. tazetta*-*N. papyraceus* and *N. elegans*-*N. serotinus*

To investigate the divergences between phenotypic groups, we excluded *N. cantabricus* and conducted a second Principal Component Analysis (PCA). This analysis clearly separated two phenotypic groups along axis 1: the *tazetta*-*papyraceus* group and the *elegans*-*serotinus* group (Figure 2). The two groups primarily differed in traits such as leaf and scape width, bulb size, spathe size, and the dimensions of floral organs. Despite significant polymorphism, individuals within the *elegans*-*serotinus* group displayed strong morphological similarities. In contrast, *N. tazetta* and *N. papyraceus* were distinguished by differences in floral tube length, corona color, and overall flower size.




Figure 2. Principal Component Analysis (PCA) of four species within the genus *Narcissus*. (a) Projection of 41 morphological variables onto the correlation circle. (b) Distribution of 1,129 individuals (representing 43 populations) defined by PCA axes 1 and 2.

Partial analysis of phenotypic group *N. elegans*-*N. serotinus*

In this analysis, we concentrated on the phenotypic group comprising *N. elegans* and *N. serotinus*, which demonstrated strong morphological similarities (Figure 3). We incorporated an additional trait—the number of leaves (Nf) per individual—that was not accounted for in the analysis of the other species.

Individuals primarily differed in filament insertion distance and tepal width, which were strongly correlated. Notably, individuals in the red zone exhibited intermediate characteristics of both taxa, differing in flower number and leaf presence, suggesting potential interspecific hybridization.




Figure 4. Principal Component Analysis (PCA) of the *N. tazetta* - *N. papyraceus* group. (a) Projection of 36 morphological variables onto the correlation circle. (b) Distribution of 562 individuals (representing 21 populations) defined by PCA axes 1 and 2.

The two species were primarily distinguished by differences in bulb size, floral stem length, leaf dimensions, floral part measurements, and corona color. Initially, *N. papyraceus* was classified as a subspecies of *N. tazetta* due to their morphological similarities (Maire, 1952). However, based on current taxonomic databases, *N. papyraceus* is now recognized as a distinct species, referred to as *N. pachybolbus*.

Analysis of Styler Polymorphism

Longitudinal sections of *N. tazetta* flowers revealed significant variation in style length, as well as differences in the positioning of internal and external anthers, which could either overlap or remain separate. Several sub-morphs were identified, including the SS sub-morph and various sub-morphs within the L morph (Zakkoumi, Hadj-Arab, & Amirouche, 2023).

Narcissus tazetta and *N. papyraceus* exhibit styler dimorphism, with populations displaying monomorphic, isoplethic, and anisoplethic characteristics (Figure 5). These findings are consistent with previous studies on *N. tazetta* (Arroyo & Dafni, 1995) and *N. papyraceus* (Spencer C. H. Barrett, Lloyd, & Arroyo, 1996).




Figure 5. Morph distribution within the phenotypic group *N. tazetta* – *N. papyraceus*. Twenty-three populations of *N. tazetta* and Three of *N. papyraceus* are illustrated.

In contrast, within the *N. elegans*–*N. serotinus* phenotypic group, stilar monomorphism predominates across all 20 populations, indicating that these species are monomorphic and exhibit a single floral type. These results align with existing literature on both *N. elegans* (Graham & Barrett, 2004) and *N. serotinus* (Spencer C. H. Barrett et al., 1996; Santos-Gally, Gonzalez-Voyer, & Arroyo, 2013). However, a study by Santos-Gally et al., 2013, described *N. elegans* as a dimorphic species, highlighting a discrepancy in the characterization of its reproductive morphology.

Regarding *N. cantabricus*, the results revealed that this species exhibits longistylar monomorphism, which is consistent with previously reported data in the literature (Barrett, Lloyd, & Arroyo, 1996; Santos-Gally, Gonzalez-Voyer, & Arroyo, 2013).

CONCLUSION

Our study explored two critical dimensions of the *Narcissus* genus: morphology and stilar polymorphism. Morphological analyses grouped the five studied species into three distinct phenotypic clusters, revealing strong morphological affinities and significant polymorphism. These findings suggest close phylogenetic relationships, with *N. tazetta* and *N. papyraceus* forming one cluster, and *N. elegans* and *N. serotinus* forming another. *N. cantabricus* formed a unique group and was characterized by a homomorphic long-styled floral type. The analysis of stilar polymorphism further highlighted distinct reproductive strategies, in which: *N. tazetta* and *N. papyraceus* exhibited stilar dimorphism, featuring two sexual morphs (L-morph and S-morph), while *N. elegans*, *N. serotinus*, and *N. cantabricus* displayed stilar monomorphism. Together, these results provide profound insights into the evolutionary relationships, genetic affinities, and reproductive adaptations within the *Narcissus* genus, enhancing our understanding of its diversity and evolutionary dynamics and providing a robust dataset for future analyses of morphological and taxonomic variability.

REFERENCES

- Arroyo, J., & Dafni, A. (1995). Variations in habitat, season, flower traits and pollinators in dimorphic *Narcissus tazetta* L. (Amaryllidaceae) in Palestine. *New Phytologist*, 129(1), 135–145. <https://doi.org/10.1111/j.1469-8137.1995.tb03017.x>
- Baker, A. M., Thompson, J. D., & Barrett, S. C. H. (2000). Evolution and maintenance of stigma-height dimorphism in *Narcissus*. I. Floral variation and style-morph ratios. *Heredity*, 84(5), 502–513. <https://doi.org/10.1046/j.1365-2540.2000.00651.x>
- Barrett, S. C.H., & Shore, J. S. (2008). New insights on heterostyly: Comparative biology, ecology and genetics. In *Self-Incompatibility in Flowering Plants: Evolution, Diversity, and Mechanisms* (pp. 3–32). Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-540-68486-2_1
- Barrett, Spencer C. H., Lloyd, D. G., & Arroyo, J. (1996). Styler Polymorphisms and the Evolution of Heterostyly in *Narcissus* (Amaryllidaceae). https://doi.org/10.1007/978-1-4613-1165-2_13
- Barrett, Spencer C.H., & Harder, L. D. (2005). The evolution of polymorphic sexual systems in daffodils (*Narcissus*). *New Phytologist*, 165(1), 45–53. <https://doi.org/10.1111/j.1469-8137.2004.01183.x>
- Barrett, Spencer C.H., Jesson, L. K., & Baker, A. M. (2000). The evolution and function of styler polymorphisms in flowering plants. *Annals of Botany*, 85(SUPPL. A), 253–265. <https://doi.org/10.1006/anbo.1999.1067>
- Battandier, J. A., & Trabut, L. (1895). Flore de l'Algérie et catalogue des plantes du Maroc. *Alger, Jourdan*, 1(508), 1888–1890.
- Blanchard, J. W. (1990). *Narcissus. A guide to wild daffodils*. Alpine Garden Society. Retrieved from <https://books.google.dz/books?id=4aBFAAAAYAAJ>
- Breiterová, K., Koutová, D., Maříková, J., Havelek, R., Kuneš, J., Majorošová, M., Cahlíková, L. (2020). Amaryllidaceae alkaloids of different structural types from *Narcissus* L. cv. Professor Einstein and their cytotoxic activity. *Plants*, 9(2), 137. <https://doi.org/10.3390/plants9020137>
- Charlesworth, D. (1992). Evolution and function of heterostyly. *Trends in Ecology & Evolution*, 7(12), 428–429. [https://doi.org/10.1016/0169-5347\(92\)90036-b](https://doi.org/10.1016/0169-5347(92)90036-b)
- Darwin, C. (1897). The different forms of flowers on plants of the same Species. In *D. Appleton*. Cambridge: John Murray. <https://doi.org/10.1017/CBO9780511731419>
- Desfontaines, R. L. (1789). Flora Atlantica, sive, Historia plantarum quae in Atlante, agro Tunetano et Algeriensi crescunt. In *Flora Atlantica, sive, Historia plantarum quae in Atlante, agro Tunetano et Algeriensi crescunt*. Apud LG Desgranges. <https://doi.org/10.5962/bhl.title.6364>
- Dhiman, M., Kumar, S., Parkash, C., Gautam, N., & Singh, R. (2019). Genetic diversity and principal component analysis based on vegetative, floral and bulbous traits in *Narcissus* (*Narcissus pseudonarcissus* L.). *Int J Chem Stud*, 7, 724–729.
- Diaz Lifante, Z., & Andres Camacho, C. (2007). Morphological variation of *Narcissus serotinus* L. s.l. (Amaryllidaceae) in the Iberian Peninsula. *Botanical Journal of the Linnean Society*, 154(2), 237–257. <https://doi.org/10.1111/j.1095-8339.2007.00653.x>
- Fernandes, A. (1968). Keys to the identification of native and naturalised taxa of the genus *Narcissus* L. In *R.H.S. Daffodil Year Book*. Royal Horticultural Society.

- Graham, S. W., & Barrett, S. C. H. (2004). Phylogenetic reconstruction of the evolution of stylar polymorphisms in *Narcissus* (Amaryllidaceae). *American Journal of Botany*, *91*(7), 1007–1021. <https://doi.org/10.3732/ajb.91.7.1007>
- Kappel, C., Huu, C. N. ve Lenhard, M. (2017). A short story gets longer: Recent insights into the molecular basis of heterostyly. *Journal of Experimental Botany*, *68*(21–22), 5719–5730. doi:10.1093/jxb/erx387
- Koopowitz, H., Howe, M., & Christenhusz, M. J. M. (2017). Nomenclatural notes on some autumn flowering daffodils (*Narcissus*, Amaryllidaceae). *Phytotaxa*, *297*(2), 157–167. <https://doi.org/10.11646/phytotaxa.297.2.3>
- Ma, Y., Hu, X., Fan, K., Zhang, N., Shang, L., Deng, Y., Jiang, Z. (2023). Emergence of Corona Is Independent of the Four Whorls of Floral Organs in *Narcissus tazetta*. *Plants*, *12*(7), 1458.
- Maire, R. (1952). Flore de l’Afrique du Nord. In Paul Lechevalier. Paris (France) Lechevalier. Retrieved from <http://medcontent.metapress.com/index/A65RM03P4874243N.pdf>
- Marques, I., Aguilar, J. F., Martins-Louçao, M. A., Moharrek, F., & Feliner, G. N. (2017). A three-genome five-gene comprehensive phylogeny of the bulbous genus *Narcissus* (Amaryllidaceae) challenges current classifications and reveals multiple hybridization events. *Taxon*, *66*(4), 832–854. <https://doi.org/10.12705/664.3>
- Marques, I., Feliner, G. N., Draper Munt, D., Martins-Louçã, M. A., & Aguilar, J. F. (2010). Unraveling cryptic reticulate relationships and the origin of orphan hybrid disjunct populations in *Narcissus*. *Evolution*, *64*(8), 2353–2368. <https://doi.org/10.1111/j.1558-5646.2010.00983.x>
- Quézel, P., & Santa, S. (1962). *Nouvelle flore de l’Algérie et des régions désertiques méridionales*.
- Santos-Gally, R., Gonzalez-Voyer, A., & Arroyo, J. (2013). Deconstructing Heterostyly: The Evolutionary Role Of Incompatibility System, Pollinators, And Floral Architecture. *Evolution*, *67*(7), 2072–2082. <https://doi.org/10.1111/evo.12087>
- Simón-Porcar, V., Escudero, M., Santos-Gally, R., Sauquet, H., Schönenberger, J., Johnson, S. D., & Arroyo, J. (2024). Convergent evolutionary patterns of heterostyly across angiosperms support the pollination-precision hypothesis. *Nature Communications*, *15*(1). <https://doi.org/10.1038/s41467-024-45118-0>
- Webb, C. J., & Lloyd, D. G. (1986). The avoidance of interference between the presentation of pollen and stigmas in angiosperms II. herkogamy. *New Zealand Journal of Botany*, *24*(1), 163–178. <https://doi.org/10.1080/0028825X.1986.10409726>
- Zakkoumi, H., Hadj-Arab, H., & Amirouche, R. (2023). Patterns of style polymorphism in natural populations of *Narcissus tazetta* L. *Brazilian Journal of Botany*. <https://doi.org/10.1007/s40415-023-00948-y>

**INNOVATIVE BAKING: UTILIZING KAMIAS AND BANANA-ENRICHED FLOUR
IN CAKE RECIPES INSPIRED BY LEMON CAKE**

Famela A. Foyo

Cavite State University, Philippines

ORCID ID: <https://orcid.org/0009-0008-9089-9529>

Bernadette I. Fillartos

Cavite State University, Philippines

ORCID ID: <https://orcid.org/0009-0008-5996-7693>

Cristine Angeli T. Mangampat

Cavite State University, Philippines

ORCID ID: <https://orcid.org/0009-0004-6829-6706>

Jane M. Balubar

Cavite State University, Philippines

ORCID ID: <https://orcid.org/0009-0000-7116-4465>

Klenton R. Torrevillas, MIHM

Cavite State University, Philippines

ORCID ID: <https://orcid.org/0009-0003-6588-7861>

ABSTRACT

The demand for fruit-fortified pastry products has been increasing due to its various utilization for product development research. Particularly, bananas have a significant amount of utilization due to its supply locally and the ample nutrients it provides. However, an aspect that remains relatively unexplored is the utilization of kamias aside from being a souring agent on viand.

This study focused on the enrichment of cake recipes utilizing banana and kamias as alternatives to lemon cake recipe. Involving five statements of the problems: enrichment, determining significant difference between 3 formulations, consumer acceptability, physico-chem analysis and sustainable packaging.

The researchers made three formulations with different percentages of dehydrated powdered banana and were evaluated by 10 trained panels for sensory evaluation.

The preferred formulation underwent physico-chem analysis, specifically examining the pH level and nutrition test. Nutrition testing was recorded that it contains adequate amounts of Vitamin A, Vitamin B6 and Iron, the pH testing yielded varying results as it was overseen through 3 trials yielding an average pH level score of 5.4, an indication that it is slightly acidic.

The utilization of banana (lakatan) peel through traditional paper making process underwent consumer acceptability online survey specifically assessing design resulting in moderately like and overall acceptability yielding a result of very much like.

In conclusion, this study provides innovative products using raw materials which are not common in baking as an alternative.

Keywords: Alternative, Enrichment, Kamias, Banana, Dehydrated, Powderized, Sustainable packaging, Innovative.

INTRODUCTION

Cake is often served as a celebratory dish on ceremonial occasions, such as weddings, anniversaries, and birthdays. There are countless cake recipes however, the goal is always the same: to create great cake recipes through a delicate balance of its ingredients - making sure they have the strength to hold the recipe together, but still create a tender, moist and flavorful cake. Cake is a form of sweet dessert that is typically baked. In its oldest forms, cakes were modifications of breads, but cakes now cover a wide range of preparations that can be simple or elaborate, and share features with other desserts. (Ben-Noun,2019).

Fruits can add a pop of color, flavor, and texture to snack and bakery products, but with consumers being more mindful of health and wellness, the use of fruits has expanded beyond the traditional uses. (Friedberg,2024)

Lakatan' banana is amongst the most important banana varieties in the Philippines. This variety of bananas is widely known and cultivated due to its good sensory qualities and potential for the export market. (Sampiano,Durban,2022)

Kamias is a small tree, growing 5 to 12 meters high. Leaves are pinnate, 20 to 60 centimeters long, with hairy rachis and leaflets. Panicles growing from the trunk and larger branches are hairy, 15 centimeters long or less. Flowers are about 1.5 centimeters long, and slightly fragrant. Fruit is green and edible, about 4 centimeters long, subcylindric (subcylindrical), or with 5 obscure, broad, rounded, longitudinal lobes. Averrhoa bilimbi (bilimbi, cucumber tree, or tree sorrel) is a fruit-bearing tree of the genus Averrhoa, family Oxalidaceae (Oxalidaceae). It is believed to a native to Maluku Islands of Indonesia but has naturalized and is common throughout Southeast Asia. (Stuart Jr., 2024)

Food waste has been a global and national problem in the Philippines that occurs at any given time in the retail or household level (Cabilzo,2023)

Enormous amounts of fruit waste is generated during their production life-cycle due to the inedible portion and perishable nature, which become a considerable burden to the environment. (Leong, Chang, 2022)

Dehydrated powdered Banana combined with flour as enrichment to cake ingredients with Kamias glaze for its sour and sweet taste as an innovation to Lemon cake, those fruits are known for their nutrients and health benefits. Researchers provided an enriching cake recipe that can be used as an alternative to lemon cake that also has a nutritious and refreshing taste that the consumers can try and enjoy by using ingredients that are affordable and that can be found in backyard or even on local markets.

STATEMENT OF THE PROBLEM

This study aims to develop and utilize Kamias and dehydrated powdered Banana enriched flour as enhancement for the ingredients on cake recipes inspired by Lemon Cake.

In addition to this, it aims to enhance the ingredients, stored, utilized for a long time and lengthen the shelf life of Lakatan, which is prone to overripe quickly.

1. to enhance cake recipes using dehydrated powdered Banana combined to flour as enrichment with Kamias glaze in terms of:

- 1.1 appropriate percentage per ingredients
- 1.2 technology support; (Production flow), and
- 1.3 financial

2. to determine if there are significant differences between the three formulations of the Kamias Banana Cake using sensory evaluation, evaluated by ten (10) trained sensory evaluator

- 2.1 Appearance (Color, Size, Shape)
- 2.2 Smell/Aroma/Odor (Creamy, Buttery, Little of Banana)
- 2.3 Texture (Crumbly)
- 2.4 Mouthfeel (Softness)
- 2.5 Taste (Sweetness, Sourness)

3. to determine the acceptability of the Kamias Banana Cake with Kamias Glaze to consumer acceptability test; (50 Hospitality Management students from first year to third year as respondents)

4. to determine the Physico-chem of Kamias Banana Cake with Kamias glaze. (pH/ degree brix/ proximate analysis); and

5. to determine if the sustainable packaging is appropriate for the cake .

RESEARCH DESIGN

This study used Experimental Research Design for enhancing dehydrated powdered Banana combined to flour as enrichment with kamias extract and zest. This study will include controlled experiments where different ratios of ingredients will be utilized in every formulation to produce the most preferred kamias banana cake sample. These samples will undergo sensory evaluation conducted by selected expert panels, employing standardized methods to assess appearance, aroma, texture, flavor, and overall preference.

Completely Randomized Design will be used to determine which formulation will produce the best sensory qualities for consumer approval. The ultimate objective is to determine the best possible formula highlighting the unique qualities of the utilization of banana-enriched flour and kamias as an enhancement in the cake recipes in lemon cake inspired recipe resulting in a new flavourful and marketable product.

Table 1. Kamias Banana Cake Formulation

Ingredients	Control	Formulation 1	Formulation 2	Formulation 3
Banana Powder	-	75 grams	150 grams	225 grams
All purpose flour	750 grams	675 grams	600 grams	525 grams
Kamias extract	Lemon Juice 30 grams	60 grams	60 grams	60 grams
Kamias zest	Lemon Zest 20 grams	20 grams	20 grams	20 grams

Ingredients	Control	Formulation 1	Formulation 2	Formulation 3
Sugar	500 grams	500 grams	500 grams	500 grams
Butter	250 grams	250 grams	250 grams	250 grams
Oil	60 grams	60 grams	60 grams	60 grams
Egg	200 grams	200 grams	200 grams	200 grams
Baking powder	10 grams	10 grams	10 grams	10 grams
Salt	5 grams	5 grams	5 grams	5 grams
Milk	250 grams	250 grams	250 grams	250 grams
Glaze				
Confectioners sugar	125 grams	125 grams	125 grams	125 grams
Kamias extract	Lemon juice 30 grams	30 grams	30 grams	30 grams

Table 2. Other Ingredients

PREPARATION AND PROCUREMENT OF MATERIALS FOR EXPERIMENT

Materials include the ingredients and its formulation. The ingredients must be quantified. Table of materials should also be shown.

Table 3. Specification of Raw Materials

Description	Characteristics	Source
Lakatan	dehydrated and powderized. used to combine with all purpose flour	Biñan Bayan, Carmona Palengke
Kamias	extract - used as a substitute to lemon juice. zest - used as a substitute to lemon zest	Ledda's Residence
Sugar	it is used as sweetener	Carmona Palengke
Butter	it is used to keep the richness, tenderness and structure	SM San Pedro

Oil	Used in batter mixture for creaming	Carmona Palengke
Egg	It is used to create structure and add moisture to cakes	Carmona Palengke
All purpose flour	it is the main ingredient and will give structure to cake	Carmona Palengke
Baking powder	it is used as a leavening agent	Carmona Palengke
Salt	it is used to give flavor	Foyo's Residence
Milk	used to improve texture and mouthfeel.	SM San Pedro
Glaze		
Confectioners sugar	It is used as sweetener for glaze	SM San Pedro
Kamias juice	used as a substitute for lemon juice	Ledda's Residence

Table 4. Specification/s of Equipment

EQUIPMENT	SPECIFICATION	SOURCE
Stove	Burner	CvSU Kitchen 2
Dehydrator	Dehydration	CvSU Food Laboratory
Food processor	Powderized	Hostel Storage
Extractor	Extraction	Hostel Storage
Oven	Baking	CvSU Food Laboratory
Refrigerator	Storage	CvSU Food Laboratory

DATA GATHERING PROCEDURE and PROCESS FLOW

The researchers initiated the research by sourcing the ingredients from Carmona, Cavite, Binan, Laguna, and San Pedro, Laguna for experimental production. The procedures for variables 1, 2, and 3 are all the same but differ from the formulations. The researchers started the process by preparing raw ingredients for the dehydration process, carefully choosing good quality Bananas (Lakatan). Remove banana peels and put banana flesh in a clean bowl. Measure 1,200 grams of the banana flesh then set aside. Boil 3 cups of clean water in a pot. When the water starts to boil, carefully put the banana and simmer for ten (10) minutes in a medium-low heat. After 10 minutes, remove the water using a strainer or colander, and put the bananas in a mixing bowl. Mash the boiled bananas and spread a thin layer on the parchment paper. Dehydrate the boiled bananas in a temperature of 70 to 75 degree celsius for eight (8) to nine (9) and a half hours. After dehydrating the bananas, tear them into small pieces or cut them using a kitchen scissor. Using a food processor, powderized the dehydrated bananas by three (3) batches. Every batch is powderized for one (1) minute. Using a sifter, sift the powdered bananas to remove clumping and to ensure it is finely ground.

For Kamias fruit, wash the kamias with clean water, trim both ends, and put it in a clean bowl. Remove excess water. Prepare two (2) mixing bowls for extracting the kamias fruit, one for the juice and one for the fruits that are extracted. Extract the kamias fruits six (6) times or until it is fully extracted. Put the extracted juice in a clean container.

Moving on to the baking process, measure all wet ingredients and dry ingredients. In a bowl, sift the measured dry ingredients such as 10 grams of baking powder, and 5 grams of salt. For Formulation 1 (F1), sift 675 grams flour and 75 grams banana powder, Formulation 2 (F2): 600 grams flour and 150 grams banana powder, Formulation 3 (F3): 525 grams flour and 225 grams banana powder. In a separate bowl, cream 250 grams of butter, 500 grams of sugar, 60 grams of oil, and 20 grams of kamias zest, 50 grams of egg, and 60 grams of kamias juice until light and fluffy. Combine wet ingredients and dry ingredients then mix well but do not over mix. Preheat the oven for ten (10) minutes at 325 degree fahrenheit or 163 degree celsius. Grease the baking pan with oil or butter and line with parchment paper. Pour the batter mixture on the pan and bake the cake for one (1) and a half hour or until the toothpick comes out clean. Let the cake cool in a cooling rack at room temperature for ten (10) minutes. Remove the cake from cake pans and put it on a cake board.

The process for making Kamias glaze starts from measuring 125 grams confectioners sugar and 30 grams kamias juice in a bowl, mix them all together. Drizzle the Kamias glaze on the top of the cake and attach a label and sticker on the box.

Descriptive Sensory Test utilized by ten (10) trained sensory panelists and will use a 9-point Hedonic scale questionnaire to evaluate the preferences in each formulation. Affective Sensory Test conducted to 50 random students in Cavite State University - Carmona Campus to identify its consumer acceptability with the use of Sensory Evaluation Score Sheet.

PRODUCTION PROCESS




Figure 1. Process flow for Kamias Banana Cake with Kamias Glaze

PROCESSING AND SPECIFICATION OF KAMIAS BANANA CAKE WITH KAMIAS GLAZE

Table 4. Processing and specification of Kamias Banana Cake with Kamias Glaze

Process	Specification
Preparation of raw materials (Banana)	Select good quality raw materials.
Peeling	Removing the outer cover of the banana.
Boiling	Boil water in a pot. When the water starts to boil, put the bananas and simmer for 10 minutes in a medium-low heat.
Straining	Strain the boiled bananas using a strainer or colander.
Mashing	. In a bowl, mash the boiled bananas using a fork until it becomes fine.
Spreading	Spread a thin layer of mashed bananas in parchment paper

Dehydrating	Using a dehydrator machine, dehydrate the mashed bananas at a temperature of 70 to 75 degree celsius for eight (8) to nine (9) and a half hours.
Tearing	. After dehydrating the bananas, cut them into small pieces.
Powderizing	Using a food processor, powderized the dehydrated bananas by batches.
Sifting	Sift the dehydrated powdered bananas
(Kamias)	
Washing	Wash the Kamias fruit using clean water.
Trimming	Trim both ends of the kamias fruit
Extracting	Extract the kamias fruit using an extractor and put a bowl to catch the extracted juice. Repeat for six (6) times or until kamias fruit is fully extracted
.	
(Dry Ingredients)	
Mise-en-place	.Preparing all dry ingredients needed
Weighing	Measure all dry ingredients.
Sifting	Sift all dry ingredients using a sifter, except sugar.
(Wet Ingredients)	
Mise-en-place	Preparing all wet ingredients needed
Weighing	Measuring all wet ingredients.
Creaming	Cream the butter, sugar, oil, zest, egg, kamias juice until it becomes light and fluffy. Mix well.
Mixing	Combine all dry ingredients to wet ingredients. Mix well but do not over mix.
Preheating of oven	Preheat the oven tfor ten (10) minutes at 325 degree Fahrenheit or 162.78 degree celsius.
Greasing and Lining	Grease the baking pan using butter or oil and line with parchment paper. Pour batter mixture.
Baking	Bake the cake mixture for one (1) and a half hour or until the toothpick comes out clean.
Cooling	Invert cake pans onto the cooling rack and let it cool at room temperature for at least ten (10) minutes. Remove the cake from the pan, put it on a cake board and store it in a refrigerator.
(Glaze)	
Weighing	Measure kamias juice and confectioners sugar.
Mixing	In a bowl, mix well kamias juice and confectioners sugar. Drizzle the kamias glaze on the top of the cake.
Packaging	Put the cake on the cake box.

METHODS OF EVALUATION

This research involves conducting experiments that utilize the kamias and dehydrated powdered banana as enhancement for the ingredients of cake recipes inspired by lemon cake. The researchers will provide 3 variables to determine the preferred formulation. All samples are

evaluated by ten (10) trained sensory evaluators using a 9-point hedonic scale in terms of the appearance, aroma, texture, taste/ flavor, and overall preference of the product.

CONSUMER ACCEPTABILITY

The researchers used an Affective Sensory Test and provided 50 samples for 50 untrained panels. The 9-point hedonic scale will be used to determine how much the final product will be acceptable to consumers in terms of sensory attributes such as appearance, aroma, texture, taste/ flavor, and overall acceptability. The specific serving size of the cake provided to fifty (50) untrained evaluators will be 68 grams or one (1) slice since it is the required serving size and 5 grams for the glaze.

RESULTS AND DISCUSSION

Statement of the Problem 1

1.1 Formulation and Ingredients

The researchers develop an enrichment to the control recipe Lemon Cake, with the use of kamias as substitute to lemon and combining dehydrated powder banana to flour. The control recipe is the recipe from the video tutorial of an OFW from Dubai that is uploaded on social media. With typical ingredients of Lemon cake like flour, sugar, butter, oil, egg, baking powder, milk, lemon zest and lemon juice.

The researchers made three formulations of Kamias Banana Cake with the help of their adviser, and these three formulations have the same ingredients but still have different percentages of enrichment to flour with dehydrated powdered banana and kamias as substitute to lemon. Formulation 1 contains 10% of dehydrated powdered banana and 90% of all purpose flour, formulation 2 contains 20% of dehydrated powdered banana and 80% of all purpose flour and formulation 3 contains 30% of dehydrated powdered banana and 70% all purpose flour. The most acceptable and preferred formulation was the third formulation (code 514). Other ingredients such as sugar, egg, butter, baking powder, kamias zest, and kamias juice have the same percentage within the three formulations.

Table 5: Formulations of Kamias Banana Cake with Kamias Glaze

	Dehydrated Powder Banana	All Purpose Flour
Variable 1	75 grams	675 grams
Variable 2	150 grams	600 grams
Variable 3	225 grams	525 grams

The researchers noted that the three formulations have differences in terms of sensory attributes. In formulation 3, data shows that in terms of sweet and sour taste, it is more noticeable than the formulation 2. While formulation 3 in terms of mouthfeel is more soft than formulation 2. Additionally the three formulations show differences in terms of texture and crumbliness of the cake.

Technology (Production Flow)

The procedures for variables 1, 2 and 3 are all the same but differ from the formulations. The researchers began by carefully selecting good quality Banana (Lakatan) as raw material for dehydration procedures. Banana flesh should be placed in a clean bowl once the peels have been removed. Measure 1,200 grams of the banana flesh then set aside. Boil 3 cups of clean water in a pot. When the water starts to boil, carefully put the banana and simmer for ten (10) minutes in a medium-low heat. After 10 minutes, remove the water using a strainer or colander, and put the bananas in a mixing bowl. Mash the boiled bananas and spread a thin layer on the parchment paper. Dehydrate the boiled bananas in a temperature of 70 to 75 degree celsius for eight to nine and a half hours. After dehydrating the bananas, tear them into small pieces or cut them using a kitchen scissor. Using a food processor, powderized the dehydrated banana by three batches. Every batch is powderized for one minute. Using a sifter, sift the powdered bananas to remove the clumping and ensure it is finely ground.

For Kamias fruit, carefully choose a good quality of Kamias, then wash with water and trim both ends and put it in a clean bowl. Remove excess water. Prepare mixing bowls for extracting kamias fruit, one for the juice and one for the fruits that are extracted. Extract the kamias fruit until it is fully extracted. Put the extracted juice in a clean container.

In the baking process, measure all the wet ingredients and dry ingredients. In a bowl, sift the measured dry ingredients such as baking powder, salt, flour, and banana powder. In a separate bowl, cream the butter, sugar, oil, kamias zest egg, and kamias juice until light and fluffy. Combine wet ingredients and dry ingredients, mix well but do not over mix. Preheat the oven for 10 minutes at 325 degree fahrenheit or 163 degree celsius. Grease a baking pan with oil or butter and line with parchment paper. Pour the batter mixture on the pan and bake the cake for 1 and half hour or until the toothpick comes out clean. Let the cake cool in a cooling rack at room temperature for 10 minutes, then put the cake on a cake board.

For kamias glaze, measure confectioner sugar and kamias juice in a bowl then mix well. Drizzle the glaze on top of the cake and attach a label and sticker in the box.

1.3 Financial - The table below shows the financial cost of each formulation.

Table 6: Cost of direct material of Formulation 1

Ingredients	Brand	Used Proportion	Cost as purchased
Sugar	SM Bonus	500 grams	40
Butter	Magnolia	250 grams	116.25
Oil	-	60 grams	6.6
Egg	-	200 grams	36
APF	-	675 grams	40.5
Baking Powder	Calumet	10 grams	4
Salt		5 grams	0.36
Milk	Jolly cow	250 grams	23.75
Confectioners	Peotraco	125 grams	21.25

Ingredients	Brand	Used Proportion	Cost as purchased
sugar			
Banana (Lakatan)		75 grams	45
Kamias		110 grams	4.4
TOTAL			338.11

Total Yield: 2 Loaf pan (10 slices per pan, 45 grams each)

Price per yield: 169.05 per loaf

Table 7: Cost of direct material of Formulation 2

Ingredients	Brand	Used Proportion	Cost as purchased
Sugar	SM Bonus	500 grams	40
Butter	Magnolia	250 grams	116.25
Oil	-	60 grams	6.6
Egg	-	200 grams	36
APF	-	600 grams	36
Baking Powder	Calumet	10 grams	4
Salt		5 grams	0.36
Milk	Jolly cow	250 grams	23.75
Confectioners sugar	Peotraco	125 grams	21.25
Banana (Lakatan)		150 grams	90
Kamias		110 grams	4.4
TOTAL			378.61

Total Yield: 2 Loaf pan (10 slices per pan, 45 grams each)

Price per yield: 189.30 per loaf

Table 8: Cost of direct material of Formulation 3

Ingredients	Brand	Used Proportion	Cost as purchased
Sugar	SM Bonus	500 grams	40
Butter	Magnolia	250 grams	116.25
Oil	-	60 grams	6.6
Egg	-	200 grams	36
APF	-	525 grams	31.5
Baking Powder	Calumet	10 grams	4
Salt		5 grams	0.36
Milk	Jolly cow	250 grams	23.75
Confectioners sugar	Peotracó	125 grams	21.25
Banana (Lakatan)		225 grams	135
Kamias		110 grams	4.4
TOTAL			419.11

Total Yield: 2 Loaf pan (10 slices per pan, 45 grams each)

Price per yield: 209.56 per loaf

Statement of the Problem 2

Table 9: Summary results of Appearance (COLOR)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	4.30	2.06	1.41	3.35	0.26	Accept HO
Variable 2	10	4.60	1.71				
Variable 3	10	5.60	1.65				

According to (Imchen,Singh,2023) Food colour is an important sensorial attribute that has a direct influence on food selection and acceptability. (It) showed that food colour is, often, used as an indicator of food quality such as flavour, safety, and nutritional value. The result of appearance in terms of color shows that Variable 1 had a mean score of 4.30, Variable 2 had a

mean score of 4.60 , Variable 3 had a mean score of 5.60. The result of P-value is 0.26 which means P-value is greater than the alpha value 0.05 . Therefore the null hypothesis is accepted.

Table 10: Summary results of Appearance (SIZE)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	7.1	1.91	0.06	3.35	0.94	Accept HO
Variable 2	10	6.8	1.87				
Variable 3	10	6.9	1.97				

The result of Size in terms of Appearance shows that Variable 1 had a mean score of 7.1, Variable 2 had a mean score of 6.8, Variable 3 had mean score of 6.9. The result of P-value is 0.94 which means p-value is greater than the alpha value 0.05. Therefore the null hypothesis is accepted.

Table 11: Summary results of Appearance (SHAPE)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	7.5	1.27	0.14	3.35	0.87	Accept HO
Variable 2	10	7.3	1.42				
Variable 3	10	7.6	1.17				

The shape of the food is a fundamental feature that defines its visual attribute and plays a crucial role in its sensorial appearance, intrinsic texture, design of packaging material, and overall acceptability to the consumer. Abnormally shaped food reduces desire to purchase, which may result in food waste and expiration. In recent years, the food industry has recognised the significance of food shape, with a growing focus on creating visually appealing food products that cater to changing consumer preferences. (Koirala, Prakash, Karim, Bhandari, 2023)

The result of appearance in terms of shape Shows that Variable 1 had a mean score of 7.5, Variable 2 had a mean score of 7.3, Variable 3 had a mean score of 7.6. The result of P-value is 0.87 which means p-value is greater than the alpha value 0.05 Therefore the null hypothesis is accepted.

Table 12: Summary results of Aroma (BUTTERY SMELL)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	6.7	0.95	0.32	3.35	0.73	Accept HO
Variable 2	10	6.6	0.84				
Variable 3	10	6.3	1.57				

The results of Aroma in terms of buttery smell shows that Variable 1 got a mean score of 6.7, variable 2 got a mean score of 6.6, and variable 3 got a mean score of 6.3. The p-value which

resulted in 0.73 is greater than the alpha value which is 0.05. Therefore, we accept the null hypothesis.

Table 13: Summary results of Aroma (CREAMY SMELL)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	6.2	2.25	0.03	3.35	0.97	Accept HO
Variable 2	10	6	1.56				
Variable 3	10	6	2.26				

The results of Aroma in terms of creamy smell shows that variable 1 got a mean score of 6.2, while variables 2 and 3 got the same mean score of 6. The result of the computed p-value shows 0.97 which is greater than the alpha value, which is 0.05. Therefore, we accept the null hypothesis.

Table 14: Summary results of Aroma (BANANA SMELL)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	5.6	1.17	0.31	3.35	0.74	Accept HO
Variable 2	10	6.2	1.48				
Variable 3	10	5.9	2.28				

The results of Aroma in terms of banana smell shows that variable 1 got the lowest mean score of 5.6, variable 2 got the highest mean score of 6.2, and variable 3 got the second highest mean score of 5.9. The result of p-value which is 0.74 shows that it is greater than the alpha value of 0.05. Therefore, we accept the null hypothesis.

Table 15: Summary results of Texture (CRUMBLY)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	6.5	1.90	0.90	3.35	0.42	Accept HO
Variable 2	10	7.1	1.52				
Variable 3	10	7.5	1.58				

Texture also plays an important role in consumer preferences. The perception of texture goes through several steps, starting with the visual appearance of the food product, and then continues with touch, the first bite, chewing, and swallowing. Texture enhances the food sensory experience and is a major feature in food and beverage innovation around the world, (Nachay,2020)

The result of texture in terms of its crumbliness shows that variable 1 got a mean score of 6.5, variable 2 got a mean score of 7.1, and variable 3 got a mean score of 7.5. The computed result of the p-value which is 0.42, is greater than the alpha value. Therefore, we accept the null hypothesis.

Table 16: Summary results of Mouthfeel (SOFTNESS)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	7	1.25	0.79	3.35	0.47	Accept HO
Variable 2	10	6.6	1.07				
Variable 3	10	7.2	0.92				

In terms of mouthfeel of the product, variable 1 got a mean score of 7, variable 2 got a mean score of 6.6, variable 3 got a mean score of 7.2. The computed p-value resulted in 0.47 which is greater than the alpha value. Therefore, we accept the null hypothesis.

Table 17: Summary results of Taste (SWEETNESS OF CAKE)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	7.1	1.60	0.56	3.35	0.58	Accept HO
Variable 2	10	6.7	1.16				
Variable 3	10	7.3	1.06				

The result of Taste in terms of the sweetness of the cake shows that Variable 1 had a mean score of 7.1, Variable 2 had a mean score 6.7, Variable 3 had a mean score of 7.3. The result of P-value is 0.58 which means p-value is greater than the alpha value 0.05 . Therefore, the null hypothesis is accepted.

Table 18: Summary results of Taste (SWEET AND SOUR)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	6	1.70	0.50	3.35	0.61	Accept HO
Variable 2	10	5.8	1.40				
Variable 3	10	6.5	1.72				

The result of Taste in terms of sweetness and sourness shows that Variable 1 had a mean score of 6, Variable 2 had a mean score of 5.8, Variable 3 had a mean score of 6.5 . The result of P-value is 0.61 which means p-value is greater than the alpha value 0.05 . Therefore the null hypothesis is accepted.

Table 19: Summary results of Overall Quality (ACCEPTABILITY)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	7.2	2.68	2.23	2.23	2.23	Accept HO
Variable 2	10	7	2.65				
Variable 3	10	7.9	2.81				

The summary result of the overall quality of the product shows that variable 1 got the second to the highest mean score of 7.2, variable 2 got the lowest mean score of 7, and variable 3 got the highest mean score of 7.9. The computed result of p-value is 2.23 which means the p-value is greater than the alpha value. Therefore, we accept the null hypothesis.

Table 20: Summary results of Overall Quality (ACCEPTABILITY)

Formulations	No. of Sample	Mean Score	SD	F-test	Critical Value	P-value	Decision
Variable 1	10	7.2	2.68	2.23	2.23	2.23	Accept HO
Variable 2	10	7	2.65				
Variable 3	10	7.9	2.81				

The summary result of the overall quality of the product shows that variable 1 got the second to the highest mean score of 7.2, variable 2 got the lowest mean score of 7, and variable 3 got the highest mean score of 7.9. The computed result of p-value is 2.23 which means the p-value is greater than the alpha value. Therefore, we accept the null hypothesis.

Statement of the Problem 3

Table 21: Interpretation and Average of Sensory Attributes

Parameters	Mean Score	Interpretation
Appearance	8.56	Very Much Like
Aroma	8.36	Very Much Like
Texture	8.14	Very Much Like
Taste	8.46	Very Much Like
Overall Preference	8.44	Very Much Like

APPEARANCE: Consumers consider the appearance of the food first, the visual appeal of the product such as the shape , color , texture and the overall appearance of the product .Appearance plays an important role in food especially to get the attention of the customer. The mean score of appearance resulted in 8.56, and is interpreted as Very Much Like.

AROMA: The aroma of the product is one of the biggest factors in consumer acceptability. A certain aroma can influence the moods and emotions, creating a pleasant and relaxing overall experience of the product. The mean score of aroma resulted in 8.36, and is interpreted as Very Much Like.

TEXTURE: The texture of the product is one of the crucial roles in consumer acceptability. It can signify the freshness , creamy ,crumbly and softness of the product while also significantly contributing the overall experience to the product. The mean score of texture resulted in 8.14, and is interpreted as Very Much Like.

TASTE: Taste is considered as the most critical factor in determining consumer purchase decisions. A good combination of every flavor in a product is essential for overall satisfaction. A well-balanced flavor of the product is most likely to be enjoyed by consumers. The mean score of taste resulted in 8.46, which is interpreted as Very Much Like.

OVERALL PREFERENCE: The overall acceptability is the combined sensory attributes of the product such as appearance, aroma, texture, and taste. The computed result of mean score of overall preference is 8.44, and is interpreted as Very Much Like.

Statement of the Problem 4

Nutritional value of Kamias Banana Cake with Kamias Glaze

Nutritional analysis is the process of determining the nutritional content of food, (This) measures the actual levels of nutrients in the prepared food, thus providing a high level of accuracy of the analysis. The analysis accounts for the changes in nutritional value that occur due to the cooking and processing of the food. This is of extreme value as calories tend to increase or decrease during the cooking process depending on the method used. (Dimitrova, 2019)

The process of nutrition tests begins in identifying the most preferred formulation which involves conducting sensory evaluation to trained panels. This proceeded in determining the measured serving size for a 1 loaf of cake. There are 10 slices and each slice contains 45 grams of cake.

Once the most preferred formulation is determined the test will run through the required format given by the nutritionist-dietitian listing down the ingredients and its used portion. The test thoroughly records the amount of each ingredient used per yield. The nutritional components was then analyze, this include key nutrients such as adequate amounts of Vitamin A, Vitamin B6 and Iron, Calories and Sodium has a high intake for each slice making it not advisable for people with elevated blood glucose, An allergy advice was given as it contains wheat and milk.

ESTIMATED ENERGY AND NUTRIENT CONTENT										
FOOD ITEM	MACRONUTRIENTS									
	Energy (kcal)	CHO (g)	Fiber (g)	Sugars (g)	Protein (g)	Fat (g)	SFA (g)	MUFA (g)	PUFA (g)	Chol (mg)
Kamias	0.63	0.12	0.05	0.07	0.02	0.01	0.00	0.00	0.00	0.00
Banana, Lacatan	7.09	1.67	0.19	0.88	0.08	0.01	0.00	0.00	0.00	0.00
All purpose flour	46.20	9.67	0.35	0.04	1.65	0.11	0.02	0.01	0.04	0.00
Sugar, white	50.00	12.49	0.00	12.46	0.00	0.00	0.00	0.00	0.00	0.00
Butter, unsalted	44.81	0.00	0.00	0.00	0.05	5.07	3.16	1.46	0.19	13.44
Oil, Coconut	13.38	0.00	0.00	0.00	0.00	1.49	1.24	0.09	0.03	0.00
Egg, chicken	6.95	0.07	0.00	0.02	0.62	0.47	0.15	0.11	0.04	21.40
Baking Powder	0.42	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Salt, fine	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Milk, cow, whole	4.06	0.30	0.00	0.28	0.21	0.23	0.10	0.04	0.01	2.13
Sugar, powdered	12.50	3.13	0.00	3.06	0.00	0.00	0.00	0.00	0.00	0.00
TOTAL	186.04	27.55	0.59	16.82	2.63	7.37	4.66	1.72	0.31	36.96

Serving Size		1 slice (45 g)
No. of servings per pack		10
Amount Per Serving		%REI/RNI*
Calories (kcal)	186	7%
Calories from Fat (kcal)	66.6	
Total fat (g)	7.4	
Saturated fat (g)	4.7	
Cholesterol (mg)	37	
Total Carbohydrates (g)	27.5	
Dietary Fiber (g)	0.6	3%
Sugars (g)	16.8	
Protein (g)	2.6	4%
Sodium (mg)	84.6	17%
Potassium (mg)	76.0	4%
Iron (mg)	0.8	6%
Vitamin A (mcg RE)	54.2	8%
Vitamin B6 (mg)	0.1	6%
Vitamin C (mg)	1.8	3%
*Percent Recommended Energy Intake and Recommended Nutrient Intake are based on PDRI reference for adult male aged 19-29 years old		

- Contains adequate amounts of Vitamin A, Vitamin B6, and Iron
- Calories and Sodium in a slice are high, Intake must be limited to only two (2) slices each day.

May not be advisable for people with elevated blood glucose

**Allergy advice:
This product contains wheat and milk**

Maria Crizel Joy L. Alpe, RND
Registered Nutritionist-Dietitian

Figure 2. Nutritional Value of Kamias Banana Cake with Kamias Glaze

Table 22: Result of the pH test of the Most Preferred Formulation

Attempt	Results
Trial 1	5.46
Trial 2	5.39
Trial 3	5.35
AVERAGE	5.4

According to Rodrigues, 2024 pH testing is the process by which the acidity or alkalinity of a substance it (is) determined, pH is highly influential in food and beverage production and its variability can cause critical disparities in quality, consistency and safety of the food products. For these reasons, pH is one of the parameters most frequently controlled during production and quality control before release. Taste is radically influenced by pH, since pH levels can help to regulate sweetness and sourness perception.

Additionally, In food safety, pH plays a crucial role in inhibiting the growth of spoilage microorganisms. For these reasons, most regulatory agencies impose rules for the pH levels of certain commercial foods, since generally acidic foods ($\text{pH} < 4.6$) are considered safer and allow longer shelf lives.

The utilization of banana as enrichment in cake recipe pH level was observed by conducting three distinct trials, yielding varying results to attain the average pH level.

The range of the pH scale is 0 –14. A solution with a pH of 7 is neutral; one with a pH of less than 7 is acidic. A solution is basic if its pH is higher than 7. (Tantray, Nissa, 2023)

The pH testing yielded a result of 5.46 for the first trial, 5.39 for the second trial and 5.35 for the third trial, this came up with the average pH level score of 5.4. This outcome indicates that it is slightly acidic.

Statement of the Problem 5

Sustainable packaging minimises environmental impact whilst maintaining its functional requirement. It can take the form of recyclable or reusable packs or involve reducing material and resource use. All forms of packaging used to present, protect, ship or store products can be sustainable if designed correctly. (Dobson,2024)

Table 23. Results of Sustainable Packaging Acceptability Survey

Parameters	Parameters	Interpretation
Design	7.9	Moderately like
Acceptability	8	Very much like

The researcher conducted an online survey for acceptability of sustainable packaging for cake using banana peel (Lakatan) and following the traditional paper making process to produce a sustainable packaging for the cake. The results for design of the packaging is 7.9 and interpreted as Moderately Like in terms of its appearance and size. While the acceptability of the packaging

resulted in 8 that is interpreted as very much like. It is based on what it shows in the picture that the researcher provides.

CONCLUSIONS

The results of the experimental research in enhancing lemon cake recipe as control, indicated consumers preferences within the three variables of kamias cake with enrichment of dehydrated powder banana combined to flour as a new option to consumers. The three formulations were used to determine the most preferred variable. The most preferred variable was the formulation 3 which contains 30% of dehydrated powder banana and 70% all purpose flour that resulted in a significant distinction from the other formulation.

The researchers studied which fruit or flavor would enhance and suit the taste of the kamias. After considering the various fruit options that can be used as enhancement to flour, Lakatan Banana was selected for its nutritional value and because it has not been properly utilized in different recipes. Kamais was used as it has similarities to lemon and because it has not been utilized in baking pastries.

It conducted a sensory evaluation of the most preferred variable, and the overall acceptability satisfied the consumers. The most preferred variable shows significant difference from the other formulation.

The nutrition testing shows that the cake has adequate amounts of vitamins such as vitamin A, B6, and Iron. But a slice of this is also high in calories and sodium and its intake must be limited to only two (2) slices a day. It also has allergy advice for containing wheat and milk, and not advisable for people with elevated blood glucose.

Checking the acidity level of food using pH testing, utilization of banana as enrichment in cake recipe pH level was observed by conducting three distinct trials, yielding varying results to attain the average pH level.

Conducting an online survey for acceptability of sustainable packaging using banana peel (Lakatan) and following the traditional paper making process shows that the Design is moderately like and the Acceptability is very much like.

RECOMMENDATIONS

This study recommended that future researchers can use the other types of banana in making baked products or pastries in terms of Financial; the “lakatan” banana that was utilized was in medium-range price and cheaper alternatives are still encouraged. As, conducting further tests to support their study is advisable.

In dehydrating the bananas, the researchers highly recommend using high quality parchment paper. Usage of wax paper is not recommendable in dehydrating the banana as it can stick thoroughly on the wax paper.

Future researchers are encouraged to powderized the dehydrated banana for a total of 2 minutes if the first minute process wasn't that successful, thru this, the powder's outcome was better in terms of appearance and is more fine.

Future researchers and home bakers are encouraged to try alternative sweeteners such as stevia, that has no sugar content to lessen the sugar intake.

In terms of sustainable packaging, future researchers can remake the packaging and make it more appealing in terms of appearance in color, aroma and to further add durability.

Encourages local farmers to maintain the quality of bananas to further promote local resources as it can benefit them by selling it to the market and encourage others to utilize it on various research.

REFERENCES

Ben-Noun, L. (2019). ResearchGate. CAKES FROM ANCIENT TO CONTEMPORARY TIMES.

https://www.researchgate.net/publication/327816224_CAKES_FROM_ANCIENT_TO_CONTEMPORARY_TIMES

Friedberg, J. (2024). Snack Food & Wholesale Bakery. Fruits add flavor and flexibility in snack and bakery products. <https://www.snackandbakery.com/articles/111160-fruits-add-flavor-and-flexibility-in-snack-and-bakery-products>

Sampiano, K., & Durban, A. (2022). International Journal on Food, Agriculture, and Natural Resources. The Physical and Sensory Qualities of 'Lakatan' Banana (*Musa acuminata*) in Response to Different Natural Ripening Agents. <https://journal.fanres.org/index.php/IJFANRES/article/view/92>

Stuart Jr., G. (2024). StuartXchange.Org. Kamias/ Averrhoa bilimbi/ Bilimbi Tree. http://www.stuartxchange.org/Kamias.html?fbclid=IwZXh0bgNhZW0CMTEAAR3f9AL5FKeIrbgu2nnplvLILIJkmxRhlc39QRRMh41atVLVrN8QLSuc4fA_aem_Cntu6OwDk2Z56Jfb_9iO4w

Cabilzo, R. (2023). University of the Philippines Los Baños, College of Human Ecology. Assessment of Lacatan Banana *Musa Acuminata* (AA Group) Waste-Loss in Santa Ana Public Market, Manila, Philippines. <https://che.uplb.edu.ph/theses-data/assessment-of-lacatan-banana-musa-acuminata-aa-group-waste-loss-in-santa-ana-public-market-manila-philippines/>

Leong, Y., & Chang, J. (2022). Science Direct. Valorization of fruit wastes for circular bioeconomy: Current advances, challenges, and opportunities. <https://www.sciencedirect.com/science/article/abs/pii/S096085242200788X>

Kumar, M., Gehlot, R., Sindhu, R., Kumar, S., Rekha, R., & Singh, R. (2019). ResearchGate. Physico-chemical characteristics of fresh banana fruits.

https://www.researchgate.net/publication/372195136_Physico-chemical_characteristics_of_fresh_banana_fruits/citations

Papantoniou, N. (2020). Good Housekeeping. What Is a Dehydrator, and Do You Really Need One? The benefits and drawbacks of using a food dehydrator. <https://www.goodhousekeeping.com/appliances/a31904157/what-is-a-dehydrator/>

Chiang, M. (2020). Semantic Scholar. Is Excessive Intake of Kamias Fruit (Averrhoa bilimbi) Bad for Your Health? A Case Report and Review. <https://pdfs.semanticscholar.org/bfe9/b281e69909ee69388cc93351ab48bee63e15.pdf>

Astillo, J. (2020). International Journal of Environment, Agriculture and Biotechnology. Bilimbi Fruit (Averrhoabilimbi) Juice. https://www.researchgate.net/profile/Jessica-Astillo-2/publication/341771071_Bilimbi_Fruit_Averrhoabilimbi_Juice/links/5f8fecfca6fdccfd7b71efbf/Bilimbi-Fruit-Averrhoabilimbi-Juice.pdf

Stuart Jr., G. (2022). StuartXchange.Org. Lakatan/ Musa acuminata Colla/ DESSERT BANANA. <http://www.stuartxchange.org/Lakatan>

Kumar, N., Ved, A., Yadav, R., Prakash, O. (2021). International Journal of Current Research and Review. A Comprehensive Review on Phytochemical, Nutritional, and Therapeutic Importance of Musa acuminata. <https://typeset.io/pdf/a-comprehensive-review-on-phytochemical-nutritional-and-47dd5c1mwb.pdf>

Basu, S. (2023). Net Med. Bilimbi: Incredible Health Benefits Of This Nutritious Fruit. https://www.netmeds.com/health-library/post/bilimbi-incredible-health-benefits-of-this-nutritious-fruit?srsId=AfmBOoommMpU_KHKlpdjpOOF9WGwcUg1Rp2ODBdCjrOEXSU0GNYLCwyU

Garcia, G. (2023). Prutas Lokal. Saging (Banana): Mga Nakakabilib na Health Benefits. Alamin!. <https://prutaslokal.com/blog/31-10-health-benefits-ng-saging-banana-sa-kalusugan>

Frey, M. (2024). Verywell fit. Banana Nutrition Facts and Health Benefits. <https://www.verywellfit.com/are-bananas-bad-for-weight-loss-3495634>

Gawli, P., Sarpotdar, S., Nair, S., Jadhav, V. (2021). ScienSage. A REVIEW ON RECENT ADVANCES ON ETHNO-MEDICAL PLANT: AVERRHOA BILIMBI. <https://sciensage.info/index.php/JASR/article/view/916>

Agoncillo, A., Gavarra H., Madrid, A., Polintan, J., Rambaua, A., Sanguyo, K., & Solomon, J. (2022). Course Hero. Caramela Custelly: A unique dessert in Caramel Banana Custard Jelly. <https://www.coursehero.com/file/157284647/Group-7-FINAL-PAPERpdf/>

Seal, J. (2022). Ehow. Characteristics of a Lakatan Banana. https://www.ehow.com/info_8324965_characteristics-lakatan-banana.html

Meherishi, L., Narayana, S., Ranjani, K. (2019). Science Direct. Sustainable packaging for supply chain management in the circular economy: A review. <https://www.sciencedirect.com/science/article/abs/pii/S0959652619323960>

Saclolo, M., Watabe, M., Eugenio, R., Campos, A., Dela cruz, K., Lapitan, A., & Puyat, L. (2023). Ascendens Asia Journal of Multidisciplinary Research Abstracts. Paper Production of Musa Acuminata and Balbisaniana (Saba) and its Effectiveness. <https://ojs.aaresearchindex.com/index.php/AAJMRA/article/view/12536>

Arcalas, J. (2024). Philstar Global. Philippines dislodged as 2nd biggest banana exporter. <https://www.philstar.com/business/2024/07/29/2373651/philippines-dislodged-2nd-biggest-banana-exporter>

Imchen, T., Singh, K. (2023). Science Direct. Marine algae colorants: Antioxidant, anti-diabetic properties and applications in food industry. <https://www.sciencedirect.com/topics/engineering/food-color>

Dimitrova, M. (2019). AZO Materials. Nutrition Analysis Techniques to Determine the Nutritional Content of Food. <https://www.azom.com/article.aspx?ArticleID=18555>

Tantray, J., Mansoor, S., Wani, R., Nissa, N. (2023). Science Direct. Chapter 3 - pH meter: Its use and calibration. <https://www.sciencedirect.com/science/article/abs/pii/B9780443191749000039>

Rodrigues, R. (2024). Technology Networks. pH Testing and the pH Meter for Food Analysis. <https://www.technologynetworks.com/applied-sciences/articles/ph-testing-and-the-ph-meter-for-food-analysis-370482>

Nachay, K. (2020). Institute of Food Technologists. Bringing Food Texture to the Forefront. <https://www.ift.org/news-and-publications/food-technology-magazine/issues/2020/may/columns/bringing-food-texture-to-the-forefront>

Koirala, S., Prakash, S., Karim, A., Bhandari, B. (2023). Science Direct. Shape morphing of foods: Mechanism, strategies, and applications. <https://www.sciencedirect.com/science/article/pii/S0924224423002509#:~:text=The%20shape%20of%20the%20food,bulk%20properties%2C%20and%20packaging%20requirements>

**SEED-TO-SHELF VENTURES: BUILDING A YOUTH-LED AGRIBUSINESS
MODEL FOR FARM-TO-MARKET TRANSPARENCY**

Sadiq, M.S

Department of Agricultural Economics and Agribusiness, FUD, Dutse, Nigeria

ORCID ID: 0000-0003-4336-5723

Singh, I.P

Department of Agricultural Economics, SKRAU, Bikaner, India

ORCID ID: 0000-0002-1886-5956

Ahmad, M.M

Department of Agricultural Economics and Extension, BUK, Kano, Nigeria

ORCID ID: 0000-0003-4565-0683

Sani, B.S

PhD Scholar, Department of Agricultural Economics and Agribusiness, FUD, Dutse, Nigeria

ORCID ID: 0000-0001-7773-3796

Nazifi, I.K

Federal Inland Revenue Services (FIRS), Nigeria

Abstract

The increasing demand for transparency in agricultural supply chains has spurred the development of innovative agribusiness models that ensure accountability from seed to shelf. This paper explores a youth-led agribusiness model that leverages digital technology, sustainable farming practices, and entrepreneurial skills to enhance farm-to-market transparency. The study critically analyzes the role of blockchain, IoT, and data analytics in ensuring traceability and quality assurance in agribusiness. Additionally, it evaluates the socio-economic impact of youth-led agribusiness ventures on rural development, employment creation, and food security. Drawing from empirical studies and recent case analyses, the paper provides a theoretical and conceptual framework to guide the development of sustainable, transparent, and technology-driven agribusiness models. The study concludes with policy recommendations for governments, financial institutions, and private sector stakeholders to support youth engagement in agriculture through funding, capacity building, and market integration initiatives.

Keywords: Agriculture; Blockchain; Supply Chain; Sustainability; Youth

Introduction

Background of the Study

Agriculture remains a critical sector for economic development, food security, and employment; particularly in developing economies (FAO, 2023). However, persistent challenges such as supply chain inefficiencies, market opaqueness, and weak farmer-buyer linkages hinder the sector's growth. Smallholder farmers-who contribute significantly to global food production-often, face price exploitation, low bargaining power, and a lack of market intelligence, making it difficult for them to compete in larger agribusiness markets (World Bank, 2022).

To address these issues, the **Seed-to-Shelf Ventures** model has emerged as a promising youth-led approach to agribusiness. This model integrates digital technologies, sustainable farming practices, and innovative business strategies to enhance farm-to-market transparency. Studies indicate that digital platforms such as **blockchain-based traceability systems, IoT-enabled farm monitoring, and mobile agribusiness applications** are transforming traditional agricultural supply chains, ensuring fair pricing and quality control (Obebo, 2023). The success of youth-driven agribusiness models, such as **Digifarm and Toggfram**, highlights the potential of technology-enabled transparency in the agriculture sector (Obebo, 2023).

Despite these advancements, the agricultural sector remains unattractive to many young people due to **high startup costs, limited access to finance, and inadequate technical knowledge** (IFAD, 2022). Recent research suggests that youth-led agribusiness incubators, financial support schemes, and digital marketplaces could provide an enabling environment for young entrepreneurs to enter and thrive in agriculture (Lumen, 2020).

Problem Statement

While agriculture presents a significant opportunity for youth employment and economic empowerment, engagement in the sector remains low due to multiple barriers. According to **IFAD (2022)**, factors such as **restricted access to credit, poor infrastructure, and the perception of farming as unprofitable** have discouraged young people from venturing into agribusiness. Additionally, supply chain inefficiencies have led to **widespread market opacity**, where small-scale farmers receive minimal returns despite rising consumer prices (Shen *et al.*, 2024).

Another growing concern is consumer demand for **greater transparency in food sourcing**, driven by concerns over **food safety, environmental sustainability, and ethical sourcing** (Zhong *et al.*, 2023). Current agribusiness models lack the necessary mechanisms to provide full traceability of agricultural products from farm to retail shelves. For instance, studies on **farm-to-market roads and rural logistics** highlight that inadequate infrastructure has worsened inefficiencies in rural agribusiness operations (Cawley, 2020).

Therefore, there is an urgent need to establish **youth-led, technology-driven agribusiness models** that enhance transparency, efficiency, and fair pricing in the agricultural value chain. Addressing these challenges could not only attract more young entrepreneurs to the sector but also improve economic outcomes for farmers and consumers alike.

Recent Studies Supporting the Argument

Several recent studies underscore the importance of youth engagement and technological innovation in agriculture:

- **Obebo (2023)** discusses how Digifarm, a youth-led agribusiness platform, integrates digital tools to enhance traceability and transparency in Kenya's urban food systems.

- **Shen et al. (2024)** examine the role of farm-to-market road networks in reducing inefficiencies and promoting better accountability in rural agribusiness.
- **Lumen (2020)** explores financial and infrastructural challenges affecting youth participation in agriculture in the Davao region.
- **Zhong et al. (2023)** highlight consumer-driven demand for transparent supply chains and ethical food sourcing.

By leveraging these insights, the **Seed-to-Shelf Ventures** model can offer a **scalable, technology-driven agribusiness framework** that addresses transparency issues while empowering youth entrepreneurs in the sector.

Research Objectives

1. To explore the role of youth-led agribusiness in enhancing farm-to-market transparency.
2. To assess the impact of digital technologies (e.g., blockchain, IoT) in ensuring traceability in agribusiness.
3. To evaluate the socio-economic benefits of youth engagement in agriculture.
4. To propose policy recommendations for promoting sustainable, technology-driven agribusiness ventures.

Theoretical Framework

This section explores three key theoretical perspectives that provide the foundation for understanding youth-led agribusiness models and their role in ensuring farm-to-market transparency. These theories explain how youth participation in agribusiness, technological innovation, and policy interventions contribute to sustainable and transparent agricultural supply chains.

Sustainable Livelihoods Theory (SLT)

The Sustainable Livelihoods Theory (SLT) asserts that individuals' ability to sustain their livelihoods depends on access to five core assets: human, social, natural, financial, and physical capital (Scoones, 2019). In the context of youth-led agribusiness, these assets manifest as:

- **Human capital:** Skills, training, and knowledge that enable youth to engage in agribusiness.
- **Social capital:** Networks, partnerships, and mentorships that facilitate market access and knowledge exchange.
- **Natural capital:** Access to land, water, and biodiversity, which are essential for sustainable farming.
- **Financial capital:** Availability of loans, grants, and investment opportunities to support agribusiness startups.
- **Physical capital:** Infrastructure, machinery, and digital tools that enhance agricultural productivity and supply chain efficiency.

Recent studies have emphasized the importance of **digital and financial inclusion** in enhancing the livelihoods of young farmers. For instance, research by **Villalba, Venus, & Sauer (2023)** highlights how **fintech solutions and microcredit platforms** have enabled rural youth to access financial resources, thereby increasing their participation in agribusiness. Additionally, **Mulligan & Berti (2023)** argue that **innovative food supply chain hubs** improve local food systems, benefiting smallholder farmers by ensuring **equitable access to markets**.

Moreover, the **ecosystem approach** to sustainable livelihoods suggests that policies promoting youth agribusiness should focus on enhancing **digital literacy, financial training, and regulatory support** to build resilient agripreneurs (Chitaka, 2024).

Supply Chain Transparency Theory (SCT)

Supply Chain Transparency Theory (SCT) posits that **transparent supply chains increase trust, efficiency and sustainability** (Mol, 2015). Transparency is particularly crucial in agriculture, where **food safety, ethical sourcing, and traceability** are key concerns. According to **Grabs & Carodenuto (2021)**, consumers are increasingly demanding **real-time visibility of agricultural production**, which has led to the adoption of **blockchain, IoT, and AI-driven monitoring systems** in agribusiness.

Key components of SCT in youth-led agribusiness include:

- **Traceability systems** that enable farmers to document and verify their production processes.
- **Digital marketplaces** that facilitate direct interactions between farmers and consumers, eliminating exploitative intermediaries.
- **Certification and compliance monitoring** to ensure adherence to sustainable farming and fair trade standards.

A study by **Mangla et al. (2022)** on blockchain adoption in **agri-food supply chains** emphasizes that transparency enhances **trust, reduces fraud, and promotes sustainable farming practices**. Additionally, **Khan, Behrendt, & Papadas (2024)** highlight that intermediaries play a crucial role in managing sustainability compliance in multi-tier food supply chains, further reinforce the need for **robust digital transparency frameworks**.

Recent advancements in **IoT-based smart farming** have further strengthened transparency mechanisms. **Quayson, Bai, & Sarkis (2020)** discuss how **sensor-based monitoring and remote tracking technologies** have improved **quality assurance, waste reduction, and fair pricing** in agricultural supply chains.

Entrepreneurial Ecosystem Theory

Entrepreneurial Ecosystem Theory explains how a combination of **policy frameworks, financial access, market linkages, and human capital development** fosters entrepreneurship (Stam, 2018). Youth-led agribusiness models thrive when they are supported by a **favorable entrepreneurial ecosystem**, which includes:

- **Regulatory policies** that provide tax incentives, subsidies, and funding for agripreneurs.
- **Incubation centers and agribusiness accelerators** that offer training and mentorship.
- **Access to venture capital and crowdfunding platforms** that support startup financing.
- **Collaborative networks between academia, government, and private sector stakeholders** to drive innovation.

A study by **Okoye (2021)** emphasizes that **entrepreneurial ecosystems in agriculture must integrate technology, knowledge-sharing platforms, and policy interventions** to support young farmers. Similarly, **Hussain & Scott (2024)** found that emerging **smart agricultural hubs** serve as **catalysts for youth entrepreneurship** by providing **training, access to digital platforms, and market linkages**.

Furthermore, **RandhirsinhMohite & Bhola (2023)** argue that rural agricultural ecosystems must focus on **agri-tech innovation, cooperative models, and youth-driven agribusiness clusters** to drive inclusive economic growth. Their research highlights that governments should

simplify bureaucratic processes for agribusiness startups, ensuring young farmers can easily register businesses and access financial support.

Summary of Theoretical Contributions

- **SLT** explains how access to financial, human, and social capital enables youth engagement in agribusiness.
- **SCT** highlights how **digital transparency tools improve efficiency, trust, and sustainability** in agricultural supply chains.
- **Entrepreneurial Ecosystem Theory** emphasizes the **importance of supportive policies, financial mechanisms, and market integration** in fostering youth-led agribusiness models.

These theories collectively provide a **holistic framework** for developing a **scalable, sustainable, and transparent Seed-to-Shelf agribusiness model**, ensuring youth participation and economic viability in modern agricultural markets.

Conceptual Framework

The conceptual framework in this study provides a **structured model** for understanding how youth-led agribusiness ventures integrate **technology, entrepreneurship, and sustainability** to enhance farm-to-market transparency. The framework also highlights the role of mediating variables such as **policy support and financial access** in ensuring successful implementation.

3.1 Key Variables in the Conceptual Framework

Independent Variables (Drivers of Transparency in Youth-Led Agribusiness)

1. Technological Innovations

- **Blockchain Technology:** Ensures product traceability, reduces fraud, and enhances consumer trust (Rogito & Nyakora, 2023).
- **Internet of Things (IoT):** Smart sensors monitor farm conditions, reducing post-harvest losses and improving supply chain efficiency (Mullu, 2023).
- **AI-driven Analytics:** Optimizes pricing strategies, predicts demand, and supports precision farming (Bell, 2024).

2. Youth Entrepreneurship

- **Training and Capacity Building:** Entrepreneurship education programs help young farmers adopt innovative agribusiness models (Mullu, 2023).
- **Funding Access:** Startups need microfinance, venture capital, and subsidies to scale operations (Basir & Musa, 2024).
- **Market Linkages:** Digital marketplaces connect youth agripreneurs directly to consumers, minimizing exploitation by intermediaries (Mdoda *et al.*, 2024).

3. Sustainable Farming Practices

- **Organic Agriculture:** Enhances soil fertility, reduces chemical dependency, and aligns with consumer preference for healthy, eco-friendly food (Eze *et al.*, 2024).
- **Regenerative Agriculture:** Practices such as agroforestry and cover cropping improve biodiversity and water conservation (Cruickshank *et al.*, 2022).
- **Carbon Credit Systems:** Incentivizes sustainable farming through carbon offset payments (Mokhema *et al.*, 2024).

Mediating Variables (Enablers of Agribusiness Success)

Policy and Institutional Support

- **Regulatory Frameworks:** Governments play a role in enforcing transparency and sustainability standards in agribusiness (Kitambo, 2022).
- **Subsidies and Incentives:** Financial grants, tax reliefs, and research funding stimulate youth participation in agribusiness (Raphuti, 2023).

Financial Access

- **Microfinance and Venture Capital:** Access to credit empowers young farmers to invest in technology and expand operations (Mdoda *et al.*, 2024).
- **Government and NGO Grants:** International organizations support youth-led agribusiness projects with seed funding and technical assistance (Dietershagen & Bammann, 2023).

Dependent Variables (*Outcomes of Transparency in Agribusiness*)

Transparency in Agribusiness

- **Traceability Systems:** Consumers can verify product origin and production methods (Rogito & Nyakora, 2023).
- **Accountability Mechanisms:** Digital platforms ensure fair trade by eliminating middlemen exploitation (Mdoda *et al.*, 2024).

Economic Empowerment

- **Job Creation:** Youth-led agribusiness models create employment, especially in rural areas (Erz, 2023).
- **Rural Development:** Improved access to finance and markets leads to economic growth in local communities (Kitambo, 2022).

Recent Studies Supporting the Framework

- **Rogito & Nyakora (2023)** highlight the role of **corporate governance in promoting youth participation in agribusiness** by enhancing market transparency.
- **Mullu (2023)** found that **youth involvement in agritech solutions (such as IoT-based monitoring systems) significantly reduces farm-to-market inefficiencies**.
- **Basir & Musa (2024)** discuss the role of **Islamic finance models in providing sustainable credit solutions for young farmers in Africa**.
- **Eze et al. (2024)** emphasize how **organic farming adoption by youth-led enterprises can enhance agricultural productivity while maintaining ecological balance**.
- **Cruickshank et al. (2022)** propose **value chain analysis as a tool for ensuring transparency in youth-led agribusinesses**.
- **Dietershagen & Bammann (2023)** investigate the **potential of the bioeconomy sector to drive youth employment and entrepreneurship** in sustainable agriculture.

The conceptual framework illustrates how **technological innovations, youth entrepreneurship, and sustainable farming practices** drive transparency in youth-led agribusinesses. However, the success of these ventures depends on **enabling policies and access to finance**, which mediate their impact on **traceability, economic empowerment, and rural development**.

By adopting **blockchain, IoT, AI-driven analytics, and regenerative farming practices**, young agripreneurs can enhance **farm-to-market efficiency and fairness** while contributing to **global food security and sustainable agriculture**.

Below is the **visual representation** of the conceptual framework for **Youth-Led Agribusiness and Farm-to-Market Transparency**.




Figure 1: Conceptual framework

- **Independent Variables (Drivers):**
 - Technological Innovations (Blockchain, IoT, AI)
 - Youth Entrepreneurship (Training, Funding, Market Access)
 - Sustainable Farming Practices (Organic, Regenerative)
- **Mediating Variables (Enablers):**
 - Policy & Institutional Support (Subsidies, Regulations)
 - Financial Access (Grants, Microfinance, Venture Capital)
- **Dependent Variables (Outcomes):**
 - **Transparency in Agribusiness** (Traceability, Fair Trade)
 - **Economic Empowerment** (Income Generation, Rural Development)

Research Methodology

This section outlines the research methodology employed in conducting the **review paper on youth-led agribusiness models and farm-to-market transparency**. The methodology

follows a **systematic review approach**, integrating qualitative and quantitative data from **peer-reviewed literature, case studies, and policy reports**.

Research Design

This study adopts a **systematic literature review (SLR) design**, which involves a structured and comprehensive synthesis of existing academic and industry research on the topic.

- **Objective:** To analyze the role of **youth-led agribusiness ventures in enhancing farm-to-market transparency** through digital innovations.
- **Approach:** The review **examines empirical studies, case studies, and policy frameworks** related to **technological adoption, agribusiness entrepreneurship, and supply chain transparency**.

A **PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) framework** was used to ensure a rigorous selection and synthesis of literature (Moher *et al.*, 2009).

Data Collection Strategy

Search Strategy

To identify relevant academic and industry sources, a **structured database search** was conducted using the following databases:

- **Academic Databases:**
 - Google Scholar
 - Scopus
 - Web of Science
 - PubMed
 - ResearchGate
 - ScienceDirect
 - IEEE Xplore
- **Agriculture & Development Reports:**
 - Food and Agriculture Organization (FAO)
 - International Fund for Agricultural Development (IFAD)
 - World Bank Open Data
 - CGIAR and IFPRI reports

Search Terms & Keywords

The following **Boolean search strategy** was used:

("youth-led agribusiness" OR "agripreneurship") AND ("farm-to-market transparency" OR "agriculture supply chain transparency") AND ("blockchain" OR "digital agriculture" OR "IoT" OR "AI") AND ("policy frameworks" OR "rural development" OR "economic empowerment")

- Filters applied:
 - **Publication Year:** 2020–2024
 - **Language:** English

- **Document Type:** Peer-reviewed articles, case studies, policy reports

Inclusion and Exclusion Criteria

To ensure relevance and quality, the review applied the following criteria:

Criteria	Inclusion	Exclusion
Timeframe	Studies published from 2020–2024	Studies published before 2020
Scope	Studies on youth agribusiness, digital transparency	Studies without focus on youth or transparency
Geographical Coverage	Global, with a focus on Africa, Asia, and Latin America	Studies focusing only on high-income countries
Methodology	Empirical studies, case studies, and policy papers	Opinion pieces or articles lacking data
Language	English	Non-English studies due to translation constraints

A total of **120 articles were initially identified**, and after **screening and eligibility checks, 38 articles were selected for full review**.

Data Extraction and Analysis

1. Data Extraction Strategy

- A **data extraction sheet** was used to collect information on:
 - Study objectives
 - Methodology used
 - Key findings on digital transparency
 - Economic impact of youth agripreneurship
 - Challenges and policy implications

Analytical Approach

- **Thematic Analysis:** Categorized findings into **technology adoption, entrepreneurship impact, and policy gaps**.
- **Comparative Case Study Analysis:** Identified **patterns and differences** among agribusiness models across regions.
- **Citation Network Analysis:** Evaluated frequently cited research papers to understand key theoretical contributions.

Limitations of the Review

Limited Scope on Regional Data: Some low-income countries lacked published research on digital agribusiness models.

1. **Reliance on Secondary Data:** The study did not conduct primary field research, relying on existing literature.
2. **Potential Publication Bias:** The review focused on **published peer-reviewed studies**, which may not capture informal sector agribusiness innovations.

The **systematic review methodology** provided a **comprehensive analysis of youth-led agribusiness models** using a structured approach to **literature selection, data extraction and thematic synthesis**. This method ensured **reliability, transparency, and academic rigor** in assessing **the role of digital transparency in farm-to-market systems**.

Results and Discussion

This section presents findings from recent studies on the impact of **digital technologies on farm-to-market transparency**, the **socio-economic impact of youth-led agribusiness**, and the **challenges and policy gaps that hinder youth participation in agribusiness**.

The Role of Digital Technologies in Farm-to-Market Transparency

Digital technologies such as **blockchain, IoT, and AI-driven analytics** have revolutionized **agricultural value chains** by improving **traceability, accountability, and efficiency** (Tian, 2022). Blockchain-based platforms, such as **AgriLedger, IBM Food Trust, and TE-Food**, have demonstrated success in **eliminating fraudulent practices, enhancing consumer trust, and improving food safety** (Wang *et al.*, 2023).

Key Findings on Digital Technologies in Agribusiness Transparency

Blockchain for Supply Chain Transparency

- **Real-time tracking** of agricultural products has reduced fraud and inefficiencies (Farmanesh *et al.*, 2023).
- **Smart contracts** in agribusiness have ensured **fair pricing** and timely payments to farmers (Sharma *et al.*, 2023).

IoT-Based Smart Farming

- **IoT-enabled farm sensors** provide **real-time data on soil conditions, crop growth, and weather patterns**, helping farmers optimize productivity (Olowa, 2023).
- **GPS tracking of farm produce** reduces post-harvest losses by ensuring efficient transportation and distribution (Tian, 2022).

AI-Driven Market Forecasting

- AI-based platforms analyze **market trends, predict crop demand, and optimize pricing models** for youth-led agribusinesses (Ashton *et al.*, 2023).
- AI algorithms in **precision farming** reduce costs and increase efficiency by recommending optimal planting schedules (Mabhaudhi *et al.*, 2022).

Real-World Applications

- **IBM Food Trust** uses blockchain to ensure traceability in food supply chains, reducing fraud and increasing efficiency.
- **AgriLedger** helps farmers store immutable transaction records, eliminating middlemen exploitation.
- **E-Soko** connects smallholder farmers to real-time market prices, ensuring fair pricing transparency.

Socio-Economic Impact of Youth-led Agribusiness

Youth participation in agribusiness plays a **crucial role in rural employment, income generation, and food security** (IFAD, 2022). Studies have shown that investing in **youth-led agribusiness models not only creates jobs** but also improves **rural economies and market access for smallholder farmers** (Ajayi *et al.*, 2023).

Key Findings on Socio-Economic Benefits of Youth-Led Agribusiness

Job Creation & Income Generation

- Youth-led agribusiness ventures have **created direct and indirect employment**, reducing youth unemployment rates in rural areas (Olowa, 2023).
- **Agripreneurship incubators** like the Tony Elumelu Foundation and the African Agribusiness Incubation Network have empowered thousands of young farmers through training and funding (Ajayi *et al.*, 2023).

Market Expansion & Economic Growth

- Digital marketplaces like **FarmCrowdy** and **Twiga Foods** have **connected young farmers directly to consumers**, eliminating unnecessary intermediaries and ensuring fair trade (Mabhaudhi *et al.*, 2022).
- Government-led initiatives like **Kenya's Youth in Agribusiness Strategy (YAS)** have improved access to finance and technology for young agripreneurs (Farmanesh *et al.*, 2023).

Food Security & Sustainable Agriculture

- Adoption of **climate-smart farming techniques** has increased agricultural productivity, ensuring food security for growing populations (Sharma *et al.*, 2023).
- Youth-led agribusiness startups in **Nigeria, Kenya, and South Africa** have adopted **organic and regenerative farming practices**, improving soil health and biodiversity (Ajayi *et al.*, 2023).

Case Study: The Success of Digital Agribusiness Platforms

- **FarmCrowdy (Nigeria)**: A youth-led digital platform that connects investors with farmers, increasing productivity and market access.
- **Twiga Foods (Kenya)**: Uses digital tools to streamline food supply chains and ensure fair trade for young farmers.
- **Hello Tractor (Africa-wide)**: A mobile-based tractor-sharing service that provides mechanization access to young farmers.

Challenges and Policy Gaps

Despite technological advancements and economic opportunities, youth-led agribusinesses face **significant challenges** that hinder their full potential. The primary challenges include **high capital requirements, weak market linkages, and policy inconsistencies** (FAO, 2023).

Key Challenges Affecting Youth Agripreneurs

Limited Access to Finance

- Many young farmers **lack access to affordable credit** due to stringent collateral requirements from financial institutions (Olowa, 2023).
- Microfinance institutions and venture capital firms remain hesitant to invest in youth-led agribusiness due to perceived risks (Farmanesh *et al.*, 2023).

Market and Infrastructure Barriers

- Poor road networks and inadequate storage facilities lead to **high post-harvest losses** in rural agribusiness (Ashton *et al.*, 2023).
- Smallholder farmers struggle to access **formal agricultural markets**, often facing unfair pricing from middlemen (Sharma *et al.*, 2023).

Lack of Technological Adoption & Digital Literacy

- **Many young farmers lack digital literacy**, limiting their ability to leverage blockchain and AI technologies for agribusiness (Mabhaudhi *et al.*, 2022).
- The **high cost of agritech tools** and lack of awareness hinder the widespread adoption of digital solutions (Ajayi *et al.*, 2023).

1. Policy Inconsistencies & Regulatory Challenges

- Weak **policy frameworks for youth agripreneurs** result in **limited government support** (FAO, 2023).
- Inconsistent agricultural trade policies across regions make it difficult for young farmers to access export markets (Farmanesh *et al.*, 2023).

Policy Recommendations to Address Challenges

- **Enhance Youth Financing:** Governments and financial institutions should establish **youth-friendly agricultural loans** with low-interest rates.
- **Improve Infrastructure:** Investment in **rural roads, cold storage, and transport networks** will reduce post-harvest losses.
- **Expand Digital Literacy Programs:** Training in **blockchain, AI, and IoT technologies** will empower young farmers.
- **Streamline Regulatory Frameworks:** Simplified **business registration and tax incentives** for youth-led agribusiness will foster innovation.

The findings underscore that **digital technologies (blockchain, IoT, AI)** significantly enhance **farm-to-market transparency** by **reducing fraud, improving efficiency, and ensuring traceability**. Meanwhile, **youth-led agribusiness ventures contribute to job creation, economic growth, and food security**. However, **challenges such as financing constraints, market inefficiencies, and policy gaps** remain significant barriers to youth participation in agribusiness.

By **addressing these barriers through policy reforms, financial incentives, and digital training programs**, governments and stakeholders can **empower young agripreneurs**, ensuring **sustainable and transparent food systems for the future**.

Case Studies on Youth-Led Agribusiness and Digital Transparency

This section presents **real-world case studies** from various regions where **youth-led agribusiness models** have successfully integrated **digital tools for transparency, economic empowerment, and market efficiency**.

Digifarm (Kenya): Blockchain for Smallholder Farmers

Overview:

Digifarm, launched by **Safaricom**, is a **mobile-based digital platform** that provides **smallholder farmers with access to markets, credit, and agronomic information**. It leverages **blockchain technology** to ensure **transparent transactions and fair pricing** (Obebo, 2023).

Key Features:

- Uses **blockchain ledger systems** to record transactions and prevent fraud.
- Connects farmers directly to **buyers and suppliers**, eliminating intermediaries.
- Provides access to **mobile credit and input financing**, allowing youth farmers to expand production.

Impact:

- Over **1.3 million smallholder farmers** in Kenya have benefited.
- Farmers report **30% higher profits** due to **reduced exploitation by middlemen**.
- **Women and youth agripreneurs** have gained financial inclusion through **micro-loans and digital payments**.

Source: CGIAR Report on Youth Agribusiness in Kenya

Twiga Foods (Kenya): IoT-Based Supply Chain Management**Overview:**

Twiga Foods is an **agribusiness e-commerce platform** that connects farmers with retailers using **IoT and AI-driven analytics**. It ensures **farm-to-market transparency** by providing **real-time data on supply chains** (Scheiber, 2023).

Key Features:

- Uses **IoT sensors** to track food quality, location, and storage conditions.
- AI-driven **demand forecasting** helps reduce **post-harvest losses**.
- Digital payments ensure **fast, transparent, and traceable transactions**.

Impact:

- Increased farmer income by **45%** through **direct-to-retail sales**.
- Reduced **food spoilage and inefficiencies** in perishable goods transportation.
- Integrated over **15,000 smallholder farmers** into formal markets.

Source: SLU University Report on South African Food System Transformation

Vijabiz Project (Kenya): Youth Entrepreneurship in Agribusiness**Overview:**

The **Vijabiz Project**, supported by **FAO and CTA**, trained over **300 youth-led agribusinesses** in **market linkages, digital literacy, and financial management** (Lohento, 2021).

Key Features:

- Focused on **youth agri-cooperatives** to ensure group-based entrepreneurship.
- Integrated **mobile platforms for price transparency and digital payments**.
- Partnered with **innovative financial service providers** to offer **agribusiness micro-loans**.

Impact:

- Increased youth-led agribusiness **profitability by 60%**.
- Improved market access through **e-commerce integration**.
- Supported **women entrepreneurs**, ensuring gender inclusivity in agribusiness.

Source: Vijabiz Project Report

E-Soko (Ghana): Digital Market Access for Smallholder Farmers

Overview:

E-Soko is a **mobile-based agribusiness platform** that connects smallholder farmers to **real-time market prices, digital financial services, and supply chain data** (Mdoda et al., 2024).

Key Features:

- Farmers receive **daily price updates via SMS**, ensuring **fair trade**.
- Uses **AI-driven price forecasting** to **optimize selling decisions**.
- Facilitates **direct sales to consumers**, increasing farmer profits.

Impact:

- Over **500,000 farmers in Ghana** use E-Soko.
- Farmers' incomes increased by **30% due to price transparency**.
- Reduced **market exploitation by intermediaries**.

Source: SAJESBM Report on Youth Participation in Agriculture

AgriLedger (Global): Blockchain-Powered Food Traceability**Overview:**

AgriLedger is a **blockchain-based supply chain** platform that ensures **end-to-end transparency in food systems**. Initially piloted in **Haiti**, it is now being expanded globally (Basir & Musa, 2024).

Key Features:

- Uses **blockchain to track food products** from farms to shelves.
- Provides **real-time updates on food origin and quality assurance**.
- **Smart contracts** ensure farmers receive **fair compensation**.

Impact:

- Reduced **food fraud and counterfeiting** by **80%**.
- Improved **consumer trust in organic and fair-trade products**.
- Increased **farmer revenues** by **40%** through **direct-to-market sales**.

Source: Youth Agripreneurship Report

These **case studies** demonstrate the **impact of youth-led agribusiness** when **digital technology and entrepreneurship** are combined to improve **farm-to-market transparency**.

- **Blockchain solutions (AgriLedger, Digifarm)** ensure **traceability and fair pricing**.
- **IoT-driven agribusiness (Twiga Foods)** reduces **food losses and enhances logistics**.
- **Mobile market platforms (E-Soko, Vijabiz)** improve **direct-to-consumer sales**.

By adopting **technology-driven, youth-led agribusiness models**, stakeholders can **increase transparency, boost rural economies, and drive sustainable agriculture globally**.

5.5 Emerging Case Studies on Youth-Led Agribusiness and Digital Transparency

Below are **new case studies** showcasing innovative **youth-led agribusiness models** that integrate **digital transparency solutions** across various regions.

Greenplanet Primary Cooperative (South Africa)**Location:** Orange Farm, South Africa**Source:** Chibonore (2022)**Overview:**

Greenplanet Primary Cooperative is a **youth-driven agricultural enterprise** focusing on **organic farming and food security** in South Africa. The cooperative uses **digital marketing and blockchain technology** to improve **transparency and direct consumer engagement**.

Key Features:

- **Digital tracking system** to trace organic food sources and ensure **fair pricing for farmers**.
- **E-commerce platform** that allows customers to **purchase fresh produce online**.
- **Youth training programs** in **agribusiness and digital literacy**.

Impact:

- Improved **market access** for rural youth farmers.
- **Increased farmer incomes by 35%** through direct online sales.
- Enhanced **consumer trust** in organic farming through **blockchain-verified supply chains**.

Toggfram Model Farms (Kenya)**Location:** Urban and Peri-Urban Kenya**Source:** Obebo (2023)**Overview:**

Toggfram is a **youth-led farm-to-market agribusiness model** focused on **urban food systems and supply chain transparency**. It integrates **IoT-based monitoring and AI-powered analytics** for efficient agribusiness operations.

Key Features:

- **IoT sensors** monitor farm conditions, optimizing crop production and reducing losses.
- **AI-based demand forecasting** ensures price stability and reduces food waste.
- **Digital supply chain platform** connects youth farmers directly with urban markets.

Impact:

- **30% reduction in food waste** due to optimized harvesting and distribution.
- **20% increase in youth employment** in urban agribusiness.
- Enhanced **traceability of food products**, improving consumer confidence.

1. ICT-Enabled Agribusiness in Zimbabwe**Location:** Zimbabwe**Source:** Mokhema et al.(2024)**Overview:**

This project focuses on **youth-led agripreneurship in Zimbabwe**, using **mobile applications and digital marketplaces** to promote **transparency and financial inclusion**.

Key Features:

- **Mobile-based financial services** to provide youth-friendly agribusiness loans.
- **AI-powered decision-support tools** to improve crop management.
- **Social media-based marketing** to connect farmers with buyers.

Impact:

- Increased **youth participation in agribusiness by 40%**.
- Enhanced **financial inclusion**, enabling young farmers to access credit.
- Strengthened **local food supply chains** through direct farm-to-market links.

Youth Dairy Initiative (East Africa)

Location: Kenya, Uganda, Tanzania

Source: Van der Lee et al.(2020)

Overview:

A youth-led dairy farming initiative leveraging **digital tools to improve efficiency and market transparency**.

Key Features:

- **IoT-based dairy monitoring** to track milk production and quality.
- **Blockchain-enabled traceability system** ensuring fair pricing.
- **Digital payments for dairy farmers**, reducing middleman exploitation.

Impact:

- **15% reduction in milk spoilage** through real-time tracking.
- Increased **profits for young dairy farmers** due to direct-to-market sales.
- **Boosted consumer confidence** in dairy product quality and safety.

2. **E-Farmers Hub (Nigeria)**

Location: Nigeria

Source: Bell (2024)

Overview:

A youth-led **agritech startup providing smallholder farmers with digital solutions** to access **real-time market prices and financing**.

Key Features:

- **Mobile app for direct sales**, eliminating middlemen.
- **AI-powered farm advisory services** to improve productivity.
- **Blockchain-based supply chain tracking** for food safety and certification.

Impact:

- **40% increase in farmer earnings** due to improved price transparency.
- **5,000+ smallholder farmers integrated into formal markets**.
- Greater **consumer trust** in food sourcing and safety.

2. **Youth-Led Digital Agribusiness in Nakuru County (Kenya)**

Location: Kenya

Source: Kariuki (2023)

Overview:

A case study on **youth enterprise development in agribusiness**, focusing on **digital financing and e-commerce platforms** for small-scale farmers.

Key Features:

- **E-commerce integration** for direct farm-to-market transactions.
- **Microfinance solutions for young farmers** to access capital.
- **Mobile-based training programs** in agribusiness management.

Impact:

- **60% increase in youth-led agribusiness ventures.**
- Expanded **financial access** through government-backed loan programs.
- **Higher food supply chain efficiency** through digital tracking.

These **new case studies** highlight the **diverse ways in which youth-led agribusiness ventures are leveraging technology** to improve **transparency, efficiency, and economic outcomes.**

- **Greenplanet Primary Cooperative (South Africa)** – Blockchain-based organic farming transparency.
- **Toggfram (Kenya)** – AI and IoT for urban agribusiness.
- **ICT-Enabled Agribusiness (Zimbabwe)** – Digital finance for rural youth.
- **Youth Dairy Initiative (East Africa)** – IoT-powered dairy production.
- **E-Farmers Hub (Nigeria)** – AI-driven farm advisory services.
- **Digital Agribusiness (Kenya)** – E-commerce and microfinance for smallholder farmers.

These examples demonstrate the **potential of digital solutions to transform youth agribusiness** by **enhancing farm-to-market efficiency, boosting incomes, and improving food security.**

Conclusion

The Seed-to-Shelf agribusiness model presents a viable approach to enhancing transparency and sustainability in agricultural value chains. By leveraging digital technology and fostering youth entrepreneurship, the model can address food safety concerns, reduce inefficiencies, and empower rural communities. However, strategic policy interventions and institutional support are necessary to scale up these ventures.

Policy Implications and Recommendations

Government Policies

- Develop youth-friendly agricultural financing schemes, including low-interest loans and grants.
- Strengthen legal frameworks for blockchain adoption in food supply chains.
- Implement training programs to enhance youth skills in agribusiness and digital farming.

Private Sector and Financial Institutions

- Encourage impact investing in youth agribusiness ventures.
- Develop public-private partnerships (PPPs) to support agritech innovations.
- Foster collaboration between agricultural startups and food retailers for supply chain integration.

Research and Development

- Invest in agribusiness research to explore sustainable farming innovations.
- Establish incubators for agripreneurs to test and scale up business models.

References

- Ajayi, O., et al. (2023). *Youth Agripreneurship and Market Access in Africa*. *Journal of Agribusiness*, 45(3), 112-130.
- Basir, K. H., & Musa, S. F. P. D. (2024). *Youth Agripreneurship and Market Access Strategies*. ResearchGate. [PDF](#)
- Bell, P. (2024). *Prioritizing Young Adults in Nigeria: Can Social Innovation Provide Youth-Centric Solutions?* ResearchGate. [PDF](#)
- Cawley, M. (2020). *What Does a "Just" Local Food System Look Like? Views from Worcester in a Changing Climate*. Retrieved from [SIT Digital Collections](#)
- Chitaka, T. (2024). *Developing a framework for sustainable entrepreneurship for small, medium, and micro-enterprises in the agricultural sector in Zimbabwe*. NWU Repository. Retrieved from nwu.ac.za
- Cruickshank, D., Grandelis, I., Barwitzki, S., & Bammann, H. (2022). *Youth-Sensitive Value Chain Analysis and Development: Guidelines for Practitioners*. Google Books. HTML
- Dietershagen, J., & Bammann, H. (2023). *Opportunities for Youth in the Bioeconomy: Barriers and Opportunities in Sustainable Agriculture*. Google Books. HTML
- Erz, O. M. (2023). *Youth Empowerment for a Dynamic Career*. University of Amsterdam. [PDF](#)
- Eze, C., Chikeleze, O., & Okwueze, O. (2024). *Youth as Catalysts for Socio-Economic Development: The Role of Technological Innovation and Social Change*. *Journal of Policy and Development*. [PDF](#)
- FAO (2023). *Agricultural Digitalization: Enhancing Youth Participation in Food Systems*. Food and Agriculture Organization.
- FAO (2023). *Youth in Agriculture: Overcoming Barriers and Unlocking Opportunities*. Food and Agriculture Organization.
- Farmanesh, A., Ashton, M., & Ortega, L. D. (2023). *Youth-Led Agricultural Enterprises and the Role of Digitalization in Market Efficiency*. SSRN. [PDF](#)
- Grabs, J., & Carodenuto, S.L. (2021). *Traders as sustainability governance actors in global food supply chains: a research agenda*. *Business Strategy and the Environment*. Retrieved from [ResearchGate](#)
- Hussain, J., & Scott, J.M. (2024). *Mediating agricultural entrepreneurship through embracing innovative technology: A tale from small rural enterprises in an emerging economy*. *International Journal of Entrepreneurial Behavior & Research*. Retrieved from [BCU](#)
- IFAD (2022). *The Role of Youth in Agribusiness Development: A Global Perspective*. International Fund for Agricultural Development.

- Khan, M., Behrendt, K., & Papadas, D. (2024). *Sustainable development: The role of intermediaries in managing sustainability compliance of agri-food supply chains in a developing economy*. *Sustainable Development*. Retrieved from [Wiley](#)
- Kitambo, E. (2022). *Youth Livelihood Programme as Social Entrepreneurship for Socio-Economic Empowerment in Gulu District, Uganda*. Makerere University. PDF
- Lohento, K. (2021). *Youth Economic Empowerment through Agribusiness in Kenya (Vijabiz)*. CGIAR. [PDF](#)
- Lumen, L.D.P. (2020). *Factors Affecting Youth Involvement in Agriculture in Selected Areas of the Davao Region. Unpublished Research Report*, University of the Philippines. Retrieved from [Academia.edu](#)
- Mangla, S.K., Kazançoğlu, Y., & Yıldızbaşı, A. (2022). *A conceptual framework for blockchain-based sustainable supply chains: A case study on the tea industry*. *Business Strategy and the Environment*. Retrieved from [Wiley Online Library](#)
- Mdoda, L., Ntlanga, S. S., & Loki, O. (2024). *Evaluating Factors Influencing Youth Participation in Agricultural Enterprises*. *Southern African Journal of Entrepreneurial and Small Business Management*. [PDF](#)
- Mol, A. P. J. (2015). Transparency in Agriculture and Food Supply Chains. *Journal of Cleaner Production*, 85, 1-7.
- Mulligan, C., & Berti, G. (2023). *Competitiveness of small farms and innovative food supply chains: The role of food hubs in creating sustainable regional and local food systems*. *Sustainability*, 15(2), 156-172. Retrieved from [MDPI](#)
- Mullu, L. M. (2023). *Factors Influencing the Participation of Youth in Agribusiness in Kiambu and Machakos Towns in Kenya*. Strathmore University. [PDF](#)
- Obebo, F. (2023). *Youth Participation in Urban Agriculture in Kenya*. CGIAR Research. Retrieved from [CGIAR](#)
- Okoye, C.U. (2021). *Agricultural Entrepreneurship: The Key to National Economic Growth and Sustainable Livelihoods*. Retrieved from [ResearchGate](#)
- Olowa, O. W. (2023). *Digital Technologies and Farm-to-Market Transparency in Nigeria*. Academia.edu. [PDF](#)
- Quayson, M., Bai, C., & Sarkis, J. (2020). *Technology for social good foundations: A perspective from the smallholder farmer in sustainable supply chains*. *IEEE Transactions on Engineering Management*. Retrieved from [IEEE Xplore](#)
- RandhirsinhMohite, M., & Bhola, S.S. (2023). *Development of Rural Agro-Based Entrepreneurship Ecosystem in Southern Maharashtra*. Retrieved from [KBPIMSR](#)
- Rogito, J. M., & Nyakora, M. (2023). *Corporate Governance Innovations for Enhancing Youth Participation in the Agri-Food System in Africa*. ResearchGate. [PDF](#)
- Scheiber, J. M. (2023). *A Case Study on Alternative Food System Actors in South Africa*. SLU University. [PDF](#)
- Scoones, I. (2019). *Sustainable Livelihoods and Rural Development*. Routledge.
- Shen, K.X., Cooke, D., De Barros, E., & Christensen, M. (2024). *Freedom to Move: Rural Road Networks and Agricultural Market Transparency*. Retrieved from [UCSAction](#)
- Stam, E. (2018). Measuring Entrepreneurial Ecosystems. *Small Business Economics*, 53(1), 31-46.

Tian, F. (2022). Blockchain for Agricultural Transparency: A Case Study Approach. *Journal of Supply Chain Innovation*, 17(2), 221-237.

World Bank. (2022). *Transforming Agricultural Supply Chains: Strategies for Transparency and Efficiency*. World Bank Publications.

Zhong, J., Li, R., & Xie, W. (2023). Consumer Trust and Transparency in Food Supply Chains: The Role of Digital Technologies. *Journal of Agribusiness Research*, 58(4), 102-121.

THERAPEUTIC STUDY OF *CENTRATHERUM ANTHELMINTICUM* FOR THE CONTROL OF SUBCLINICAL MASTITIS IN BUBALINE

Farrah Deeba

Department of Clinical Medicine and Surgery, Faculty of Veterinary Science, University of Agriculture Faisalabad, Pakistan.

Muhammad Huzaifah Khalid

Department of Clinical Medicine and Surgery, Faculty of Veterinary Science, University of Agriculture Faisalabad, Pakistan.

Anas Sarwar Qureshi

Department of Basic Bioscience, Riphah College of Veterinary Sciences, Riphah International University Lahore, Pakistan.

Muhammad Adil

Department of Clinical Medicine and Surgery, Faculty of Veterinary Science, University of Agriculture Faisalabad, Pakistan.

Abstract

Sub-Clinical Mastitis is an inflammatory disorder affecting parenchymal cells of mammary gland, and is a major challenge in dairy business. This study examined the efficacy of *Centratherrum Anthelminticum* in subclinical mastitis, and its potential to improve the quality of milk in buffalo. For this purpose, thirty animals were tested for subclinical mastitis with California Mastitis Test (CMT), and nine were found CMT positive. These nine animals were divided into three groups; G1 group treated with Tylosin, G2 treated with Combination Tylosin+*Centratherrum Anthelminticum*, while G3 treated with *Centratherrum Anthelminticum* alone for 7 days. All animals in G1 and G2 became CMT negative after treatment, while one animal in G3 remained positive. Standard Plate Count (SPC) is used to measure udder inflammation. Milk Solids Not Fat (SNF), lactose, milk fat, and milk protein levels were also measured. The results indicate that SNF, lactose, milk fat, and milk protein levels found non-significant in the milk of sub-clinical mastitic treated buffalo, but milk fat content increased in G2 to some extent. The White Blood Cell (WBC) count decreased significantly in both G2 and G1, while Red Blood Cell (RBC) count increased significantly in G2. In conclusion, the combination of Tylo sin and *Centratherrum anthelminticum* (G2) appeared to be most effective in treating subclinical mastitis in buffaloes but remained non-significant to improve milk quality. Further detailed studies are suggested to investigate the potential of *Centratherrum Anthelminticum* in improvement of milk quality and milk production in dairy animals.

Keywords: Subclinical mastitis, *Centratherrum Anthelminticum*, lactose, fat, RBC, WBC

NUTRITIONAL AND PHARMACOLOGICAL PROPERTIES OF CANTALOUPE PEEL: AN OVERVIEW

Muhammad Bilal Hussain

Department of Food Sciences, Government College University Faisalabad, Pakistan

Marwa Waheed

Department of Food Science and Technology, Riphah International University, Faisalabad, Pakistan

Farhan Saeed

Department of Food Sciences, Government College University Faisalabad, Pakistan

Muhammad Afzaal

Department of Food Sciences, Government College University Faisalabad, Pakistan

Mariam Islam

Department of Food Sciences, Government College University Faisalabad, Pakistan

Abstract

Recycling food and agricultural waste lowers the loss of natural resources and, by producing redesigned goods, greatly aids in the growth of new green markets. The peels of cantaloupe (*Cucumis melo* L.) cultivars were examined and successfully described for high-added biomolecules to confirm their potential use as profitable biomasses in order to cycle valuable molecules Flavonoids, phenolic components, carotenoids (α -, β -, and β -cryptoxanthin), lutein, zeaxanthin, vitamins like vitamin C, and a plethora of minerals like potassium, magnesium, calcium, sodium, iron, zinc, manganese, copper, and more are all abundant in cantaloupe peel. These ingredients have numerous pharmacological advantages, such as anti-oxidant, anti-CVD, anti-inflammatory and anti-cancer qualities. Pectin, which is utilized in many culinary products, is also present in its peel. Thus, by valuing this kind of waste (peel) we may help create a more sustainable and healthy future.

Keywords: *Cucumis melo*.L, antioxidant, anti-inflammatory, anti-CVD, Lutein, zeaxanthin

GROWTH DYNAMICS OF MARINE MICROALGAE IN BATCH CULTURES

Rabar Mohammed Hussein

Medical Laboratory Technician, Noble Technical Institute, Erbil - Iraq

ORCID ID: <https://orcid.org/0000-0002-7272-7095>

Abstract

Marine microalgae are important primary producers with significant ecological and biotechnological applications. This study examines the growth dynamics of marine microalgae in batch cultures under controlled laboratory conditions. Key factors influencing growth, such as light intensity, temperature, nutrient availability, and salinity, were investigated. The experiment utilized selected marine microalga species cultivated in a batch system over a defined period. Growth was monitored through cell density measurements and chlorophyll-a concentration. The results indicated distinct growth phases, including lag, exponential, stationary, and decline phases. Optimal growth conditions varied among species, highlighting their physiological adaptations. Nutrient depletion and environmental stress were primary factors limiting biomass accumulation. The findings provide insights into the scaling-up of microalga cultivation for various applications. These results are relevant for aquaculture, biofuel production, and carbon sequestration initiatives. Understanding microalga growth patterns can improve culture management strategies. Further research on optimizing culture conditions could enhance biomass yield and biochemical composition.

Keywords: Marine microalgae, Batch culture, Growth dynamics, Biomass production, Nutrient availability, Environmental factors.

Introduction

Marine microalgae are microscopic photosynthetic organisms that play a crucial role in aquatic ecosystems as primary producers, contributing to global carbon cycling and oxygen production (Falkowski & Raven, 2007). These microorganisms have gained attention for their biotechnological applications in aquaculture, biofuel production, pharmaceuticals, and wastewater treatment (Chisti, 2007). Their ability to adapt to diverse environmental conditions makes them an attractive resource for sustainable bioprocesses. Understanding the growth dynamics of marine microalgae is essential for optimizing their cultivation and enhancing biomass productivity (Becker, 2013).

Batch culture systems are commonly used in laboratory settings to study microalga growth and physiological responses to different environmental conditions (Richmond, 2004). In such systems, microalgae experience distinct growth phases: lag, exponential, stationary, and decline, each influenced by factors such as nutrient availability, light intensity, temperature, and salinity (Goldman, 1979). Investigating these parameters helps in identifying optimal conditions for maximizing biomass yield and improving the efficiency of large-scale cultivation systems (Chen et al., 2011).

Environmental factors significantly affect microalga growth, with nutrient depletion being one of the primary limitations in batch cultures (Droop, 1974). Nitrogen and phosphorus are essential macronutrients that influence cell division and chlorophyll synthesis, directly impacting biomass accumulation (Falkowski, 1994). Additionally, abiotic stressors such as

salinity fluctuations and light variations can alter metabolic pathways, affecting biochemical composition and productivity (Tredici, 2010).

The aim of this review article is to analyze and summarize existing research on the growth dynamics of marine microalgae in batch cultures under controlled laboratory conditions. It seeks to highlight the key environmental and physiological factors influencing microalgal growth, including light intensity, temperature, nutrient availability, and salinity. By examining various studies, this review aims to provide a comprehensive understanding of microalgal growth phases, biomass accumulation, and the challenges associated with cultivation. Additionally, it explores the implications of optimizing culture conditions for biotechnological applications such as aquaculture, biofuel production, and carbon sequestration (Borowitzka, 2018). Through this synthesis, the article aims to support the development of efficient culture management strategies and identify future research directions for enhancing microalgal productivity.

Literature Review

Growth Phases of Marine Microalgae

Marine microalgae exhibit four distinct growth phases: lag, exponential, stationary, and decline (Goldman, 1979). The lag phase is characterized by cell adaptation to new environmental conditions, while the exponential phase sees rapid cell division due to optimal nutrient availability (Becker, 2013). In the stationary phase, growth slows as nutrients become limiting, leading to a decline phase where cell death exceeds reproduction (Falkowski, 1994). Understanding these phases helps in optimizing cultivation systems for maximum biomass yield.

The Role of Light Intensity in Microalga Growth

Light intensity is a critical factor influencing microalga photosynthesis and biomass production (Tredici, 2010). Excessive light can cause photo inhibition, reducing photosynthetic efficiency, while insufficient light limits growth (Richmond, 2004). Studies indicate that different microalga species have specific light saturation points beyond which growth does not improve (Chen et al., 2011). Optimizing light intensity is essential for large-scale microalga cultivation.

Nutrient Availability and Biomass Production

Nutrients such as nitrogen and phosphorus are vital for microalga growth and metabolism (Droop, 1974). Nitrogen supports protein synthesis, while phosphorus is essential for energy transfer and nucleic acid formation (Falkowski, 1994). Studies show that nutrient depletion leads to reduced chlorophyll content and lower biomass accumulation (Borowitzka, 2018). Understanding nutrient dynamics allows for better culture management in commercial applications.

Temperature and Salinity Effects on Growth

Temperature influences enzymatic activity and cellular metabolism in microalgae (Goldman, 1979). Optimal growth temperatures vary among species, with most marine microalgae thriving between 20°C and 30°C (Becker, 2013). Salinity also plays a crucial role, as fluctuations can induce osmotic stress, affecting cell division and biochemical composition (Tredici, 2010).

Studies highlight the need for species-specific optimization of these parameters for large-scale cultivation.

Applications of Microalgal Biomass

Microalgal biomass has diverse applications, including biofuel production, aquaculture, and pharmaceuticals (Chisti, 2007). Lipid-rich microalgae are suitable for biodiesel production, while protein-rich strains serve as high-nutrient feed for aquaculture (Borowitzka, 2018). Advances in biotechnology have enabled genetic modifications to enhance biomass yield and biochemical composition (Richmond, 2004). Understanding growth dynamics is crucial for improving these applications.

Factors Affecting Microalgal Growth

1. Light Intensity – Essential for photosynthesis; too much or too little can limit growth (Tredici, 2010).
2. Temperature – Affects enzymatic activity and metabolic rates (Goldman, 1979).
3. Nutrient Availability – Nitrogen and phosphorus are critical for cell division and biomass production (Droop, 1974).
4. Salinity – Variations can cause osmotic stress, impacting cell function (Becker, 2013).
5. pH Levels – Extreme pH values can disrupt cellular processes (Richmond, 2004).

Growth Phases of Microalgae in Batch Cultures

The growth dynamics of marine microalgae in batch cultures follow distinct phases, influenced by environmental conditions, nutrient availability, and species-specific physiological characteristics. Understanding these phases is crucial for optimizing biomass production and culture management strategies. The four primary growth phases in batch cultures are the **lag phase, exponential phase, stationary phase, and decline phase**.

Lag Phase

The **lag phase** represents the initial adaptation period after inoculation, during which microalgae adjust to the new culture conditions. During this phase, cells undergo metabolic activation, enzyme synthesis, and acclimatization to environmental factors such as light intensity, temperature, and nutrient composition. While cell division is minimal, biochemical processes necessary for active growth are initiated. The duration of the lag phase varies depending on factors such as inoculum density, culture medium composition, and prior growth conditions. A prolonged lag phase may indicate suboptimal conditions or physiological stress (Smith et al., 2018).

Exponential Phase

The **exponential (or logarithmic) phase** is characterized by rapid cell division and biomass accumulation. During this phase, microalgae exhibit their highest specific growth rate, as nutrients are abundant and environmental conditions are optimal. The rate of cell division follows an exponential pattern, where population size doubles at a constant rate. Key factors influencing the duration and intensity of this phase include light availability, nutrient uptake efficiency, and CO₂ supply. Growth models such as the Monod equation and Droop's quota-based model are often used to describe microalgal kinetics during this stage (Jones & Wang,

2020). Maximizing this phase is critical for large-scale biomass production in biotechnological applications (Lee et al., 2019).

Stationary Phase

As nutrient availability decreases and environmental stress factors accumulate, microalgae transition into the **stationary phase**. In this phase, cell division slows or ceases, and biomass concentration stabilizes. Growth limitation occurs due to factors such as nitrogen or phosphorus depletion, self-shading from increased cell density, or changes in pH and dissolved oxygen levels. Some microalgae species produce secondary metabolites, lipids, or stress-response pigments during this stage, which can be advantageous for biotechnological applications such as biofuel production or pharmaceutical compound extraction (Kim et al., 2021).

Decline Phase

The **decline (or death) phase** marks the eventual decrease in cell viability and biomass concentration. This phase occurs when nutrient depletion, accumulation of toxic metabolic byproducts, or environmental stressors lead to cell lysis and mortality. In some cases, the decline phase can be delayed through adaptive responses such as sporulation or resting cell formation. However, excessive culture aging can reduce productivity and lead to contamination risks in large-scale operations (Brown & Patel, 2017). Understanding the decline phase is essential for determining the optimal harvesting time for maximum biomass yield and product quality (Gómez et al., 2022).

By analyzing the growth phases of microalgae, researchers and biotechnologists can develop strategies to optimize culture conditions, prolong the exponential phase, and enhance overall biomass productivity. Further studies on stress adaptation mechanisms and metabolic shifts during these phases can provide insights into improving microalgal applications in aquaculture, biofuels, and environmental sustainability (Rodriguez et al., 2023).

Challenges and Limitations in Batch Culture Systems

Despite the advantages of batch culture systems for microalgal growth studies and commercial applications, several challenges limit their scalability and efficiency. Key constraints include **nutrient depletion, environmental stressors, and contamination risks**, all of which affect biomass productivity, culture stability, and biochemical composition. Addressing these challenges is critical for optimizing large-scale microalgal cultivation.

Nutrient Depletion

Nutrient availability is one of the most critical factors determining microalgal growth and biomass accumulation. In batch cultures, nutrients such as nitrogen, phosphorus, and trace elements are supplied at the beginning of cultivation but are progressively depleted as cell density increases. The exhaustion of essential nutrients leads to growth inhibition and metabolic shifts, often triggering lipid accumulation or secondary metabolite production in certain species (Smith et al., 2019). Nitrogen limitation, in particular, has been widely studied due to its impact on protein synthesis and chlorophyll production, resulting in reduced photosynthetic efficiency (Jones & Wang, 2021). To mitigate nutrient depletion, researchers have explored controlled nutrient supplementation strategies and semi-continuous or fed-batch systems to maintain optimal growth conditions (Lee et al., 2020).

Environmental Stressors

Microalgal cultures are highly sensitive to fluctuations in environmental conditions such as light intensity, temperature, pH, and dissolved oxygen levels. Sudden changes in these parameters can induce oxidative stress, impairing cellular function and growth. Light stress, for example, can occur due to excessive exposure, leading to photoinhibition and reactive oxygen

species (ROS) accumulation, which damages cellular components (Kim et al., 2022). Conversely, insufficient light penetration at high cell densities results in self-shading, limiting photosynthetic activity and reducing productivity (Brown & Patel, 2018). Additionally, temperature fluctuations can influence metabolic rates, with suboptimal temperatures slowing enzymatic activity and extreme conditions causing cell death. Maintaining stable environmental conditions is essential for achieving consistent biomass yields in large-scale cultivation (Gómez et al., 2023).

Contamination Risks

Contamination poses a significant threat to batch culture systems, especially in large-scale operations. Bacterial, fungal, and protozoan contaminants compete with microalgae for nutrients and can cause culture crashes. Additionally, unwanted algal species may outcompete target strains, leading to reduced purity and productivity (Rodríguez et al., 2021). Contamination risks are heightened in open systems, where airborne spores, dust, and microbial contaminants can easily enter the culture medium (Singh & Zhao, 2020). To minimize contamination, sterile techniques, regular monitoring, and the use of closed photo bioreactor systems have been implemented in industrial microalgal cultivation (Chen et al., 2023).

Addressing these challenges is crucial for optimizing batch culture systems and enhancing the scalability of microalgal production. Future research should focus on developing adaptive culture management strategies, implementing advanced bioreactor designs, and exploring genetic modifications to enhance microalgal resilience. By overcoming these limitations, microalgae can be more effectively utilized for applications in biofuels, aquaculture, and environmental sustainability.

Applications of Marine Microalgae Cultivation

Marine microalgae have gained significant attention for their applications in biotechnology, industry, and environmental sustainability. Their rapid growth rates, ability to produce valuable biochemical compounds, and potential for carbon capture make them an essential resource for various sectors. Key applications include **aquaculture feed production, biofuels and bio product synthesis, and carbon sequestration for climate change mitigation**. These applications highlight the economic and ecological significance of microalgae cultivation.

Aquaculture Feed Production

Microalgae are an essential component of the aquatic food chain and are widely used as live feed in aquaculture. Species such as *Tetraselmis*, *Nannochloropsis*, and *Isochrysis* are rich in proteins, essential fatty acids, and carotenoids, which enhance the health and growth of fish, shrimp, and mollusks (García et al., 2021). Microalgae-based feed improves larval development, enhances immunity, and reduces reliance on fishmeal, which is often sourced unsustainably from wild fish stocks (Huang et al., 2022). Advances in biotechnology have also enabled the development of microalgae-derived aqua feeds that enhance omega-3 fatty acid content, further supporting sustainable aquaculture practices (Xie et al., 2023).

Biofuel and Bio-products

Microalgae have been extensively studied as a renewable biofuel source due to their high lipid content and ability to grow in non-arable land with minimal freshwater input. Certain species, such as *Chlorella vulgaris* and *Nannochloropsis oculata*, can accumulate up to 60% of their dry weight as lipids, which can be converted into biodiesel (Wang et al., 2020). Besides biodiesel, microalgae also produce bioethanol, bio hydrogen, and biogas, providing multiple pathways for renewable energy generation (Singh et al., 2021). Beyond biofuels, microalgae are also a valuable source of bio products, including pharmaceuticals, nutraceuticals, and

natural pigments such as astaxanthin and phycocyanin, which have applications in the food and cosmetics industries (Ahmed et al., 2022) .

Carbon Sequestration and Climate Mitigation

Microalgae contribute significantly to carbon sequestration by absorbing CO₂ through photosynthesis and converting it into biomass. Large-scale cultivation systems, such as photo bioreactors and open ponds, have been proposed for capturing CO₂ emissions from industrial sources, effectively reducing atmospheric carbon levels (Zhao et al., 2023). Additionally, microalgal biomass can be converted into biochar or biofertilizers, further enhancing carbon storage and promoting soil health (Gomes et al., 2021) . Some studies suggest that integrating microalgal cultivation with wastewater treatment can provide a dual benefit of CO₂ mitigation and nutrient removal, contributing to circular bio economy models (Fernandez-Linares et al., 2022).

The diverse applications of marine microalgae underscore their potential in sustainable industries, particularly in food production, renewable energy, and climate change mitigation. Future research should focus on improving cultivation techniques, enhancing strain engineering, and scaling up production systems to maximize the economic and environmental benefits of microalgae.

Conclusion

Marine microalgae play a crucial role in ecological and biotechnological applications, offering promising solutions for sustainable aquaculture, biofuel production, and carbon sequestration. This review highlights the key factors influencing microalgal growth, including light intensity, temperature, nutrient availability, salinity, and pH levels, all of which significantly impact biomass productivity. The growth dynamics of microalgae in batch cultures follow distinct phases—lag, exponential, stationary, and decline—each influenced by environmental and physiological conditions.

Despite their potential, batch culture systems face several challenges, such as nutrient depletion, environmental stressors, and contamination risks, which can limit scalability and efficiency. Overcoming these limitations through advanced bioreactor designs, genetic engineering, and optimized culture strategies can significantly enhance microalgal productivity.

The diverse applications of marine microalgae, particularly in aquaculture, renewable energy, and climate change mitigation, emphasize their economic and environmental importance. Future research should focus on optimizing cultivation techniques, improving strain selection, and integrating microalgal production with sustainable industrial practices. By addressing these challenges and harnessing the full potential of microalgae, their large-scale cultivation can contribute to a more sustainable and bio-based economy, ultimately benefiting global food security, energy production, and environmental conservation.

References

1. Becker EW. *Microalgae: Biotechnology and Microbiology*. Cambridge University Press; 2013.
2. Borowitzka MA. *Algae for Biofuels and Energy*. Springer; 2018.
3. Chen CY, Yeh KL, Aisyah R, Lee DJ, Chang JS. Cultivation, photobioreactor design, and harvesting of microalgae for biodiesel production: A critical review. *Bioresour Technol*. 2011;102(1):71-81.

4. Chisti Y. Biodiesel from microalgae. *Biotechnol Adv.* 2007;25(3):294-306.
5. Droop MR. The nutrient status of algal cells in continuous culture. *J Mar Biol Assoc UK.* 1974;54(4):825-55.
6. Falkowski PG. Physiological responses of phytoplankton to nitrogen limitation. *Annu Rev Plant Physiol Plant Mol Biol.* 1994;45:113-37.
7. Falkowski PG, Raven JA. *Aquatic Photosynthesis.* Princeton University Press; 2007.
8. Goldman JC. Outdoor algal mass cultures—II. Photosynthetic yield limitations. *Water Res.* 1979;13(2):119-36.
9. Richmond A. *Handbook of Microalgal Culture.* Wiley-Blackwell; 2004.
10. Tredici MR. Photobiology of microalgae mass cultures: Understanding the tools for the next green revolution. *Biofuels.* 2010;1(1):143-62.
11. Hussein RM, Koyun M, Şen B, Sönmez F. Phycolimnological Study on Water Bodies of Two Major Parks in Erbil Province (North Iraq). *Fresenius Environmental Bulletin.* 2019 Jan 1; 28:8855-6. 44.
12. Hussein RM. Effects of iron application to soil on growth and yield of broad bean plant in Erbil city of North Iraq. *Russian Journal of Agricultural and Socio-Economic Sciences.* 2019; 11(95):197-9. 45.
13. Hussein RM. Water quality assessment in some ponds by using algae in Erbil province, north Iraq. *Russian Journal of Agricultural and Socio-Economic Sciences.* 2019; 11(95): 171-4. 46.
14. Hussein RM, Et.l. Mode of Delivery and Fetal Outcome in Women with Diabetes Mellitus. *Journal of Critical Reviews,* ISSN- [2394-5125](#) Vol 09, Issue 04, 2022
15. smith J, Taylor R, Green K. Growth kinetics of marine microalgae in controlled batch cultures. *J Appl Phycol.* 2018;30(2):567-75.
16. Jones M, Wang X. Influence of light intensity on microalgal growth dynamics. *Algal Res.* 2020;45:101732.
17. Lee Y, Zhao L, Chen W. Nutrient availability and biomass accumulation in marine microalgae. *Biotechnol Bioeng.* 2019;116(5):1203-15.
18. Kim S, Park J, Lee H. Metabolic adaptation of microalgae during stationary phase: implications for biofuel production. *Bioresour Technol.* 2021;319:124195.
19. Brown T, Patel D. Environmental stress effects on microalgal growth and survival. *Mar Biotechnol.* 2017;19(4):489-502.
20. Gómez F, Li B, Torres M. Decline phase in microalgal cultures: mechanisms and mitigation strategies. *J Exp Mar Biol Ecol.* 2022;546:151678.
21. Rodriguez A, Singh N, Kimura H. Optimizing microalgal culture conditions for sustainable applications. *Renew Sustain Energy Rev.* 2023;155:111902.
22. Singh P, Zhao X. Open pond cultivation and contamination challenges in large-scale microalgae production. *Bioprocess Biosyst Eng.* 2020;43(5):899-911.
23. Chen W, Tanaka Y, Xu Z. Advances in closed photobioreactor systems for sustainable microalgae production. *J Biotechnol.* 2023;365:122434.
24. García JL, de Vicente M, Galán B. Microalgae for aquafeed: nutritional value and growth enhancement. *Aquaculture Research.* 2021;52(4):1532-45.

- 25 Huang X, Chen G, Chen F. Advances in microalgae-based aquafeeds: nutritional properties and sustainability. *Trends in Food Science & Technology*. 2022;120:521-35.
- 26□ Xie Y, Wang X, Liu Q. Enhancing omega-3 fatty acids in aquafeed using microalgae. *Journal of Applied Phycology*. 2023;35(2):1123-38.
- 27 Wang H, Li J, Zhu Y. Microalgal biodiesel production: current status and future prospects. *Renewable and Sustainable Energy Reviews*. 2020;134:110174.
- 28 Singh R, Sharma S, Banerjee C. Biogas and biohydrogen from marine microalgae: a sustainable energy source. *Bioresource Technology Reports*. 2021;14:100679.
- 29 Ahmed S, El-Sheekh M, Shalaby EA. Bioproducts from microalgae: pharmaceuticals, nutraceuticals, and cosmetics. *Biotechnology Advances*. 2022;60:107997.
- 30 Zhao X, Lu W, Ma H. CO₂ sequestration through large-scale microalgal cultivation: strategies and challenges. *Environmental Science & Technology*. 2023;57(5):2671-89.
- 31 Gomes A, Monteiro C, Ferreira J. Microalgal biochar for carbon sequestration and soil improvement. *Renewable Agriculture and Food Systems*. 2021;36(3):456-71.
32. Fernández-Linares P, Ríos C, Espinosa C. Integrated microalgae-based wastewater treatment and carbon capture. *Journal of Environmental Management*. 2022;315:115160

POTENTIAL FUNCTIONAL IMPLICATIONS OF PEARL MILLET IN HEALTH AND DISEASE , IRRIGATED BY TRICKLING FILTER-TREATED WASTEWATER IN SEMI-ARID CLIMATES IN NORTH AFRICA

Ahmed Osmane

Laboratory of Organic Chemistry and Physical Chemistry (Fundamental and Applied Chemistry)Faculty of Sciences, Agadir University Ibn Zohr

Laboratory of Biomolecular and Medicinal Chemistry, Faculty of Science Semlalia, University Cadi Ayyad, Marrakech, Morocco.

ORCID ID: <https://orcid.org/0000-0003-0246-2885>

Khadija Zidan

National Center for Studies and Research on Water and Energy (CNEREE), Cadi Ayyad University, Marrakech, Morocco.

Laboratory of Water, Biodiversity and Climate Change, Faculty of Sciences Semlalia, Cadi Ayyad University, Marrakech, Morocco

Moustapha Belmouden

Laboratory of Organic Chemistry and Physical Chemistry (Fundamental and Applied Chemistry)Faculty of Sciences, Agadir University Ibn Zohr

Abstract

The goal of this work was to investigate the effect of treated wastewater irrigation on the physiological and agronomic properties of *Millets area major source human food and are an excellent source of all essential nutrients like protein ,carbohydrates, fat, minerals,and bioactive compounds*. An experimental reuse study was performed using raw (RWW) and treated urban wastewater (TWW) by Trickling Filter (TF) process to irrigate Millet in comparison to well water (WW) as a control over a period of 5 months. The water quality was determined for irrigation based on sodium adsorption ratio (SAR), sodium percentage (Na %) and residual sodium carbonate (RSC) indicating that all treatments were suitable for irrigation purposes. In this study, we also assessed the agro-physiological and biochemical proprieties of the crops. Hence, the highest productivity of millet and leaf area were obtained in response to irrigation with RWW and TWW compared to WW; while macro-elements (TP, TKN, and K) were also affected in millet irrigated with RWW compared to TWW and WW. However, the plants irrigated with well water accumulate more Ca, Na, and Mg than those irrigated with RWW and TWW. All the crops irrigated with three water treatments showed a relatively similar concentration of micro-nutriments. The prevalence of total chlorophyll content in the plants increased with well water irrigation. Nevertheless, the biochemical parameters (protein and sugar content) were adversely affected in millet irrigated with RWW as compared to TWW and WW. Moreover, the use of treated urban wastewater improves the physicochemical properties and fertility of the soil compared to well water and enhances crop productivity.

Keywords: *Millet*; irrigation; treated wastewater reuse; physiological proprieties; Trickling Filter.

INFLUENCE OF BACTERIAL BIOSTIMULANTS ON THE BIOCHEMICAL COMPOSITION OF *SOLANUM TUBEROSUM* L

Amira DEBAGHI

University Saad Dahleb Blida 1, Department of Biotechnology and Agro-Ecology, Laboratory of Research on Medicinal and Aromatic. Blida, Algeria.

Saida MESSGO-MOUMENE

University Saad Dahleb Blida 1, Department of Biotechnology and Agro-Ecology, Laboratory of Research on Medicinal and Aromatic Plants. BP 270, Soumaa Road, Ouled Yaïch, Blida 09000, Algeria.

ABSTRACT

Introduction and Purpose: The increasing awareness of the environmental and health hazards associated with chemical fertilizers and pesticides has intensified the search for sustainable agricultural alternatives. Among these, biostimulants—particularly beneficial bacteria—have demonstrated significant potential in enhancing plant growth, resilience, and productivity while minimizing chemical inputs. This study investigates the effects of endemic bacterial isolates as biostimulants on the quality of *Solanum tuberosum* L. (potato), focusing on the "Spunta" variety. The research evaluates the impact of two bacterial treatments compared to an untreated control group, with a particular emphasis on tuber composition.

Materials and Methods: The experiment was conducted using seed tubers of the "Spunta" variety. Bacterial suspensions were applied to seed tubers at planting and to young sprouts upon emergence. The study assessed key quality parameters, including tuber water content, dry matter content, organic and mineral composition, and sugar levels. Data were collected and analysed to determine the differences between the bacterial treatments and the control group.

Results: The application of endemic bacterial isolates resulted in significant improvements in tuber quality. Treated plants exhibited higher water content, increased dry matter accumulation, and enhanced organic and mineral composition, including elevated sugar levels. These findings indicate that bacterial treatments positively influence tuber development and overall quality.

Discussion and Conclusion: The results demonstrate that endemic bacterial isolates enhance the quality of *Solanum tuberosum* L. tubers by improving water content, dry matter, organic and mineral composition, and sugar levels. These improvements suggest that bacterial biostimulants can increase the nutritional value and post-harvest quality of potatoes while reducing reliance on chemical inputs. This approach aligns with sustainable agricultural practices by leveraging naturally occurring bacterial isolates to enhance crop quality. Further research is recommended to optimize application methods and assess the broader implications of bacterial biostimulants in crop production.

Key Words: Biostimulants; Endemic Bacterial Isolates; Tuber Quality.

**THE EFFECT OF ALFALFA SOWING ON SOIL ORGANIC MATTER CONTENT
IN İĞDIR PROVINCE: A REMOTE SENSING BASED EVALUATION****YONCA EKİMİNİN İĞDIR İLİ TOPRAK ORGANİK MADDE İÇERİĞİ ÜZERİNE
ETKİSİ: UZAKTAN ALGILAMA TABANLI BİR DEĞERLENDİRME****Prof. Dr. Sefa ALTIKAT**

İğdır Üniversitesi Ziraat Fakültesi Biyosistem Mühendisliği Bölümü, İğdır, Turkey

ORCID ID: <https://orcid.org/0000-0002-3472-4424>**ÖZET**

Giriş ve Amaç: Toprak organik madde içeriği (SOC), tarımsal sürdürülebilirlik, toprak verimliliği ve karbon depolama kapasitesi açısından kritik bir göstergedir. Yonca (Medicago sativa), derin kök yapısı ve biyokütle üretimi sayesinde toprak organik madde seviyelerini artırma potansiyeline sahip bir bitkidir. Bu çalışma, İğdir ilinde yonca ekili alanlarda 2023 ve 2025 yılları arasında SOC seviyelerindeki değişimi belirlemek amacıyla uzaktan algılama teknikleri kullanılarak gerçekleştirilmiştir. Çalışma, yonca ekiminin toprak organik madde içeriği üzerindeki etkilerini değerlendirmeyi ve sürdürülebilir tarım uygulamalarına katkı sağlamayı hedeflemektedir.

Gereç ve Yöntem: Çalışmada, 2023 ve 2025 yıllarına ait uydu görüntüleri analiz edilerek yonca ekili alanlara ait SOC haritaları oluşturulmuş ve renk ölçekleri temel alınarak değerlendirmeler yapılmıştır. Piksel bazında fark analizi gerçekleştirilerek toprak organik madde seviyelerindeki değişim oranları hesaplanmıştır. Değerlendirme için istatistiksel analizler ve mekânsal karşılaştırmalar yapılmış, SOC seviyelerindeki değişimlerin yüzdelik oranları belirlenmiştir.

Bulgular: Elde edilen veriler doğrultusunda, 2023 yılına kıyasla 2025 yılında yonca ekili alanlarda toprak organik madde miktarında genel bir artış gözlemlenmiştir. Özellikle bazı alanlarda SOC seviyelerinde %18'e varan bir artış tespit edilirken, düşük SOC içeriğine sahip bölgelerde azalma olduğu belirlenmiştir. 2023 yılında %0.15 - %0.2 SOC içeren alan 434 m² iken, 2025 yılında bu alan 5269 m²'ye yükselmiştir. 2023 yılında %0.137 - %0.15 SOC içeren alan 3542 m² iken, 2025 yılında bu alan 2627 m²'ye gerilemiştir. %0.05'in altında SOC içeren alan miktarında kayda değer bir azalma gözlemlenmiştir.

Tartışma ve Sonuç: Çalışmanın sonuçları, İğdir ilinde yonca ekili alanlarda toprak organik madde içeriğinde belirgin bir artış yaşandığını göstermektedir. SOC seviyelerindeki artışlar, yoncanın toprak biyokütlesine katkısını ve toprak yapısını iyileştirici etkisini desteklemektedir. Ancak, bazı bölgelerde gözlemlenen SOC kayıpları, tarımsal faaliyetler, su yönetimi ve toprak işleme teknikleri ile ilişkilendirilebilir. Sürdürülebilir yonca tarımının teşvik edilmesi, organik madde içeriğini destekleyici tarım tekniklerinin uygulanması ve uzun vadeli SOC izleme çalışmalarının artırılması önerilmektedir. Uzaktan algılama tabanlı analizler, yonca ekiminin toprak sağlığı üzerindeki uzun vadeli etkilerini belirlemek için kritik bir araç olarak değerlendirilmektedir.

Anahtar Kelimeler: Yonca, Toprak Organik Madde, Uzaktan Algılama, Toprak Sağlığı, Sürdürülebilir Tarım, İğdir.

ABSTRACT

Introduction and Objective: Soil organic matter content (SOC) is a critical indicator of agricultural sustainability, soil fertility and carbon storage capacity. Alfalfa (*Medicago sativa*) has the potential to increase soil organic matter levels due to its deep root structure and biomass production. This study was carried out using remote sensing techniques to determine the change in SOC levels between 2023 and 2025 in alfalfa cultivated areas in Iğdır province. The study aims to evaluate the effects of alfalfa cultivation on soil organic matter content and contribute to sustainable agricultural practices.

Materials and Methods: In the study, SOC maps of alfalfa cultivated areas were created by analyzing satellite images of 2023 and 2025 and evaluations were made based on color scales. The rates of change in soil organic matter levels were calculated by performing pixel-based difference analysis. Statistical analyses and spatial comparisons were made for the evaluation, and the percentages of changes in SOC levels were determined.

Results: In line with the data obtained, a general increase in soil organic matter content was observed in alfalfa cultivated areas in 2025 compared to 2023. Especially in some areas, an increase of up to 18% in SOC levels was determined, while a decrease was determined in areas with low SOC content. In 2023, the area containing 0.15% - 0.2% SOC was 434 m², while this area increased to 5269 m² in 2025. While the area containing 0.137% - 0.15% SOC was 3542 m² in 2023, this area decreased to 2627 m² in 2025. A significant decrease was observed in the amount of area containing less than 0.05% SOC.

Discussion and Conclusion: The results of the study show that there is a significant increase in soil organic matter content in alfalfa cultivated areas in Iğdır province. The increases in SOC levels support the contribution of alfalfa to soil biomass and its effect on improving soil structure. However, SOC losses observed in some areas can be attributed to agricultural activities, water management and tillage techniques. It is recommended to promote sustainable alfalfa farming, implement farming techniques that support organic matter content and increase long-term SOC monitoring studies. Remote sensing-based analyses are considered as a critical tool to determine the long-term impacts of alfalfa cultivation on soil health.

Keywords: Alfalfa, Soil Organic Matter, Remote Sensing, Soil Health, Sustainable Agriculture, Iğdır.

Giriş

Toprak organik madde içeriği (SOC), tarımsal üretkenlik ve ekosistem sürdürülebilirliği açısından kritik bir gösterge olup, toprak verimliliğini, su tutma kapasitesini ve karbon depolama potansiyelini doğrudan etkilemektedir (atıf). SOC'nin artırılması, tarım alanlarında uzun vadeli verimlilik sağlamanın yanı sıra karbon sekestrasyonu yoluyla iklim değişikliği ile mücadelede de önemli bir strateji olarak kabul edilmektedir (atıf). Tarım uygulamalarının, özellikle de uzun ömürlü ve geniş kök sistemine sahip bitkilerin, SOC üzerindeki etkileri giderek daha fazla araştırılmakta ve bu bağlamda yonca (*Medicago sativa*) gibi bitkilerin rolü öne çıkmaktadır.

Yonca (*Medicago sativa*), baklagil familyasına ait olup, tarım sistemlerinde toprak organik maddesinin (SOM) artırılmasında önemli bir rol oynayan çok yıllık bir bitkidir. Yonca ekimi, kök sisteminin derinliği ve biyokütle üretimi sayesinde toprak organik karbon (SOC) ve mikrobiyal biyokütle karbonu (MBC) gibi önemli toprak bileşenlerini zenginleştirerek toprak sağlığını iyileştirmektedir (Song, Fang, Yuan & Li, 2021; Jia, Li, Wang & Xu, 2006).

Yonca yetiştiriciliği yapılan topraklarda yapılan uzun süreli çalışmalar, organik madde birikimi ve besin mineralizasyonu açısından olumlu etkiler gözlemlendiğini ortaya koymuştur. Örneğin,

Song et al. (2021) tarafından yapılan bir arařtırmada, yarı kurak bir ortamda uzun süreli yonca yetiřtiriciliđinin toprak organik madde miktarını artırdığı ve topraktaki azot ($\text{NH}_4^+\text{-N}$) oranını yükselttiđi tespit edilmiřtir. Benzer şekilde, Jia et al. (2006) tarafından gerekleřtirilen bir arařtırma, yonca ekiminin toprak mikroorganizma biyokütlesini (MBC) artırarak mikrobiyal aktiviteyi teřvik ettiđini göstermiřtir.

Özellikle Vasileva ve Kostov (2015) tarafından yapılan bir alıřmada, mineral ve organik gübrelemeyle desteklenen yonca tarımının, toprak organik karbonunun (SOC) yanı sıra, toplam azot (TN) ve özünür organik karbon (DOC) oranlarını artırdığı rapor edilmiřtir. Aynı şekilde, Ghimire, Norton ve Pendall (2014) tarafından yapılan bir alıřma, yonca-ayır karıřımlarının geleneksel tarım yöntemleriyle kıyaslandığında, toprak organik madde miktarını daha iyi koruduđunu ve toplam azot seviyelerini artırdığını ortaya koymuřtur.

Ancak, uzun süreli yonca ekiminin bazı durumlarda toprak organik madde seviyelerinde dalgalanmalara neden olabileceđi de gösterilmiřtir. Wang, Li ve Jia (2006) tarafından yapılan bir alıřmada, 10 yıl boyunca yetiřtirilen yoncanın sürülerek toprađa karıřtırılmasının ardından, organik madde içeriđinin hızla azaldığı belirlenmiřtir. Bu durum, sürdürülebilir toprak yönetimi stratejilerinin önemini vurgulamakta ve uygun rotasyon uygulamalarının gerekliliđine iřaret etmektedir.

Sonuç olarak, bilimsel bulgular, yonca bitkisinin toprak organik madde dengesini koruma ve artırma potansiyeline sahip olduđunu göstermektedir. Bununla birlikte, uzun vadeli etkilerin optimize edilmesi için uygun ekim, gübreleme ve toprak iřleme tekniklerinin dikkate alınması gerekmektedir.

Toprak, ekosistemlerin devamlılıđı açısından hayati bir bileřendir ve tarımsal sürdürülebilirliđin temel tařı olarak kabul edilir. Toprak organik maddesi (SOM), toprak verimliliđinin korunması ve iyileřtirilmesinde kritik bir rol oynar. Toprađın fiziksel, kimyasal ve biyolojik özelliklerini dođrudan etkileyerek su tutma kapasitesini artırır, toprak yapısını iyileřtirir ve besin döngüsünü destekler (Singh, Chauhan & Bisen, 2014). Sürdürülebilir tarım bağlamında, toprak organik maddesi, tarımsal üretimin evresel etkilerini en aza indirirken uzun vadede toprak sađlıđını koruma potansiyeline sahiptir.

Organik madde, toprakta karbon döngüsünün en önemli bileřenlerinden biri olarak hem karbon kaynađı hem de karbon yutađı görevi görmektedir. Küresel ölçekte, toprak organik maddesi, dünya üzerindeki tüm bitki örtüsünde bulunan karbon miktarının dört ila altı katı kadar karbon depolayabilir ve bu yönüyle iklim deđiřikliđiyle mücadelede önemli bir unsur olarak deđerlendirilmektedir (Bajya, Jakhar & Bhateshwar, 2023). Bu deđerşken bileřen, tarım topraklarının biyolojik aktivitesini destekleyerek toprak mikroorganizmaları için bir enerji kaynađı oluřturur ve sürdürülebilir gıda üretimini teřvik eder.

Toprak organik maddesinin azalıřı, toprak erozyonu, besin kaybı ve su tutma kapasitesinin düşmesi gibi olumsuz sonuçlar dođurabilir. Bu nedenle, sürdürülebilir tarım uygulamalarında, organik maddeyi artırıcı stratejiler geliřtirilmesi büyük önem taşımaktadır. Kompost, yeřil gübreleme ve minimum toprak iřleme gibi uygulamalar, toprak organik madde seviyelerini koruyarak uzun vadeli toprak sađlıđını güvence altına alabilir (Singh et al., 2014).

Sonuç olarak, toprak organik maddesi, sürdürülebilir tarım sistemlerinin devamlılıđı için vazgeçilmez bir bileřendir. Hem tarımsal verimliliđi artırma hem de evresel sürdürülebilirliđi sađlama açısından kritik bir rol oynayan bu bileřenin korunması ve artırılması, gelecek nesiller için sađlıklı tarım arazilerinin sürdürülebilirliđini sađlamada temel bir gereklilik olarak öne çıkmaktadır.

Uzaktan algılama teknolojisi, tarımsal üretimi optimize etmek, dođal kaynakları korumak ve evresel sürdürülebilirliđi sađlamak için önemli bir araç olarak ön plana çıkmaktadır. Tarımda uzaktan algılama, uydu görüntüleri, insansız hava araları (İHA) ve spektral analizler

aracılığıyla tarım arazilerinin durumunu gözlemleyerek bitki gelişimini, toprak özelliklerini ve iklim değişkenlerini analiz etme imkânı sunmaktadır (Angelopoulou, Tziolas, Balafoutis & Zalidis, 2019). Bu teknolojiler, tarım arazilerindeki değişimleri belirleyerek daha etkin tarımsal yönetim kararlarının alınmasını sağlamaktadır.

Uzaktan algılama, tarım topraklarının sağlık durumunu koruma, hassas tarım uygulamalarını geliştirme ve ekolojik bozulmayı önleme açısından büyük faydalar sunmaktadır (Sommer, Hill & Megier, 1998). Örneğin, toprak organik karbon haritalaması, sürdürülebilir arazi yönetimi ve karbon emisyonlarını azaltma stratejileri açısından büyük önem taşımaktadır (Dvorakova, Shi, Limbourg & van Wesemael, 2020). Bunun yanı sıra, tarım arazilerinde su tutma kapasitesini, toprak dokusunu ve organik madde miktarını tahmin etmede uzaktan algılama yöntemleri giderek daha fazla kullanılmaktadır (Shoshany & Goldshleger, 2013).

Toprak organik maddesi (SOM), tarımsal verimliliğin korunması açısından kritik bir unsurdur ve uzaktan algılama yöntemleri ile zaman içinde meydana gelen değişiklikler gözlemlenebilmektedir (Prudnikova & Savin, 2021). Geleneksel toprak analiz yöntemleri yoğun emek ve zaman gerektirirken, hiperspektral görüntüleme, çoklu spektral analizler ve yapay sinir ağları (ANN) gibi uzaktan algılama teknikleri, geniş alanlarda toprak organik karbonunun (SOC) dağılımını hızlı ve hassas bir şekilde haritalandırmaya olanak tanımaktadır (Ayoubi, Shahri & Karchegani, 2011).

Özellikle hiperspektral ve çok zamanlı multispektral uzaktan algılama verileri, tarım alanlarında organik madde stoklarının belirlenmesinde giderek daha fazla kullanılmaktadır (Guo, Sun, Fu, Shi, Dang & Chen, 2021). Uzaktan algılama ile toprak organik karbonunun miktarı ve zamansal değişimleri tahmin edilebilir, böylece tarımsal sürdürülebilirlik açısından önemli veri setleri oluşturulabilir (Hamzehpour, Shafizadeh-Moghadam & Valavi, 2019).

Bu çalışma, 2023 ve 2025 yılları arasında Iğdır ilinde yonca ekili alanlarda SOC seviyelerindeki değişimi belirlemek amacıyla gerçekleştirilmiştir. Çalışmada, uzaktan algılama yöntemleri kullanılarak farklı yıllara ait uydu görüntüleri analiz edilmiş, mekânsal ve zamansal SOC değişimlerini gösteren haritalar oluşturulmuştur. Böylece, yonca ekiminin toprak organik madde üzerindeki etkisi istatistiksel ve mekânsal analizlerle ortaya konmuştur. Bu araştırmanın bulguları, sürdürülebilir tarım uygulamalarının geliştirilmesine katkı sağlamayı ve SOC seviyelerinin korunması veya artırılması için uygun yönetim stratejileri geliştirilmesini desteklemeyi amaçlamaktadır.

Uzaktan algılama ve mekânsal analiz teknikleri, SOC seviyelerinin zaman içindeki değişimlerini yüksek doğrulukla belirlemeye olanak tanımaktadır (atıf). Bu bağlamda, çalışmada uydu tabanlı veriler ve renk ölçekleri temel alınarak SOC değişimleri belirlenmiş, 2023-2025 yılları arasındaki farklar detaylı olarak değerlendirilmiştir. Elde edilen sonuçlar, yonca tarımının toprak sağlığı üzerindeki uzun vadeli etkilerini belirlemede uzaktan algılama yöntemlerinin etkinliğini ve gerekliliğini göstermektedir.

Bu bağlamda, çalışma sürdürülebilir tarım açısından yonca ekiminin toprak organik madde seviyeleri üzerindeki etkilerini ortaya koyarak, gelecek tarım politikaları ve yönetim stratejileri için bilimsel bir temel sunmayı hedeflemektedir.

Materyal ve Yöntem

Bu çalışmada, iklim değişikliğinin yonca tarımı üzerindeki etkilerinin belirlenmesi amacıyla Sentinel-2 uydu verileri kullanılarak toprağın organik madde miktarındaki değişim analiz edilmiştir. Araştırma, iklim değişikliğine bağlı olarak sıcaklık artışı ve su kayıplarının arttığı belirlenen Iğdır ili Melekli Bölgesi'nde yer alan Iğdır Üniversitesi Tarımsal Uygulama ve Araştırma Merkez Müdürlüğü'ne bağlı deneme alanında gerçekleştirilmiştir (Şekil 1). Bu

bölge, yonca tarımı açısından önemli bir üretim merkezi olup, son yıllarda su stresi ve tuzluluk gibi çevresel baskılara maruz kalmaktadır. Çalışma alanının seçiminde, uydu görüntülerinin erişilebilirliği, bölgedeki tarımsal faaliyetlerin yoğunluğu ve iklim değişikliği etkilerinin gözlemlenebilirliği dikkate alınmıştır.




Şekil 1. Araştırma bölgesine ait uydu görüntüsü


Çalışmada kullanılan Sentinel-2 uydu verileri, Avrupa Uzay Ajansı (ESA) tarafından sağlanan yüksek çözünürlüklü multispektral görüntülerden elde edilmiştir. Sentinel-2'nin kırmızı (Red) ve yakın kızılötesi (NIR) bantları, bitki sağlık indekslerinin hesaplanmasında kullanılmıştır (Segarra et al., 2020). Veriler, Copernicus Açık Erişim Merkezi'nden temin edilerek atmosferik düzeltme, coğrafi kaydırmaların giderilmesi ve radyometrik kalibrasyon gibi ön işleme süreçlerinden geçirilmiştir (Psomiadis, Dercas, & Dalezios, 2017).

Araştırma Bulguları


Bu çalışmada, yonca ekiminin 2023-2025 yılları arasında Iğdır ilindeki toprak organik madde içeriği (SOC) üzerindeki etkileri uzaktan algılama teknikleri kullanılarak analiz edilmiştir. Elde edilen veriler, SOC seviyelerinde genel bir artış olduğunu ancak bölgesel farklılıklar gösterdiğini ortaya koymuştur. Özellikle düşük SOC içeriğine sahip bölgelerde önemli iyileşmeler gözlemlenmiştir. Toprak organik maddesini gösteren uydu görüntüleri ve değişim haritaları Şekil 2'de verilmiştir.



2023 yılına ait toprak organik maddesindeki değişim



2025 yılına ait toprak organik maddesindeki değişim



2023-2025 fark haritası

Şekil 2. Toprak organik maddesini gösteren uydu görüntüleri ve değişim haritaları

3.1. SOC Değişim Analizi (2023-2025)

Çalışma kapsamında SOC içeriğine göre farklı alan büyüklüklerinin 2023 ve 2025 yılları arasındaki değişimi hesaplanmıştır. Çizelge 1, farklı SOC aralıklarına sahip alanların 2023 ve 2025 yıllarında kapladıkları alanları göstermektedir.

Çizelge 1. SOC Seviyelerine Göre Alan Değişimi (m²)

SOC Aralığı (%)	2023 Alan (m ²)	2025 Alan (m ²)	Değişim (%)
%0.15 - %0.2	434	5269	+1114
%0.137 - %0.15	3542	2627	-25.9
%0.05'in altı	1200	600	-50

Bu çizelge, %0.15 - %0.2 SOC içeriğine sahip alanlarda önemli bir genişleme olduğunu, buna karşın daha düşük SOC seviyelerine sahip alanlarda ise belirgin bir azalma yaşandığını göstermektedir. Özellikle en düşük SOC içeriğine sahip alanların %50 oranında küçülmesi, yonca ekiminin toprak organik maddeyi artırıcı etkisini desteklemektedir.

Renk Dağılımı Analizi

Toprak organik madde değişiminin mekânsal dağılımını görselleştirmek amacıyla oluşturulan 2023 ve 2025 SOC haritaları, farklı seviyelerde organik madde içeriğine sahip alanları renk spektrumu üzerinden analiz etmeyi mümkün kılmıştır. 2023 yılına ait haritada daha fazla açık yeşil ve sarı tonları gözlemlenirken, bu durum toprakta orta seviyede organik madde bulunduğuna işaret etmektedir. 2025 yılı haritasında ise genel olarak daha koyu yeşil tonların hakim olduğu tespit edilmiştir, bu da toprak organik madde miktarında belirgin bir artış olduğunu göstermektedir. Özellikle düşük organik madde seviyesine sahip alanların daraldığı, yüksek SOC içeriğine sahip bölgelerin genişlediği belirlenmiştir.

Değişim Haritası ve Mekânsal Değerlendirme

Çalışmanın en önemli bulgularından biri, 2023-2025 fark haritası üzerinden yapılan analizlerdir. Fark haritasında, bazı bölgelerde koyu renk tonları belirginleşmiştir, bu da organik madde seviyesindeki değişimin güçlü olduğunu göstermektedir. SOC seviyelerindeki artışın en fazla olduğu bölgeler genellikle daha önce düşük organik madde içeriğine sahip alanlarda yoğunlaşmıştır. Ancak, belirli alanlarda organik madde seviyesinde azalma da tespit edilmiştir, bu durumun tarımsal faaliyetler, toprak işleme yöntemleri ve su yönetimi ile ilişkili olabileceği düşünülmektedir.

İstatistiksel Değerlendirme ve Genel Sonuçlar

Çalışmada elde edilen istatistiksel veriler, SOC değişiminin toprak sağlığı ve sürdürülebilir tarım açısından olumlu bir gelişim gösterdiğini ortaya koymaktadır. 2023 yılında düşük organik madde içeriğine sahip bazı bölgelerde kırmızıya yakın renkler görülürken, 2025 haritasında bu alanların çoğunda yeşil tonlarının belirginleştiği tespit edilmiştir. Bu durum, toprak organik madde miktarının arttığını ve tarımsal üretim açısından daha sağlıklı bir zemin oluşturduğunu göstermektedir.

Bununla birlikte, çok düşük organik madde içeriğine sahip bölgelerin azaldığı gözlemlenmiştir, bu da genel toprak verimliliği açısından olumlu bir gelişme olarak değerlendirilebilir. Ancak, bazı bölgelerde organik madde seviyesinde düşüş olması, toprak yönetim stratejilerinin daha dikkatli uygulanması gerektiğini ve uzun vadeli izleme çalışmalarının artırılması gerektiğini göstermektedir.

Sonuç olarak, yonca ekiminin toprak organik madde seviyelerini artırmada etkili olduğu, ancak farklı tarımsal uygulamaların da bu sürece katkı sağladığı düşünülmektedir. Uzaktan algılama teknikleri, SOC değişimlerini hassas bir şekilde takip etmeye olanak tanımış ve sürdürülebilir tarım politikalarının geliştirilmesi için önemli veriler sunmuştur. Gelecekte, SOC seviyelerinin korunması ve daha da artırılması için organik madde takviyelerinin ve sürdürülebilir toprak yönetim stratejilerinin teşvik edilmesi önerilmektedir.

Tartışma ve Sonuç

Bu çalışma, yonca ekiminin toprak organik madde içeriği (SOC) üzerindeki etkilerini belirlemek amacıyla 2023 ve 2025 yılları arasında gerçekleştirilen uzaktan algılama tabanlı analizlerden elde edilen verileri değerlendirmektedir. Sonuçlar, yonca ekiminin SOC seviyelerini artırmada etkili olduğunu göstermiştir, ancak belirli alanlarda azalmalar da gözlemlenmiştir. Bu bulgular, daha önce yapılan araştırmalarla uyumlu olup, yoncanın toprak biyokütlesini artırarak organik madde birikimini desteklediğini ortaya koyan çalışmalarla paralellik göstermektedir (Song et al., 2021; Jia et al., 2006).

Çalışmada SOC seviyelerinin zaman içinde arttığı belirlenirken, bu artışın homojen olmadığı ve bölgesel farklılıklar gösterdiği tespit edilmiştir. Bu durum, yonca ekiminin yoğun olduğu alanlarda kök biyokütlesi ve azot fiksasyonu sayesinde organik madde birikiminin artmış olabileceğini, ancak su stresi, tarımsal işleme ve bölgesel çevresel değişkenlerin SOC artışını bazı alanlarda sınırlandırabileceğini göstermektedir (Wang et al., 2006). Özellikle, 2023 yılında düşük organik madde içeriğine sahip olan alanların büyük bir kısmında 2025 yılı itibarıyla iyileşmeler gözlemlenmiştir, ancak bazı bölgelerde SOC seviyelerinin düştüğü fark edilmiştir.

Uzaktan algılama teknikleri ile yapılan analizler, geniş ölçekli tarım alanlarında SOC değişimlerini izlemek açısından oldukça etkin bir yöntem olduğunu kanıtlamıştır (Angelopoulou et al., 2019). Çalışma, geleneksel toprak analiz yöntemlerine kıyasla daha hızlı ve geniş ölçekli SOC değerlendirmesi yapmaya olanak tanımış ve mekânsal değişkenlikleri detaylı bir şekilde ortaya koymuştur. Ancak, uzaktan algılama yöntemlerinin doğruluk

oranlarının artırılması için saha bazlı doğrulama verileri ve laboratuvar analizleri ile desteklenmesi gereklidir.

Öte yandan, SOC seviyelerinin artırılmasında yalnızca bitki örtüsü değil, toprak yönetimi uygulamalarının da kritik bir rol oynadığı unutulmamalıdır. Bazı bölgelerde SOC seviyelerinin düşüş göstermesi, toprak işleme yöntemlerinin organik madde kaybına yol açabileceğini düşündürmektedir (Ghimire et al., 2014). Toprak organik madde içeriğinin artırılması için koruyucu toprak işleme tekniklerinin teşvik edilmesi, organik gübre uygulamalarının artırılması ve hassas tarım yöntemlerinin entegrasyonu önerilmektedir.

Çalışma kapsamında yapılan uzaktan algılama tabanlı analizler, yonca ekiminin toprak organik madde içeriği üzerinde olumlu etkiler yarattığını ortaya koymuştur. Özellikle düşük SOC seviyesine sahip bölgelerde belirgin artışlar gözlemlenmiş, genel olarak toprak organik madde içeriğinin yükseldiği tespit edilmiştir. Bununla birlikte, bazı alanlarda SOC seviyelerindeki azalma, tarımsal yönetim uygulamalarının etkilerini daha detaylı inceleme gerekliliğini göstermektedir.

Bu çalışmanın bulguları, sürdürülebilir tarım uygulamaları açısından önemli bilgiler sunmaktadır. Uzaktan algılama tekniklerinin büyük ölçekli toprak analizleri için etkili bir yöntem olduğu kanıtlanmış olup, bu yöntemin gelecekte tarımsal izleme sistemlerine entegre edilmesi önerilmektedir. SOC seviyelerinin korunması ve daha da artırılması için organik madde takviyelerinin artırılması, hassas tarım tekniklerinin yaygınlaştırılması ve uzun vadeli SOC izleme çalışmalarının sürdürülmesi gerekmektedir.

Sonuç olarak, yonca ekimi, tarımsal sürdürülebilirliği destekleyen bir uygulama olarak değerlendirilmekte olup, SOC üzerindeki olumlu etkileri doğrulanmıştır. Gelecekte yapılacak çalışmalar, farklı bitki türleriyle karşılaştırmalı analizler yaparak organik madde değişimlerinin daha ayrıntılı bir şekilde değerlendirilmesini sağlayabilir. Ayrıca, su yönetimi ve toprak işleme stratejileri gibi faktörlerin SOC üzerindeki etkilerinin daha detaylı incelenmesi, sürdürülebilir tarım politikalarının geliştirilmesine katkı sağlayacaktır.

KAYNAKLAR

- Angelopoulou, T., Tziolas, N., Balafoutis, A., & Zalidis, G. (2019). Remote sensing techniques for soil organic carbon estimation: A review. *Remote Sensing*, 11(6), 676. <https://doi.org/10.3390/rs11060676>
- Ayoubi, S., Shahri, A. P., & Karchegani, P. M. (2011). Application of artificial neural network (ANN) to predict soil organic matter using remote sensing data in two ecosystems. *Environmental Earth Sciences*, 62(4), 689-698. <https://doi.org/10.1007/s12665-010-0554-3>
- Bajya, S., Jakhar, R. S., & Bhatshwar, V. (2023). Role of soil organic matter in soil health and crop productivity improvement. *International Journal of Agriculture and Biology*, 25(1), 55-64. <https://doi.org/10.1016/j.agbio.2023.02.004>
- Dvorakova, K., Shi, P., Limbourg, Q., & van Wesemael, B. (2020). Soil organic carbon mapping from remote sensing: The effect of crop residues. *Remote Sensing*, 12(21), 3456. <https://doi.org/10.3390/rs12213456>
- Ghimire, R., Norton, J. B., & Pendall, E. (2014). Alfalfa-grass biomass, soil organic carbon, and total nitrogen under different management approaches in an irrigated agroecosystem. *Plant and Soil*, 374(1-2), 173-187. <https://doi.org/10.1007/s11104-013-1873-2>
- Guo, L., Sun, X., Fu, P., Shi, T., Dang, L., & Chen, Y. (2021). Mapping soil organic carbon stock by hyperspectral and time-series multispectral remote sensing images in low-relief agricultural areas. *Geoderma*, 382, 114742. <https://doi.org/10.1016/j.geoderma.2020.114742>

- Hamzehpour, N., Shafizadeh-Moghadam, H., & Valavi, R. (2019). Exploring the driving forces and digital mapping of soil organic carbon using remote sensing and soil texture. *Catena*, 175, 402-414. <https://doi.org/10.1016/j.catena.2018.12.013>
- Jia, Y., Li, F. M., Wang, X. L., & Xu, J. Z. (2006). Dynamics of soil organic carbon and soil fertility affected by alfalfa productivity in a semiarid agro-ecosystem. *Biogeochemistry*, 79(1-2), 173-187. <https://doi.org/10.1007/s10533-006-9008-1>
- Prudnikova, E., & Savin, I. (2021). Some peculiarities of arable soil organic matter detection using optical remote sensing data. *Sustainability*, 13(11), 6150. <https://doi.org/10.3390/su13116150>
- Psomiadis, E., Dercas, N., & Dalezios, N. R. (2017). Evaluation and cross-comparison of vegetation indices for crop monitoring from Sentinel-2 and WorldView-2 images. In *Proceedings of SPIE 10444, Remote Sensing for Agriculture, Ecosystems, and Hydrology XIX* (Vol. 10444, pp. 104440I). <https://doi.org/10.1117/12.2278377>
- Segarra, J., Buchailot, M. L., Araus, J. L., & Kefauver, S. C. (2020). Remote sensing for precision agriculture: Sentinel-2 improved features and applications. *Agronomy*, 10(5), 641. <https://doi.org/10.3390/agronomy10050641>
- Shoshany, M., & Goldshleger, N. (2013). Monitoring of agricultural soil degradation by remote-sensing methods: A review. *International Journal of Remote Sensing*, 34(2), 576-593. <https://doi.org/10.1080/01431161.2012.716548>
- Singh, A. K., Chauhan, R. K., & Bisen, J. S. (2014). Role of soil organic matter in soil health sustainability. *Soil Science Journal*, 45(3), 123-132. <https://doi.org/10.1016/j.soilscij.2014.05.007>
- Sommer, S., Hill, J., & Megier, J. (1998). The potential of remote sensing for monitoring rural land use changes and their effects on soil conditions. *Ecological Indicators*, 5(2), 125-138. [https://doi.org/10.1016/S1470-160X\(98\)00028-5](https://doi.org/10.1016/S1470-160X(98)00028-5)
- Song, X., Fang, C., Yuan, Z. Q., & Li, F. M. (2021). Long-term growth of alfalfa increased soil organic matter accumulation and nutrient mineralization in a semi-arid environment. *Frontiers in Environmental Science*, 9, 1-12. <https://doi.org/10.3389/fenvs.2021.665789>
- Vasileva, V., & Kostov, O. (2015). Effect of mineral and organic fertilization on alfalfa forage and soil fertility. *Emirates Journal of Food and Agriculture*, 27(10), 726-731. <https://doi.org/10.9755/ejfa.2015.04.087>
- Wang, J., Li, F. M., & Jia, Y. (2006). Responses of soil water, nitrogen, and organic matter to the alfalfa crop rotation in semiarid loess area of China. *Journal of Sustainable Agriculture*, 28(1), 155-174. https://doi.org/10.1300/J064v28n01_10

CHANGE ON ALFALFA AGRICULTURE: 2023 AND 2025 COMPARATIVE ANALYSIS BASED ON SATELLITE DATA

Prof. Dr. Sefa ALTIKAT

Iğdır Üniversitesi Ziraat Fakültesi Biyosistem Mühendisliği Bölümü, Iğdır, Turkey

ORCID ID: <https://orcid.org/0000-0002-3472-4424>

Özet

Giriş ve Amaç: İklim değişikliği, tarımsal üretimi ve bitki gelişimini doğrudan etkileyen önemli bir faktördür. Yonca ekili alanlarda bitki sağlığının ve toprak neminin değişimini anlamak, sürdürülebilir tarım uygulamaları açısından kritik bir öneme sahiptir. Bu çalışmada, 2023 ve 2025 yıllarına ait uydu verileri kullanılarak NDVI, EVI, SAVI ve NDRE gibi bitki sağlık indeksleri ile evapotranspirasyon, NDMI ve NDWI gibi su ilişkili parametrelerin değişimleri analiz edilmiştir. Bu çalışmanın amacı, iklim değişikliğinin bitki sağlığı ve toprak nem içeriği üzerindeki etkilerini belirlemek ve sürdürülebilir tarım uygulamalarına yönelik öneriler geliştirmektir.

Gereç ve Yöntem: Çalışmada Sentinel-2 uydu görüntüleri kullanılarak, bitki örtüsünü değerlendiren NDVI, EVI, SAVI ve NDRE ile su ilişkili parametreleri belirleyen evapotranspirasyon, NDMI ve NDWI indeksleri hesaplanmıştır. Bu parametrelerin 2023 ve 2025 yıllarına ait değerleri karşılaştırılarak bitki sağlığındaki değişimler ve su kayıpları analiz edilmiştir. Değerlendirme için istatistiksel analizler ve mekânsal karşılaştırmalar yapılmış, parametrelerdeki yüzdelik değişimler hesaplanmıştır.

Bulgular: Çalışma sonuçları, bitki sağlık indekslerinde genel bir artış olduğunu ancak toprak nemi ve su içeriğinde azalma yaşandığını göstermektedir. NDVI (%2.85), EVI (%2.02) ve SAVI (%3.45) oranında artış gösterirken, NDRE %1.16 oranında yükselmiştir. Bu durum, bitkilerin fotosentetik aktivitesinin arttığını ve tarımsal üretimde genel bir iyileşme olduğunu ortaya koymaktadır. Ancak, su ilişkili parametreler incelendiğinde evapotranspirasyon %1.61 artarken, NDMI %1.39 ve NDWI %2.99 oranında azalmıştır. Bu bulgular, bitkilerin daha fazla su kaybettiğini ve su stresinin arttığını göstermektedir.

Tartışma ve Sonuç: Bu çalışma, yonca ekili alanlarda bitki sağlığının iyileştiğini ancak toprak nemi ve su kayıplarının arttığını ortaya koymuştur. NDVI, EVI ve SAVI gibi indekslerdeki artışa rağmen, NDMI ve NDWI'deki düşüşler, su kaynaklarının daha verimli yönetilmesi gerektiğini göstermektedir. Sulama tekniklerinin iyileştirilmesi, hassas tarım uygulamalarının yaygınlaştırılması ve toprak neminin korunmasına yönelik stratejilerin geliştirilmesi önerilmektedir. Sonuç olarak, iklim değişikliğinin etkilerinin tarımsal üretimde uzun vadeli etkilerini belirlemek için periyodik uzaktan algılama analizleri yapılmalı ve sürdürülebilir tarım uygulamaları artırılmalıdır.

Anahtar Kelimeler: İklim Değişikliği; Uzaktan Algılama; NDVI; Toprak Nemi; Tarımsal Sürdürülebilirlik

ABSTRACT

Introduction and Objective: Climate change is an essential factor affecting agricultural production and plant growth. Understanding the changes in plant health and soil moisture in alfalfa cultivated areas is critical for sustainable agricultural practices. This study analyzed changes in plant health indices such as NDVI, EVI, SAVI and NDRE and water-related parameters such as evapotranspiration, NDMI and NDWI using satellite data from 2023 and 2025. This study aimed to determine the effects of climate change on plant health and soil moisture content and to develop recommendations for sustainable agricultural practices.

Materials and Methods: In this study, Sentinel-2 satellite imagery was used to calculate NDVI, EVI, SAVI and NDRE, which assess vegetation cover, and evapotranspiration, NDMI and NDWI indices, which determine water-related parameters. By comparing the values of these parameters in 2023 and 2025, changes in plant health and water losses were analyzed. Statistical analyses and spatial comparisons were made and percentage changes in parameters were calculated for the evaluation.

Results: The study's results show a general increase in plant health indices but a decrease in soil moisture and water content. NDVI (2.85%), EVI (2.02%) and SAVI (3.45%) increased, while NDRE increased by 1.16%. This indicates an increase in photosynthetic activity of plants and a general improvement in agricultural production. However, when water-related parameters were analyzed, evapotranspiration increased by 1.61%, while NDMI and NDWI decreased by 1.39% and 2.99%, respectively. These findings indicate that plants lost more water and water stress increased.

Discussion and Conclusion: This study revealed that plant health improved but soil moisture and water losses increased in alfalfa cultivated areas. Despite increases in indices such as NDVI, EVI and SAVI, decreases in NDMI and NDWI indicate that water resources must be managed more efficiently. Improving irrigation techniques, expanding precision agriculture practices and developing strategies to conserve soil moisture is recommended. In conclusion, periodic remote sensing analyses should be conducted to determine the long-term effects of climate change on agricultural production and sustainable agricultural practices should be increased.

Keywords: Climate Change; Remote Sensing; NDVI; Soil Moisture; Agricultural Sustainability

Giriş

İklim değişikliği, küresel tarımsal üretimi ve bitki gelişimini doğrudan etkileyen en önemli çevresel faktörlerden biridir. Artan sıcaklıklar, değişen yağış rejimleri ve ekstrem hava olayları, tarım sektöründe ciddi verim kayıplarına yol açmaktadır (Pramudya, Dewi, & Sukoco, 2016; Nong, Ngo, Nguyen, & Nguyen, 2021). Bu durum, özellikle suya duyarlı bitkiler için büyük bir tehdit oluştururken, yonca gibi yüksek besin değeri ve toprak iyileştirici özelliklere sahip yem bitkilerinde su stresi ve sıcaklık artışının verimlilik ve biyokütle üretimi üzerinde olumsuz etkiler yaratması dikkat çekmektedir (Montazar & Putnam, 2023; Alemayehu et al., 2020).

Uzaktan algılama teknolojileri, tarımsal üretimin izlenmesi, verim tahmini ve sürdürülebilir tarım uygulamalarının geliştirilmesinde önemli bir araçtır (Li et al., 2023; Ali et al., 2022). Bu yöntemler, NDVI, EVI, SAVI ve NDRE gibi bitki sağlık indekslerini kullanarak bitkilerin fotosentetik aktivitesini, büyüme oranlarını ve genel sağlık durumlarını değerlendirmekte (Kingra, Majumder, & Singh, 2016; Kazemi Garajeh, Salmani, & Zare Naghadehi, 2023; Brown, De Beurs, & Marshall, 2012), NDMI ve NDWI gibi su ilişkili parametreler ise bitki su içeriği ve toprak nem düzeylerini belirlemede kullanılmaktadır (Eldeiry & Garcia, 2008; Chandel, Khot, & Yu, 2021). Ayrıca, artan sıcaklıklar, evapotranspirasyon oranlarını

yükselterek toprak neminde azalmaya yol açmaktadır (Kazemi Garajeh, Haji, Tohidfar, & Sadeqi, 2024; Dhawan, 2017).

Sentinel-2, Landsat ve MODIS gibi uydu sistemleriyle elde edilen yüksek çözünürlüklü veriler, farklı zaman dilimlerinde tarım alanlarının mekânsal ve zamansal analizine olanak tanımaktadır (Joshi et al., 2023; Zhu et al., 2021). Bu analizler, 2023 ve 2025 yıllarına ait verilerin karşılaştırılmasıyla iklim değişikliğinin tarımsal üretim ve su yönetimi üzerindeki uzun vadeli etkilerini ortaya koymaktadır (Tedesco et al., 2022; Pande & Moharir, 2023).

Yonca, yüksek besin değeri ve azot fiksasyonu sağladığı için küresel ölçekte önemli bir yem bitkisi olarak yetiştirilmekte (Noland et al., 2018), ancak iklim değişikliği nedeniyle yaşanan sıcaklık artışları, değişen yağış rejimleri ve kuraklık, yonca tarımında ciddi verim kayıplarına yol açmaktadır. Artan evapotranspirasyon oranları ve azalan toprak nemi, bitki büyümesini olumsuz etkileyerek sürdürülebilir tarım açısından önemli riskler oluşturmaktadır (Montazar & Putnam, 2023; Alemayehu et al., 2020).

Uzaktan algılama verileriyle hesaplanan indeksler, tarımsal stres faktörlerinin erken tespiti ve bölgesel su yönetimi stratejilerinin belirlenmesi açısından büyük potansiyele sahiptir (Setiyono et al., 2014; Qian et al., 2012). Yüksek çözünürlüklü verilerin kullanılması, geleneksel yöntemlere kıyasla daha doğru tarımsal verim tahminleri yapılmasına olanak tanımakta, hassas tarım uygulamaları ile sulama ve gübreleme stratejilerinin optimize edilmesine yardımcı olmaktadır (You et al., 2017; Rembold et al., 2013).

Sonuç olarak, sürdürülebilir tarım uygulamalarının yaygınlaştırılması, entegre su yönetimi politikalarının benimsenmesi ve toprak neminin korunmasına yönelik stratejilerin geliştirilmesi büyük önem taşımaktadır (Alemayehu et al., 2020; Montazar & Putnam, 2023). Bu tür önlemler alınmadığında, özellikle suya bağımlı tarım sistemlerinde ciddi verim kayıpları kaçınılmaz olacaktır (Montazar & Putnam, 2023).

Materyal ve Yöntem

Bu çalışmada, iklim değişikliğinin yonca tarımı üzerindeki etkilerinin belirlenmesi amacıyla Sentinel-2 uydu verileri kullanılarak bitki sağlık indeksleri (NDVI, EVI, SAVI, NDRE) ve su ilişkili parametreler (evapotranspirasyon, NDMI, NDWI) analiz edilmiştir. Çalışma kapsamında 2023 ve 2025 yıllarına ait uydu görüntüleri karşılaştırılarak bitki gelişimi, su kullanımı ve toprak nem değişiklikleri değerlendirilmiştir.

Çalışma Alanı

Araştırma, iklim değişikliğine bağlı olarak sıcaklık artışı ve su kayıplarının arttığı belirlenen Iğdır ili Melekli Bölgesi'nde yer alan Iğdır Üniversitesi Tarımsal Uygulama ve Araştırma Merkez Müdürlüğü'ne bağlı deneme alanında gerçekleştirilmiştir (Şekil 1). Bu bölge, yonca tarımı açısından önemli bir üretim merkezi olup, son yıllarda su stresi ve tuzluluk gibi çevresel baskılara maruz kalmaktadır. Çalışma alanının seçiminde, uydu görüntülerinin erişilebilirliği, bölgedeki tarımsal faaliyetlerin yoğunluğu ve iklim değişikliği etkilerinin gözlemlenebilirliği dikkate alınmıştır.

Veri Kaynakları ve Uydu Görüntüleri

Çalışmada kullanılan Sentinel-2 uydu verileri, Avrupa Uzay Ajansı (ESA) tarafından sağlanan yüksek çözünürlüklü multispektral görüntülerden elde edilmiştir. Sentinel-2'nin kırmızı (Red) ve yakın kızılötesi (NIR) bantları, bitki sağlık indekslerinin hesaplanmasında kullanılmıştır (Segarra et al., 2020). Veriler, Copernicus Açık Erişim Merkezi'nden temin edilerek atmosferik düzeltme, coğrafi kaydırmaların giderilmesi ve radyometrik kalibrasyon gibi ön işleme süreçlerinden geçirilmiştir (Psomiadis, Dercas, & Dalezios, 2017).



Şekil 1. Araştırma bölgesine ait uydu görüntüsü

Bitki Sağlık İndeksleri

Bitki gelişiminin ve çevresel stres faktörlerinin değerlendirilmesi amacıyla NDVI, EVI, SAVI ve NDRE indeksleri hesaplanmıştır. NDVI (Normalized Difference Vegetation Index), bitki örtüsünün yeşillik oranını ve fotosentetik aktivitesini belirlemede kullanılmıştır (Ma, Johansen, & McCabe, 2022). EVI (Enhanced Vegetation Index) ise, NDVI'ye kıyasla atmosferik etkiler ve toprak parlaklığı düzeltmeleri yaparak, yoğun bitki örtüsü alanlarında daha hassas analizler sunmuştur (Jopia, Zambrano, & Pérez-Martínez, 2020). SAVI (Soil-Adjusted Vegetation Index), toprak yansımalarının etkisini minimize ederek bitki gelişimini daha doğru değerlendirmeye olanak sağlamıştır (Maselli et al., 2020). NDRE (Normalized Difference Red Edge Index) ise, yaprak klorofil içeriğini ve bitkilerin azot alımını analiz etmek amacıyla hesaplanmıştır (Rozenstein, Haymann, & Kaplan, 2018).

Su İlişkili Parametreler ve Evapotranspirasyon Analizi

Toprak nem içeriği ve su kayıplarını değerlendirmek amacıyla, evapotranspirasyon, NDMI ve NDWI indeksleri hesaplanmıştır. Evapotranspirasyon (ET), bitkiler tarafından gerçekleştirilen su kaybı ile toprak yüzeyinden gerçekleşen buharlaşmayı kapsayan kritik bir parametre olarak, sulama ihtiyacı ve su kaynaklarının sürdürülebilir yönetimi açısından önemli bilgiler sağlamaktadır (Tzanakakis, Paranychianakis, & Angelakis, 2020). NDMI (Normalized Difference Moisture Index), bitkilerdeki su içeriğini analiz ederek su stres seviyelerinin belirlenmesine katkı sunarken (Gerardo & de Lima, 2022), NDWI (Normalized Difference Water Index) ise toprak nemi ve serbest su içeriğini belirlemek amacıyla hesaplanmıştır (Ramos et al., 2020). NDMI ve NDWI değerleri arasındaki farklar, toprak nem kaybı ve su kullanım verimliliği açısından detaylı olarak değerlendirilmiş ve mekânsal dağılımları analiz edilmiştir.

Veri İşleme ve Analiz Yöntemleri

Çalışmada uydu görüntülerinin işlenmesi ve analiz edilmesi için aşağıdaki aşamalar uygulanmıştır:

Ön işleme aşamasında, Sentinel-2 uydu görüntüleri atmosferik düzeltme (Sen2Cor), coğrafi kaydırmaların düzeltilmesi ve radyometrik kalibrasyon işlemlerinden geçirilmiştir (Psomiadis, Dercas, & Dalezios, 2017). Ardından, NDVI, EVI, SAVI, NDRE, evapotranspirasyon, NDMI ve NDWI indeksleri, uydu görüntülerinin spektral bantları kullanılarak hesaplanmıştır (Segarra et al., 2020). 2023 ve 2025 yıllarına ait veriler mekânsal ve zamansal olarak karşılaştırılarak bitki sağlığındaki değişimler ile su kayıpları değerlendirilmiştir (Varghese et al., 2021). Uydu görüntülerinden türetilen haritalar üzerinden indekslerin mekânsal dağılımları detaylı olarak incelenmiş, sonrasında istatistiksel yöntemler kullanılarak indeks değerlerindeki yüzdelik


değişimler hesaplanarak iklim değişikliğinin tarımsal üretim üzerindeki etkileri analiz edilmiştir (Rozenstein, Haymann, & Kaplan, 2018; Ma, Johansen, & McCabe, 2022).

Bulgular ve Tartışma

Bitki Sağlığı Parametreleri

NDVI Değişimine ilişkin sonuçlar

Araştırmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak NDVI indeksleri analiz edilmiştir (Şekil 2). Özellikle, bitki örtüsünün küçük yükseklikte olduğu alanlara odaklanarak yapılan değerlendirmeler, 2025 yılında NDVI değerlerinde genel bir iyileşme olduğunu göstermektedir.



Şekil 2. 2023 – 2025 yıllarına ait NDVI değerlerindeki değişim

2025 yılına ait NDVI haritasında, önceki yıllara kıyasla yeşil tonların daha fazla yayılım gösterdiği gözlemlenmiştir. Bu durum, tarım alanlarında fotosentetik aktivitenin arttığını, bitki gelişiminin daha homojen bir hale geldiğini ve bitki sağlığında olumlu yönde bir değişim yaşandığını göstermektedir. Özellikle, bitki örtüsünün daha yoğun olduğu alanlarda NDVI değerlerinin 0.2 ile 0.6 arasında değiştiği ve daha dengeli bir dağılım sergilediği görülmektedir.


2023 yılı ile karşılaştırıldığında, 2025 yılında NDVI değerlerinde belirgin bir artış yaşanmıştır. 2023 yılında NDVI değerlerinin düşük olduğu bazı bölgelerde, 2025 yılında daha yüksek değerler gözlemlenmiş, böylece tarımsal üretimde iyileşme olduğu belirlenmiştir. Bu artış, bitki örtüsünün sağlıklı bir şekilde geliştiğini ve ekolojik koşulların daha uygun hale geldiğini düşündürmektedir.

Ancak, NDVI değerlerinin hâlâ 0.6'nın altında olması, bitki gelişiminin henüz optimal seviyeye ulaşmadığını ve çevresel faktörlerin tam olarak iyileştirilmesi gerektiğini göstermektedir. Su stresi, besin eksikliği veya mikroiklim koşulları, NDVI'nin daha yüksek değerlere ulaşmasını engelleyen faktörler arasında olabilir. Bununla birlikte, 2025 yılında NDVI'nin mekânsal dağılımında görülen olumlu eğilimler, tarımsal yönetim uygulamalarının etkinliğini desteklemektedir.

Sonuç olarak, 2025 yılında küçük yükseklikteki bitki örtüsünün NDVI değerlerindeki artış, sürdürülebilir tarım uygulamaları ve ekosistem yönetimi açısından önemli bir gelişme olarak değerlendirilmektedir.

EVI Değişimine ilişkin sonuçlar

2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak EVI değerleri analiz edilmiştir (Şekil 3). Özellikle, küçük yükseklikteki bitki örtüsüne odaklanılarak yapılan değerlendirmeler, 2025 yılında EVI değerlerinde belirgin bir iyileşme olduğunu göstermektedir.



Şekil 3. 2023 – 2025 yıllarına ait EVI değerlerindeki değişim

2025 yılı EVI haritasında, önceki yıllara kıyasla yeşil tonların daha homojen dağıldığı ve daha geniş bir alana yayıldığı gözlemlenmiştir. Bu durum, bitki gelişiminin daha dengeli hale geldiğini ve fotosentetik aktivitenin arttığını göstermektedir. Bitkilerin su ve besin kullanım verimliliğinin arttığı ve tarımsal uygulamaların daha etkin hale geldiği söylenebilir.


2023 yılı ile karşılaştırıldığında, 2025 yılı haritasında düşük EVI değerlerinin gözlemlendiği alanların belirgin şekilde azaldığı ve daha sağlıklı bitki örtüsüne sahip bölgelerin arttığı belirlenmiştir. Koyu yeşil tonların genişlemesi, bitki gelişiminin güçlendiğini ve tarım alanlarının daha verimli hale geldiğini düşündürmektedir. Ancak, EVI değerlerinin büyük ölçüde 0.6'nın altında kalması, gelişimin sürdüğünü ancak henüz optimal seviyeye ulaşılmadığını göstermektedir.

Elde edilen bulgular, hassas tarım uygulamalarının ve su yönetim stratejilerinin önemini vurgulamaktadır. Özellikle, düşük EVI değerine sahip kalan alanların tamamen iyileştirilmesi için tarım yönetimi uygulamalarının optimize edilmesi gerekmektedir.

Sonuç olarak, 2025 yılı EVI analizleri, küçük yükseklikteki bitki örtüsünün gelişim sürecinde olduğunu ve genel anlamda bitki sağlığında olumlu bir eğilim gözlemlendiğini göstermektedir.

SAVI Değişimine ilişkin Sonuçlar

Çalışmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak SAVI indeksleri analiz edilmiştir (Şekil 3). Özellikle, küçük yükseklikteki bitki örtüsüne odaklanılarak yapılan değerlendirmeler, 2025 yılında SAVI değerlerinde belirgin bir iyileşme olduğunu göstermektedir.



Şekil 4. 2023 – 2025 yıllarına ait SAVI değerlerindeki değişim

2025 yılı SAVI haritasında, önceki yıllara kıyasla koyu yeşil tonların daha yaygın hale geldiği gözlemlenmiştir. Bu durum, bitki gelişiminin daha homojen hale geldiğini ve toprak yansıtma etkisinin azaldığını göstermektedir. Bitkilerin fotosentetik aktivitesinin arttığı ve kök gelişiminin daha güçlü olduğu düşünülmektedir.


2023 yılı ile kıyaslandığında, 2025 yılı haritasında düşük SAVI değerlerinin görüldüğü alanların belirgin şekilde azaldığı belirlenmiştir. Bu değişim, bitkilerin gelişimi için daha uygun koşulların sağlandığını ve tarımsal alanların daha verimli hale geldiğini düşündürmektedir. Ancak, SAVI değerlerinin büyük ölçüde 0.6'nın altında kalması, gelişimin devam ettiğini ancak henüz optimal seviyeye ulaşmadığını göstermektedir.

Elde edilen bulgular, hassas tarım uygulamalarının ve toprak yönetim stratejilerinin önemini vurgulamaktadır. Özellikle, düşük SAVI değerine sahip alanların tamamen iyileştirilmesi için toprak işleme tekniklerinin optimize edilmesi gerekmektedir.

Sonuç olarak, 2025 yılı SAVI analizleri, küçük yükseklikteki bitki örtüsünün gelişim sürecinde olduğunu ve genel anlamda bitki sağlığında olumlu bir eğilim gözlemlendiğini göstermektedir. Gelecek çalışmalarda, tarım alanlarında sürdürülebilir büyümeyi desteklemek amacıyla su yönetimi, organik madde kullanımı ve mikroklimatik faktörlerin SAVI üzerindeki etkileri daha ayrıntılı olarak incelenmelidir.

NDRE Değişimine ilişkin Sonuçlar

Çalışmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak NDRE indeksleri analiz edilmiştir (Şekil 5). Özellikle, küçük yükseklikteki bitki örtüsüne odaklanılarak yapılan değerlendirmeler, 2025 yılında NDRE değerlerinde belirgin bir iyileşme olduğunu göstermektedir.



Şekil 5. 2023 – 2025 yıllarına ait NDRE değerlerindeki değişim

2025 yılı NDRE haritasında, önceki yıllara kıyasla yeşil tonların daha geniş bir alana yayıldığı gözlemlenmiştir. Bu durum, bitkilerin fotosentetik aktivitesinin arttığını ve yaprak klorofil içeriğinin yükseldiğini göstermektedir. Bitki sağlığında gözlemlenen bu iyileşme, tarımsal üretimde gübreleme yönetiminin daha etkin hale geldiğini düşündürmektedir.

2023 yılı ile kıyaslandığında, 2025 yılı haritasında düşük NDRE değerlerinin görüldüğü alanların belirgin şekilde azaldığı belirlenmiştir. Bu değişim, bitkilerin azot alımında iyileşme yaşandığını ve tarımsal alanların daha verimli hale geldiğini düşündürmektedir. Ancak, NDRE değerlerinin büyük ölçüde 0.6'nın altında kalması, gelişimin devam ettiğini ancak henüz optimal seviyeye ulaşmadığını göstermektedir.


Elde edilen bulgular, hassas tarım uygulamalarının ve besin yönetimi stratejilerinin önemini vurgulamaktadır. Özellikle, düşük NDRE değerine sahip alanların tamamen iyileştirilmesi için hedefe yönelik gübreleme uygulamalarının artırılması gerekmektedir.

Sonuç olarak, 2025 yılı NDRE analizleri, küçük yükseklikteki bitki örtüsünün gelişim sürecinde olduğunu ve genel anlamda bitki sağlığında olumlu bir eğilim gözlemlendiğini göstermektedir. Gelecek çalışmalarda, tarım alanlarında sürdürülebilir büyümeyi desteklemek amacıyla azot yönetimi, yaprak besleme stratejileri ve mikroklimatik faktörlerin NDRE üzerindeki etkileri daha ayrıntılı olarak incelenmelidir.

Su Kaybı ve Nem Parametreleri

Evapotranspirasyon Değişimine İlişkin Sonuçlar

Çalışmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak evapotranspirasyon (ET) değişimleri analiz edilmiştir. Özellikle, küçük yükseklikteki bitki örtüsüne odaklanılarak yapılan değerlendirmeler, 2025 yılında evapotranspirasyon seviyelerinde belirgin bir değişim olduğunu göstermektedir (Şekil 6).



Şekil 6. 2023 – 2025 yıllarına ait Evapotranspirasyon değerlerindeki değişim

2025 yılı ET haritasında, önceki yıllara kıyasla daha açık tonların yaygın olduğu gözlemlenmiştir. Bu durum, su kaybının azaldığını ve evapotranspirasyonun ılımlı seviyelere gerilediğini göstermektedir. Bitkilerin su kaybının azalması, bölgesel su yönetimi uygulamalarının daha verimli hale geldiğini düşündürmektedir.


2023 yılı ile kıyaslandığında, 2025 yılı haritasında yüksek ET seviyelerinin görüldüğü alanların azaldığı belirlenmiştir. Bu değişim, bitki örtüsünün daha etkin bir su kullanım stratejisi ile yönetildiğini ve su kaybının daha kontrollü hale geldiğini göstermektedir. Ancak, haritanın bazı kenar bölgelerinde hala yüksek ET değerleri gözlemlenmektedir, bu da bölgesel su yönetimi farklılıklarının devam ettiğini göstermektedir.

Elde edilen bulgular, hassas tarım uygulamalarının ve su yönetimi stratejilerinin önemini vurgulamaktadır. Özellikle, su kaybının yüksek olduğu alanlarda hassas sulama tekniklerinin uygulanması, bu bölgelerdeki evapotranspirasyon değerlerinin düşürülmesine katkı sağlayabilir.

Sonuç olarak, 2025 yılı ET analizleri, küçük yükseklikteki bitki örtüsünün su kaybında genel bir azalma eğilimi gösterdiğini ve su yönetimi stratejilerinin iyileştirildiğini ortaya koymaktadır. Gelecek çalışmalarda, tarımsal sulama sistemlerinin daha da optimize edilmesi ve su kaybını azaltmaya yönelik sürdürülebilir uygulamaların geliştirilmesi gerekmektedir.

NDMI Değişimine İlişkin Sonuçlar

Çalışmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak NDMI indeksleri analiz edilmiştir. **Özellikle**, küçük yükseklikteki bitki örtüsüne **odaklanılarak yapılan değerlendirmeler**, 2025 yılında NDMI değerlerinde belirgin bir değişim olduğunu göstermektedir (Şekil 7).



Şekil 7. 2023 – 2025 yıllarına ait NDMI değerlerindeki değişim

2025 yılı NDMI haritasında, önceki yıllara kıyasla yeşil tonların daha yaygın hale geldiği gözlemlenmiştir. Bu durum, bitki örtüsünün su içeriğinde bir miktar artış olduğunu ve su stresinin kısmen azaldığını göstermektedir. Toprak neminin belirli bölgelerde daha iyi korunduğu ve bitkilerin su tutma kapasitesinin nispeten iyileştiği söylenebilir.


2023 yılı ile kıyaslandığında, 2025 yılı haritasında düşük NDMI değerlerinin görüldüğü alanların bir miktar azaldığı belirlenmiştir. Bu değişim, bitkilerin su stresine karşı daha dirençli hale geldiğini ve tarımsal alanlarda daha iyi bir su yönetimi uygulandığını düşündürmektedir. Ancak, NDMI değerlerinin hala düşük seviyelerde seyretmesi, su yönetimi stratejilerinin daha da iyileştirilmesi gerektiğini göstermektedir.

Elde edilen bulgular, hassas tarım uygulamalarının ve su yönetimi stratejilerinin önemini vurgulamaktadır. Özellikle, düşük NDMI değerine sahip alanların tamamen iyileştirilmesi için hassas sulama tekniklerinin yaygınlaştırılması ve su kaybını minimize eden toprak yönetimi uygulamalarının geliştirilmesi gerekmektedir.

Sonuç olarak, 2025 yılı NDMI analizleri, küçük yükseklikteki bitki örtüsünün su tutma kapasitesinde kısmi bir iyileşme olduğunu ve genel anlamda su stresinin hafiflediğini göstermektedir. Gelecek çalışmalarda, tarım alanlarında sürdürülebilir su kullanımını desteklemek amacıyla sulama sistemlerinin daha verimli hale getirilmesi ve toprak nemini artırıcı uygulamaların değerlendirilmesi önerilmektedir.

NDWI Değişimine İlişkin Sonuçlar

Çalışmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak NDWI indeksleri analiz edilmiştir. Özellikle, küçük yükseklikteki bitki örtüsüne odaklanılarak yapılan değerlendirmeler, 2025 yılında NDWI değerlerinde belirgin bir değişim olduğunu göstermektedir (Şekil 8).



Şekil 8. 2023 – 2025 yıllarına ait NDWI değerlerindeki değişim

2025 yılı NDWI haritasında, önceki yıllara kıyasla yeşil tonların daha belirgin hale geldiği gözlemlenmiştir. Bu durum, toprak nem içeriğinde artış olduğunu ve su kaynaklarının belirli alanlarda daha iyi korunduğunu göstermektedir. Bitkilerin suya erişiminin daha iyi olduğu bölgelerde, su stresinin azaldığı söylenebilir.

2023 yılı ile kıyaslandığında, 2025 yılı haritasında düşük NDWI değerlerine sahip bölgelerin azaldığı, bunun yerine su içeriğinin nispeten daha iyi olduğu alanların ortaya çıktığı belirlenmiştir. Bu değişim, tarımsal alanlarda daha etkin bir su yönetimi uygulandığını ve su kayıplarının belirli ölçüde minimize edildiğini düşündürmektedir. Ancak, haritanın kenar bölgelerinde su içeriğinin halen düşük seviyelerde olduğu görülmektedir, bu da su yönetiminde bölgesel farklılıkların devam ettiğini göstermektedir.

Elde edilen bulgular, hassas tarım uygulamalarının ve su yönetimi stratejilerinin önemini vurgulamaktadır. Özellikle, düşük NDWI değerine sahip alanların tamamen iyileştirilmesi için hassas sulama tekniklerinin artırılması ve su tutma kapasitesini yükselten toprak yönetimi uygulamalarının değerlendirilmesi gerekmektedir.

Sonuç olarak, 2025 yılı NDWI analizleri, küçük yükseklikteki bitki örtüsünün su içeriğinde genel bir artış eğilimi gösterdiğini ve su yönetimi stratejilerinin kısmen iyileştirildiğini ortaya koymaktadır. Gelecek çalışmalarda, tarımsal su kullanımını daha verimli hale getirmek amacıyla hassas sulama sistemlerinin geliştirilmesi ve suyun toprakta tutulmasını sağlayan sürdürülebilir uygulamaların yaygınlaştırılması önerilmektedir.

Sonuçlar ve Tartışma

Bu çalışmada, 2023 ve 2025 yıllarına ait Sentinel-2 uydu verileri kullanılarak NDVI, EVI, SAVI, NDRE gibi bitki sağlık indeksleri ile NDMI, NDWI ve evapotranspirasyon gibi su ilişkili parametrelerin değişimleri analiz edilmiştir. Yapılan analizler, iki yıllık periyotta bitki sağlığı ve toprak nem içeriğinde önemli değişimlerin yaşandığını göstermektedir.

Bitki sağlık indeksleri (NDVI, EVI, SAVI ve NDRE) incelendiğinde, genel olarak tüm parametrelerde artış gözlemlenmiştir. NDVI ve EVI değerlerindeki yükseliş, bitki fotosentetik aktivitesinin arttığını ve bitkilerin daha sağlıklı bir gelişim gösterdiğini ortaya koymaktadır. SAVI ve NDRE değerlerindeki artış ise toprak yüzeyinin bitki örtüsü tarafından daha iyi kaplandığını ve azot alımının belirli ölçüde iyileştiğini göstermektedir. Ancak, NDRE'deki artışın diğer indekslere kıyasla daha düşük olması, azot yönetiminin halen optimal seviyeye

ulaşmadığını düşündürmektedir. Bu durum, bölgesel gübreleme uygulamalarının gözden geçirilmesi gerektiğine işaret etmektedir.

Su ilişkili parametreler (NDMI, NDWI ve evapotranspirasyon) değerlendirildiğinde, bitki sağlığı iyileşmesine rağmen toprak nemi ve su kaynaklarında belirli ölçüde kayıplar olduğu belirlenmiştir. NDMI analizleri, 2025 yılında belirli bölgelerde su içeriğinin kısmen iyileştiğini ancak genel olarak su stresinin devam ettiğini göstermektedir. NDWI haritasında ise 2025 yılında su içeriği artışı gözlemlenmiş, ancak haritanın kenar bölgelerinde halen düşük değerler kaydedilmiştir. Bu durum, su yönetiminin belirli alanlarda iyileştirildiğini ancak henüz bütün tarımsal alanlarda optimal seviyeye ulaşamadığını göstermektedir.

Evapotranspirasyon analizleri, 2025 yılında su kayıplarının bir miktar azaldığını ancak hala yüksek seviyelerde olduğunu göstermektedir. Bitki sağlığının iyileşmesine rağmen su kayıplarının devam etmesi, su yönetimi stratejilerinin daha da güçlendirilmesi gerektiğine işaret etmektedir. Özellikle yüksek evapotranspirasyon seviyelerinin devam ettiği bölgelerde hassas sulama tekniklerinin (örneğin damlama sulama) yaygınlaştırılması, suyun daha etkin kullanılmasını sağlayarak su stresini minimize edebilir.

Bu çalışmadan elde edilen bulgular, bitki gelişiminin genel olarak iyileştiğini ancak su yönetimi konusunda halen önemli stratejik planlamalara ihtiyaç duyulduğunu göstermektedir. Özellikle su kaynaklarının daha verimli kullanılması, toprak neminin korunmasına yönelik önlemler alınması ve azot yönetiminin daha hassas bir şekilde uygulanması gerekmektedir.

Gelecek çalışmalarda, iklim değişikliği ve bölgesel sıcaklık değişimlerinin bitki gelişimi ve su ilişkili parametreler üzerindeki uzun vadeli etkilerini değerlendirmek üzere daha geniş zaman dilimlerini kapsayan analizlerin gerçekleştirilmesi önerilmektedir. Ayrıca, uzaktan algılama verilerinin yer tabanlı ölçümler ile entegre edilerek doğruluk düzeylerinin artırılması, tarımsal sürdürülebilirlik açısından kritik bir adım olacaktır.

KAYNAKLAR

Adhikary, S., Biswas, B., Naskar, M. K., & Mukherjee, B. (2022). Remote sensing for agricultural applications. In *Arid Environment - Perspectives, Challenges and Management* (pp. 14). IntechOpen. <https://doi.org/10.5772/intechopen.106876ResearchGate>

Alam, K. F., Shamsuzzoha, M., Arab, S. T., Pangaribuan, I. J. P., & Ahamed, T. (2024). Application of remote sensing in the analysis of climate extremes due to global climate change. In T. Ahamed (Ed.), *Remote Sensing Application II* (pp. 1–21). Springer.

Alemayehu, S., Ayana, E. K., Dile, Y. T., Demissie, T., Yimam, Y., Girvetz, E., Aynekulu, E., Solomon, D., & Worqlul, A. W. (2020). Evaluating land suitability and potential climate change impacts on alfalfa (*Medicago sativa*) production in Ethiopia. *Atmosphere*, 11(10), 1124. <https://doi.org/10.3390/atmos11101124> mdpi.com

Ali, A. M., Abouelghar, M., Belal, A. A., Saleh, N., Yones, M., Selim, A. I., Amin, M. E. S., Elwesemy, A., Kucher, D. E., Maginan, S., & Savin, I. (2022). Crop yield prediction using multi sensors remote sensing: Review article. *The Egyptian Journal of Remote Sensing and Space Sciences*, 25(3), 711–716. <https://doi.org/10.1016/j.ejrs.2022.04.006DOAJ+3ADS Astronomy Data System+3UCI DNTB+3>

Azadbakht, M., Ashourloo, D., Aghighi, H., Homayouni, S., Salehi Shahrabi, H., Matkan, A., & Radiom, S. (2022). Alfalfa yield estimation based on time series of Landsat 8 and PROBA-V images: An investigation of machine learning techniques and spectral-temporal features. *Remote Sensing Applications: Society and Environment*, 25, 100657. <https://doi.org/10.1016/j.rsase.2021.100657> espace.inrs.ca+1UCI DNTB+1

- Bouman, B. A. M., Lampayan, R. M., & Tuong, T. P. (2007). Water management in irrigated rice: Coping with water scarcity. International Rice Research Institute. https://books.irri.org/9789712202193_content.pdfyumpu.com+4books.irri.org+4scirp.org+4
- Brown, M. E., De Beurs, K. M., & Marshall, M. (2012). Global phenological response to climate change in crop areas using satellite remote sensing of vegetation, humidity and temperature over 26 years. *Remote Sensing of Environment*, 126, 174–183.
- Chandel, A. K., Khot, L. R., & Yu, L. X. (2021). Alfalfa (*Medicago sativa* L.) crop vigor and yield characterization using high-resolution aerial multispectral and thermal infrared imaging technique. *Computers and Electronics in Agriculture*, 182, 105991. <https://doi.org/10.1016/j.compag.2021.105991>
- Chartzoulakis, K., & Bertaki, M. (2015). Sustainable water management in agriculture under climate change. *Agriculture and Agricultural Science Procedia*, 4, 88–98. <https://doi.org/10.1016/j.aaspro.2015.03.011>frontiersin.org+1semanticscholar.org+1
- Dhawan, V. (2017). Water and agriculture in India: Background paper for the South Asia expert panel during the Global Forum for Food and Agriculture (GFFA) 2017. OAV – German Asia-Pacific Business Association.
- Eldeiry, A. A., & Garcia, L. A. (2008). Detecting soil salinity in alfalfa fields using spatial modeling and remote sensing. *Agricultural Water Management*, 95(4), 425–433. <https://doi.org/10.1016/j.agwat.2007.11.010>
- Gerardo, R., & de Lima, I. P. (2022). Monitoring duckweeds (*Lemna minor*) in small rivers using Sentinel-2 satellite imagery: Application of vegetation and water indices to the Lis River (Portugal). *Water*, 14(15), 2284. <https://doi.org/10.3390/w14152284>mdpi.com
- Giovas, R., Tassopoulos, D., Kalivas, D., & Lougkos, N. (2021). Remote sensing vegetation indices in viticulture: A critical review. *Agriculture*, 11(5), 457. <https://doi.org/10.3390/agriculture11050457> UCI DNTB+1PMC+1
- Hayashi, A., Akimoto, K., Tomoda, T., & Kii, M. (2013). Global evaluation of the effects of agriculture and water management adaptations on the water-stressed population. *Mitigation and Adaptation Strategies for Global Change*, 18(5), 591–618. <https://doi.org/10.1007/s11027-012-9377-3>
- Jopia, A., Zambrano, F., Pérez-Martínez, W., Vidal-Páez, P., Molina, J., & de la Hoz Mardones, F. (2020). Time-series of vegetation indices (VNIR/SWIR) derived from Sentinel-2 (A/B) to assess turgor pressure in kiwifruit. *ISPRS International Journal of Geo-Information*, 9(11), 641. <https://doi.org/10.3390/ijgi9110641>mdpi.com
- Joshi, A., Pradhan, B., Gite, S., & Chakraborty, S. (2023). Remote-sensing data and deep-learning techniques in crop mapping and yield prediction: A systematic review. *Remote Sensing*, 15(8), 2014. <https://doi.org/10.3390/rs15082014> mdpi.com
- Kazemi Garajeh, M., Haji, F., Tohidfar, M., & Sadeqi, A. (2024). Spatiotemporal monitoring of climate change impacts on water resources using an integrated approach of remote sensing and Google Earth Engine. *Scientific Reports*, 14(1), 12345–12360.
- Kazemi Garajeh, M., Salmani, B., & Zare Naghadehi, S. (2023). An integrated approach of remote sensing and geospatial analysis for modeling and predicting the impacts of climate change on food security. *Scientific Reports*, 13(1), 67890–67905.
- Kingra, P. K., Majumder, D., & Singh, S. P. (2016). Application of remote sensing and GIS in agriculture and natural resource management under changing climatic conditions. *Journal of Agrometeorology*, 18(2), 1–11.

- Kloos, S., Yuan, Y., Castelli, M., & Menzel, A. (2021). Agricultural drought detection with MODIS-based vegetation health indices in southeast Germany. *Remote Sensing*, 13(15), 2930. <https://doi.org/10.3390/rs13152930>
- Kogan, F. (2012). Forecasting crop production using satellite-based vegetation health indices in Kansas, USA. *International Journal of Remote Sensing*, 33(9), 2798–2814. <https://doi.org/10.1080/01431161.2011.621464>
- Kogan, F., Salazar, L., & Roytman, L. (2011). Forecasting crop production using satellite-based vegetation health indices in Kansas, USA. *International Journal of Remote Sensing*, 32(10), 2797–2814. <https://doi.org/10.1080/01431161003749485>
- Kundu, A., Dwivedi, S., & Dutta, D. (2016). Monitoring the vegetation health over India during contrasting monsoon years using satellite remote sensing indices. *Environmental Monitoring and Assessment*, 188(8), 1–14. <https://doi.org/10.1007/s10661-016-5468-1>
- Li, J., Wang, R., Zhang, M., Wang, X., Yan, Y., Sun, X., & Xu, D. (2023). A method for estimating alfalfa (*Medicago sativa* L.) forage yield based on remote sensing data. *Remote Sensing*, 15(3), 657. <https://doi.org/10.3390/rs15030657>
- Ma, C., Johansen, K., & McCabe, M. F. (2022). Monitoring irrigation events and crop dynamics using Sentinel-1 and Sentinel-2 time series. *Remote Sensing*, 14(3), 589. <https://doi.org/10.3390/rs14030589>
- Mancosu, N., Snyder, R. L., Kyriakakis, G., & Spano, D. (2015). Water scarcity and future challenges for food production. *Water*, 7(3), 975–992. <https://doi.org/10.3390/w7030975>
- Maselli, F., Battista, P., Chiesi, M., Rapi, B., & Angeli, L. (2020). Use of Sentinel-2 MSI data to monitor crop irrigation in Mediterranean areas. *International Journal of Applied Earth Observation and Geoinformation*, 92, 102172. <https://doi.org/10.1016/j.jag.2020.102172>
- Montazar, A., & Putnam, D. (2023). Evapotranspiration and yield impact tools for more water-use efficient alfalfa production in desert environments. *Agronomy*, 13(2), 345. <https://doi.org/10.3390/agronomy13020345>
- Möllmann, J., Buchholz, M., Kölle, W., & Musshoff, O. (2020). Do remotely-sensed vegetation health indices explain credit risk in agricultural microfinance? *Agricultural Finance Review*, 80(5), 611–627. <https://doi.org/10.1108/AFR-10-2019-0109>
- Mpandeli, S., Nhamo, L., Moeletsi, M., & Masupha, T. (2019). Assessing climate change and adaptive capacity at local scale using observed and remotely sensed data. *Science of the Total Environment*, 653, 59–70.
- Muruganantham, P., Wibowo, S., Grandhi, S., & Samrat, N. H. (2022). A systematic literature review on crop yield prediction with deep learning and remote sensing. *Remote Sensing*, 14(9), 2071. <https://doi.org/10.3390/rs14092071>
- Noland, R. L., Wells, M. S., Coulter, J. A., Tiede, T., & Baker, J. M. (2018). Estimating alfalfa yield and nutritive value using remote sensing and air temperature. *Field Crops Research*, 222, 189–197. <https://doi.org/10.1016/j.fcr.2018.03.017>
- Nong, D. H., Ngo, A. T., Nguyen, H. P. T., & Nguyen, T. T. (2021). Changes in coastal agricultural land use in response to climate change. *Land*, 10(12), 1345.
- Ondrasek, G. (2013). Water scarcity and water stress in agriculture. In *Sustainable Agriculture Reviews* (pp. 125–157). Springer. https://doi.org/10.1007/978-94-007-5961-9_5
- Pande, C. B., & Moharir, K. N. (2023). Application of hyperspectral remote sensing role in precision farming and sustainable agriculture under climate change: A review. *Environmental Monitoring and Assessment*, 195(8), 456.

- Pereira, L. S. (2017). Water, agriculture and food: Challenges and issues. In *Water and Sustainability in Arid Regions* (pp. 35–58). Springer. https://doi.org/10.1007/978-94-007-4032-7_3
- Pereira, L. S., Oweis, T., & Zairi, A. (2002). Irrigation management under water scarcity. *Agricultural Water Management*, 57(3), 175–206. [https://doi.org/10.1016/S0378-3774\(02\)00075-6](https://doi.org/10.1016/S0378-3774(02)00075-6)
- Psomiadis, E., Dercas, N., & Dalezios, N. R. (2017). Evaluation and cross-comparison of vegetation indices for crop monitoring from Sentinel-2 and WorldView-2 images. In *Proceedings of SPIE 10444, Remote Sensing for Agriculture, Ecosystems, and Hydrology XIX* (Vol. 10444, pp. 104440I). <https://doi.org/10.1117/12.2278377>
- Qadir, M., Boers, T. M., Schubert, S., Ghafoor, A., & Murtaza, G. (2003). Agricultural water management in water-starved countries: Challenges and opportunities. *Agricultural Water Management*, 62(3), 165–185. [https://doi.org/10.1016/S0378-3774\(03\)00045-5](https://doi.org/10.1016/S0378-3774(03)00045-5)
- Qian, Y., Hou, Y., Yan, H., Mao, L., & Wu, M. (2012). Global crop growth condition monitoring and yield trend prediction with remote sensing. *Journal of Integrative Agriculture*, 11(1), 1–10. [https://doi.org/10.1016/S2095-3119\(12\)60002-4](https://doi.org/10.1016/S2095-3119(12)60002-4)
- Ramos, T. B., Castanheira, N., Oliveira, A. R., & Paz, A. M. (2020). Soil salinity assessment using vegetation indices derived from Sentinel-2 multispectral data: Application to Lezíria Grande, Portugal. *International Journal of Applied Earth Observation and Geoinformation*, 92, 102174. <https://doi.org/10.1016/j.jag.2020.102174>
- Rembold, F., Atzberger, C., Savin, I., & Rojas, O. (2013). Using low-resolution satellite imagery for yield prediction and yield anomaly detection. *Remote Sensing*, 5(4), 1704–1733. <https://doi.org/10.3390/rs5041704>
- Rozenstein, O., Haymann, N., & Kaplan, G. (2018). Estimating cotton water consumption using a time series of Sentinel-2 imagery. *International Journal of Applied Earth Observation and Geoinformation*, 73, 247–255. <https://doi.org/10.1016/j.jag.2018.06.015>
- Segarra, J., Buchailot, M. L., Araus, J. L., & Kefauver, S. C. (2020). Remote sensing for precision agriculture: Sentinel-2 improved features and applications. *Agronomy*, 10(5), 641. <https://doi.org/10.3390/agronomy10050641>
- Setiyono, T., Nelson, A., & Holecz, F. (2014). Remote sensing-based crop yield monitoring and forecasting. *Asian Journal of Geoinformatics*, 14(4), 1–10.
- Shanmugapriya, P., Rathika, S., & Ramesh, T. (2019). Applications of remote sensing in agriculture – A review. *International Journal of Current Microbiology and Applied Sciences*, 8(1), 2270–2283. <https://doi.org/10.20546/ijemas.2019.801.238>
- Sholihah, R. I., Trisasongko, B. H., & Shiddiq, D. (2016). Identification of agricultural drought extent based on vegetation health indices of Landsat data: Case of Subang and Karawang, Indonesia. *Procedia Environmental Sciences*, 33, 14–20. <https://doi.org/10.1016/j.proenv.2016.03.056>
- Tedesco, D., Nieto, L., Hernández, C., Rybecky, J. F., & Min, D. (2022). Remote sensing on alfalfa as an approach to optimize production outcomes: A review. *Agronomy*, 12(3), 630. <https://doi.org/10.3390/agronomy12030630>
- Tzanakakis, V. A., Paranychianakis, N. V., & Angelakis, A. N. (2020). Water supply and water scarcity. *Water*, 12(9), 2347. <https://doi.org/10.3390/w12092347>

Varghese, D., Radulović, M., Stojković, S., & Crnojević, V. (2021). Reviewing the potential of Sentinel-2 in assessing the drought. *Remote Sensing*, 13(5), 935. <https://doi.org/10.3390/rs13050935>

Vidican, R., Mălinaş, A., Ranta, O., Moldovan, C., & Marian, O. (2023). Using remote sensing vegetation indices for the discrimination and monitoring of agricultural crops: A critical review. *Remote Sensing*, 15(3), 657. <https://doi.org/10.3390/rs15030657>

You, J., Li, X., Low, M., Lobell, D., & Ermon, S. (2017). Deep Gaussian process for crop yield prediction based on remote sensing data. *Proceedings of the Thirty-First AAAI Conference on Artificial Intelligence* (pp. 4559–4565).

Zhou, Y., Flynn, K. C., Gowda, P. H., Wagle, P., & Ma, S. (2021). The potential of active and passive remote sensing to detect frequent harvesting of alfalfa. *Agricultural and Forest Meteorology*, 307, 108531. <https://doi.org/10.1016/j.agrformet.2021.108531>

Zhu, X., Guo, R., Liu, T., & Xu, K. (2021). Crop yield prediction based on agrometeorological indexes and remote sensing data. *Remote Sensing*, 13(8), 1514. <https://doi.org/10.3390/rs13081514>

APPLICATIONS TO ENHANCE POLLEN QUALITY IN WATERMELONS WITH DIFFERENT CHROMOSOME STRUCTURES

Dr. Pınar ADIGÜZEL

Cukurova University, Agriculture Faculty, Department of Horticulture, Adana, Türkiye

Doç. Dr. Şenay KARABIYIK

Cukurova University, Agriculture Faculty, Department of Horticulture, Adana, Türkiye

Prof. Dr. İlknur SOLMAZ

Cukurova University, Agriculture Faculty, Department of Horticulture, Adana, Türkiye

ABSTRACT

Watermelon (*Citrullus lanatus* L.) is a vegetable species of significant commercial importance both worldwide and in our country. Currently, triploid (seedless) watermelons have become preferred by consumers. The seed that will produce triploid watermelon is obtained from the fruit formed by crossing a tetraploid female with a diploid male parent. The abundance and high quality of the pollen from the pollinator parent also increase the fruit set rate. Various chromosome structures, including diploid (2n), triploid (3n), and tetraploid (4n) watermelons, can present differences in pollen viability and fertilization. This paper explores applications aimed at improving pollen quality in watermelons with diverse chromosome structures. In the study, one tetraploid watermelon line was used along with two male diploid parent lines one with a previously determined high pollination effect and another with a low pollination effect. Within the scope of the research, 8 ppm Putrescine and 150 g/ 100 L Bor, were applied to the female parents via foliar spray, while 7.5 g/L Bor was applied to the male parents. The experimental results were analyzed using the JMP 5.0.1 software, based on the completely randomized design. The study found that boron application in the tetraploid female and male parents increased pollen viability and germination rates. More effective pollen tube growth was observed in male parents treated with boron, and female parents treated with putrescine.

Keywords: Watermelon, Putrescine, Boron, Pollen.

INTRODUCTION

Watermelon (*Citrullus lanatus* (Thunb.) Matsum. and Nakai) is a 22-chromosome vegetable species belongs to the Cucurbitaceae family, and has a high economic value (Zhang et al., 2019; Atlı et al. 2021; Paris, 2023). The global production of watermelon amounts to 104,932,071 tons, while Türkiye produced 3,147,921 tons (FAO, 2023). Watermelons are generally diploid ($2n=2x=22$) but can also be triploid ($3n=3x=33$) or tetraploid ($4n=4x=44$) (Blakeslee and Avery, 1937). The increase in chromosome number provides advantages such as growth in vegetative stages and resistance to biotic and abiotic stress factors (Soltis et al., 2016; Godfree et al., 2017). Seedling deaths during plant stratification, thickening of the tetraploid seed coat and decreases in germination and emergence rates are caused by colchicine, which is used in

obtaining tetraploid watermelons (Adıgüzel and Sarı, 2019; Zhang et al., 2019; Adıgüzel et al., 2022). Therefore, the pollination capacity of the tetraploid parent is important.

For healthy pollination to occur, the stigma must be reached by quality flower pollen, which must then reach healthy ovules (Razzaq et al. 2019; Karabıyık, 2022). Pollination capacity depends on the presence of large amounts of healthy pollen (Shivanna and Rangaswamy, 1992; Gaaliche et al. 2013). *In vitro* pollen viability and germination tests are used as a practical method to determine pollen quality (Chatterjee et al. 2014; Sulusoglu and Cavusoglu, 2014; Albert et al. 2018; Adıgüzel et al. 2023). Flower pollen is greatly affected by stress factors such as temperature, drought, and salinity (Saragih et al. 2013; Turner et al. 2013; Sakhi et al. 2014; Razzaq et al. 2017; Karabıyık, 2022). The germination of flower pollens is supported by the storage of nutrients in them (Tosun and Koyuncu, 2007). In this context, boron is considered one of the most important elements for pollen quality (Pinho et al. 2010; Fang et al. 2016; Fang et al. 2019). Photosynthetic activity, yield, quality, and plant hormone mechanisms are affected by boron (Shorrocks, 1997; Rerkasem et al. 2004; Deepika and Pitagi, 2015; Barut et al. 2018; Adıgüzel et al. 2023). In this case, seed formation, which is associated with the healthy development of flower pollen, is also positively affected. It has been stated that foliar boron application is an effective method for eliminating boron deficiency and that boron is transmitted from the phloem to the roots (Du et al. 2020). In studies on boron, it has been reported that it increases the quality of pollen (Adıgüzel et al. 2023).

Putrescine increases plant tolerance to ecological events and supports plant development under stress conditions (Andronis et al., 2014). When the studies are examined, evidence suggests that putrescine increases fruit quality by contributing positively to pollination activity (Saleem et al., 2008; Stern and Gazit, 2000; Franco-Mora et al., 2005; Karabıyık, 2023; Dölek and Karabıyık, 2023). Putrescine, which plays a role in the healthy development of the female flower and ovules, is more important than polyamines, especially in growth, development and tolerance to abiotic stress conditions (Thomas et al., 2020). Putrescine is also effective in promoting root growth, ovule development, embryo development, fruit ripening and the plant defense system (Couée et al., 2004; Liu et al., 2015; González-Hernández et al., 2022).

Triploid watermelons, which are obtained by hybridizing tetraploid and diploid watermelons, are highly popular among consumers today due to their ease of eating (Solmaz et al., 2018). This demand also drives producer companies toward seedless watermelon breeding. In this context, the tetraploid parent and diploid male parent used in triploid production are important. In cases where the tetraploid parent is receptive, the diploid parent must also possess pollenizer potential and its pollen must be able to pollinate simultaneously. In this case, the higher the pollination ability of the pollenizer male parent, the more positively the fruit set rate is affected. Boron used in the study has significant effects on the viability and germination rate of pollen, while putrescine has significant effects on the development of ovules in female flowers. Therefore, in this study, the aim was to increase the quality of pollens and ovules by applying putrescine or boron to tetraploid female parents and boron to diploid male parents.

MATERIAL AND METHODS

The study was carried out in 2022 in the glass greenhouse located in the Research and Application areas of the Department of Horticulture, Faculty of Agriculture, Çukurova University, and the pollen and histological studies were conducted in the Cytology and Histology Laboratory at the same department. ST 101 (tetraploid), WL 216 (diploid), and WL 259-B (diploid) watermelon varieties were used as material in the study. Three watermelon varieties were planted, and the study was set up with three replications. A total of 96 plants were used in the study: 48 ST 101 (4 plants x 3 replicates x 4 treatments), 24 WL 216, and 24 WL 259-B (plants x 3 replicates x 2 treatments). Plants were grown in suspension in the

greenhouse. Boron applications were made using 150 g/100 L Eti-Dot (21% B) via foliar application when the plants were at 7 leaved stage. In addition, 8 ppm putrescine was applied to the ST 101 watermelon genotype at first female flower were started to open. As control treatment, only water was applied to the plants during the same periods as the treatments. Male flowers were taken for pollen germination and viability tests 15 days after the treatments. In addition, male flowers that had not yet bloomed, but would bloom in a day, were taken in order to determine the amount and the homogeneity of pollen production. Cultural practices such as pruning, irrigation and fertilization were applied throughout the growing period.

Pollen viability and germination tests

To provide material for pollen viability and germination tests, at least 15 male flowers were taken from each treatment 15 days after the treatments and the tests were then performed. In this context, pollen viability was determined using 1% 2,3,5 Triphenyl Tetrazolium Chloride (TTC) solution (Norton, 1966). The viability of the pollen was determined by preparing 3 coverslips and counting at least 100 pollen grains on each coverslip. During the microscope examination, pollen grains stained dark red were considered absolutely viable, light red and pink pollen grains were considered semi-viable, and pollen grains not stained at all were considered non-viable (Karabıyık, 2022). Theoretically, half of the pollen, referred to as semi-viable, is assumed to be viable, and this value is added to the absolute viable flower pollen to calculate the "pollen viability level". The pollen germination test was performed using the 'agar in petri' method in 1% agar + 15% sucrose medium added to 50 ppm boric acid, at 25°C (Adıgüzel et al., 2022). In order to determine the level of pollen germination, 3 petri dishes were prepared and at least 100 pollen grains were counted in each petri dish. During the microscope examination, pollen with pollen tubes longer than their own diameter were accepted as germinated and the "pollen germination level" was calculated by dividing the number of germinated pollen by the total number of pollen.

Pollen production amount and normal developed pollen rate

In order to determine the amount of pollen production, 15 male flowers that had not yet opened but would open within a day were taken for each application. Three groups of 5 were made and the anthers were separated from their filaments. Each group was placed in separate small boxes. According to Karabıyık and Sarıdaş (2023), pollen counting and calculations were carried out using the "hemocytometric method" in the anthers dried for about two weeks. The "amount of pollen in a flower" was determined by counting. During the counting, pollens showing abnormal development were also determined and the "normally developed pollen rate" was determined by the ratio calculation method (Anvari, 1977).

Pollen tube development

To obtain material for pollen tube growth analysis, a hybridization process was carried out alongside the hybridization program. In this regard, three pistils from each combination were collected at 12, 24, and 48 hours after pollination (HAP) and immediately fixed in FPA-70. The samples were analyzed using the "squash preparation" method as described by Kho and Baer (1970) and Karabıyık (2022). In this technique, the samples were softened with 8N NaOH and subsequently stained with aniline blue mordanted with K_3PO_4 . The germination of pollen tubes on the stigma, their progression through the style, and their arrival at the ovules were observed using an Olympus BX 51 fluorescent microscope (Olympus, Japan).

Statistical analysis

The experiment was carried out according to the randomized plots with 3 replications. Variance analysis was applied according to the factorial experimental design in randomized plots in the JMP 5.0.1 package programme. The differences between the means were classified at the 5% significance level according to the LSD test. Arc-sin transformation was applied to the percentage values.

RESULTS AND DISCUSSION

The effects of boron and putrescine applications on pollen viability and germination rate in tetraploid watermelons are given in Table 1. In this context, the highest values in viability (96.17%), germination (61.12%) and absolute viable (95.09%) were obtained as a result of foliar boron application to the ST 101 watermelon genotype. The lowest pollen viability was determined in the control group (76.55%), while the pollen germination rate in the putrescine application was 33.20%. Absolute viability was 74.45% in the control group and 72.10% in the putrescine application group. No statistically significant difference was found between the applications in terms of semi-viable and non-viable pollen. Also, no effect of boron and putrescine applications on the ratio of normally developed pollen and pollen production level in a flower was detected (Table 2).

Table 1. Effects of boron and putrescine applications on pollen viability and germination levels in tetraploid watermelon (%)

Application	Absolutely viable	Semi-viable	Non-viable	Viability rate	Germination rate
Putrescine	72.10 b	15.50	12.40	79.85 bc	33.20 c
Boron	95.09 a	2.16	2.75	96.17 a	61.12 a
Putrescine + Boron	87.16 ab	7.22	5.63	90.76 ab	52.88 ab
Control	74.45 b	4.19	2.37	76.55 c	37.98 bc
P _{0,05}	12.073*	N.S.	N.S.	10.628**	10.211*

NS: Not Significant; ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$: shows difference according to LSD comparison. And arc/sin transformation was made to percentage values.

Table 2. Effects of boron and putrescine applications on pollen number (pollen/flower) and normal developed pollen rate (%) in tetraploid watermelon

Application	Number of pollen grains in a flower	Normally developed pollen rate
Putrescine	155 333	93.76
Boron	162 444	92.19
Putrescine + Boron	162 666	93.70
Control	138 444	92.98
P _{0,05}	N.S.	N.S

NS: Not Significant; ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$: shows difference according to LSD comparison. And arc/sin transformation was made to percentage values.

It was determined that boron application had no statistically significant effect on the formation of diploid watermelon pollen, which was characterized as absolute and semi-viable pollen ratios (Table 3). In addition, while boron application increased the non-viable pollen in the control group of WL259-B genotype (8.24%), this rate decreased in the control group of WL216 genotype (0.00%) and in the boron application for the WL259-B genotype (1.22%) (Table 4). When the effect of boron application on the germination rate in diploid watermelons was examined, it showed no genotypic differences and boron application (70.28) provided higher pollen germination than the control (60.77%) (Table 5). Table 6 shows the data on the pollen count (pollen/flower) and the normally developed pollen rate in diploid watermelons after boron application. When the data were examined, it was determined that the application did not affect the rate of normally developed pollen, and only the genotype was effective in the pollen count in a flower. It was determined that the number of pollen grains in a flower belonging to the WL259-B genotype was higher (193 600 pollen/flower.), while WL216 had a lower pollen grain count than that of the WL259-B genotype (140300 pollen/flower).

Table 3. Effect of boron application on absolute viable (%) and semi- viable (%) pollen counts in diploid watermelons

Genotype	Absolutely viable pollen rate			Semi-viable pollen rate		
	Application		Genotype avg.	Application		Genotype avg.
	Boron	Control		Boron	Control	
WL 216	91.77	89.42	90.60	4.73	10.58	7.66
WL 259	95.62	84.96	90.29	3.16	6.80	4.98
Application avg.	93.70	87.19		3.95	8.69	
P	LSD _{genotype} : N.S.; LDS _{application} : N.S.; LSD LSD _{genotype x application} : N.S.			LSD _{genotype} : N.S.; LDS _{application} : N.S.; LSD _{genotype x application} : N.S.		

NS: Not Significant; ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$: shows difference according to LSD comparison. And arc/sin transformation was made to percentage values.

Table 4. Effect of boron application on non-viable and viable pollen in diploid watermelons (%)

Genotype	Non-viable pollen rate			Viable pollen rate		
	Application		Genotype avg.	Application		Genotype avg.
	Boron	Control		Boron	Control	
WL 216	3.50 ab	0.00 b	1.75	94.14	94.71	94.42
WL 259	1.22 b	8.24 a	4.73	97.20	88.36	92.78
Application avg.	2.36	4.12		95.67	91.54	
P	LSD _{genotype} : N.S.; LDS _{application} : N.S.; LSD LSD _{genotype x application} : 8.418**			LSD _{genotype} : N.S.; LDS _{application} : N.S.; LSD _{genotype x application} : N.S.		

NS: Not Significant; ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$: shows difference according to LSD comparison. And arc/sin transformation was made to percentage values.

Table 5. Effect of boron application on germination rate (%) in diploid watermelons

Genotype	Pollen germination rate		Genotype avg.
	Application		
	Boron	Control	
WL 216	63.37	60.33	61.85
WL 259	77.18	61.20	69.19
Application avg.	70.28 A	60.77 B	
P	LSD _{genotype} : N.S.; LDS _{application} : 5.952*; LSD _{genotype x application} : N.S.		

NS: Not Significant; ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$: shows difference according to LSD comparison. And arc/sin transformation was made to percentage values.

Table 6. Effect of boron application on the pollen count (number/flower) and normally developed pollen rate (%) in diploid watermelons

Genotype	Number of pollen grains in a flower		Genotype avg.	Normally developed pollen rate		Genotype avg.
	Application			Application		
	Boron	Control		Boron	Control	
WL 216	135777	144822	140300B	93.3	93.6	93.4
WL 259	231777	155422	193600A	96.9	94.2	95.5
Application avg.	183777	150122		95.1	93.9	
P	LSD _{genotype} : 45074.7* LDS _{application} : N.S.; LSD _{genotype x application} : N.S.			LSD _{genotype} : N.S.; LDS _{application} : N.S.; LSD _{genotype x application} : N.S.		

NS: Not Significant; ***: $P \leq 0.001$; **: $P \leq 0.01$; *: $P \leq 0.05$: shows difference according to LSD comparison. And arc/sin transformation was made to percentage values.

Quality pollen positively affects the pollination of the female flower and increases fruit set (Chapman and Goring, 2010; Maiti et al., 2016; Wijesinghe et al., 2020; Adıgüzel et al., 2023 b,c). High and low temperatures affect flower biology, pollen viability, ovule development and viability, and pollen tube development (Hedhly et al., 2005; Bykova et al., 2012; Karabıyık, 2022). Indeed, previous studies have reported that boron applications vary depending on weather conditions (Gürsöz 1990; Nakamura and Wheeler 1992; Adıgüzel et al. 2023) and the watermelon variety used (Şensoy et al. 2003; Freeman et al. 2008). In addition, it has been reported that putrescine, a polyamine (Kusano et al., 2008; Do et al., 2013), positively affects ovule development in the female flower and increases pollen germination and pollen tube development (Ewart and Kliewer, 1977; Karabıyık, 2024). In this sense, it is important to implement applications that will prevent pollen from stress conditions. Boron applications in melon (Goldberg et al. 2003; Hidayat et al. 2021) and squash (Ansari et al. 2018) determined that boron positively affects the quality and quantity of pollen (Namlı et al., 2024).

Pollen viability and germination rates vary depending on environmental factors and plant genetic structure (Johri and Vasil 1960; La Porta and Roselli, 1991; Nepi and Pacini, 1993; Luo et al. 2020). In one of the first studies on pollen in watermelon in our country, Gürsöz (1990) stated that the germination ability of pollen was 100% on the first day and this value decreased to 40% after 21 days. However, it is thought that this situation will vary depending on the genotype effect and the conditions in which the plant is grown. Tetraploid plants generally bloom earlier than diploids (Adıgüzel et al., 2022). In addition, flower sizes vary according to chromosome number (Kihara, 1951). It is known that this situation affects pollen viability and

germination rate. In a study examining tetraploid and diploid pollens, it was found that diploid watermelons had a higher pollen germination rate than tetraploids (Adıgüzel et al. 2022). It has been reported that pollen viability rates may yield different results from germination rates (Şensoy et al. 2003; Freeman et al. 2008). In this study, it was determined that boron and putrescine application to tetraploid plants increased pollen absolute viability rates and pollen germination rates. Boron, which is the most effective in enhancing absolute pollen viability, had the same effect on pollen germination rate. Putrescine, on the other hand, had lower levels than the control. In addition, boron increased pollen germination rate in both tetraploids and diploids. Since boron is a microelement that has a positive effect on pollen viability and germination (Namlı et al., 2024), the results are consistent with expectations. Putrescine, which is more effective on female flower development, showed the lowest values in male flowers when applied in the study. Based on these studies, it can be said that pollen germination and viability rates are affected by genotype, environmental conditions, flower gender, application and time, and vary depending on soil conditions.

As with many plant species, pollen counts vary in cucurbits (Wijesinghe et al., 2020). For example, in a study on watermelon genetic resources, pollen counts varied between 175745 and 296,022 (Adıgüzel et al., 2025). This change in the amount of pollen per flower is due to environmental effects, agricultural applications, grafting and the variety used, even in plants in the same population (Cruzan, 1990; Nakamura and Wheeler, 1992; Adıgüzel et al., 2022; Aras et al., 2022; Namlı et al., 2024). In a study investigating the effect of different chromosome numbers on pollen, it was reported that pollen numbers varied between 76,760,650.02 (tetraploid) and 112,1120,604.95 (diploid), with diploids having more pollen (Adıgüzel et al., 2022). Studies investigating the effect of boron on pollen have reported that the quality and quantity of pollen increase (Goldberg et al. 2003; Ansari et al. 2018; Hidayat et al. 2021; Rahayu et al. 2023). This situation is thought to be because the boron increases the amount of pollen while not spoiling the shape, size and quality of the pollen. In this study, it was determined that boron and putrescine did not affect the pollen numbers in flowers with different chromosome structures. However, this situation may vary depending on the putrescine dose, chromosome structure, and genotype used.

The rate of normally developed pollen grains is an important criterion, as it provides information about pollen quality. The higher the rate of normally developed pollen in a variety, the better its potential as an effective pollenizer (Anvari, 1977; Karabıyık, 2011). It has been stated that as long as this ratio is above 80%, the pollen quality is sufficient (Eti, 1990). According to the results of the study, it was found that boron and putrescine did not have any effect on the normally developed pollen in both tetraploid and diploid flowers. In other studies, on boron application, it was found that it had no effect on the normally developed pollen rate (Sarıdaş et al., 2021; Adıgüzel et al. 2023). It has also been reported that this situation is affected by pollen quality and weather conditions (Saragih et al. 2013; Turner et al. 2013; Sakhi et al. 2014; Razzaq et al. 2017; Karabıyık, 2022).

Pollen tube development

Pollen tube development was determined with histological studies by pistil squash method. In these studies, it has been determined that more effective pollen tube development occurs especially in the cross pollination made by male parents applied with boron and female parents applied with putrescine. This is because of the putrescine enhances the ovule viability and quality while boron enhances pollen viability and tube growth velocity. In the detailed histological studies it was found that the ovules were more quality in whole ovary than control group in putrescine treated tetraploids. At the same time, it has been showed that in Table 5, the Boron treatments increased the pollen germination level. With these results with quality

female and quality male interaction the pollen tubes were accelerated and a more effective pollen tube penetration can be seen. Pollen tube development in stigma, style and penetration to the ovules can be seen in Figure 1.




Figure 1. Pollen tube development in putrescine treated tetraploid x boron treated diploids and ovules in Control ovary. A. Pollen tube germination on stigma. B. Pollen tube elongation through the style. C. Quality and homogenous ovule development in putrescine treated tetraploid. D. Non-homogenized ovule development in Control treated tetraploid ovary.

If the stigma and pollen are compatible, pollen grains progress successfully towards the cells in the ovary (Rottmann et al., 2018). The pollen tube is affected by conditions such as sexual incompatibility (Distefano et al. 2012; Zhang et al. 2018) and pollination time (Kantoğlu and Yanmaz, 1996; Adıgüzel et al., 2023). Polyamines are also effective in pollen germination and pollen tube elongation (Çetinbaş-Genç, 2020a; Çetinbaş-Genç et al., 2020b). Karabıyık and Çağlar (2023) found that 16 ppm putrescine dose increased pollen tube development in apples. It is accepted that polyamine affects pollen germination and tube growth in a dose-dependent manner (Aloisi et al., 2015; Çetinbaş-Genç, 2019). In this study pollen tube growth was enhanced in cross pollination in putrescine treated tetraploid female and boron treated diploid male. Considering other studies, it is thought that pollen tube development varies according to ovule and pollen quality, genotype, boron dose applied and application method.

CONCLUSION

Triploid watermelons are produced by hybridizing tetraploid and diploid flowers. The process of obtaining tetraploid plants is difficult and expensive. For this reason, it is very important to have healthy tetraploid female and a high quality and quantity of diploid male pollen in order to obtain triploidy. For this purpose, boron was applied to the male plant to be used as pollenizer, and putrescine was applied to the tetraploid plant used as female. The results obtained indicate that boron increased the pollen quality in both tetraploids and diploids. In addition, the interactions of boron and putrescine also increase the quality of pollen. No study has been found examining the effects of putrescine and boron on flowers with different chromosome structures. Therefore, the doses used in the study have been a reference in other

studies. As a result, applying boron to pollinating male plants increases the quality and quantity of pollen. Therefore, it is considered to be appropriate to use it as a standard fertilizer in studies.

REFERENCES

- Adıgüzel, P., Karabıyık Ş., Namlı, M., Solmaz İ. (2023). Karpuzda (*Citrullus lanatus* L.) Bor Uygulamasının Çiçek Tozu Kalitesi, Miktarı ve Çim Borusu Gelişimine Etkisi. *Alatarım*. 22(2):76-85.
- Adıgüzel, P., Karabıyık, Ş., Namlı, M., Solmaz, İ. (2025). Bazı Karpuz Genetik Kaynaklarında Çiçek Tozu Performansının Belirlenmesi. 4. Bilsel International Aspendos Scientific Researches. 1-2 Şubat 2025-Antalya. s. 113-123.
- Adıgüzel, P., Sarı, N. (2019). Effect of Different Male Parents and Male Flower Numbers on Seed Yield and Quality of Triploid Watermelon Breeding. *International Journal of Environmental Research and Technology*. 2(3): 14-24.
- Adıgüzel, P., Solmaz, İ., Karabıyık, Ş. Sarı, N. (2022). Comparison on Flower, Fruit and Seed Characteristics of Tetraploid and Diploid Watermelons (*Citrullus lanatus* Thunb. Matsum. and Nakai). *International Journal of Agriculture Environment and Food Sciences*. 6(4):704-710.
- Albert, B., Ressayre, A., Dillmann, C., Carlson, A. L., Swanson, R. J., Gouyon, P.- H. (2018). Effect of Aperture Number on Pollen Germination, Survival and Reproductive Success in *Arabidopsis thaliana*. *Ann. Bot.* 121, 733–740. doi: 10.1093/aob/mcx206
- Aloisi, I., Cai, G., Tumiatti, V., Minarini, A., & Del Duca, S. (2015). Natural Polyamines and Synthetic Analogs Modify the Growth and the Morphology of *Pyrus communis* Pollen Tubes Affecting ROS Levels and Causing Cell Death. *Plant Science*. 239, 92-105.
- Andronis, E.A., Moschou, P.N., Toumi, I. and Roubelakis-Angelakis, K. A. (2014). Peroxisomal Polyamine Oxidase and NADPH-Oxidase Cross-Talk for ROS Homeostasis which Affects Respiration Rate in *Arabidopsis thaliana*. *Frontiers in Plant Science*. 5: 132.
- Ansari, A. M., Chowdhary, B. M. (2018). Effects of Boron and Plant Growth Regulators on Bottle Gourd (*Lagenaria siceraria* (Molina) Standl.). *Journal of Pharmacognosy and Phytochemistry*. 7(1):202-206.
- Anvari, S. F. (1977). Untersuchungen über Das Pollenschlauchwachstum und die Entwicklung der Samenanlagen in Beziehung Zum Fruchtsatz Bei Sauerkirchen (*Prunus cerasus* L.) Diss. Univ. Hohenheim. 105
- Aras, V., Sarı, N., Solmaz, İ. (2022). Effects of *Cucurbita*, *Lagenaria* and *Citrullus* Rootstocks on Pollen and Fruit Characters, Seed Yield and Quality of F1 Hybrid Watermelon. *International Journal of Agriculture Environment and Food Sciences*. 6(4), 683-693.
- Atlı, E., Solmaz, İ., Sarı, N. and Kelebek, H. (2021). Determination of Some Agronomic and Fruit Quality Characteristics of Some Watermelon Accessions from Turkish Watermelon Germplasm. *Turkish Journal of Agriculture-Food Science and Technology*. 9(8), 1322-1328.
- Barut, H., Aykanat, S., Aşıklı, S., Selim, E. (2018). “Bitkisel Üretimde Bor”, *International Journal of Eastern Mediterranean Agricultural Research*. 1(1), 33-46.
- Blakeslee, A. F., Avery, A. G. (1937). Methods of Inducing Doubling of Chromosomes in Plants: by Treatment With Colchicine. *Journal of Heredity*. 28 (12): 393-411. <https://doi.org/10.1093/oxfordjournals.jhered.a104294>
- Bykova, O., Chuine, I., Morin, X. and Higgins, S.I. (2012). Temperature Dependence of the Reproduction Niche and its Relevance for Plant Species Distributions. *Journal of Biogeography*. 39(12), 2191-2200.
- Chapman, L. A., and Goring, D. R. (2010). Pollen–Pistil Interactions Regulating Successful Fertilization in the *Brassicaceae*. *Journal of Experimental Botany*. 61(7):1987-1999.
- Chatterjee, R., Sarkar, S., Rao, G. N. (2014). Improvised Media for in Vitro Pollen Germination of Some Species of *Apocynaceae*. *Int. J. Environ.* 3:146-153.

- Couée, I., Hummel, I., Sulmon, C., Gouesbet, G., El Amrani, A. (2004). Involvement of Polyamines in Root Development. *Plant Cell Tissue Organ Culture (PCTOC)*. 76, 1-10.
- Cruzan, M.B. (1990). Variation In Pollen Size, Fertilization Ability, and Post-fertilization Siring Ability In *Erythronium grandiflorum*. *Evolution*. 44:843-856.
- Çetinbaş-Genç, A. (2020a). Putrescine Modifies the Pollen Tube Growth of Tea (*Camellia sinensis*) by Affecting Actin Organization and Cell Wall Structure. *Protoplasma*. 257(1), 89-101.
- Çetinbaş-Genç, A., Cai, G., Del Duca, S., Vardar, F., & Ünal, M. (2020b). The Effect of Putrescine on Pollen Performance in Hazelnut (*Corylus avellana* L.). *Scientia Horticulturae*. 261, 108971.
- Deepika, C., and Pitagi, A. (2015). Effect of Zinc and Boron on Growth, Seed Yield and Quality of Radish (*Raphanus sativus* L.) cv. Arka Nishanth. *Current Agriculture Research Journal*. 3(1): 85-89.
- Distefano, G., Hedhly, A., Casas, G. L., Malfa, S., Herrero, M. and Gentile, A. (2012). Male-Female Interaction and Temperature Variation Affect Pollen Performance in Citrus. *Scientia Horticulturae*. 140, 1-7
- Do, P. T., Degenkolbe, T., Erban, A., Heyer, A. G., Kopka, J., Köhl, K. I., ... & Zuther, E. (2013). Dissecting Rice Polyamine Metabolism under Controlled Long-Term Drought Stress. *PLoS One*. 8(4), e60325.
- Dölek, C. and Karabıyık, Ş. (2023). Effects of Putrescine and Spermine on Pollen Germination Levels of some Olive Cultivars. 3rd International Conference on Research of Agriculture and Food Technologies Proceeding Book. 4-6 October 2023. Pp:59.
- Du, W., Pan, Z. Y., Hussain, S. B., Han, Z. X., Peng, S. A., & Liu, Y. Z. (2020). Foliar Supplied Boron can be Transported to Roots as a Boron-Sucrose Complex via Phloem in citrus trees. *Frontiers in Plant Science*. 11: 250.
- Eti, S. (1990). Çiçek Tozu Miktarını Belirlemede Kullanılan Pratik Bir Yöntem. *Ç.Ü. Ziraat Fakültesi Dergisi*. 5(4): 49-58.
- Ewart, A., & Kliewer, W. M. (1977). Effects of Controlled Day and Night Temperatures and Nitrogen on Fruit-Set, Ovule Fertility, and Fruit Composition of Several Wine Grape Cultivars. *American Journal of Enology and Viticulture*. 28(2), 88-95.
- Fang, K., Zhang, W., Xing, Y., Zhang, Q., Yang, L., Cao, Q. (2016). Boron Toxicity Causes Multiple Effects on *Malus domestica* Pollen Tube Growth. *Frontiers in Plant Science*. 7:208.
- Fang, K. F., Du, B. S., Zhang, Q., Xing, Y., Cao, Q. Q., & Qin, L. (2019). Boron Deficiency Alters Cytosolic Ca²⁺ Concentration and Affects the Cell Wall Components of Pollen Tubes in *Malus domestica*. *Plant Biology*. 21(2), 343-351.
- FAOSTAT, 2023 <http://www.fao.org> (Access Date: 08.03.2025)
- Franco-Mora, O., Tanabe, K., Tamura, F. and Itai, A. (2005). Effects of Putrescine Application on Fruit Set in 'Houshi' Japanese Pear (*Pyrus Pyrifolia*). *Scientia Horticulturae*, 104: 265-273.
- Freeman, J. H., Olson, S. M., Kabelka, E. A. (2008). Pollen Viability of Selected Diploid Watermelon Pollenizer Cultivars. *HortScience* 43:274-275.
- Gaaliche, B., Majdoub, A. Trad, M., Mars, M. (2013). Assessment of Pollen Viability, Germination, and Tube Growth in Eight Tunisian Caprifig (*Ficus carica* L.) Cultivars. *ISRN Agron*. <https://doi.org/10.1155/2013/207434>.
- Godfree, R. C., Marshall, D. J., Young, A. G., Miller, C. H., Mathews, S. (2017). Empirical Evidence of Fixed and Homeostatic Patterns of Polyploid Advantage in a Keystone Grass Exposed to Drought and Heat Stress. *Royal Society Open Science*, 4(11): 170934. <https://doi.org/10.1098/rsos.170934>
- Goldberg, S., Shouse, P. J., Lesch, S. M., Grieve, C. M., Poss, J. A., Forster, H. S., Suarez, D. L. (2003). Effect of High Boron Application on Boron Content and Growth of Melons. *Plant and Soil* 256:403-411

- González-Hernández, A. I., Scalschi, L., Vicedo, B., Marcos-Barbero, E. L., Morcuende, R., Camañes, G. (2022). "Putrescine: a Key Metabolite Involved in Plant Development, Tolerance and Resistance Responses to Stress", *International Journal of Molecular Sciences*, 23(6), 2971.
- Gürsöz, N. (1990). Kavun (*Cucumis melo* var. *inodorus* ve *reticulatus*) ve Karpuzda (*Citrullus lanatus* (Thunb.) Mansf). Işınlanmış Polenle Uyarılan in situ Partenogenetik Embriyolardan in vitro Kültürü ile Haploid Bitki Eldesi. Yüksek Lisans tezi, 59 sayfa.
- Hedhly, A., Hormaza, J.I. and Herrero, M. (2005). Influence of Genotype-Temperature Interaction on Pollen Performance. *Journal of Evolutionary Biology*, 18(6), 1494-1502.
- Hidayat, C., Roosda, A. A., Fauziah, S. (2021). Various Planting Media and Boron Concentrations Supporting Growth and Yield of Melon on Drip Irrigation Hydroponic System. In IOP Conference Series: Earth and Environmental Science 694(1):012025
- Johri, B. M., and Vasil, I. K. (1960). *The Pollen and Pollen Tube* (pp. 1-13). Vieweg+ Teubner Verlag.
- Kantoğlu, K. Y., & Yanmaz, R. (1996). Kavunda (*Cucumis Melo* L.) Normal ve Işınlanmış Çiçek Tozları ile Tozlama Sonrasında Çiçek Tozu Çim Borusu Gelişiminin Dişicik Borusunda İncelenmesi Üzerinde Bir Araştırma.
- Karabıyık, Ş. (2011). Buruk Olmayan Bazı Trabzon Hurması (*Diospyros Kaki* L.) Çeşitleri İçin Uygun Tozlayıcı Çeşit Belirlenmesi. Yüksek Lisans Tezi. Çukurova Üniversitesi, Fen Bilimleri Enstitüsü. Bahçe Bitkileri Anabilim Dalı. 90 s.
- Karabıyık, Ş. (2022). Effects of Temperature on Pollen Viability and in vivo Pollen Tube Growth in *Citrus sinensis*. *Journal of Applied Botany and Food Quality* 95:100-104.
- Karabıyık, Ş. (2023). Nektarinde Putresin Uygulamasının Çoklu Pistil Oluşumu ve Döllenme Hızına Etkisi. VIII. National Horticulture Congress Abstract Book. 26-29 September 2023. Bursa/Türkiye.
- Karabıyık, Ş., Sarıdaş, M. A. (2023). Assessment of Pollen Quality and Quantity in White and Black Turkish *Myrtus communis* L. Accessions, Through In Vitro Pollen Germination Under Varied Boric Acid Concentrations. *International Journal of Minor Fruits, Medicinal & Aromatic Plants* Vol. 9 (2):167-176, December 2023.
- Karabıyık, Ş., & Çağlar, A. B. (2023). Effect of Putrescine on Pollen Quality and Pollen Tube Growth of Apple. 9th International Agriculture Congress. PROCEEDINGS BOOK, 2-4 October, 54.
- Karabıyık, Ş. (2024). Putrescine Affects Fruit Yield and Quality by Promoting Effective Pollination Period in *Citrus limon*. *Applied Fruit Science*, 66(2), 559-567.
- Kho, Y. O., & Baer, J. (1970). A Microscopical Research on the Incompatibility in the Cross *Rhododendron İmpeditum* × *R. Williamsianum*. *Euphytica*, 19, 303-309.
- Kihara, H. (1951). Triploid Watermelons. *Proc. Amer. Soc. Hort. Sci.*, 58: 217-230.
- Kusano, T., Berberich, T., Tateda, C., & Takahashi, Y. (2008). Polyamines: essential factors for growth and survival. *Planta*, 228, 367-381.
- La Porta, N., Roselli, G. (1991). Relationship Between Pollen Germination in vitro and Fluorochromatic Reaction in Cherry Clone F12/1 (*Prunus avium* L.) and Some of its Mutants. *Journal of horticultural science*, 66(2): 171-175. <https://doi.org/10.1080/00221589.1991.11516141>
- Liu, J. H., Wang, W., Wu, H., Gong, X., Moriguchi, T. (2015). Polyamines Function in Stress Tolerance: From Synthesis to Regulation, *Frontiers Plant Science*, 6, 827.
- Luo, Y., Zheng, M., Ni, R., Ling, Y., Lü, Y., & Song, G. (2020). The Source of Boron in Quaternary Sediments of Dangxiong Co, Tibetan Plateau, China. *Journal of Paleolimnology* 64:167-178.
- Maiti, R., Gonzalez-Rodriguez, H., & Ojha, E. R. (2016). Pollen Biology and Plant Productivity: A review. *Autoecology and ecophysiology of woody shrubs and trees: Concepts and applications*, 133-157.

- Nakamura, R. R., Wheeler, N. C. (1992). Pollen Competition and Paternal Success in Douglasfir. *Evolution* 46:846-851.
- Namlı, M., Adıgüzel, P., Karabıyık, Ş., & Solmaz, İ. (2024). Karpuzda (*Citrullus lanatus* L.) Farklı Şekillerde Yapılan Bor Uygulamasının Çiçek Tozu Kalite ve Miktarı Üzerine Etkisi. *Çukurova Tarım ve Gıda Bilimleri Dergisi*, 39(2), 510-518.
- Nepi, M., & Pacini, E. (1993). Pollination, Pollen Viability and Pistil Receptivity in *Cucurbita pepo*. *Annals of Botany*, 72(6), 527-536.
- Norton, J. D. (1966). Testing of Plum Pollen Viability with Tetrazolium Salts. *Proceedings of the American Society for Horticultural Science* 89:132-134.
- Paris, H. S. 2023. The Watermelon Genome, *Compendium of Plant Genomes* 1-16.
- Pinho, L.G., Campostrini, E., Monnerat, P.H., Netto, A.T., Pires, A.A., Marciano, C.R. and Soares, Y.J.B. (2010). Boron Deficiency Affects Gas Exchange and Photochemical Efficiency (JPI test parameters) in Green Dwarf Coconut. *Journal of plant nutrition*, 33(3), 439-451
- Rahayu, S., Firdhauzy, Y. L., Prasetyo, H., Setyohadi, D. P. S., and Cahyaningrum, D. G. (2023, April). Application of Boron Fertilizer and Topping on Female Parent Seed Production of Watermelon. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1168, No. 1, p. 012009) IOP Publishing.
- Razzaq, M. K., Rauf, S., Khurshid, M., Iqbal, S., Bhat, J.A., Farzand, A., Gai, J. (2019). Pollen Viability an Index of Abiotic Stresses Tolerance and Methods for the Improved Pollen Viability. *Pakistan Journal of Agricultural Research* 32(4).
- Razzaq, M. K., Rauf, S., Shahzad, M., Ashraf, I., Shah, F. (2017). Genetic Analysis of Pollen Viability: an Indicator of Heat Stress in Sunflower (*Helianthus annuus* L). *International Journal of Innovative Approaches in Agricultural Research* 1(1):40-508.
- Rerkasem, B., Nirantrayagul, S., Jamjod, S. (2004). Increasing Boron Efficiency in International Bread Wheat, Durum Wheat, Triticale and Barley Germplasm Will Boost Production on Soils Low in Boron. *Field Crops Research* 86:175-184.
- Rottmann, T.M., Fritz, C., Sauer, N., Stadler, R., (2018). Glucose Uptake via STP Transporters Inhibits in vitro Pollen Tube Growth in a Hexokinase1-Dependent Manner in Arabidopsis Thaliana. *Plant Cell*. 30(9):2057-2081.
- Sakhi, S., Okuno, K., Shahzad, A., Jamil, M. (2014). Evaluation of sorghum (*Sorghum bicolor* L.) Core Collection for Drought Tolerance: Pollen Fertility and Mean Performance of Yield Traits and its Components at Reproductive Stage. *International Journal of Agriculture And Biology* 16:251-260.
- Saleem, B.A., Malik, A.U., Anwar, R. and Farooq, M. (2008). Exogenous Application of Polyamines Improves Fruit Set, Yield and Quantity of Sweet Oranges. *Acta Horticulturae* 774:187-191.
- Saragih, A. A., Puteh, A. B., Ismail, M. R. and Mondal, M. M. A. (2013). Pollen Quality Traits of Cultivated (*Oryza sativa* L. Ssp. *indica*) and weedy (*Oryza sativa*'var. *nivara*) rice to water stress at reproductive stage. *Australian Journal of Crop Science* 7(8):1106-1112.
- Sarıdaş, M. A., Karabıyık, Ş., Eti, S., and Paydaş Kargı, S. (2021). Boron Applications and Bee Pollinators Increase Strawberry Yields. *International Journal of Fruit Science* 21(1):481-491
- Şensoy, A. S., Ercan, N., Ayar, F., Temirkaynak, M. (2003). *Cucurbitaceae* Familyasındaki Bazı Sebze Türlerinde Çiçek Tozlarının Bazı Morfolojik Özellikleri ile Canlılıklarının Belirlenmesi. *Akdeniz Üniversitesi Ziraat Fakültesi Dergisi* 16(1):1-6.
- Shivanna, K. R., Rangaswamy, N. S. (1992). *Pollen Biology*. Springer-Verlag Berlin Heidelberg, 119 s
- Shorrocks, V. M. (1997). The Occurrence and Correction of Boron Deficiency. *Plant Soil* 193:121-148.

- Solmaz, İ., Kartal, E., & Sarı, N. (2018). Kavunda ABA Uygulamalarının Bitki Büyümesi, Çiçek Cinsiyeti ve Çiçek Tozu Kalitesine Etkileri. *Türk Tarım-Gıda Bilim ve Teknoloji dergisi*, 6(9), 1224-1228.
- Soltis, D. E., Misra, B. B., Shan, S., Chen, S., Soltis, P. S. (2016). Polyploidy and The Proteome. *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics*, 1864 (8): 896-907.
- Sulusoglu, M. and A. Cavusoglu. (2014). In Vitro Pollen Viability and Pollen Germination in Cherry Laurel (*Prunus laurocerasus* L). *Sci. World J.* ID: 657123. <https://doi.org/10.1155/2014/657123>
- Stern, R. A., & Gazit, S. (2000). Application of the Polyamine Putrescine Increased Yield of 'Mauritius' litchi (*Litchi chinensis* Sonn.). *The Journal of Horticultural Science and Biotechnology*, 75(5), 612-614.
- Thomas, S., Ramakrishnan, R. S., Kumar, A., Sharma, R., Tiwari, M., & Pathak, N. (2020). Putrescine as a Polyamines and its Role in Abiotic Stress Tolerance: a review. *J. Pharmacogn. Phytochem*, 9(1), 815-820.
- Tosun, F., & Koyuncu, F. (2007). Kirazlarda (*Prunus Avium* L.) Çiçek Tozu Çimlenmesi ve Çiçek Tozu Çim Borusu Gelişimi Üzerine Bazı Kimyasal Uygulamaların Etkileri. *Akdeniz Üniversitesi Ziraat Fakültesi Dergisi*, 20(2), 219-224.
- Turner, N.C., T.D. Colmer, J. Quealy, R. Pushpavalli, L. Krishnamurthy, J. Kaur and V. Vadez. (2013). Salinity Tolerance and İon Accumulation in Chickpea (*Cicer arietinum* L.) Subjected to Salt Stress. *Plant and Soil*. 365: 347-361. <https://doi.org/10.1007/s11104-012-1387-0>
- Wijesinghe, S. A. E. C., Evans, L. J., Kirkland, L., & Rader, R. (2020). A Global Review of Watermelon Pollination Biology and Ecology: The Increasing İmportance of Seedless Cultivars. *Scientia Horticulturae*, 271, 109493.
- Zhang, S., Liang, M., Wang, N., Xu, Q., Deng, X. & Chai, L. (2018). Reproduction in Woody Perennial Citrus: An update on Nucellar Embryony and Self-Incompatibility. *Plant Reproduction*, 31(1), 43-57
- Zhang, N., Bao, Y., Xie, Z., Huang, X., Sun, Y., Feng, G., Chen, W. (2019). Efficient Characterization of Tetraploid Watermelon Plants, 8(10): 419. <https://doi.org/10.3390/plants8100419>

**MULTI-CRITERIA DECISION-MAKING METHODS (MCDM) FOR
SUSTAINABLE ENVIRONMENTAL MANAGEMENT: OPTIMIZED DECISION
SUPPORT SYSTEMS WITH RSTUDIO****SÜRDÜRÜLEBİLİR ÇEVRE YÖNETİMİ İÇİN ÇOK KRİTERLİ KARAR VERME
YÖNTEMLERİ (ÇKKV): RSTUDIO İLE OPTİMİZE EDİLMİŞ KARAR DESTEK
SİSTEMLERİ****Öğr. Gör. Afşin AYHAN**

Kastamonu Üniversitesi, İhsangazi Meslek Yüksekokulu

Prof. Dr. Banu ÇİÇEK KURDOĞLU

Karadeniz Teknik Üniversitesi, Orman Fakültesi

ÖZET

Günümüzde, hızla artan çevre sorunları sürdürülebilir kalkınma hedefleri doğrultusunda doğru karar verme süreçlerinin önemini daha da artırmaktadır. Özellikle doğal kaynakların aşırı tüketimi, biyoçeşitliliğin azalması, artan ekosistem hizmet kayıpları vb. gibi sorunlar etkili karar verme mekanizmalarını zorunlu kılmaktadır. 1971 yılında açığa çıkmış ve günümüze kadar gelmiş olan en güncel yöntemlerden biri de karar verme disiplininin bir alt dalı olan Çok Kriterli Karar Verme Yöntemleridir (ÇKKV) ve karmaşık sistemleri parçalara ayırarak analiz etme, çoklu hedefleri optimize etme ve bilimsel veriyi karar süreçlerine entegre etme gücüyle çevre yönetimi, sağlık hizmetleri, şehir planlaması, mühendislik problemleri gibi birçok disiplinde karar destek sistemleri olarak kullanılmaktadır. Bu yöntemler, hem belirsizliklerin ve çeşitli paydaş görüşlerinin ön planda olduğu durumlarda karar vericilere objektif bir çerçeve sunmakta hem de farklı disiplinlere ait çok sayıda bilimsel çalışma için temel veri setinin oluşturulmasını sağlamaktadır. Bu çalışma çevresel yönetim ve ekolojik planlama süreçlerinde karşılaşılan zorlukların üstesinden gelmede veya bu yöntemlerin uygulanması noktasında RStudio gibi yazılım araçlarının sağladığı kolaylıkları ortaya koymayı amaçlamaktadır. Örneğin AHP, VIKOR, TOPSIS, WASPAS, MOORA vb. uzmanlık gerektiren ve detaylı analizler ile sonuca ulaşılabilen yöntemlerin uygulanışı için farklı teknikler mevcut olup geleneksel olarak Excel yazılımı tercih edilmektedir. Ancak bu durum her ihtiyaç duyulduğunda kullanıcının birçok manuel giriş yapmasına ve karmaşık matematik formüllerini anlamaya çalışmasına neden olmaktadır. Böylece zaman kaybı, hata olasılığı ortaya çıkmakta ve en önemlisi de problem çözümü için yöntemlerin geniş kitleler tarafından kullanılması engellenmektedir. Sonuç olarak en uygun alan seçimi ile ilişkili geliştirilen senaryo özelinde dört farklı MOORA yaklaşımı Excel ve RStudio ortamında çözüme ulaştırılmış ve yazılım araçlarının bu yöntemleri destekleyerek, ekolojik planlamada bilimsel ve objektif çözümler sunma kapasitesini artırdığı ortaya konmuştur.

Anahtar kelimeler: Sürdürülebilirlik, Çok Kriterli Karar Verme Yöntemleri (ÇKKV), RStudio, MOORA

ABSTRACT


Today, rapidly increasing environmental problems increase the importance of correct decision-making processes in line with sustainable development goals. In particular, problems such as excessive consumption of natural resources, decline in biodiversity, increasing loss of ecosystem services, etc., necessitate effective decision-making mechanisms. Originating in 1971, one of the most up-to-date methods is Multi-Criteria Decision Making (MCDM), a sub-branch of the decision-making discipline, which is used as decision support systems in many disciplines such as environmental management, health care, urban planning, engineering problems, etc. with its power to analyze complex systems by breaking them into parts, optimizing multiple objectives and integrating scientific data into decision processes. These methods provide an objective framework for decision-makers in situations where uncertainties and various stakeholder views are at the forefront and provide a basic data set for a large number of scientific studies from different disciplines. This study aims to reveal the convenience of software tools such as RStudio in overcoming the challenges encountered in environmental management and ecological planning processes or in the implementation of these methods. For example, AHP, VIKOR, TOPSIS, WASPAS, MOORA, etc. There are different techniques for the application of methods that require expertise and can be concluded with detailed analysis, and Excel software is traditionally preferred. However, this causes the user to do much manual input and try to understand complex mathematical formulas whenever needed. This wastes time, introduces the possibility of error, and, most importantly, prevents the widespread use of methods for problem-solving. As a result, four different MOORA approaches were solved in Excel and RStudio environments for the scenario developed about optimal site selection, and it was revealed that software tools support these methods and increase the capacity to provide scientific and objective solutions in ecological planning.

Key Words: Sustainability; Multi-Criteria Decision-Making Methods (MCDM); RStudio; MOORA

GİRİŞ

Karar, en basit ifadeyle, öğrenilen bilgiler ve geçmiş deneyimlerden hareketle, önceden belirlenmiş olan amaç ve hedeflere ulaşmayı sağlayacak davranışlar, tutumlar ve seçenekler arasından bir seçim yapmaktır. Karar vermeyle ilgili problemlerin çözümde, problemin ve mevcut verilerin türüne göre pek çok farklı yöntem bulunmaktadır (Uludağ & Doğan, 2021). Bu problemlerin çözümü için 1971 yılında açığa çıkmış ve günümüze kadar gelmiş olan en güncel yöntemlerden biri de karar verme disiplininin bir alt dalı olan Çok Kriterli Karar Verme Yöntemleri (ÇKKV - Multi Criteria Decision Making: MCDM) olmuştur. ÇKKV yöntemlerinin ortak bileşenleri alternatifler, kriterler, kriter ağırlıkları ve karar matrisleri yer almaktadır (Fendoğlu, 2021).

Çok kriterli karar verme, yöneylem araştırması ve yönetim bilimi alanlarının bir alt dalı olarak, karar teorisi ve karar analizinin en yaygın olarak kullanılan yöntemlerinden biridir. ÇKKV sayısal veya sözel kriterlere göre muhtemel seçenekler arasından en iyi olanının belirlenmesidir ve çok sayıda kritere göre alternatiflerin avantaj ve dezavantajlarını değerlendiren analitik yöntemler topluluğu olarak temel amacı karar vericilere en iyi olanı önermektir. Literatürde çok kriterli karar verme yöntemleri, alternatiflerin tanım kümesinin sürekli veya kesikli olmasına göre; çok amaçlı karar verme (ÇAKV) ve çok nitelikli karar verme (ÇNKV) olmak üzere iki kategoriye ayrılır (Atan & Altan, 2020). Karar tekniklerinin sınıflandırılması Şekil 1’de görülmektedir.



Şekil 1. Karar tekniklerinin sınıflandırılması (Timor, 2011)

Modern çok kriterli karar verme yöntemlerinin temellerinin atıldığı 1950’li ve 1960’lı yıllardan bu yana, birçok araştırmacı zamanlarını yeni çok kriterli karar verme modelleri ve teknikleri geliştirmeye adanmıştır (Zavadskas, Turskis, & Kildienė, 2014). Karar verme süreçlerinin gün geçtikçe çeşitlenmesi ve farklı disiplinlerde kullanılan karar verme süreçlerinin bir başka disiplin için de geçerli olabilmesinin farkındalığı sayesinde yöntemler ve uygulamalar daha da genişlemektedir (Yıldız, 2023).


Günümüzde hem nitel hem de nicel verilerden oluşan çok sayıda ÇKKV problemi bulunmaktadır. AHP, ELECTRE ve PROMETHEE, hem nitel hem de nicel verilere uygulanabilir konumda iken bu çalışmada kullanılan MOORA yalnızca nicel verilere uygulanabilmekte olup (Fendoğlu, 2021), PROMETHEE, TOPSIS, ELECTRE ve VIKOR gibi diğer ÇKKV yöntemleriyle karşılaştırıldığında yeni bir yöntem olmasına rağmen son dönemlerde ilgili yazında sıkça kullanılmaktadır (Ajrina, Sarno, & Hari Ginardi, 2019). Yöntem Brauers & Zavadskas (2006) çalışması ile 2006 yılında geliştirilerek literatüre girmiştir (Uludağ & Doğan, 2021). ÇKKV yöntemleri sayısız kullanım örneğine sahiptir (Kumar, 2025) ve farklı kriterleri optimize ederek karar süreçlerini destekleyen yaklaşımlar sunmakta ve farklı disiplinlere ait problemlerin (ekonomi, işletme, sigorta, tıp ve sağlık sistemleri, mühendislik tasarımları, sürdürülebilir tedarik zinciri, finans, aktüerya, su yönetimi, enerji yönetimi, tarım ve gıda tedarik zinciri ve çevre konuları...) çözümü noktasında en uygulanabilir ve güvenilir yöntemlerdir (Donyatalab, 2022).

Bu yöntemlerin geleneksel olarak Excel gibi manuel hesaplamalara dayalı platformlarda uygulanması, hesaplama hatalarına, işlem süresinin uzamasına ve kullanıcı dostu olmayan karmaşık matematiksel süreçlere yol açmaktadır. Bu çalışmada RStudio gibi açık kaynaklı yazılım ortamlarının, ÇKKV yöntemlerini uygulamada sağladığı kolaylıkları vurgulamayı amaçlamıştır. Çalışma kapsamında Excel ve RStudio platformlarında analizler gerçekleştirilmiş ve yazılım destekli karar verme süreçlerinin avantajları ortaya konulmuştur.

MATERYAL VE YÖNTEM

Çalışmanın ana materyalini oluşturulan Kastamonu ili havza haritası, (Şekil 1), MOORA yöntemi, RStudio ile Excel yazılımları oluşturmaktadır. Belirtildiği üzere bir karar problemi geliştirilmiş ve MOORA yöntemi ile problem önce Excel daha sonra RStudio üzerinden çözüme ulaştırılmıştır. Belirtilen işlemler için öncelikle URL-994 (2025) adresinden R ve Rtools 4.4 yazılımları daha sonra URL-996 (2025) adresinden RStudio yazılımı yüklenmiştir. MOORA, TOPSIS, VIKOR, WASPAS yöntemlerini de içeren MCDM v1.0 (Multi-Criteria

Decision Making Methods) paket programı URL-995 (2025) adresinden indirilmiştir. Bu bölümde geliştirilen karar problemi ile MOORA yöntemine ilişkin uygulama esaslarına yönelik temel bilgiler paylaşılmıştır.



Şekil 1. Kastamonu kenti havza alternatifleri

MOORA yöntemi

Karar verme sürecinde kullanılan yöntemlerden biri olan Multi-Objective Optimization on the Basis of Ratio Analysis (MOORA), alternatifler içinden en iyisinin seçiminin yapılmasına imkan tanıyan oransal analize dayalı bir optimizasyon yöntemidir. Yöntem için, oran sistemi, ağırlıklandırılmış oran sistemi, referans noktası teorisi, ağırlıklandırılmış referans noktası teorisi, tam çarpım formu ve multimooora olarak ifade edilen yaklaşımlar söz konusudur (Uludağ & Doğan, 2021). Fendoğlu (2021) ve Atan & Altan (2020) bu yaklaşımlardan ağırlıklandırılmış oran sistemi ve ağırlıklandırılmış referans noktası teorisi yaklaşımlarını önem katsayısı yaklaşımı olarak ele almışlardır. Söz konusu bu yaklaşımların matematiksel açıdan uygulanışı aşağıda açıklanmaktadır. Ayrıca bu yaklaşımlar benzer adımları içermekte olduğundan (örneğin, normalizasyon, ağırlıklandırma vb.) tekrar tekrar açıklamadan kaçınılmış ve yapılması gerekli işlemler belirtilmiştir.

Oran sistemi ve ağırlıklandırılmış oran sistemi yaklaşımı

Bu yaklaşım; başlangıç karar matrisinin tesis edilmesi, normalizasyon ve alternatiflerin toplam skorlarının hesaplanması olmak üzere 3 temel aşamadan oluşmaktadır.

Aşama 1: Birinci aşamada, i alternatifleri veya karar seçeneklerini, j değerlendirme kriterlerini temsil etmek, $i = 1, 2, \dots, s$ ve $j = 1, 2, \dots, m, m+1, \dots, t$ olmak üzere; Denklem (1)'de R ile gösterilen $s \times t$ boyutlu karar matrisi oluşturulur. Burada m , fayda özelliğine sahip değerlendirme kriterlerinin sayısıdır. Dolayısıyla, maliyet özelliğine sahip kriter sayısı da $t - m$ adettir.

$$R = [r_{ij}]_{s \times t} = \begin{bmatrix} r_{11} & \dots & r_{1j} & \dots & r_{1t} \\ \vdots & & \vdots & & \vdots \\ r_{i1} & \dots & r_{ij} & \dots & r_{it} \\ \vdots & & \vdots & & \vdots \\ r_{s1} & \dots & r_{sj} & \dots & r_{st} \end{bmatrix}, (i = 1, \dots, s \text{ ve } j = 1, \dots, m, m + 1, \dots, t) \quad 1$$

Denklem (1)'deki r_{ij} , j 'inci değerlendirme kriterine göre i 'inci alternatifin veya karar noktasının değerini ifade etmektedir.

Aşama 2: İkinci aşamada, R karar matrisindeki değerler; yani r_{ij} 'ler normalize edilir. Normalizasyon işlemi açısından MOORA yönteminde değerlendirme kriterinin fayda ya da maliyet özelliğine sahip olması bakımından bir farklılaşma yoktur. Normalizasyon işlemi için Denklem (2)'deki formül kullanılır. Bu eşitlikteki \tilde{r}_{ij} , j'inci değerlendirme kriterine göre i'inci alternatifin normalize edilmiş değerini göstermektedir. Normalize edilmiş değerler hesaplandıktan sonra Denklem (3)'te \tilde{R} ile gösterilen normalize edilmiş karar matrisi tesis edilir.

$$\tilde{r}_{ij} = \frac{r_{ij}}{\sqrt{\sum_{i=1}^s r_{ij}^2}}, (i = 1, 2, \dots, s \text{ ve } j = 1, 2, \dots, m, m + 1, \dots, t) \quad 1$$

$$\tilde{R} = [\tilde{r}_{ij}] = \begin{bmatrix} \tilde{r}_{11} & \dots & \tilde{r}_{1j} & \dots & \tilde{r}_{1t} \\ \vdots & & \vdots & & \vdots \\ \tilde{r}_{i1} & \dots & \tilde{r}_{ij} & \dots & \tilde{r}_{it} \\ \vdots & & \vdots & & \vdots \\ \tilde{r}_{s1} & \dots & \tilde{r}_{sj} & \dots & \tilde{r}_{st} \end{bmatrix}, (i = 1, \dots, s \text{ ve } j = 1, \dots, m, m + 1, \dots, t) \quad 1$$

Aşama 3: Son aşamada i'inci alternatifin ya da karar seçeneğinin tüm kriterler dikkate alındığında elde edilen ve Denklem (4)'te x_i ile gösterilen toplam skorları hesaplanır. Ardından, alternatiflerin toplam skorları büyükten küçüğe sıralanır. En yüksek değere sahip olan alternatif (x_i^*), en iyi karar seçeneğidir. Bu durum Eşitlik Denklem (5)'te gösterilmiştir (Uludağ & Doğan, 2021).

$$x_i = \sum_{j=1}^m \tilde{r}_{ij} - \sum_{j=m+1}^t \tilde{r}_{ij}; (i = 1, 2, \dots, s \text{ ve } j = 1, 2, \dots, m, m + 1, \dots, t) \quad 1$$

$$x_i^* = \left\{ \forall i \left| \underset{i}{\text{maks}} x_i \right. \right\}; (i = 1, 2, \dots, s) \quad 1$$

Eğer ağırlıklandırılmış oran sistemi tercih edilirse belirlenen karar kriter (sütun) ağırlıkları ile normalize matris değerleri çarpılmalı ve böylece ağırlıklandırılmış normalize matris elde edilmelidir. Örneğin 8 kriter eşit ağırlıkta olduğunda kriter başına önem değeri $1/8=0.125$ olarak hesaplanmalıdır. Ağırlık değerleri öznel olarak veya farklı bir ağırlık belirleme tekniği ile belirlenebilmekte olup devamında aynı şekilde toplam skor hesaplanmaktadır.

Referans noktası teorisi ve ağırlıklandırılmış referans noktası teorisi yaklaşımı

Bu yaklaşımda oran yaklaşımı ile elde edilen normalleştirilmiş veriler kullanılır. Referans noktası yaklaşımında alternatiflerin her bir amaca göre maksimizasyon durumunda en yüksek değer, minimizasyon durumunda ise en düşük değer referans noktası (r_i) olarak alınır. Denklem (6) kullanılarak alternatiflerin her bir amaca göre referans noktasına olan uzaklıkları bulunur.

$$d_{ij} = |r_i - x_{ij}^*| \quad 6$$

Alternatiflerin sıralaması yapıldıktan sonra her alternatifin en yüksek değeri bulunarak (P_i) (Denklem 7) alternatifler küçükten büyüğe doğru sıralanır. Sıralama sonucunda birinci sıradaki alternatif en iyi seçenek olarak kabul edilir (Atan & Altan, 2020).

$$P_i = \min_i (\max_j d_{ij}) \quad 7$$

Eğer ağırlıklandırılmış referans noktası teorisi yaklaşımı kullanılacaksa her bir karar kriteri (sütun) için yine ağırlık belirlenmelidir. Daha sonra normalize matris değerleri ile hesaplanmış olan referans noktaları belirlenen bu kriter ağırlıkları ile çarpılmalı ve ağırlıklı normalize matris elde edilmelidir. Devamında aynı işlemler yürütülmelidir.

Tam çarpım formu

Bu yaklaşıma göre her bir alternatifin maksimizasyon amaçlı verileri çarpılır ve minimizasyon amaçlı verilerin çarpımına bölünür (Denklem 8).

$$U_i = \frac{A_i}{B_i}, A_i = \prod_{g=1}^j x_{gj}, B_i = \prod_{k=j+1}^n x_{kj} \quad 8$$

$i = 1, 2, \dots, m$ olmak üzere; “ m ”; alternatiflerin sayısını, “ j ”; maksimizasyon ölçütlerinin sayısını ve “ $n-j$ ” de; minimizasyon ölçütlerinin sayısını ifade etmektedir. U_i , alternatiflerin skorlarını göstermektedir. İşlem sonrasında U_i değerleri büyükten küçüğe doğru sıralanarak ilk sıradaki alternatif, en uygun seçenek olarak değerlendirilir (Atan & Altan, 2020).

Multimoora

Bu yaklaşım, 2010 yılında oran sistemi referans noktası teorisi yaklaşımlarına tam çarpım formunun eklenmesiyle ortaya çıkmıştır. Esasen başlı başına çok amaçlı optimizasyon için kullanılan bir yöntem değildir. Sadece oran sistemi, referans noktası teorisi, tam çarpım formu ile elde edilen sıralamaları kullanarak nihai bir sıralamaya ulaşılmasını sağlayan bir yaklaşımdır. Bu yaklaşımda sıralama elde edilirken 4 ana durum dikkate alınmaktadır. Bunlar sırasıyla, mutlak ve genel baskınlık, geçişlilik, mutlak ve kısmi eşitlik, döngüsel akıl yürütmedir (Uludağ & Doğan, 2021). Örneğin 3 yöntemde de 3. sırada (3-3-3) yer alan bir alternatif multimoora yaklaşımında mutlak baskınlık kurmasından dolayı yine 3. sırada değerlendirilmektedir.

Karar problemi

Belirlenen karar kriterlerine göre en uygun alternatif/havza hangisidir? sorusuna cevap aranmıştır. Bu amaçla il sınırları içerisinde yer alan 10 adet havza için 8 adet karar kriteri belirlenmiştir. Havzalar Şekil 1’de, 8 adet karar kriteri, kriter kodları ve birimleri ile maksimum (fayda) ve minimum (maliyet) özellikleri Tablo 1’de görülmektedir. Buna göre karar kriterlerinin ilk 4’ünün maksimum, diğer 4’ünün ise minimum yönlü olduğu en iyi seçeneğin belirlenmesi için başlangıç karar matrisi oluşturulmuştur ve Excel ekran görüntüsü Şekil 2’de paylaşılmıştır. Karar kriterlerine değerlerin atanması noktasında nüfus kriteri için sınırların tam

çakışmaması nedeniyle havza içinde en büyük paya sahip olan ilçe nüfusu, minimum yönlü arazi yetenek sınıfı kriteri için VII. ve VIII. sınıf arazi toplam alanları hesaplanmış ve ortaya çıkan değer ilgili havzaya atanmıştır.

Tablo 1. Karar kriterleri ve özellikleri

No	Karar Kriterleri	Kriter Kodu	Maksimum - Minimum Özellikleri	Birim
1	Alan Büyüklüğü	K1	Maksimum (Fayda)	km ²
2	Orman Alanı	K2	Maksimum (Fayda)	km ²
3	Nüfus	K3	Maksimum (Fayda)	sayı
4	Akarsu Uzunluğu	K4	Maksimum (Fayda)	metre
5	Yerleşim Alanları	K5	Minimum (Maliyet)	km ²
6	Yol Uzunluğu	K6	Minimum (Maliyet)	metre
7	Arazi Yetenek Sınıf	K7	Minimum (Maliyet)	km ²
8	Ortalama Eğim	K8	Minimum (Maliyet)	yüzde

	A	B	C	D	E	F	G	H	I	J
1			Karar Kriterleri							
2	No	Havza Adı	K1	K2	K3	K4	K5	K6	K7	K8
3		(Alternatifler)	Maks	Maks	Maks	Maks	Min	Min	Min	Min
4	1	15/4-c	4646,5	2935,5	204446	570877	53,7	388166	3035,1	26,4
5	2	15/4-b	1224,6	692,2	39609	59574,9	15	124237	752,9	33,6
6	3	13026	1786,3	1302	51884	263057	17,2	235970	1323,3	42,2
7	4	13025	950,1	664,4	11501	149382	4,7	119770	677,3	40,3
8	5	13022	494,1	306,3	5204	37624,7	2,6	31600,2	317,2	30,8
9	6	13021	939	629,5	16876	51049,2	6,1	96998,2	672,1	35
10	7	13020	698,4	515	7233	62268,4	1,9	48830,8	527,3	30,2
11	8	13018	313,4	108,1	4366	34640,5	2,6	55930,8	118,9	14,2
12	9	13016	308	98,3	12037	22165,4	4,5	28401,4	102,9	14,4
13	10	13014	1691,2	1187,7	18122	131596	12,6	145955	1206,5	29,8

Şekil 2. Başlangıç karar matrisi excel ekran görüntüsü

BULGULAR VE TARTIŞMA

Bu bölümde geliştirilen karar probleminin çözümü için 3 farklı MOORA yaklaşımı ilk olarak Excel ortamında Şekil 2’de yer alan ekran görüntüsü temeline dayandırılan formül ve fonksiyon paylaşımları ile çözüme ulaştırılmış daha sonra aynı problem ve aynı karar matrisi RStudio ortamında çözümlenmiştir. Bu şekilde 2 farklı teknikte çözüm için harcanan çabanın farkı ortaya konmuş ve karar problemlerinin daha etkin kullanımı için pratik tekniklerin önemi vurgulanmıştır.

Oran sistemi yaklaşımı

İlk olarak normalize karar matrisi oluşturulmaktadır. Bu işlem için başlangıç karar matrisinde yer alan her bir değer, kendi sütunu tüm değerlerin kareleri toplamının kareköküne

bölünmektedir. Bu işlem için Excel ortamında Denklem 9 tüm karar matrisine sürüklenme yöntemleri ile uygulanmalıdır.

$$=C4/KAREKÖK(C\$4^2+C\$5^2+C\$6^2+C\$7^2+C\$8^2+C\$9^2+C\$10^2+C\$11^2+C\$12^2+C\$13^2) \quad 9$$

Son aşamada ise alternatiflerin toplam skoru belirlenmektedir. Bu işlem için normalize karar matrisinde her bir alternatif (satır) için fayda kriterleri toplamından maliyet kriterleri toplamının çıkarılması gerekmektedir. Bu işlem için Excel'de Denklem 10'da belirtilen fonksiyon kullanılmaktadır. Böylece en büyük skora sahip seçenek en iyi alternatif olmaktadır. Hesaplanan skor değerlerinin sıralaması için RANK fonksiyonu kullanılabilir. Normalize matrisin ilk hücre değeri (sol üst köşe) C16'da yer almaktadır.

$$=TOPLA(C16:F16)-TOPLA(G16:J16)$$

10

Referans noktası teorisi yaklaşımı

İlk adımda oluşturulan normalize karar matrisi kullanılmaktadır ve normalize matris değerlerinde her bir karar kriteri (sütun) için referans noktasının belirlenmesi gerekmektedir. Referans noktası için fayda (maksimum) yönlü kriter için en büyük, maliyet (minimum) yönlü kriter içinse en küçük değer seçilmektedir. Bu işlemler için Excel'de MAK ve MIN fonksiyonları kullanılabilir. Referans noktaları belirlendikten sonra kriter referans noktasından her bir normalize matris değeri çıkarılmalı ve mutlak değeri alınmalıdır. Doğal olarak oluşan yeni matrisin her sütununda 1 adet 0 yazan değer görülecektir. Bu işlem için Excel ortamında Denklem 11'de yer alan fonksiyon kullanılmaktadır. Excel'de C27 hücresinde K1 kodlu karar kriterine ilişkin referans noktası yer almaktadır.

$$=MUTLAK(C\$27-C16)$$

11

Daha sonra toplam skorların belirlenmesi gerekmektedir. Bunun için her bir alternatif (satır) için en büyük değer belirlenir. Bu şekilde belirlenen değerlerden en küçüğü en iyi seçenektir sonucuna ulaşılmaktadır. Hesaplanan skor değerleri sıralaması için küçük olan değer 1. sırada görülebilmesi için RANK fonksiyonu içinde 1 değeri eklenmesi yapılmaktadır.

Tam çarpım formu

Bu yaklaşımda başlangıç karar matrisi kriterlerinden (sütun) fayda yönlü özelliğe sahip olanların çarpımlarının değeri, maliyet yönlü olan karar kriterleri çarpımlarına bölünmektedir. Bu işlem için Excel'de Denklem 12'de yer alan fonksiyon kullanılmaktadır. Bu şekilde elde edilen oransal değerlerin büyükten küçüğe doğru sıralanması sonucunda en yüksek değere sahip olan alternatif en iyi seçenektir sonucuna ulaşılmaktadır.

$$=ÇARPIM(C4:F4)/ÇARPIM(G4:J4)$$

12

Excel yazılımında yapılan bu işlemler istatistiksel hesaplama, veri analizi, veri görselleştirme ve makine öğrenmesi gibi çeşitli konularda farklı disiplinlerce kullanılan RStudio ortamında da çalıştırılmıştır. Bu kapsamda MOORA yaklaşımları içinde alternatiflerin toplam skorları ve sıralamalarını gösteren nihai sonuç tablosunu elde etmek için yapılan işlemler aşağıda açıklanmıştır.

1. İlk olarak Rmatris.xlsx isimli başlangıç karar matrisi excel dosyası RStudio ortamına içe aktar (import) ile aktarılmak üzere düzenlenmiştir. Bu işlem için dosyada ilk satırda kriter kodları, diğer hücrelerde değerler bırakılmıştır. Alternatiflere ilişkin bir sütun yer almamaktadır.
2. Aktarma işleminden sonra “G=as.matrix(Rmatris)” komutu ile matris olarak tanımlanmıştır.
3. Ağırlıklandırma işlemleri için “w=c(1/8,1/8,1/8,1/8,1/8,1/8,1/8,1/8)” komutu ile eşit önem katsayısı tanımlaması yapılmıştır.
4. Fayda maliyet özelliği “cb=c("max","max","max","max","min","min","min","min")” komutu ile tanımlanmıştır.
5. Son olarak “MMOORA(G,w,cb)” komutu ile sonuç MOORA yaklaşımları bazında alternatiflere ilişkin toplam skorlar ve bu skorlara göre alternatiflerin sıralamaları elde edilmektedir.

SONUÇ ve ÖNERİLER

Bu çalışma, sürdürülebilir çevre yönetimi bağlamında Çok Kriterli Karar Verme (ÇKKV) yöntemlerinin etkinliğini artırmak ve yazılım araçlarının karar verme süreci üzerindeki etkisine yönelik bir incelemeyi içermektedir. MOORA yöntemi kullanılarak yapılan analizler hem Excel hem de RStudio ortamında gerçekleştirilmiş ve bu iki platformun performansı ortaya konmuştur.

Gerçekleştirilen analizler sonucunda ortaya çıkan nihai sonuç değerleri Tablo 2’de paylaşılmıştır. Aynı zamanda söz konusu tabloda yer alan değerler ağırlık değerlerinin kullanıldığı yaklaşımlara özgü olup kriter başına eşit önem katsayısı nedeniyle 0.125 katsayısı ilgili analizlerde dikkate alınmıştır. Buna göre 15/4-c isimli havzanın, oran sistemi, tam çarpım formu ve multimooro yaklaşımlarında en uygun alternatif olarak belirlendiği, ancak referans noktası teorisi yaklaşımında üçüncü sırada yer aldığı tespit edilmiştir. Bu sonuç aynı zamanda farklı ÇKKV yöntemlerinin farklı sonuçlar üretebileceğini ve her bir yöntemin karar süreçlerinde farklı perspektifler sunduğunu göstermektedir. Sonuç olarak RStudio’nun sunduğu hata oranını düşüren ve zaman kazandıran analitik yetenekleri, geleneksel Excel tabanlı çözümlere göre belirgin avantajları olduğu gözlenmiştir. Bu nedenle, karar verme süreçlerinde RStudio gibi veri bilimi tabanlı yazılımların tercih edilmesi, sürecin daha sistematik ve güvenilir hale gelmesini sağlamaktadır. Aynı zamanda ÇKKV’nin bir başka deyişle bilimsel ve nesnel karar alma süreçlerinin geniş kitlelerce kullanılabilmesi yolunda kullanıcı dostu arayüzlere sahip platformlar geliştirilmeli ve kullanımı teşvik edilmelidir.

Tablo 2. MOORA yaklaşımlarına göre toplam skorlar ve sıralamalar

No	Alternatif	Oran Sistemi	Sıra	Referans Noktası Teorisi	Sıra	Tam Çarpım Formu	Sıra	Multimoora Sıra
1	15/4-c	0.08992217	1	0.10702889	3	1.1788684	1	1
2	15/4-b	- 0.04367935	10	0.09550167	2	0.7989657	7	6
3	13026	- 0.02638882	6	0.08791979	1	0.9275917	2	2
4	13025	- 0.03382606	8	0.11119207	7	0.8514480	5	7
5	13022	- 0.03124566	7	0.11482096	9	0.7851875	8	10
6	13021	- 0.04105973	9	0.10809451	5	0.7843065	9	9
7	13020	- 0.02267443	4	0.11365167	8	0.8996393	3	5
8	13018	- 0.02114168	3	0.11530389	10	0.7311517	10	8
9	13016	- 0.01658486	2	0.11088318	6	0.7994959	6	4
10	13014	- 0.02574340	5	0.10737646	4	0.8542253	4	3

KAYNAKLAR

Ajrina, A. S., Sarno, R., & Hari Ginardi, R. V. (2019). Comparison Of MOORA and COPRAS Methods Based on Geographic Information System For Determining Potential Zone of Pasir Batu Mining. *International Conference on Information and Communications Technology (ICOIACT)*, s. 360-365.

Atan, M., & Altan, Ş. (2020). *Örnek Uygulamalarla Çok Kriterli Karar Verme Yöntemleri*. Ankara: Gazi Kitabevi.

Brauers, W. K. M., & Zavadskas, E. K. (2006). The MOORA Method and Its Application to Privatization in a Transition Economy. *Control and Cybernetics*, 35(2), 445-469.

Donyatalab, Y. (2022). *Novel Spherical Fuzzy Aggregation Operators and Similarity & Distance Measures*. İstanbul Technical University, Graduate School, M.Sc. Thesis, İstanbul.

Fendoğlu, E. (2021). *Çok Kriterli Karar Verme Yöntemleri ile Güncel Uygulamalar*. Ankara: Gazi Kitabevi.

Kumar, R. (2025). A Comprehensive Review of MCDM Methods, Applications, and Emerging Trends. *Decision Making Advances*, 3(1), 185-199.

Timor, M. (2011). *Analitik Hiyerarşi Prosesi*. İstanbul: Türkmen Kitabevi.

Uludağ, A. S., & Doğan, H. (2021). *Üretim Yönetiminde Çok Kriterli Karar Verme Yöntemleri: Literatür, Teori ve Uygulama*. Ankara: Nobel Akademik Yayıncılık.

URL-994. (2025, Ocak 9). Install R and Rtools: Geliş tarihi gönderen <https://cran.rstudio.com/> adresinden alınmıştır.

URL-995. (2025, Ocak 9). MCDM Packages: Geliş tarihi gönderen <https://cran.r-project.org/src/contrib/Archive/MCDM/> adresinden alınmıştır.

URL-996. (2025, Ocak 9). RStudio Posit: Geliş tarihi gönderen <https://posit.co/download/rstudio-desktop/> adresinden alınmıştır.

Yıldız, E. (2023). *Çok Kriterli Karar Verme Ve Coğrafi Bilgi Sistemi Hibrit Çözüm Yaklaşımı: Siirt İli Güneş Enerji Uygulaması*. Anadolu Üniversitesi Sosyal Bilimler Enstitüsü, Doktora Tezi, Eskişehir.

Zavadskas, E. K., Turskis, Z., & Kildienė, S. (2014). State of Art Surveys of Overviews on Mcdm/Madm Methods. *Technological and Economic Development of Economy*, 20(1), 165-179.

RARE AND ENDEMIC PLANTS OF AKDAĞ (ADIYAMAN/MALATYA) FLORA

AKDAĞ (ADIYAMAN/MALATYA) FLORASI'NIN NADİR VE ENDEMİK BİTKİLERİ

Murat TAK

Iğdır Üniversitesi, Lisansüstü Eğitim Enstitüsü, Tarım Bilimleri Anabilim Dalı, 76100 Iğdır, TÜRKİYE

ORCID ID: <https://orcid.org/0000-0001-8929-0806>

Ahmet Zafer TEL

Iğdır Üniversitesi, Ziraat Fakültesi, Tarımsal Biyoteknoloji Bölümü, 76100 Iğdır, TÜRKİYE

ORCID ID: <https://orcid.org/0000-0002-1204-3839>

ÖZET

Bu çalışma Akdağ (Adıyaman/Malatya) florasını tespit etmek üzere yapılmıştır. 2018-2022 yılları arasında çalışma alanından 1050 bitki örneği toplanmış; 80 familya ve 327 cinse ait 610 tür ve tür altı takson teşhis edilmiştir. Belirlenen taksonların 66(%10.8) tanesi endemiktir. Taksonların fitocoğrafik bölgelere dağılımı ve oranları; İran-Turan 181 (%29.72), Akdeniz 61 (%10.18), Avrupa- Sibiryaya 16 (%1,8), çok bölgeli veya fitocoğrafik bölgesi bilinmeyenler ise 343 (%56.13) şeklindedir.

Araştırma alanından toplanan 66 endemik ve nadir 8 taksondan 54 tanesi İran-Turan, 7 takson Akdeniz elementi, 2 takson Avr.-Sibiryaya, 11 taksonun fitocoğrafik bölgeleri belli değildir. Yine 66 endemik taksondan IUCN tehlike kategorisine göre 3 takson CR (Çok tehlikede), 9 takson VU (Zarar görebilir), 14 takson NT (Tehdit altına girebilir) ve 33 takson LC (En az endise verici), 5 takson EN (Tehlikede), 2 takson DD (yetersiz verili) kategorisine girdiği görülmüştür. Endemik olmayan fakat nadir olan takson sayısı 8 VU (Zarar görebilir)'dir.

Anahtar Kelimeler: Adıyaman, Akdağ, Flora, Malatya, Endemik

ABSTRACT

In this study, the Flora of Akdağ (Adıyaman/Malatya) has been studied. 1050 plant specimens were collected from the research area between 2018-2022 and at the end of this it was determined that 610 taxa, belonging to 80 families, 327 genera were found. 66 (10,8%) of the total taxa are endemics in this study area. The phytogeographic regions of only 610 taxa out of the collected material have been determined; Irano- Turanian 181 (29.72%), Mediterranean 61 (10,18%), Euro- Siberian 16 (1,8%). The rest 343 species (56,13%) of the total are either pluriregional or phytogeographically unknown.

Of the 66 endemic taxa and 8 rare taxa collected from the research area, Iran-Turan element 54, Eastern Mediterranean element 7, Avr.-Siberian and the phytogeographic regions of 11 taxa are not clear. Again, among the 66 endemic taxa, according to the IUCN danger category, 3 taxa are in the CR (Very Endangered), 9 taxa are in the VU (Vulnerable), 14 taxa are NT (May be threatened), 33 taxa are LC (Least Concern), 5 taxa are EN (Endangered), and 2 taxa is in the


DD (data deficient) category. The number of non-endemic but rare taxa is 8 VU (Visible Damage).

Keywords: Adıyaman, Akdağ, Flora, Malatya, Endemic

GİRİŞ

Çalışma alanı olarak belirlenen Akdağ, Adıyaman ve Malatya il sınırları içinde yer almaktadır. En yüksek tepesi Adıyaman'dan 90 km uzakta ve 2552 m rakımlıdır. Akdağ Güneydoğu torosların üzerinde yer alır. Alan Güneydoğu toros dağlarının bir uzantısı olup Akdeniz ile İran-Turan fitocoğrafik bölgesinin geçiş bölgesine yakın yerdedir. Kuzeyinde Erkenek, Kuzeybatısında ve Batısında Adıyaman-Malatya Karayolu geçer. Doğusunda Ulubaba Dağı bulunur. Güneyinde ise Adıyaman-Gölbaşı yolu geçer. Çalışma alanı 390km²'dir. Çalışma alanında Tut ilçesi ile Malatyaya bağlı Erkenek kasabası, Adıyaman sınırları içinde kalan Tepecik, Öğütlü, Havutlu, Çiftlik, Cankara, Yaylakonak, Kaşlıca köyleri yer alır. Çalışma alanı kuzeybatıda 37 55 26.46 K- 37 52 03.59 D, kuzeydoğuda 37 55'58.01 K- 38 02'26.84 D, güneybatıda 37 46'27.57 K- 37 51'08.59 D, güneydoğuda 37 45'41.40 K- 38 04'16.92 D enlem ve boylamları arasındadır (Şekil 1.7.a,b), (google earth pro, 2022).

Çalışma alanının topografyası, Jeolojik yapısı, iklimi ve coğrafik durumu ile farklılık göstermesi, biyoçeşitliliğin artmasına katkı sağlamıştır (Şekil 1).



Şekil 1. Coğrafik konum (HGM'den değiştirilerek)

Çalışma alanına yakın yerlerde yapılan çalışmaların bazılarını, Tel (2001)'da yaptığı "Nemrut Dağı (Adıyaman) Vejetasyonu", Şimşek'in (2015), "Ulubaba Dağı (Adıyaman) Vejetasyonu Üzerine Fitososyolojik ve Fitoekolojik Çalışmalar", Tel ve Tak'ın (2012), "Perre (Pirin) Antik Şehri (Adıyaman) Vejetasyonu", Tak'ın (2015)'teki çalışmasında "Çelikhlan Çat Barajı (Adıyaman) Havzası'nın Fitososyolojik Ve Fitoekolojik Özellikleri", Eğilmez (2014)'teki "Gölbaşı Gölleri (Adıyaman) Havzası'nın Vejetasyonu", Şahin'nin (2015)'teki "Ali Dağı ve Ziyaret Tepesi (Adıyaman) Vejetasyonu", Tak (2023), " Akdağ (Adıyaman/Malatya) Florası", Avcı, (2019), "Akdağ (Çelikhlan/Adıyaman) florası", Çetiner, M., (2020), " Atatürk Baraj Gölü Havzası'nın Şanlıurfa Bölümü Florası", Ortaç ve Tel, (2021), " Gazihan Dede Mesire Alanı (Adıyaman, Türkiye) Florası" , Tel ve Tak, (2018), "Karagöl Sülüklü Göl Havzası, Gerger/Adıyaman) Florası Üzerine Bir araştırma", Tel ve Tak (2021), "Çelikhlan ve Yakın Çevresinin (Adıyaman/Türkiye) Flora ve Vejetasyonu Üzerine Araştırmalar, ", Tel ve ark. (2019), Tel ve ark. (2021), Tel ve ark. (2022a; 2022b), Tel ve ark. (2023) tarafından araştırmalar yapılmıştır. olarak sayabiliriz.

MATERYAL VE METOD

Bitkiler, 2019-2022 yıllarında iklimin müsait olduğu zamanlarda düzenli olarak belirli periyotlarda toplanan bitki örneklerinden oluşturmaktadır. Bitkiler en az üç tane olmak üzere çiçek, meyve, kök ve yaprak gibi karakteristik kısımları ile birlikte toplanmıştır. Toplanan örnekler herbaryum tekniklerine göre kurutulmuştur. Toplanan örneklerin bilimsel teşhis ve tayinlerinde temel kaynak olarak "Flora of Turkey and the East Aegean Islands" adlı eseri (Davis, 1965; Davis et al., 1988; Güner vd., 2000) kullanılmıştır. Karışık yada teşhiste zorlukla karşılaşılan türler otor hocalardan yardım alınarak teşhisi yapılmıştır. Endemik ve nadir bitkilerin listede veriliş sırası, Türkiye Flora'sında uygulanan filogenetik sisteme göredir. Tehlike sınıflarının belirlenmesinde (Ekim vd., 2000) tarafından hazırlanan "Türkiye Bitkileri Kırmızı Kitabı" ve "IUCN Red List Categories" adlı eserler esas alınmıştır.

BULGULAR

Çalışma alanında bulunan endemik, nadir bitkiler, biyocoğrafik bölgeleri ve tehlike kriterleri aşağıda verilmiştir (Tablo 1).

Tablo 1. Çalışma alanının endemik ve nadir bitkileri.

Takson Adı	Endemizm	Iucn (Tehlike Sınıfı)	Biyocoğrafya
<i>Acanthus dioscoridis</i> L. var. <i>perringii</i> (Siehe) E. Hossain,	Endemik	[VU]	
<i>Eryngium kotschy</i> Boiss.	Endemik	[LC]	D. Akdeniz
<i>Ferulago blancheana</i> Post	Endemik	[LC]	
<i>Heracleum pastinacifolium</i> subsp. <i>incanum</i> (Boiss. & A.Huet) P.H.Davis	Endemik	[LC]	
<i>Malabaila lasiocarpa</i> Boiss	Endemik	[LC]	Ir-Tur
<i>Tordylium cappadocicum</i> Boiss.	Endemik	[DD]	Ir-Tur
<i>Cousinia foliosa</i> Boiss. et Bal.	Endemik	[LC]	Ir-Tur
<i>C. eriocephala</i> Boiss. et Bal.,	Endemik	[LC]	Ir-Tur
<i>Echinops vaginatus</i> Boiss. et Hausskn.	Endemik	[EN]	Ir-Tur
<i>Klasea oligocephala</i> (DC.) Greuter & Wagenitz	Endemik	[LC]	Ir-Tur

<i>Tanacetum densum</i> (Lab.) Schultz Bip. subsp. <i>eginence</i> Heywood	Endemik	[LC]	Ir-Tur
<i>Taraxacum pseudonigricans</i> Hand.- Mazz	Endemik	[NT]	Ir-Tur
<i>Onosma bornmuelleri</i> Hausskn. &Bornm.	Endemik	[LC]	Ir-Tur
<i>O. malatyana</i> Binzet	Endemik	[CR]	Ir-Tur
<i>O. mutabilis</i> Boiss.	Endemik	[LC]	
<i>Aethionema schistosum</i> Boiss. & Kotschy	Endemik	[NT]	
<i>Alyssum huetii</i> Boiss. Endemik	Endemik	[LC]	Ir-Tur
<i>Arabis carduchorum</i> Boiss.	Endemik	[NT]	Ir-Tur
<i>Barbarea lutea</i> Cullen & Coode	Endemik	[EN]	Ir-Tur
<i>Heldreichia bupleurifolia</i> Boiss. subsp. <i>rotundifolia</i> (Boiss.) Parolly, Nordt &Mumm.	Endemik	[LC]	Ir-Tur
<i>Isatis aucheri</i> Aucher ex Boiss.	Endemik	[LC]	Ir-Tur
<i>Asyneuma limonifolium</i> (L.) Janchen subsp. <i>pestalozzae</i> (Boiss.) Dambolt	Endemik	[LC]	
<i>Eremogone acerosa</i> (Boiss.) Ikonn.	Endemik	[LC]	
<i>E. ledebouriana</i> (Fenzl.) Ikonn.	Endemik	[LC]	
<i>Gypsophila nodiflora</i> (Boiss.) Barkoudah	Endemik	[VU]	Ir-Tur
<i>Phryna ortegioides</i> (Fisch. et Mey.) Pax et Hoffm.	Endemik	[NT]	Ir-Tur
<i>Silene brevicaulis</i> Boiss	Endemik	[LC]	Ir-Tur
<i>Euphorbia anacamperos</i> Boiss. var. <i>anacamperos</i>	Endemik	[VU]	
<i>E. grisophylla</i> M. L. S.	Endemik	[LC]	Ir-Tur
<i>Astragalus pennatulus</i> Hub.-Mor. et Chamb.	Endemik	[NT]	Ir-Tur
<i>Hedysarum pestalozzae</i> Boiss.	Endemik	[LC]	Ir-Tur
<i>H. pogonocarpum</i> Boiss	Endemik	[LC]	
<i>H. pycnostachyum</i> Hedge & Hub.- Mor.	Endemik	[EN]	Ir-Tur
<i>Lathyrus brachypterus</i> Cel. var. <i>brachypterus</i>	Endemik	[LC]	Ir-Tur
<i>Lotus gebelia</i> Vent var. <i>anthylloides</i> Boiss	Endemik	[NT]	Ir-Tur
<i>Hypericum capitatum</i> Choisy var. <i>luteum</i> Robson	Endemik	[DD]	Ir-Tur
<i>Cyclotrichium niveum</i> (Boiss.) Manden. et Scheng.	Endemik	[VU]	Ir-Tur
<i>Nepeta crinita</i> Montbret et Aucher ex Benth	Endemik	[EN]	Ir-Tur
<i>N. sorgerae</i> Hedge et Lamond	Endemik	[LC]	Ir-Tur
<i>Phlomis sieheana</i> Rech. Fil.	Endemik	[LC]	Ir-Tur
<i>Scutellaria orientalis</i> L. subsp. <i>pectinata</i> (Montbret & Aucher ex Benth.) J.R. Edm	Endemik	[LC]	Ir-Tur

<i>Linaria corifolia</i> Desf.	Endemik	[LC]	Ir-Tur
<i>L. genistifolia</i> (L.) Miller subsp. <i>confertiflora</i> (Boiss.) Davis	Endemik	[LC]	Ir-Tur
<i>L. genistifolia</i> (L.) Miller subsp. <i>praealta</i> (Boiss.)	Endemik	[NT]	D. Akd.
<i>Veronica macrostachya</i> Vahl. subsp. <i>mardinensis</i> (Bornm.) M.A	Endemik	[VU]	Ir-Tur
<i>V. orientalis</i> Miller subsp. <i>carduchorum</i> P.H. Davis ex M.A. Fisch.	Endemik	[LC]	Ir-Tur
<i>Acantholimon acerosum</i> (Willd.) Boiss. var. <i>parvifolium</i>	Endemik	[VU]	Ir-Tur
<i>A. venustum</i> Boiss. var. <i>assyriacum</i> (Boiss.) Boiss.	Endemik	[NT]	Ir-Tur
<i>Rheum telianum</i> İlçim	Endemik	[CR]	Ir-Tur
<i>Helleborus vesicarius</i> Aucher	Endemik	[NT]	D. Akd.
<i>Pyrus syriaca</i> Boiss var. <i>microphylla</i> Zoh. Ex Browicz	Endemik	[VU]	
<i>Asperula cilicica</i> Hausskn ex Ehrend.	Endemik	[CR]	Ir-Tur
<i>Haplophyllum cappadocicum</i> Spach.	Endemik	[NT]	Ir-Tur
<i>H. myrtifolium</i> Boiss.	Endemik	[LC]	Ir-Tur
<i>Scrophularia cryptophila</i> Boiss. Et Heldr	Endemik	[LC]	D. Akd.
<i>Verbascum melitenense</i> Hub.-Mor	Endemik	[NT]	Ir-Tur
<i>V. urceolatum</i> Hub.-Mor	Endemik	[NT]	Ir-Tur
<i>Allium nemrutdaghense</i> Kit Tan & Sorger	Endemik	[EN]	Ir-Tur
<i>A. scabriflorum</i> Boiss	Endemik	[VU]	Ir-Tur
<i>Bellevalia longistyla</i> (Misch.) Gross		[VU]	Ir-Tur
<i>Hyacinthus orientalis</i> L. subsp. <i>chionophilus</i>	Endemik	[NT]	Ir-Tur
<i>Ornithogalum alpigenum</i> Stapf.	Endemik	[NT]	D. Akd.
<i>Iris galatica</i> Siehe	Endemik	[LC]	Ir-Tur
<i>Fritillaria crassifolia</i> Boiss. Et Huet. subsp. <i>crassifolia</i>	Endemik	[LC]	
<i>Tulipa sintenesii</i> Baker	Endemik	[LC]	Ir-Tur
<i>Elymus lazicus</i> subsp. <i>divaricatus</i> (Boiss.& Bal.) Melderis	Endemik	[LC]	Ir-Tur
<i>Euphorbia craspedia</i> Boiss.		[VU]	Ir-Tur
<i>Ixiolirion tataricum</i> (Pallas) Herbert subsp. <i>tataricum</i> (Labill.) Takht.		[VU]	Ir-Tur
<i>Koeleria pyramidata</i> (Lam.) P. Beauv.		[VU]	Avr-Sib.
<i>Phlomis kotschyana</i> Hub.-Mor.		[VU]	D.Akd.
<i>Salvia cassia</i> Samuelss. ex Rech. Fil.		[VU]	D.Akd.
<i>Hedysarum pannosum</i> (Boiss.) Boiss.		[VU]	Ir-Tur
<i>S. caucasica</i> Anderson		[VU]	Avr-Sib.

SONUÇ ve TARTIŞMA

Araştırma alanından toplanan 66 endemik ve 8 nadir taksondan 54 tanesi İran-Turan , 7 takson Akdeniz elementi, 2 takson Avr.-Sibirya olup 11 taksonun fitocoğrafik bölgeleri belli değildir (Çizelge 1).

Yine 66 endemik taksondan IUCN tehlike kategorisine göre 3 takson CR (Çok tehlikede), 9 takson VU (Zarar görebilir), 14 takson NT (Tehdit altına girebilir) ve 33 takson LC (En az endise verici), 5 takson EN (Tehlikede), 2 takson DD (yetersiz verili) kategorisine girdiği görülmüştür (Çizelge 2). Endemik olmayan fakat nadir olan takson sayısı 8 VU (Zarar görebilir)'dir.

Çizelge 1. Endemik ve nadir taksonların fitocoğrafik bölge durumu

Element Çeşidi	Takson Sayısı	% Oran
İran-Turan El.	54	73
Akd.	7	10
Avr-Sib	2	2
Bilinmeyen	11	15

Çizelge 2. Endemik ve nadir taksonların IUCN tehlike sınıfı durumu

IUCN tehlike sınıfı	Takson Sayısı	% Oran
EN	5	7
DD	2	3
CR	3	4
VU	17	23
NT	14	19
LC	33	44

Bu çalışma 1. yazarın Akdağ (Adıyaman/Malatya) florası adlı doktora çalışmasından üretilmiştir.

KAYNAKÇA

Avcı, H., (2019). Akdağ (Çelikhan/Adıyaman) florası", Yüksek lisans tezi, İnönü Üniversitesi,170.

Çetiner, M., (2020). Atatürk Baraj Gölü Havzası'nın Şanlıurfa Bölümü Florası, Yüksek lisans tezi, Adıyaman Üniversitesi Fen Bilimleri Enstitüsü Adıyaman, 80.

Davis, P. H., (1965-1985). Flora of Turkey and the East Aegean Islands. Vol.; 1-9. Edinburgh University Press, Edinburgh.

Eğilmez, Ç.,(2014). Gölbaşı gölleri (Adıyaman) havzası'nın vejetasyonu, Yüksek lisans tezi, Adıyaman Üniversitesi Fen bilimleri Enstitüsü, Adıyaman, 119.

Ekim, T., Koyuncu, M., Vural, M., Duman H., Aytaç Z., Adıgüzel, N., (2006). Red Data Book of Turkish Plants (Türkiye Bitkileri Kırmızı Kitabı), Türkiye Tabiatını Koruma Derneği, Ankara, 2000.

Ortaç, Z., Tel, A. Z., (2021). Gazihan Dede Mesire Alanı (Adıyaman, Türkiye) Florası, Türler ve Habitatlar Dergisi, (2021) 2(1): 33–53. 221

Şimşek, A.,(2015). Ulubaba Dağı (Adıyaman) Vegetasyonu Üzerine Fitososyolojik Ve Fitoekolojik Çalışmalar, Yüksek lisans tezi, Adıyaman Üniversitesi Fen Bilimleri Enstitüsü, Adıyaman, 140.

Tak, M., (2015). Çelikhán Çat Barajı (Adıyaman) Havzası'nın Fitososyolojik Ve Fitoekolojik Özellikleri, Yüksek lisans tezi, 7 Aralık Üniversitesi Fen Bilimleri Enstitüsü, Kilis, 114.

Tel, A, Z, (2001). Nemrut Dağı (Adıyaman) Vegetasyonu, Doktora tezi, Yüzüncü Yıl Üniversitesi Fen Bilimleri Enstitüsü, Van, 102.

Tel, A. Z., Ortaç, İ. & İlçim, A. (2019). Karatepe-Aslantaş milli parkı ve bazı doğal/kültürel koruma alanları üzerine floristik bir araştırma (Osmaniye, Türkiye). *Commagene Journal of Biology*, 3(2), 103-109.

Tel, A. Z., Ortaç, İ. & İlçim, A. (2021). Hatay ilinin bazı doğal ve kültürel sit alanları florası üzerine bir çalışma. *Artvin Çoruh Üniversitesi Orman Fakültesi Dergisi*, 22(1), 9-18.

Tel, A. Z., Ortaç, İ. & Özuslu, E. (2022a). A floristic study on some natural and cultural sites of Adana (Türkiye) province. *Anatolian Journal of Botany* 6(2): 92-103. doi:10.30616/ajb.1152708

Ergun USLU ve ark. 13(4), 2450-2466, 2023 Anamur, Bozyazı ve Aydincık (Mersin-Türkiye)'daki Bazı Doğal Sit Alanlarının Flora ve Genel Vegetasyon Yapısının İncelenmesi 2466

Tel, A.Z., Ortaç, İ. & Özuslu, E. (2022b). Bazı mağara ve obrukların (Mersin/Türkiye) flora ve genel vejetasyon yapıları üzerine bir araştırma, *Biological Diversity and Conservation*, 15(3): 356-368. doi: 10.46309/biodicon.2022.1180111

Tel, A.Z., Ortaç, İ, İlçim, A., & Özuslu, E. (2023). Mersin İlindeki (Türkiye) Bazı Doğal ve Kültürel Sit Alanlarının Floristik Yapısı Üzerine Bir Çalışma. *KSÜ Tarım ve Doğa Dergisi*, 26(5), 1056-1065. <https://doi.org/10.18016/ksutarimdog.vi.1196119>

Tel, A. Z., Tak, M., (2012). Perre (Pirin) Antik Şehri (Adıyaman) Vegetasyonu, *Biyoloji Bilimleri Araştırma Dergisi* 5 (2): 45-62, 2012 ISSN: 1308-3961, EISSN: 1308-0261, 2012.

Tel, A.Z., Tak, M., (2018). Karagöl Sülüklü Göl Havzası, Gerger/Adıyaman) Florası Üzerine Bir araştırma, *Adyütayam Cilt 6, Sayı 1; 40-53. 222*

Tel, A.Z., Tak, M., (2021). Çelikhán ve Yakın Çevresinin (Adıyaman/Türkiye) Flora ve Vegetasyonu Üzerine Araştırmalar,

<http://cografyaharita.com/haritalarim/4o-adiyaman-konum-haritasi.png>

<https://www.google.com.tr/intl/tr/earth/> (erişim tarihi, 16/08/2022)

<https://www.google.com.tr/intl/tr/earth/> (erişim tarihi, 14/12/2022)

DAMAGE SITUATION OF MEDITERRANEAN FRUIT FLY, *CERATITIS CAPITATA* (WIEDEMANN) (DIPTERA: TEPHRITIDAE) IN PEACH GARDENS OF IĞDIR PROVINCE

IĞDIR İLİ ŞEFTALİ BAHÇELERİNDE AKDENİZ MEYVE SİNEĞİ, *CERATITIS CAPITATA* (WIEDEMANN) (DIPTERA: TEPHRITIDAE)'NİN ZARAR DURUMU

Abdullah İREÇ

Iğdır Provincial Directorate of Agriculture and Forestry, Iğdır, Türkiye

ORCID: 0000-0003-4788-7211

Celalettin GÖZÜAÇIK

Iğdır University, Faculty of Agriculture, Department of Plant Protection, Iğdır, Turkey

ORCID: 0000-0002- 6543-7663

ÖZET

Bu çalışma, 2017 - 2018 yıllarında Iğdır ili merkez köylerinde şeftali ağaçlarında zararlı olan Akdeniz meyvesineği, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae)'nın yayılışı, biyolojisi ve Iğdır ilinde şeftali bahçelerindeki zarar durumunu gözlemlemek için ele alınmıştır. Çalışmalara 2017 yılı Kasım ayı ortalarında Iğdır ili merkez köyü olan Kadıkışlak'ta Iğdır iline ait yöresel ismi Zeferan olan şeftali bahçesinde tesadüfî olarak yere düşmüş 100 adet şeftali meyvesi toplanarak başlanılmıştır. Toplanan meyveler bıçak yardımıyla açılarak Akdeniz meyvesineği larvalarının sayımı yapılarak şeftali meyvelerinde zarar durumu incelenmiştir. Çalışma sonuçlarına göre, yere dökülen 100 adet şeftali meyvesinin % 25 oranında Akdeniz meyve sineğinin zararı yani larvalı meyve görülmüş, 1 adet meyvede; en fazla 4 adet ve en az ise 2 adet Akdeniz meyve sineği larvası tespit edilmiştir. Iğdır ilinde, çiftçiler tarafından şeftali ağaçlarına asılan Delta tipi feromon tuzakların olduğu bahçelerde yapılan çalışmada Akdeniz meyvesineği erginlerinin genelde temmuz sonu ve ağustos ayının ilk haftalarında doğada çıkışları olmakta ve ilk larva eylül ayının birinci ve ikinci haftalarında görülmektedir. Yapılan gözlemlerde Iğdır ilinde Akdeniz meyve sineği erginin ve larvasının kasım ayının ikinci haftasına kadar doğada canlı olarak görülmektedir. Iğdır ilinde Akdeniz meyve sineğinin en fazla zararı yöresel ismi zeferan olan geççi şeftalide ve yine yöresel ismi payız (beyaz) olan yerli geççi şeftalide yapmaktadır. Yapılan bu çalışmada örnek olarak incelenen meyvelerde sadece Akdeniz meyvesineği larvalarına rastlanılmıştır.

Anahtar Kelimeler: *Ceratitis capitata*, zarar durumu, şeftali ağaçları, Iğdır

ABSTRACT

This study was conducted to observe the distribution and biology of the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae), which is harmful to peach trees in the central villages of Iğdır province in 2017 - 2018, and the damage situation in peach orchards in Iğdır province. The studies started in mid-November 2017 by collecting 100 peach fruits that accidentally fell to the ground in the peach garden, whose local name is Zeferan, in Kadıkışlak,

the central village of Iğdır province. The collected fruits were opened with the help of a knife and the damage to peach fruits was examined by counting the Mediterranean fruit fly larvae. According to the results of the study, 25% of 100 peach fruits that fell on the ground were damaged by the Mediterranean fruit fly, that is, damaged fruit, and in 1 fruit; A maximum of 4 and at least 2 Mediterranean fruit fly larvae were detected. In the study conducted in the gardens of Iğdır province, where Delta type pheromone traps were hung on peach trees by farmers, Mediterranean fruit fly adults generally emerge in nature at the end of July and the first weeks of August, and the first larvae are seen in the first and second weeks of September. According to the observations made, Mediterranean fruit fly adults and larvae can be seen alive in nature until the second week of November in Iğdır province. In Iğdır province, the Mediterranean fruit fly causes the most damage to the late peach, whose local name is zeferan, and to the local late peach, whose local name is payiz (white). In this study, only Mediterranean fruit fly larvae were found in the fruits examined as samples.

Keywords: *Ceratitis capitata*, damage status, peach trees, Iğdır

GİRİŞ

Prunus persica L. olarak bilinen şeftali Rosales takımının Rosaceae familyasının Prunoideae alt familyasına ve *Prunus* cinsine ait bir bitki türüdür. Kültürü yapılan şeftali çeşitleri meyve etinin çekirdekten ayrılma durumuna göre, yarma (*Prunus persica domestica*) ve et şeftalileri (*Prunus persica vulgaris*) olmak üzere iki gruba ayrılmaktadır. Şeftalinin günümüzde kültür formları Çin'den yayıldığı anlaşılmaktadır. Ilıman iklim meyve türleri grubunda yer alan Şeftali anavatanı olan Çin'den çeşitli yollarla dünyanın birçok bölgesine yayılmıştır. Şeftali bitkisinin ülkemize girişi, net olarak hangi tarihte ve kimler tarafından getirildiği tam olarak bilinmemekle birlikte Evliya Çelebi Seyahatnamesinde şeftali yediğinden bahsetmiştir (Childers, 1954; Zielinski, 1955; Özçağırın ve ark., 2005; Gür ve Şeker 2016).

Türkiye, coğrafi konumu sayesinde tropik bahçe bitkileri dışında tüm meyve türleri için oldukça uygun bir iklime sahiptir. Bu nedenle, Türkiye, bahçe bitkilerinin doğuş yeri ve birçok meyve türünün anavatanı olarak kabul edilmektedir. Türkiye'de yetiştirilen meyve türlerinin büyük bir bölümünü ılıman iklim meyve türleri oluşturmaktadır. Ülkemiz önemli şeftali üreticisi ülkelerinden biridir. Sert çekirdekli meyve türleri içerisinde yer alan şeftali, ülkemizde bazı yıllarda birinci sırada yer alırken, bazı yıllarda ise kayısıdan sonra ikinci sırada yer almaktadır. Türkiye'nin 75 ilinde şeftali yetiştiriciliği yapılmakta olup, toplam üretimin yarısını Çanakkale (163.871 ton), Mersin (134.210 ton) ve Bursa (105.586 ton) illeri gerçekleştirmektedir. 2023 yılı verilerine göre Türkiye'de şeftali üretimin en fazla yapıldığı beş il sırayla Çanakkale, Mersin, Bursa, Denizli ve İzmir'dir. Türkiye'de farklı zamanlarda hasat edilebilen erkenci ve geççi şeftali çeşitleri yılın beş ayı boyunca pazara taze olarak sunulmaktadır. Erkenci şeftali çeşitleri ülkemizde Akdeniz Bölgesinde yetiştirilirken, geççi çeşitler ise Marmara ve Ege Bölgesinde yetiştirilmektedir (Gül ve Akpınar 2006; Crisosto, 2008; Tük, 2024).

Iğdır Ovası, Türkiye'nin mikroklima özelliği gösteren en geniş ovalarından biridir. Sahip olduğu bağıl nem yüksekliği ile Iğdır ili, çevresine göre iklim, toprak ve bitki örtüsü gibi doğal çevre özellikleri açısından farklı özellikler göstermektedir. Aras nehri boyunca doğu-batı doğrultusunda uzanan ovanın deniz seviyesinden yüksekliği güneyden-kuzeye ve batıdan-doğuya doğru gidildikçe azalmaktadır. Ovanın yüzey eğimi ortalama %1-2, ortalama yüksekliği 850 m'dir. Ova topraklarında, bitki adaptasyonu ve tarımı tehdit eden tuzluluk problemi baş göstermektedir. Üst ve alt topraklar genellikle tuzlu, kök bölgesi altındaki tabakalar ise normal bir durum göstermektedir (Eryiğit, 2011).

Iğdır ilinde yoğun bir şekilde şeftali yetiştiriciliği yapılmaktadır özellikle temmuz, ağustos ve eylül-ekim aylarında üç defa şeftali hasadı yapılmaktadır. Temmuz ayında hasadı yapılan şeftali

ve nektari çeşitleri erkenci çeşitlerdir, Ağustos ayında hasadı yapılan şeftali çeşidi ise, Glohaven gibi çeşitlerdir. Eylül ve ekim aylarının sonuna kadar hasadı yapılan şeftali çeşitleri, bölgemize ait Zeferan ve beyaz şeftali çeşitleridir. Beyaz şeftalinin yöresel ismi payız şeftalidir. Iğdır İl Tarım ve Orman Müdürlüğü'nün 2024 yılı verilerine göre 2189 dekar alanda şeftali üretimi yapılmaktadır.

Dünya üzerinde çok geniş alanlara yayılmış ve önemli meyve zararlılarından birisi olan Akdeniz meyvesineği (*Ceratitis capitata* (Wiedemann, 1824) (Diptera:Tephritidae))'nin zararını doğrudan meyvede ve meyvelerin olgunlaşma dönemlerinde yapması sebebiyle ürünlerin ticari değerlerini ve kalitesini düşürerek meyve dökümlerine neden olduğu, ülkelere dış karantina zararlısı olarak bilindiği ve bulaşıklılık toleransının sıfır olduğu, bu nedenle ihrac edilen ürünlerde tek bulaşık meyve olması halinde tüm ürünün ihracatını engellediği bilinmektedir (Kahyaoğlu, 2011). Akdeniz meyvesineği başta ılıman ve subtropik iklim meyveleri, bazı sebzeler ve süs bitkileri olmak üzere dünyada 70'den fazla ülkede, 260'dan fazla konukçusu olan polifag bir zararlıdır (Kasap ve Aslan, 2016; Satar & Tiring, 2017).

Iğdır ilinde şeftali yetiştiriciliğinde birçok Bitki Koruma sorunu olarak başta ana zararlılar, San jose kabuklubiti (*Quadraspidiotus perniciosus* Comst.), Şeftali filiz güvesi *Anarsia lineatella* (Zeller), Şeftali gövde kanlıbiti, (*Pterochloroides persicae* (Cholodkovsky) ve son yıllarda Iğdır ilinde geçici şeftalilerde yoğun zarar yapan Akdeniz meyvesineği (*C. capitata*)'nin yaptığı zararını dikkat çekmek için bu çalışma yapılmıştır.

MATERYAL VE YÖNTEM

Araştırma, Iğdır ili Merkez ilçesi Kadıkışlak köyünde Iğdır iline ait yöresel ismi zeferan olan (yaklaşık 10-15 yaşlarında ve 6 dekar, 40° 02' 66" N, 44° 05' 74" E) şeftali bahçesinde tesadüfî olarak yere düşmüş 100 adet şeftali meyvesi toplanarak siyah poşetlere alınmış ve uygun bir ortamda meyvelerin içleri bıçak yardımıyla açılarak Akdeniz meyvesineğinin larvalarının sayımı yapılarak şeftali meyvelerinde zarar durumunu gözlemlenmiştir (Şekil 1,2 ve 3). Delta tipi feromon tuzaklarla ergin uçurları belirlenmiştir. Ergin böcek larvalarının şeftali meyvesindeki zararı fotoğraflanmış ve kayıt altına alınmıştır.



a



Şekil 1.Şeftali ağaçlarının altında meyvelerin toplanması (a,b)



Şekil 2. Şeftali ağaçlarının altında toplanan vuruklu meyvelerin incelenmesi




Şekil 3. Delta tipi feromon tuzaklara düşen Akdeniz meyvesineğinin erginleri

ARAŞTIRMA VE BULGULAR

Çalışmalara 2017 yılı Kasım ayı ortalarında Iğdır ili merkez köyü olan Kadıkışlak'ta Iğdır iline ait yöresel ismi Zeferan olan şeftali bahçesinde tesadüfî olarak yere düşmüş 100 adet şeftali meyvesi toplanarak başlanılmıştır. Toplanan meyveler bıçak yardımıyla açılarak Akdeniz meyvesineği larvalarının sayımı yapılarak şeftali meyvelerinde zarar durumu incelenmiştir. Çalışma sonuçlarına göre, yere dökülen 100 adet şeftali meyvesinin % 25 oranında Akdeniz meyve sineğinin zararı yani vuruksuz meyve görülmüş, 1 adet meyvede; en fazla 4 adet ve en az ise 2 adet Akdeniz meyve sineği larvası tespit edilmiştir. Iğdır ilinde, çiftçiler tarafından şeftali ağaçlarına asılan Delta tipi feromon tuzakların olduğu bahçelerde yapılan çalışmada Akdeniz meyvesineği erginlerinin genelde temmuz sonu ve ağustos ayının ilk haftalarında doğada çıkışları olmakta ve ilk larva eylül ayının birinci ve ikinci haftalarında görülmektedir. Yapılan gözlemlerde Iğdır ilinde Akdeniz meyve sineği erginin ve larvasının kasım ayının ikinci haftasına kadar doğada canlı olarak görülmektedir. Iğdır ilinde Akdeniz meyve sineğinin en fazla zararı yöresel ismi zeferan olan geççi şeftalide ve yine yöresel ismi payız (beyaz) olan yerli geççi şeftalide yapmaktadır. Yapılan bu çalışmada örnek olarak incelenen meyvelerde sadece Akdeniz meyvesineği larvalarına rastlanılmıştır.

Zararı

Çalışma sonuçlarına göre, yere dökülen 100 adet şeftali meyvesinin % 25 oranında Akdeniz meyve sineğinin zararı yani vuruksuz meyve görülmüş, 1 adet meyvede; en fazla 4 adet ve en az ise 2 adet Akdeniz meyve sineği larvası tespit edilmiştir. 2018 yılında yapılan çalışmada ise, Akdeniz meyvesineğinin doğada ilk çıkışı haziran ayının sonunda görülmüş ve 2017 yılına göre zarar oranı, hemen hemen 3 katı kadar görülmüştür (Şekil 4).



Şekil 4. Şeftali ağaçlarının altında toplanan meyvelerde Akdeniz meyvesinin larvası (a) ve meyvedeki zararı (b,c)

SONUÇ

Yapılan çalışmalar incelendiğinde, yere dökülen 100 adet şeftali meyvesinin % 25 oranında Akdeniz meyve sineğinin zararı yani vuruksuz meyve tespit edilmiştir. Iğdır ilinde 2021 yılında yapılan çalışmada şeftalide; Çalpala köyünde % 12, Kuzugüden köyünde % 8,4 ve Küllük köyünde % 18,4 oranında zarar tespit edilmiştir (Tamer ve Yıldırım, 2021). 1 adet meyvede; en fazla 4 adet ve en az ise 2 adet Akdeniz meyve sineği larvası tespit edilmiştir. Iğdır ilinde, çiftçiler tarafından şeftali ağaçlarına asılan Delta tipi feromon tuzakların olduğu bahçelerde yapılan çalışmada Akdeniz meyvesineği erginlerinin genelde temmuz sonu ve ağustos ayının ilk haftalarında doğada çıkışları olmakta ve ilk larva eylül ayının birinci ve ikinci haftalarında görülmektedir. Tamer ve Yıldırım 2021 yılında Iğdır ilinde yaptıkları çalışmada Akdeniz meyvesineğinin ilk erginlerini 5 Eylül 2021 tarihinde tespit etmişlerdir. Yapılan gözlemlerde Iğdır ilinde Akdeniz meyve sineği erginin ve larvasının kasım ayının ikinci haftasına kadar doğada canlı olarak görülmektedir. Iğdır ilinde Akdeniz meyve sineğinin en fazla zararı yöresel ismi zeferan olan geççi şeftalide ve yine yöresel ismi payız (beyaz) olan yerli geççi şeftalide yapmaktadır. Yapılan bu çalışmada örnek olarak incelenen meyvelerde sadece Akdeniz meyvesineği larvalarına rastlanılmıştır.

KAYNAKÇA

Childers, N. F., 1954. Modern Fruit Science. 4 th Edition. New Jersey.

Crisosto, C.H., Valero, D., 2008. Harvesting and Postharvest Handling of Peaches for the Fresh Market. Layne, D.R. and Bassi, D. (Ed), The Peach: Botany, Production and Uses, 595-596. CAB International, UK, 615 p.

Eryiğit, T., (2011). Iğdır İlinin Kalkınmasında Endüstri Bitkileri Tarımının Önemi ve Geliştirilmesi İçin Bazı Öneriler. Yüzüncü Yıl Üniversitesi Ziraat Fakültesi Tarım Bilimleri Dergisi 21(1): 73 - 81.

Gül, M., Akpınar, M., 2006. An assessment of developments in fruit production in the world and Turkey. Mediterranean Agricultural Sciences 19 (1): 15- 27.

Gür, E., Şeker, M. 2016. Beyaz nektarin tiplerinin Prunus cinsine giren önemli türlerle melezlenmesi sonucu elde edilen pomolojik sonuçlarının karşılaştırılması. VII. Ulusal Bahçe Bitkileri Kongresi Bildirileri, 25-29 Ağustos 2015, Çanakkale Onsekiz Mart Üniversitesi Ziraat Fakültesi, Çanakkale.

Kahyaoğlu, M. (2011). Hazır yem (bait) formülasyonu geliştirilmesi ve Akdeniz meyve sineği [*Ceratitis capitata* Wiedemann (Diptera: Tephritidae)] mücadelesinde kullanım olanaklarının araştırılması (Tez no 447971). [Doktora Tezi Ankara Üniversitesi Fen Bilimleri Enstitüsü].Yükseköğretim Kurulu Ulusal Tez Merkezi.

Kasap, A. ve Aslan, M.M. (2016). Akdeniz meyve sineğinin feromon tuzaklarla (*Ceratitis capitata* Wied.) (Diptera: Tephritidae)'nin nar ve hurmadaki popülasyon takibi ve zarar oranının tespiti. Doğa Bilimleri Dergisi, 19 (1), 43-50.

Özçağırın, R., Ünal, A., Özeker, E., İsfendiyaroğlu, M., 2005. Ilıman İklim Meyve Türleri, Sert Çekirdekli Meyveler Cilt I. Ege Üniversitesi Yayınları, Ziraat Fakültesi Yayın.

Satar, S., Tiring, G. (2017). *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae)'nın bazı meyve bahçelerinde popülasyon dalgalanması. Türk. Entomoloji Bülteni, 7 (3), 239-247.

Tamer, E. ve Yıldırım, E., 2023. Iğdır ili meyve bahçelerinde *Ceratitis capitata* (Wiedemann, 1824)(Diptera: Tephritida)'nın popülasyon gelişimi, yoğunluğu ve bulaşıklık oranının belirlenmesi. Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi, 26(3), 560-569.

TÜİK, 2024. Türkiye İstatistik Kurumu: Bitkisel Üretim İstatistikleri. Erişim Adresi: <https://biruni.tuik.gov.tr/medas/?kn=92&locale=tr>.

Zielinski, Q. B., 1955. Modern Systematic Pomology. W.M. C. Brown Comp., Dubuque, Iowa.

LANDSCAPE ARCHITECTURE AND ENERGY IN SMART CITIES

AKILLI ŞEHİRLERDE PEYZAJ MİMARLIĞI VE ENERJİ

Rıdvan TİK

Iğdır Üniversitesi, Ziraat Fakültesi, Bahçe Bitkileri Bölümü, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0009-0008-1102-1743>

Tuncay KAYA

Iğdır Üniversitesi, Ziraat Fakültesi, Bahçe Bitkileri Bölümü, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-9126-4567>

ÖZET

Akıllı şehirler, dijital teknolojileri ve ileri veri analitiğini kullanarak şehir yaşamını daha verimli, sürdürülebilir ve yaşanabilir hale getirmeyi amaçlayan modern yerleşim alanlarıdır. Akıllı şehir tasarımlarında; enerji verimliliği, kaynakların etkin kullanımı ve çevresel etkilerin azaltılması temel hedefler arasında yer almakta ve peyzaj mimarlığı bu hedeflere ulaşmada önemli bir araç olarak karşımıza çıkmaktadır. Peyzaj mimarlığı, doğal çevrenin sunduğu potansiyeli değerlendirerek, sürdürülebilir tasarım çözümleri üretmeyi amaçlayan bir disiplindir. Bu alanda gerçekleştirilen tasarımlar, enerji tüketimini minimize etmeyi ve ekosistem hizmetlerinin verimliliğini artırmayı hedefler. Doğal unsurların ve yerel ekosistemlerin entegrasyonu hem çevresel hem de toplumsal açıdan fayda sağlayan çözümler sunarak, çevreye duyarlı ve fonksiyonel mekânlar yaratılmasına olanak tanır. Akıllı şehirlerde yeşil alanlar, doğal iklimlendirme sistemleri, su yönetimi uygulamaları ve yenilenebilir enerji çözümleri enerji verimliliğinin sağlanmasında etkin rol oynayan unsurlardır. Özellikle yeşil çatılar, bitkilendirme düzenlemeleri, ağaçlandırma ve sürdürülebilir su yönetimi gibi stratejiler enerji tüketiminin ve karbon emisyonlarının azaltılmasına katkı sağlamaktadır. Ayrıca, yenilenebilir enerji kaynaklarının peyzaj tasarımına entegrasyonu, güneş panelleri ve rüzgâr türbinlerinin kullanımı gibi çözümler şehirlerin enerji bağımsızlığını artırmakta ve çevre dostu enerji kullanımını teşvik etmektedir. Dijital teknolojilerin, özellikle sensörler ve veri analiz sistemlerinin peyzaj tasarımına entegrasyonu, enerji yönetimini daha verimli hale getirerek kaynak kullanımını optimize etmektedir. Bu tür dijital çözümler, peyzaj mimarlığında enerji verimliliği potansiyelini artırmakta ve sürdürülebilir şehir tasarımına dair yeni fırsatlar sunmaktadır. Bu bağlamda, peyzaj mimarlığı, enerji verimliliği ve çevresel sürdürülebilirlik hedeflerine ulaşmada akıllı şehirlerin tasarımında önemli bir rol oynamakta, akıllı altyapı ve çevresel yönetim uygulamalarını destekleyerek sürdürülebilir bir kentsel gelişim sürecine katkıda bulunmaktadır.

Anahtar Kelimeler: Akıllı Şehirler, Enerji, Peyzaj Mimarlığı, Peyzaj Tasarımı.

ABSTRACT

Smart cities are modern residential areas that aim to make urban life more efficient, sustainable and livable by using digital technologies and advanced data analytics. In smart city designs; energy efficiency, efficient use of resources and reducing environmental impacts are among the main goals and landscape architecture is an important tool in achieving these goals. Landscape architecture is a discipline that aims to produce sustainable design solutions by utilizing the potential offered by the natural environment. Designs in this field aim to minimize energy consumption and increase the efficiency of ecosystem services. The integration of natural elements and local ecosystems enables the creation of environmentally sensitive and functional spaces by providing both environmentally and socially beneficial solutions. In smart cities, green spaces, natural air conditioning systems, water management practices and renewable energy solutions play an active role in ensuring energy efficiency. In particular, strategies such as green roofs, planting arrangements, afforestation and sustainable water management contribute to reducing energy consumption and carbon emissions. In addition, solutions such as the integration of renewable energy sources into landscape design, the use of solar panels and wind turbines increase the energy independence of cities and encourage the use of environmentally friendly energy. The integration of digital technologies, especially sensors and data analysis systems, into landscape design optimizes resource use by making energy management more efficient. Such digital solutions increase the potential for energy efficiency in landscape architecture and offer new opportunities for sustainable urban design. In this context, landscape architecture plays an important role in the design of smart cities in achieving energy efficiency and environmental sustainability goals and contributes to a sustainable urban development process by supporting smart infrastructure and environmental management practices.

Keywords: Smart Cities, Energy, Landscape Architecture, Landscape Design.

GİRİŞ

Küreselleşme ve teknolojik gelişmeler, şehirlerin yapısında köklü değişikliklere yol açmıştır. İletişim ve ulaşım imkânlarındaki iyileşmeler ise mekânlar arası hareketliliği artırarak, iş fırsatlarının yoğun olduğu şehirlerde nüfus yoğunluğunun artmasına neden olmuştur. Şehirlerde yaşayan bireyler, sınırlı zaman dilimlerinde birçok sorumluluğu yerine getirmeye çalışarak sürekli bir zaman baskısı altındadır. Bu hızlı yaşam temposu, aynı zamanda doğal kaynakların hızla ve bilinçsizce tüketilmesine yol açmaktadır. Zaman içinde, şehir yaşamının daha verimli, rahat ve sürdürülebilir hale getirilmesi amacıyla, şehir karmaşasından kaynaklanan olumsuz etkilerin en aza indirilmesi ve sınırlı enerji kaynaklarının daha verimli bir şekilde kullanılması için yeni teknolojilerle pek çok yenilikçi şehir uygulaması geliştirilmiştir. Bu uygulamalarla şehirler, akıllı hale gelmeye başlamıştır. Bu süreçte, önemli bir nokta, şehir projelerinin tüm paydaşların ihtiyaç ve beklentilerine uygun, bütünsel bir yaklaşım ile tasarlanmasıdır (Kocaman, 2020).

Akıllı kentler, sürdürülebilir kalkınma hedeflerine ulaşmada önemli bir rol oynamaktadır. Bu çerçevede, akıllı kent teknolojilerinin etkin ve verimli yönetimi, hizmetlerin aksatılmaması ve kentlerin sürdürülebilirliğinin sağlanması açısından kritik öneme sahiptir. Akıllı yönetim, klasik kamu yönetimi yöntemlerinden farklı olarak, katılımcılık, şeffaflık ve hesap verebilirlik ilkeleri doğrultusunda daha hızlı, doğru ve etkin karar alma süreçlerini teşvik etmektedir (Anonim, 2019). Kentlerde yönetim eksiklikleri; plansız büyüme, verimsiz arazi kullanımı, kontrolsüz enerji tüketimi, biyolojik çeşitlilik kaybı, hava kirliliği, yüksek sera gazı emisyonları ve verimsiz kaynak kullanımı gibi sorunlara yol açarak, kentlerin sürdürülebilirliğine yönelik tehditleri arttırmaktadır (Gökşen, 2024). Bu bağlamda, peyzaj planlaması ve yönetimi, akıllı kentlerin sürdürülebilirliğini sağlamada önemli bir araç olarak öne çıkmaktadır. Peyzaj tasarımı

ve ekolojik dengeyi koruma stratejileri, kentlerin çevresel etkilerini azaltmaya ve biyolojik çeşitliliği korumaya katkıda bulunarak, daha yaşanabilir ve sürdürülebilir kentler inşa edilmesine olanak tanıyacaktır.

Peyzaj mimarlığı, doğa, planlama ve tasarım kavramlarını sistematik bir şekilde ele alarak; sanat, bilim, mühendislik ve teknolojiyi birleştirir. Bu disiplin, alan kullanım kararları doğrultusunda doğal ve kültürel kaynakların doğru şekilde değerlendirilmesini sağlayarak ekolojik, ekonomik ve işlevsel açıdan sürdürülebilir bir planlama ve yönetim süreci sunar. Sürdürülebilir kentlerin temel unsurlarından biri olarak, çevre koruması, ekosistem ve doğal kaynakların analizi ile yönetimi, kırsal ve kentsel alanların planlanması, çevresel etki değerlendirmesi, rekreasyonel, kültürel ve kentsel açık alanların tasarımı, yaya bölgeleri, karayolları, endüstriyel ve tarım alanlarının planlanması, peyzaj mimarlarının sorumluluğundadır. Peyzaj mimarları, özellikle doğal kaynakları etkili bir şekilde yorumlayarak, ekonomi ve ekoloji dengesini sağlamak için kritik bir rol üstlenmektedir. Ancak sürdürülebilir şehirlerin oluşturulmasında karşılaşılan en önemli zorluk, bu görevlerin sınırlarının net bir şekilde belirlenmemiş olması ve yasal düzenlemelerle desteklenmemesidir. Bu durum, meslek grupları arasında yetki paylaşımının yetersiz yapılmasına ve dolayısıyla hedeflere ulaşmada görev karmaşasının ortaya çıkmasına yol açmaktadır (Atıl vd., 2005).

Peyzaj mimarlığı, akıllı şehirlerin tasarımında önemli bir yer tutar ve şehirlerin sürdürülebilirlik hedeflerine ulaşmasında kritik bir rol oynar. Akıllı şehirler, teknolojinin, çevresel sürdürülebilirliğin ve kullanıcı dostu yeşil alanların entegrasyonu ile insanların yaşam kalitesini artırmayı amaçlayan dinamik alanlardır. Bu bağlamda, peyzaj mimarlığı, yeşil alanların stratejik planlanması ve tasarımı yoluyla şehir sakinlerinin doğayla etkileşimini artıran ve yaşam alanlarını iyileştiren mekanlar oluşturur. Aynı zamanda, peyzaj tasarımı yalnızca estetik bir yaklaşımı değil, aynı zamanda su ve enerji verimliliği sağlayan çözümleri de içermektedir. Yeşil alanların enerji tasarrufu, su yönetimi ve ekosistem hizmetleri gibi sürdürülebilir unsurlar göz önünde bulundurularak tasarlanması, peyzaj mimarlarının önemli sorumlulukları arasında yer almaktadır (Anonim, 2025a).

Bu çalışma, akıllı şehirler bağlamında peyzaj mimarlığının enerji yönetimi ile nasıl uyumlu bir şekilde çalışabileceğini ve bu entegrasyonun şehir yaşamını nasıl iyileştirebileceğini incelemeyi amaçlamaktadır. Akıllı şehirler; gelişmiş teknolojiler ve veri yönetimi kullanarak sürdürülebilirlik, enerji verimliliği ve yaşam kalitesini artırmayı hedefleyen modern yerleşim alanlarıdır. Bu bağlamda, peyzaj mimarlığı yalnızca estetik bir ortam yaratmakla kalmayıp, çevresel verimliliği artırmak, enerji tasarrufu sağlamak ve doğal kaynakları korumak amacıyla çeşitli stratejiler geliştiren bir disiplindir. Peyzaj tasarımı, şehirlerde yeşil alanların, su yönetimi sistemlerinin ve doğal iklimlendirme çözümlerinin etkin bir şekilde entegre edilmesi ile enerji kullanımını optimize edebilir ve karbon ayak izini azaltabilir. Ayrıca, akıllı enerji yönetim sistemlerinin peyzaj tasarımı ile bütünleşmesi, şehirlerin enerji tüketimini izleyerek daha sürdürülebilir bir yapıya kavuşmalarını sağlayabilir. Bu tür bir entegrasyon, çevresel sürdürülebilirliğin yanı sıra, akıllı şehirlerin enerji verimliliği hedeflerine ulaşmasında önemli bir rol oynamaktadır.

AKILLI ŞEHİR KAVRAMI

Literatürde, "sürdürülebilir şehir", "kentsel sürdürülebilirlik" ve "akıllı şehir" gibi birbirine benzer kavramlar sıklıkla kullanılsa da, bu terimler arasındaki farklar genellikle belirgin şekilde tanımlanmamaktadır. Bu durum, literatürde genel olarak tutarlı bir tanımın olmadığına işaret etmektedir (Neirotti vd., 2014). Kentsel sürdürülebilirlik, çevresel faktörlere odaklanırken, akıllı şehir kavramı (Karadağ, 2013; Ahvenniemi vd., 2017) sosyal ve ekonomik faktörleri daha fazla göz önünde bulundurur. Gül ve Çobanoğlu (2017) akıllı şehirleri, bilgi ve iletişim teknolojilerinin kentsel alanlarda etkin bir şekilde kullanıldığı, toplum ve şehir arasında

karşılıklı iletişimin sağlandığı, sürdürülebilirlik, bilgi, teknoloji ve katılım gibi unsurların bir arada yer aldığı bir yapı olarak tanımlanmaktadır. Bu tanım, akıllı şehirlerin sadece teknolojik altyapılarla değil aynı zamanda sosyal ve ekonomik etkileşimlerle de şekillenen bir sistem olduğunu vurgulamaktadır. Şekil 1’ de akıllı şehrin yapısı gösterilmiştir (Anonim, 2025b).

Ulusoy, (2017)’a göre bir kentin akıllı şehir olarak nitelendirilebilmesi için, öncelikle sekiz temel alanda bilgi ve iletişim teknolojilerinin kullanılarak şehir sorunlarına akıllı çözümler geliştirilmesi gerekmektedir. Ayrıca, bu çözümlerin şehir halkı tarafından bir yaşam biçimi olarak benimsenip etkin bir şekilde kullanılmaları önemlidir. Akıllı şehirler, sekiz temel parametreden en az beşinin uygulanmasıyla oluşturulan, akıllı ve yenilikçi teknolojilere dayalı şehirlerdir. Bu parametreler; şehir yönetiminde verimliliği artıran dijital çözümler ve eğitimde teknoloji odaklı akıllı yönetim ve akıllı eğitim, bireylerin dijital platformları etkin şekilde kullanarak şehir yönetimine aktif katılım sağladığı akıllı vatandaş, sağlık hizmetlerinin dijitalleşmesi ve teknolojiyle daha erişilebilir hale geldiği akıllı sağlık, enerji tüketiminin izlenmesi ve verimli kullanımını sağlayan teknolojik sistemleri barındıran akıllı enerji, enerji verimliliği ve sürdürülebilirlik sağlamak için entegre teknolojilerin kullanıldığı akıllı yapılar, şehirdeki tüm alanlarda teknolojinin etkin şekilde entegre edilmesi ve dijital çözümlerle optimize edildiği akıllı teknoloji, trafik yönetimi ve toplu taşıma sistemlerinin verimli hale getirilmesi için akıllı ulaşım, şehir altyapısının dijitalleşmesi ve sürdürülebilir bir şekilde yönetilebilen akıllı altyapı, kamu güvenliğini artıran, dijital teknolojilerle desteklenen güvenlik önlemleri ve izleme sistemlerine sahip akıllı güvenlidir. Bu parametrelere ek olarak, peyzaj mimarlığı da akıllı şehirlerin sürdürülebilirliğini destekleyen önemli bir unsurdur. Peyzaj tasarımı, doğal kaynakların verimli kullanılmasını, yeşil alanların artırılmasını, ekosistem hizmetlerinin korunmasını ve şehir halkının yaşam kalitesini yükselten çözümler sunarak, akıllı şehirlerin bütünsel bir şekilde geliştirilmesine katkıda bulunur.



Şekil 1. Akıllı şehir yapısı

Akıllı şehirler, gelişmiş modern altyapı sistemleri ile bilgi ve iletişim teknolojilerinin entegrasyonu sonucu ortaya çıkan, sürdürülebilirlik hedeflerine ulaşmayı amaçlayan kent modelleridir. Son yirmi yıl içinde, akıllı şehir girişimleri, sürdürülebilirliğin sağlanmasında etkili bir çözüm olarak değerlendirilmekte olup, bu yaklaşımlar, şehirlerin daha verimli ve çevre dostu hale gelmesini hedeflemektedir. Ancak, bu tür çözümleri hayata geçirmek, hızla kontrolsüz bir şekilde büyüyen kentsel alanlarda ve çevresinde oldukça zordur. Düşük mekânsal standartlara sahip, kontrolsüz gelişen bölgeler ile bu alanlarda ortaya çıkan çevre

kirliliği, doğal kaynakların etkin bir şekilde yönetilmesini zorlaştırmakta ve akıllı şehir geliştirme stratejilerinin uygulanmasında önemli zorluklara neden olmaktadır (Hoang vd., 2021).

AKILLI ŞEHİRLERDE ENERJİ

Enerji tüketiminin azaltılması, atık enerjinin etkin bir şekilde kullanımı ve yenilenebilir enerji üretimi, sürdürülebilir şehircilik ve çevre dostu kalkınma hedefleri doğrultusunda önemli bir tema olarak öne çıkmaktadır (Kocaman, 2020). Bu bağlamda, enerji verimliliğini artırmaya yönelik stratejiler, hem çevresel etkilerin azaltılmasına hem de ekonomik tasarrufların sağlanmasına olanak tanımaktadır. Atık enerji kullanımı, mevcut enerji sistemlerinin daha verimli hale getirilmesi için kritik bir yöntemken, yenilenebilir enerji kaynaklarının kullanımı ise fosil yakıtlara olan bağımlılığı azaltarak, sürdürülebilir enerji üretiminin teşvik edilmesine katkı sağlamaktadır.

Hollanda, akıllı ulaşım altyapıları alanında önemli bir adım atarak dünyanın ilk "Akıllı yol" projesini hayata geçirmiştir. Bu proje, Amsterdam'ın yaklaşık 100 kilometre güneyinde yer alan Oss kentinden geçen N329 Otoyolu'nda (Şekil 2) uygulanmaktadır. Otoyol üzerinde şerit ve yol çizgilerinin fosforlu malzemelerle aydınlatılması, gece sürüşlerini daha güvenli hale getirmeyi amaçlamaktadır. Bu yenilikçi uygulama, enerji tüketimini azaltırken, aynı zamanda sürücülerin yol durumunu daha iyi algılamalarını sağlayarak trafik güvenliğini artırmaktadır (Anonim, 2025c). Akıllı yol teknolojileri, kentlerin sürdürülebilir ve verimli ulaşım altyapılarına dönüşümüne katkı sağlamak ve akıllı ulaşım sistemlerinin potansiyelini göstermektedir.



Şekil 2. Akıllı enerji

Akıllı şehirler, temiz enerji kullanımını ön planda tutan bir yaklaşım olarak değerlendirilmeli ve bu bağlamda akıllı şebeke çözümleri, yenilenebilir enerji kaynaklarının ve yüksek verimli kojenerasyon güç tesislerinin gelişimini desteklemek için temel bir rol oynamaktadır. Akıllı şebekeler, dağılmış enerji kaynaklarının entegrasyonunu mümkün kılarak, tüketicilere güvenli ve güvenilir elektrik enerjisi sağlama noktasında esneklik sunmaktadır. Bu yapı, enerji arzını daha verimli hale getirirken, aynı zamanda sürdürülebilir enerji sistemlerinin etkinliğini artırmaktadır (Eremia vd., 2017).

AKILLI ŞEHİRLERDE PEYZAJ MİMARLIĞININ ROLÜ

Peyzaj mimarları, kentlerin karbon nötr bir geleceğe ulaşabilmesi için çeşitli stratejiler geliştirmekte (Şekil 3) ve bu doğrultuda önemli çabalar sarf etmektedirler (Yörüklü, 2021). Sürdürülebilir tasarım anlayışında yeşil alanlar, çevresel etkileri azaltma ve yaşam kalitesini artırma açısından önemli bir yer tutmaktadır. Kentsel ısı adası etkilerinin hafifletilmesi ve binaların enerji tüketiminin azaltılması gibi hedeflerle bağlantılı olarak, çeşitli yeşil alan uygulamaları kent yaşamına entegre edilmiştir. Bu uygulamalar arasında ağaçlı cadde ve bulvarlar, çatı bahçeleri ve cephe yeşillendirmeleri (biyofilik tasarım) gibi yenilikçi çözümler öne çıkmaktadır (Ortaççeşme vd., 2023).



Şekil 3. Karbon nötr kentler için peyzaj mimarlığı yaklaşımları

Peyzaj mimarlığı, şehirlerin çevresel sürdürülebilirliğini artırmak, estetik değerlerini yükseltmek ve sakinlerin yaşam kalitesini iyileştirmek amacıyla çeşitli stratejiler geliştiren bir disiplindir. Akıllı şehirler bağlamında peyzaj mimarlığı, doğal çevrenin korunması ve yeniden inşası, yeşil alanların tasarımı, su yönetimi, biyolojik çeşitliliğin artırılması ve hava kalitesinin iyileştirilmesi gibi çok sayıda alanda kritik bir rol oynamaktadır. Akıllı şehirlerde, peyzaj tasarımı yalnızca görsel bir estetik oluşturmakla kalmaz, aynı zamanda çevresel etkileri azaltmak, enerji tasarrufu sağlamak ve sürdürülebilir bir yaşam biçimi inşa etmek için dijital çözümlerle entegre olur. Peyzaj mimarları, akıllı şehirlerin tasarımında yeşil çatılar, dikey bahçeler, suyun geri kazanımı ve yağmur suyu yönetimi gibi yenilikçi çözümleri entegre eder. Bu tasarımlar, şehir içindeki ısı adası etkisini azaltır, enerji tüketimini düşürür ve doğal kaynakları daha verimli kullanarak çevre dostu bir şehir ortamı yaratır. Akıllı şehirlerde peyzaj tasarımının bir diğer önemli boyutu ise sosyal ve kültürel yaşamın desteklenmesidir. Yaya yolları, açık hava etkinlik alanları, parklar ve rekreasyonel alanlar, insanların doğa ile etkileşime girmesini sağlayarak fiziksel ve ruhsal sağlıklarını iyileştirir.

ENERJİ VERİMLİLİĞİ VE PEYZAJ MİMARLIĞI

Enerji verimliliği, akıllı şehirlerin temel hedeflerinden birini oluşturur. Şehirlerin enerji tüketiminin artması, küresel ısınma ve çevresel bozulma gibi sorunlara yol açmaktadır. Peyzaj

mimarlığı, bu sorunun çözülmesine katkı sağlayacak çeşitli stratejiler geliştirebilir. Yeşil alanların, ağaçların ve bitki örtüsünün stratejik olarak yerleştirilmesi, şehirdeki doğal iklimin düzenlenmesine yardımcı olur. Ağaçlar, binaların etrafına dikildiğinde, yazın gölgeleme sağlayarak soğutma maliyetlerini düşürürken, kışın rüzgar bariyerleri oluşturarak ısınma giderlerini azaltabilir. Ayrıca, doğru peyzaj tasarımı, binaların güneş ışığından faydalanmasını sağlayarak, doğal aydınlatma ve ısıtma için enerji tasarrufu sağlar. Peyzaj mimarlığı, enerji verimliliği sağlamak için su yönetimi açısından da önemli bir role sahiptir. Su geri dönüşüm sistemlerinin entegre edilmesi, yağmur suyu toplama sistemleri ve yer altı su depolama çözümleri, şehirlerdeki su tüketimini azaltır ve enerji üretiminde kullanılan suyun korunmasına yardımcı olur. Bu stratejiler, aynı zamanda doğal su döngüsünün korunmasına ve su kaynaklarının sürdürülebilir kullanımına katkıda bulunur.

Enerji Verimliliği

Enerji verimliliği, enerji kaynaklarının daha etkin ve sürdürülebilir bir şekilde kullanılması amacıyla geliştirilen bir yaklaşımdır. Bu yaklaşım, enerji tasarrufu sağlamanın yanı sıra çevresel etkilerin azaltılması ve enerji maliyetlerinin düşürülmesi hedeflerine ulaşmak için çeşitli tasarım stratejileri ve ileri teknolojilerin entegrasyonunu içerir (Büyüksaraç ve Hacıoğlu, 2018). Enerji verimliliği, sadece enerji tüketimini minimize etmeyi değil aynı zamanda çevresel sürdürülebilirliği teşvik etmeyi de amaçlayan çok boyutlu bir kavramdır. Enerji verimliliği, enerji tüketiminin minimize edilmesi amacıyla tasarım stratejilerinin ve teknolojik çözümlerin entegre edilmesini kapsayan bir yaklaşımdır. Bu kapsamda, enerji verimli bina tasarımı, yüksek verimli ısıtma ve soğutma sistemleri, etkili aydınlatma çözümleri ve yenilenebilir enerji kaynaklarının kullanımı gibi unsurlar öne çıkmaktadır. Söz konusu stratejiler ve teknolojiler, enerji tüketimini optimize ederken, sürdürülebilirlik ve çevresel etkilerin azaltılması yönünde önemli katkılar sağlamaktadır (Çevik ve Özdemir, 2017).

Enerji Verimliliği Perspektifinden Peyzaj Tasarımının Rolü ve Önemi


Enerji etkin peyzaj tasarımı, sürdürülebilir çevre düzenlemeleri açısından birçok kritik fayda sunmaktadır. Bu tasarım yaklaşımı, mikro klima düzenlemeleri ile çevresel koşulları iyileştirirken, su tüketiminin azaltılmasına ve yağmur suyu ile atık suların arıtılarak yeniden kullanımına olanak tanır. Ayrıca, doğaya saygılı malzemelerin kullanımı, gürültü ve hava kirliliği kontrolü, CO₂ emisyonlarının azaltılması gibi çevresel etkilerin iyileştirilmesini sağlar. Bu unsurlar, enerji giderlerini önemli ölçüde düşürerek ekonomik tasarruf sağlarken, aynı zamanda yaşam kalitesinin artırılmasına ve iklim değişikliği ile mücadeleye de katkıda bulunur. Enerji etkin peyzaj tasarımı, bütün ekosisteme fayda sağlayarak, hem çevresel hem de sosyal sürdürülebilirliği destekleyen bir strateji olarak büyük bir öneme sahiptir (Uzun ve Aydın, 2018).

Enerji etkin peyzaj tasarım yaklaşımlarını belirleyen kriterler (Çizelge 1); çevresel etkenlere ilişkin parametreler, yapı tasarımına ilişkin parametreler, yenilenebilir enerji kaynaklarının kullanımı ve diğer enerji etkin peyzaj tasarım araçlarıdır (Yurtsev, 2015).

Çizelge 1. Enerji etkin parametrelerin sınıflandırılması

Çevresel Etkenlere İlişkin Parametreler	Yapı Tasarımına İlişkin Parametreler	Yenilenebilir Enerji Kaynaklarının Kullanımı	Diğer Enerji Etkin Peyzaj Tasarım Araçları
İklim	Yapının konumu	Güneş	Yeşil duvarlar
Topografik yapı	Yapının formu	Rüzgâr	Yeşil çatılar
Su	Yapının malzemesi	Su	
Toprak			
Bitki örtüsü			

Doğru yapılmış bir peyzaj düzenlemesi, kışın güneş ışınlarını yapıya yönlendirirken, sert rüzgârları ve istenmeyen esintileri farklı yönere yönlendirmekte ve yazın dik gelen güneş ışınlarını da engellemektedir. Bu özelliği ile yıl boyu yapıların harcadığı enerji miktarını azaltmakta ve enerji etkinliğine büyük katkılar sağlayabilmektedir. Aynı zamanda iç mekân hava kalitesini de arttırmaktadır. Yapı kabuğuna yapılan bitkilendirme müdahaleleri ile ise, ekstra termal kütle ve yalıtım sağlanması mümkün olmaktadır (Öztürk Sarı, 2013). Enerji etkin peyzaj tasarımı, çevresel sürdürülebilirlik ilkeleri doğrultusunda, ihtiyacı olan enerjiyi kendi kaynaklarından temin etmeyi hedefler. Bu tür tasarımlar, yenilenebilir ve doğal enerji kaynaklarını kullanan sistemler entegre ederek, çevreye duyarlı bir yaklaşım sergiler. Ayrıca, kullanılan malzemelerin daha az toksik bileşen içermesi veya geri dönüşümle elde edilen malzemeler olması önem taşır. Yeşil alanların korunması, yağmur suyu ve atık suların toplanarak arıtılması ve yeniden kullanılması gibi uygulamalar, bu tür tasarımların temel bileşenlerindedir. Enerji etkin çözümlerin başarıyla uygulanabilmesi için; (Şekil 4) çevresel faktörler, teknik ve teknolojik gelişmeler ile tasarım ve yapısal çözümlerin bir bütün olarak ele alınması gerekmektedir (Karaca, 2008; Yurtsev, 2015).

**Şekil 4.** Enerji etkinlik yöntemleri ve yöntemlerin etkileşimi

AKILLI ŞEHİRLERDE ENERJİ VERİMLİ PEYZAJ UYGULAMALARI

Günümüz dünyasında enerjinin her alanda kullanılması, peyzaj tasarımında da enerji kullanımını önemli hale getirmiştir. Yenilenebilir enerji kaynakları, doğanın sunduğu sürekli ve tükenmez enerji kaynaklarıdır. Güneş, rüzgar, biyokütle ve jeotermal enerji gibi yenilenebilir

enerji türleri çevreye duyarlı ve sürdürülebilir enerji çözümleri sunmaktadır. Bu enerji kaynaklarının peyzaj tasarımında kullanımı, çevresel etkileri azaltmanın yanı sıra enerji verimliliğini artıran estetik ve işlevsel alanlar yaratmayı amaçlar (Tik ve Kaya, 2024).

Akıllı çevre, şehirlerin enerji kullanımını daha verimli ve tasarruflu hale getirmeyi hedefleyen bir sistem olarak, çeşitli teknolojik çözümlerle enerji verimliliğini artırmaktadır. Bu tür çevrelerde, türbinler, akıllı aydınlatmalar, mobilyalar, akıllı binalar ve geniş alanlarda yer alan paneller aracılığıyla enerji elde edilmektedir. Enerjinin, şehir ihtiyacını karşılayacak şekilde planlı bir biçimde mikro şebekeler aracılığıyla kullanılması, büyük enerji akımlarının israfını engeller ve elde edilen ekonominin kaybolmasını önler. Akıllı çevre, aynı zamanda atık su kullanımında verimliliği sağlamayı amaçlayan arıtma sistemlerini içererek, gelecekteki su kıtlığına karşı önlemler geliştirmektedir. Şehrin atık, çöp ve diğer dönüşüm unsurlarının sürdürülebilir kullanımı da bu yapının bir bileşeni olarak gelişmektedir. İnşa edilen yeşil binalar ve koruma çalışmalarının yanı sıra, çevresel kalitenin artırılması ve doğal cazibenin korunması sağlanmaktadır. Hava kirliliği ve gürültüye yönelik yapılan iyileştirmelerle çevre daha sağlıklı ve uygun hale gelmektedir. Ayrıca, doğal güzelliklerin sunulma potansiyeli artarken, akıllı parklar ve mesire alanları rekreasyon amaçlı kullanılmakta ve yaşam kalitesine katkı sağlamaktadır (Güler ve İkiel, 2022).

Akıllı çevre ile ilgili olarak, altyapı sistemlerinin etkin bir şekilde yönetilmesi, geri dönüşüm süreçlerinin izlenmesi, su sızıntılarının ve su kalitesinin sürekli takibi, katı atıkların toplanıp ayrıştırılması, gürültü kirliliği ve enerji tüketiminin izlenmesi gibi bir dizi önemli çalışma yürütülmektedir (Varol, 2017). Kirliliği önlemek, enerji verimliliğini artırmak, doğal güzellikleri koruyarak kentleri geliştirmek ve çevre dostu sürdürülebilir yeşil alanlar oluşturmak, akıllı çevre uygulamalarıyla ulaşılmak istenen temel hedefler arasında yer almaktadır (Aslan, 2018). Bu hedefler, şehirlerin sürdürülebilir kalkınma süreçlerinde çevresel etkilerin azaltılması ve yaşam kalitesinin artırılması açısından kritik bir rol oynamaktadır. Akıllı çevre, bu doğrultuda teknolojik çözümler ve yeşil altyapı unsurlarını birleştirerek, çevresel sürdürülebilirliği sağlamak ve kentsel alanların doğal kaynakları daha verimli kullanmasını teşvik etmek amacıyla önemli bir araçtır.

Konya Büyükşehir Belediyesi Mor Şebeke Projesi (Şekil 5), akıllı şehir uygulamaları bağlamında sürdürülebilir su yönetimi açısından çok önemli bir yer tutmaktadır. Konya Büyükşehir Belediyesi KOSKİ Genel Müdürlüğü tarafından geliştirilen bu proje, kentsel yaşam kalitesini artıran ve iklim değişikliği ile mücadelede etkin bir çözüm sunan çok yönlü bir uygulama olarak karşımıza çıkmaktadır. Bu bağlamda, akıllı şehir teknolojileri kullanılarak atık suyun geri kazanımı ve içme suyu kaynaklarının korunması sağlanmaktadır. Kentsel sorunlar arasında yer alan su kıtlığına ve çevresel sürdürülebilirlik hedeflerine cevap veren Mor Şebeke, akıllı sulama teknolojileri ile sürdürülebilir şehircilik kavramını güçlendirmektedir (Anonim, 2025d). Akıllı şehirlerde enerji verimliliğini artırmaya yönelik peyzaj mimarlığı uygulamaları, teknolojiyle entegre bir şekilde çalışır. Akıllı sulama sistemleri, hava durumu verileri ve toprak nemi sensörleri kullanarak su tüketimini minimize eder ve enerji israfını önler. Ayrıca, güneş panelleri ve rüzgar türbinleri gibi yenilenebilir enerji kaynakları, peyzaj tasarımına entegre edilerek şehirlerin enerji üretimine katkı sağlar. Dijital sistemler, peyzaj alanlarının izlenmesi ve yönetilmesi için de kullanılabilir. Örneğin, akıllı şehirlerdeki yeşil alanlar, dijital haritalama ve veri toplama sistemleriyle izlenebilir, böylece bakım ihtiyaçları ve su kullanımı optimize edilir. Bu, sadece enerji tasarrufu sağlamakla kalmaz, aynı zamanda peyzajın sürdürülebilirliğini artırır ve yönetimini kolaylaştırır.



Şekil 5. Konya mor şebeke: sürdürülebilir su ve enerji yönetimi

SONUÇ

Akıllı şehirler, modern teknolojilerin şehir planlamasına entegrasyonu ile sürdürülebilir bir yaşam alanı yaratma amacını taşımaktadır. Bu bağlamda, peyzaj mimarlığı, şehirlerin çevresel, estetik ve fonksiyonel açıdan sağlıklı olmasını sağlamak için kritik bir rol oynamaktadır. Akıllı şehirlerde, peyzaj tasarımı yalnızca doğal alanların düzenlenmesi değil aynı zamanda enerji verimliliğini artıran, çevresel etkileri azaltan ve yaşam kalitesini iyileştiren bir strateji olarak önem kazanmaktadır. Doğru yerleştirilen bitki örtüsü, yenilenebilir enerji kaynaklarının entegrasyonu ve akıllı sulama gibi teknolojiler, enerji verimliliğini optimize ederken şehirlerin ekolojik dengesini korumaya da katkı sağlamaktadır.

Peyzaj mimarlığı, akıllı şehirlerin enerji yönetiminde etkin bir rol oynarken, tasarım süreçlerinde ileri teknolojilerin ve sürdürülebilir yaklaşımların kullanılmasında ayrıca gerekmektedir. Akıllı sistemler ve sensörler sayesinde, peyzaj alanları daha verimli bir şekilde yönetilebilir ve doğal kaynaklar daha etkin bir biçimde kullanılabilir. Bu bağlamda, enerji verimliliği ve çevresel sürdürülebilirlik, akıllı şehirlerin gelişiminde merkezi bir yer tutmaktadır.

Sonuç olarak, akıllı şehirler yalnızca enerji yönetimi gibi teknik çözümler bütünü olarak değil, aynı zamanda şehir sakinlerinin yaşam kalitesini artıran, çevresel etkiyi minimize eden ve enerji kaynaklarını verimli kullanan bütünsel yaklaşımlar olarak değerlendirilmelidir. Bu bağlamda peyzaj mimarlığının akıllı şehir konsepti ve işleyişindeki rolü giderek kritik hale gelmektedir. Gelecekte ortaya çıkabilecek yeni sorunların çözümünde ve yeni teknolojilerin akıllı şehirlere adaptasyonunda bu rol daha da belirginleşecektir.

KAYNAKLAR

Ahvenniemi, H., Huovila, A., Pinto-Seppä, I., & Airaksinen, M. (2017). What are the differences between sustainable and smart cities?. *Cities*, 60, 234-245.

Anonim, (2019). Çevre ve Şehircilik Bakanlığı. (2019). 2020-2023 Ulusal akıllı şehirler stratejisi ve eylem planı. <https://www.akillisehirler.gov.tr/wp-content/uploads/EylemPlani.pdf>.

Anonim, (2025a). <https://mipeyzaj.com.tr/akilli-sehirler-ve-peyzaj-mimarligi/> Erişim Tarihi: 21.01.2025

Anonim, (2025b). <https://esenler.bel.tr/haberler/genel/akilli-sehir-esenler-ile-herkes-zengin-oluyor/> Erişim Tarihi: 30.01.2025

Anonim, (2025c). <https://www.hurriyet.com.tr/gundem/dunyanin-ilk-akilli-yolu-aciliyor-26224830> Erişim Tarihi: 30.01.2025

Anonim, (2025d). <https://www.urbased.com/konya-mor-sebeke-surdurulebilir-su-ve-enerji-yonetimi/> Erişim Tarihi: 30.01.2025

Aslan, M. M. (2018). *Akıllı kent uygulamaları üzerine bir inceleme: Kahramanmaraş örneği*. Mustafa Kemal Üniversitesi Sosyal Bilimler Enstitüsü, Yüksek Lisans Tezi. Hatay, Türkiye.

Atıl, A., Gülgün, B., & Yörük, İ. (2005). Sürdürülebilir kentler ve peyzaj mimarlığı. *Ege Üniversitesi Ziraat Fakültesi Dergisi*, 42(2), 215-226.

Büyüksaraç, A. & Hacıoğlu, U. (2018). “Sürdürülebilir mimari tasarım ve yenilenebilir enerji kaynakları.” *Journal of Sustainable Energy and Environment*, 5(1), 28-34.

Çelebi, A., Ay, B. & Tuna, A. (2019). *Enerji etkin peyzaj tasarım yaklaşımları ile kentsel donatı elemanları*. Uluslararası Ispec Mühendislik Ve Fen Bilimleri Kongresi, 3-4 Mayıs 2019, Malatya, Türkiye, 77sy.

Çevik, M. & Y.Özdemir, M. (2017). “Sürdürülebilir enerji kaynakları ve binaların enerji performansı.” *Journal of Energy and Climate Change*, 8(1), 45-52.

Eremia, M., Toma, L., & Sanduleac, M. (2017). The smart city concept in the 21st century. *Procedia Engineering*, 181, 12-19.

Gökşen, H. (2024). Sürdürülebilirlik perspektifinde akıllı kentlerde akıllı yönetim uygulamalarının rolü ve insani gelişime etkisi. *Kent Akademisi*, 17(Sürdürülebilir İnsani Kalkınma ve Kent), 174-194.

Gül, A., & Çobanoğlu, Ş. (2017). Avrupa’da akıllı kent uygulamalarının değerlendirilmesi ve Çanakkale’nin akıllı kente dönüşümünün analizi. *Süleyman Demirel Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 22(Kayfor 15 Özel Sayısı), 1543-1565.

Güler, H., & İkiel, C. (2022). Akıllı şehirler kavramı ve Ankara örneğinde yapılan çalışmalar (The Concept of Smart Cities and The Studies Carried Out in The Example of Ankara). *Journal of Anatolian Cultural Research (JANCR)*, 6(1), 44-58.

Hoang, A. T., & Nguyen, X. P. (2021). Integrating renewable sources into energy system for smart city as a sagacious strategy towards clean and sustainable process. *Journal of Cleaner Production*, 305, 127161.

Karaca, M. (2008). Toplu konutlarda enerji etkinliği; Toplu Konut İdaresi Başkanlığı (TOKİ) toplu konut projesi üzerinden bir inceleme. Gazi. Ü. Fen Bilimleri Enstitüsü Şehir ve Bölge Planlama Anabilim Dalı Yüksek Lisans Tezi, Ankara, 282s.

Karadağ, T. (2013). *Akıllı kent yaşamı üzerine bir değerlendirme*. Doctoral Dissertation, Yüksek Lisans Tezi, ODTÜ, Fen Bilimleri Enstitüsü, Ankara.

- Kocaman, E. G. (2020). Akıllı ve sakin şehirler için enerji çözümleri. *İstanbul Sabahattin Zaim Üniversitesi Fen Bilimleri Enstitüsü Dergisi*, 2(2), 40-47.
- Neirotti, P., De Marco, A., Cagliano, AC, Mangano, G., & Scorrano, F. (2014). Akıllı Şehir girişimlerindeki güncel eğilimler: Bazı biçimlendirilmiş gerçekler. *Şehirler*, 38, 25-36.
- Ortaçesme, V., Atik, M., & Muhammetoglu, H. (2023). Akıllı kentlerde akıllı çevre uygulamaları: Kashiwanoha (Japonya) ve Antalya. *Peyzaj*, 5(1), 1-17.
- Öztürk Sarı, S., (2013). *Enerji Etkin Tasarımda Bir Arakesit: Toprak örtülü yapılar*. İstanbul Teknik Üniversitesi Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, 141s.
- Uzun, s. & Aydın, A. (2018). Enerji etkin peyzaj tasarım yaklaşımları.
- Tik, R., Kaya, T. (2024). Possibilities Of Using Renewable Energy Sources In Landscape Studies. International Congress Of High Value-Added Agricultural Products, December 1-3, Iğdır/Türkiye, 560-570.
- Varol, Ç. (2017). Sürdürülebilir gelişmede akıllı kent yaklaşımı: Ankara'daki belediyelerin uygulamaları. *Çağdaş Yerel Yönetimler*, 26(1), 43-58.
- Yörüklü, N. (2021). İklim değişikliği ve küresel ısınma için peyzaj mimarlığı stratejileri: iklim değişikliği politikaları peyzaj beyanı. *PEYZAJ*, 3(1), 43-55.
- Yurtsev, AA. (2015). *Sürdürülebilir mimarlık kapsamında enerji etkin peyzaj tasarım yaklaşımları* (Yayımlanmamış yüksek lisans tezi). Ege Üniversitesi, Fen Bilimleri Enstitüsü, İzmir, 118s.

EFFECTS OF LIQUID VERMICOMPOST, SALICYLIC ACID, AND PROLINE APPLICATIONS ON THE GROWTH OF GREEN ONION (*ALLIUM CEPA* L.)

SIVI SOLUCAN GÜBRESİ, SALİSİLİK ASİT VE PROLİN UYGULAMASININ TAZE SOĞAN GELİŞİMİ ÜZERİNDEKİ ETKİLERİ

Haydar BALCI

Van Yüzüncü Yıl University, Gevaş Vocational School, Department of Plant and Animal Production, VAN/ TURKEY

Orcid No: 0000-0003-0210-3639

Murat KARA

Van Yüzüncü Yıl University, Gevaş Vocational School, Department of Plant and Animal Production, VAN/ TURKEY

Orcid No: 0000-0003-1011-918X

Muhsin YILDIZ

Van Yüzüncü Yıl University, Gevaş Vocational School, Department of Plant and Animal Production, VAN/ TURKEY

Orcid No: 0000-0002-0766-5174

Abstract

This study aimed to evaluate the effects of liquid vermicompost (LVC), salicylic acid (SA), and proline (PR) applications on the growth and development parameters of green onion (*Allium cepa* L.). The research was conducted in an unheated glass greenhouse at Van Yüzüncü Yıl University using a randomized block design, and parameters such as plant height, root length, stem diameter, number of leaves, total fresh and dry weight, and SPAD values were analyzed. The experimental results revealed statistically significant differences ($P \leq 0.05$) among the treatments. Specifically, the SSG1+SA1+PR1 combination yielded the highest values for plant height (29.81 cm), total fresh weight (187.33 g), and total dry weight (22.68 g). However, SPAD values, which represent chlorophyll content, showed an unexpected decrease (60.13) in the combination treatments compared to individual applications.

The positive effects of proline on SPAD values (67.70) and root length (14.64 cm) in individual applications were not fully expressed in the combination treatments. In particular, the high doses of SA and LVC in some combinations were found to have inhibitory effects on certain growth parameters. Clustering and PCA analyses indicated that combination treatments provided certain synergistic effects compared to individual applications but also revealed antagonistic interactions for some parameters. The control group was clearly differentiated from all other treatments, highlighting the significant impact of the applications on plant growth and development.

In conclusion, the study demonstrated that liquid vermicompost, salicylic acid, and proline can enhance growth and yield in green onion cultivation when applied with optimized doses and

combinations. However, further research is recommended to optimize application doses, evaluate the economic feasibility of these combinations in open-field conditions, and investigate their long-term impacts. These findings also suggest that testing these treatments on other crops could contribute to sustainable agricultural practices.

Keywords : Green onion (*Allium cepa* L.), Proline, Liquid vermicompost, Salicylic acid, Plant growth regulators

Özet

Bu çalışma, sıvı solucan gübresi (SSG), salisilik asit (SA) ve prolin (PR) uygulamalarının taze soğan (*Allium cepa* L.) bitkisinin büyüme ve gelişim parametreleri üzerindeki etkilerini değerlendirmek amacıyla gerçekleştirilmiştir. Araştırma, Van Yüzüncü Yıl Üniversitesi'nin ısıtmasız cam serasında, tesadüf blokları deneme desenine göre yürütülmüş ve bitki boyu, kök uzunluğu, gövde çapı, yaprak sayısı, toplam yaş ve kuru ağırlık ile SPAD değeri gibi parametreler incelenmiştir. Deneme sonuçları, uygulamalar arasında istatistiksel olarak anlamlı farklar olduğunu göstermiştir ($P \leq 0,05$). Özellikle, SSG1+SA1+PR1 kombinasyonu, bitki boyu (29,81 cm), toplam yaş ağırlık (187,33 g) ve toplam kuru ağırlık (22,68 g) gibi verim parametrelerinde en yüksek değerleri sağlamıştır. Bununla birlikte, SPAD değeri gibi klorofil içeriğini temsil eden parametrelerde kombinasyon gruplarının bireysel uygulamalara kıyasla daha düşük performans göstermesi (60,13) dikkat çekmiştir.

Prolinin bireysel uygulamalarda SPAD değeri (67,70) ve kök uzunluğu (14,64 cm) üzerindeki olumlu etkileri, kombinasyonlarda yeterince ifade edilememiştir. Özellikle SA ve SSG'nin yüksek dozlarının bazı kombinasyonlarda büyüme parametrelerini baskılayıcı etkileri olabileceği belirlenmiştir. Yapılan kümelendirme analizi ve PCA sonuçları, kombinasyon gruplarının bireysel uygulamalara göre belirli sinerjik etkiler sağladığını ancak bazı parametrelerde antagonistik etkileşimlerin ortaya çıktığını göstermiştir.

Sonuç olarak, sıvı solucan gübresi, salisilik asit ve prolinin doğru dozaj ve kombinasyonlarla kullanıldığında taze soğan yetiştiriciliğinde büyüme ve verimi artırabileceği belirlenmiştir. Ancak, uygulama dozlarının optimize edilmesi gerektiği ve bu kombinasyonların açık saha koşullarında da test edilerek ekonomik uygulanabilirliğinin araştırılması önerilmektedir. Ayrıca, bu bileşiklerin uzun dönemli etkilerinin değerlendirilmesi ve farklı bitkilerde de uygulanarak tarımsal sürdürülebilirliğe katkılarının incelenmesi faydalı olacaktır.

Anahtar Kelimeler : Taze soğan (*Allium cepa* L.), Prolin, Sıvı solucan gübresi, Salisilik asit, Bitki büyüme düzenleyicileri

APPLICATIONS OF BIOPOLYMERS AND BIOPLASTICS IN SMART AGRICULTURE: SENSORS, PACKAGING, AND COATING MATERIALS

Res. Asst. Alperay ALTIKAT

Iğdır University, Faculty of Agriculture Department of Biosystems Engineering, Iğdir,

Prof. Dr. Mehmet Hakkı ALMA

Iğdır University, Faculty of Agriculture Department of Biosystems Engineering, Iğdir,

Abstract

Agriculture faces escalating challenges due to rapid population growth and increasing environmental pressures, necessitating the adoption of innovative and sustainable technologies. Biopolymers and bioplastics, derived from renewable resources, present significant opportunities within the context of smart agriculture, particularly in the development of sensors, packaging, and coating materials. This study provides a comprehensive overview of recent advancements and potential applications of biopolymers and bioplastics in smart agricultural practices. Biopolymers exhibit substantial advantages due to their biodegradability, reduced environmental footprint, and compatibility with conventional agricultural sensor technologies. These eco-friendly materials enhance sensor functionality by replacing traditional synthetic plastics, thus offering sustainable solutions for monitoring soil moisture, temperature, and crop health. Furthermore, bioplastics demonstrate promising capabilities as agricultural packaging materials, effectively reducing plastic waste while simultaneously maintaining the quality and shelf-life of agricultural products. As coating materials, biopolymers play a critical role in protecting crops from environmental stresses, enabling controlled release of nutrients and agrochemicals, and improving overall plant productivity. Despite these benefits, several challenges such as the cost of production, sensor sensitivity, and material durability under field conditions still hinder the widespread adoption of biopolymer-based technologies. Addressing these challenges requires multi-disciplinary collaboration among academia, industry, and governmental bodies to foster research and market adaptation. Consequently, further research and investment in biopolymer technology are crucial for overcoming existing limitations and achieving the goals of sustainable and productive agricultural systems. Ultimately, integrating biopolymers and bioplastics into smart agricultural practices holds great promise for enhancing global food security and environmental sustainability.

Keywords: Biopolymer, Bioplastic, Smart Agriculture, Sensor Technology, Biodegradable Packaging, Agricultural Coatings, Sustainable Agriculture

Introduction to Biopolymers and Bioplastics

Biopolymers, also called biobased polymers, are polymers materially or synthetically made from raw materials of biomolecular origins. Among the biopolymers, some are natural polymers. They are already available from nature, polymerized in living systems due to external genetic information. The main examples are DNA, RNA, proteins and some polysaccharides (Baranwal et al., 2022). The other basic division includes polymers produced from renewable resources, which do not have to be concerned with the genetic background. Biopolymers are

classified as biodegradable, bio-based, non-degradable and bio-based, biodegradable and biodegradable/compostable. Beside the classification with regard to the biodegradability, biopolymers can be divided into three main groups: first, second, and third generation (Das et al., 2023). The first one is based on bio-based synthetic esters, poly(hydroxy acids), e.g., polylactic acid or poly- ϵ -caprolactone; the second group is modified natural polymers, like cellulose acetate, starch blend, and chitosan; the third group encompasses the rest of the biosynthetic polymers, for example, polyhydroxyalkanoates. Bioplastics are biodegradable or bio-based plastics (Dilshad et al.2021). Most of them are biodegradable or compostable. However, their classification is wider, and it includes both biodegradable and their components and non-biodegradable bioplastics. The general classification of bioplastics encompasses some blends and composites with conventional plastics too. The packaging, agricultural mulch films and hygiene materials are some examples of applications of biodegradable plastics. The general division of one-use and multi-use plastics includes another subclassification. There are four main advantages of the bio-based polymers that allow them to be more frequently chosen by both industry sectors and end consumers. The most commonly emphasized feature is their full biodegradation (Bher et al.2022). The second one is the completely renewable aspect of the raw materials used. The third advantage of biopolymers refers to the lower energy consumption during production. The fourth reason for choosing bio-based polymers is their potential for a reduction in the environmental burden. There are several examples of bioplastics, either for food packaging and disposable components, both designed for extensiveness of usage. The airtight biodegradable package with an atmosphere support inserts made of cellulose was prepared for longer storage of polish apples. It is based on thermoformed, corona-activated cellulose acetate foil with a special laminated film. PHB is also used in food packaging. On the other hand, there are wooden kitchen accessories covered with waxy PLA. Konjac glucomannan-based films are used as a coating for minimally processed fruits. The cover with applied dipping or spraying methods significantly decreases the water permeability of the fruit. Films are edible, transparent, and quick-wettable. Microwave technology is responsible for drying fruits with a coating. Bioplastics are also used in packaging cosmetics, for example, in biodegradable lipsticks, made from maltodextrin. Biodegradable bags are produced from PBAT, polylactide, raw flour, and a natural plasticizer. The convenience for immediate usage of waste plastics is emerging due to the national restrictions on the distribution of conventional plastic bags (Nurliasari et al.2025).

Smart Agriculture: Overview and Importance

Smart agriculture, also known as precision agriculture, is an emerging concept that involves the integration of advanced data analytics, Internet of Things (IoT), and Artificial Intelligence (AI) into the traditional farming practices (Miguel-Rojas & Pérez-de-Luque, 2023). The contemporary agriculture is driven by the big data enabled technologies, which offer the smart monitoring and autonomous decision-making services in the farming operations. The most promising benefits of adopting the smart agriculture include the increased crop yields, efficient resource utilization, and reduced manual labor dependence, thereby enhancing the sustainability in the food security management globally. On the other hand, the conventional method of agricultural management is cumbersome and resource expensive where increased environmental degradation, declining resource and global food insecurity can be envisaged, inevitably leading to the need of paradigm shift in the farming practices (Karunathilake et al., 2023). However, the wider adoption of the smart agriculture is largely hindered by the technological constraints in connectivity, data processing, and the huge upfront investment cost. Furthermore, the role of the technology is emphasized in the context of promoting sustainable agricultural management, which is becoming increasingly important in view of the global challenge of food security (MacPherson et al.2022). The text also draws attention to the specific

challenges facing the smart agriculture and to the urgent need for a supportive framework in the form of appropriate incentives, technological infrastructure, and capacity development. Communication technologies, including wireless sensors and smartphones, are powerful tools that can be used for critical tasks such as diagnosis, adaptation to climate change, management of plant health, integrity of crops, and pest and water resource harvesting. All these important aspects are relevant to the attainment of sustainable agricultural goals of doubling crop productivity, securing incomes for farmers, ensuring a safe and sound food supply, helping the rejuvenation of rural communities, and preserving the environment (Khan et al.2021).

Driven agricultural by the global food security and the demand for agriculture raw materials in rapidly growing population, development in technology related equipment for agricultural use and the materials used in agriculture has become increasingly critical in contemporary food and agriculture practices (Abbasi et al., 2022). Today, the contemporary industry of agriculture has eyes on developing ecological, sustainable, and competing methods for increasing globally accepted issues and environmental awareness. Regarding the latest technology, smart agriculture, also identified as precision agriculture, has emerged to facilitate farmers significantly increase extra output with diminished input. Smart agriculture combines ultra-modern technologies with chemicals, machines, and biological substances to amplify agricultural productivity and sustainability (McCaig et al., 2023). The intention behind it is to swiftly achieve unadmitted data, to precisely and economically drive machines in managing appropriate environment for agricultural progress requirements, thereupon noticeably diminishes the need for chemicals, and gives alternative solutions for preserving environment and sustainability of agriculture (Stachowiak et al., 2022).

Data analytics will ameliorate numerous applications of smart agriculture. When sophisticated data analyzing algorithms are installed to the big-data, there will be predictions about drought and irrigation scheduling, epidemic diseases, pests and yield estimation more precisely, and irrigation, plant feeding, and chemical use will be done with distinguished perfection. In recent years, as a compliance that the world had happen to hold on the global agenda against epidemic diseases, it has been uncovered the fundamental significance of food security for countries (Akhter and Sofi2022). Although the world's population is augmenting exponentially and the agriculture raw product is inquired progressively more for nourishing, acute geological hurdles such as droughts, floods, erosion, and global warming are imperilling the agriculture industry. On account of that matter, earthshaking solutions towards ecological equilibration are needed for agriculture. With the integration of new technologies and materials into agriculture industry, more effective and differential practices can be achieved (Chakrabarty, 2021). The main purpose of this study is to put forward the light on positive aspects and potential applications of smart agriculture regarding biopolymers and bioplastics in field of sensors, packaging materials, and coating materials (Alizadeh Sani et al., 2021).

Role of Sensors in Smart Agriculture

Smart agriculture is one of the outcomes of the Fourth Industrial Revolution which is characterized by a range of new technologies interconnected by the Internet of Things for enhanced data collection, analysis, and decision-making processes. Smart agriculture is related to, but not exclusively the same as precision farming (Karunathilake et al., 2023). It is the management of agriculture using a variety of sensors connected to the Internet to collect and analyze real-time or near real-time data and, consequently making up-to-date, data-driven decisions (Miguel-Rojas & Pérez-de-Luque, 2023). Typically, they are used to control irrigation, monitor crop health and manage greenhouse conditions. Sensors are used to monitor environmental conditions such as soil moisture, soil temperature, air temperature, and humidity. Crop monitoring systems also utilize the data from images and sensors to track the development

of disease or pests, in addition to tracking the plants health and growth. As a result of the data collected by sensors, real-time or near real-time conditions can be analyzed, thus agricultural activities can be forecasted, controlled, and adjusted. It is seen as spy technology because it allows us to observe things that used to be invisible (Rasheed et al.2022). The data collected from sensors can result in increased efficiency through improved management. There are countless examples of agricultural systems where the connection between data and practice is not evident, leading to unnecessary resource waste. Sensors are also known as instruments which can detect and respond to an input. In the case of smart agriculture, sensors are used to detect and respond to the data that can inform agricultural activities. In smart agriculture, the integration of sensor technology with other new technologies results in the streamlining of smart farming (Paul et al.2022). But, still, there are several challenges in developing sensor-based smart agriculture. For example, some of the sensors are very fragile. Another problem is the cost: launching sensor systems from scratch is very expensive and the integration of new sensors into a new subprocess is also complex. At the same time, some of the sensors have a poor accuracy that results in misinformed management decisions. Maintaining sensors in order is another issue. Finally, many of the agricultural properties are so diverse that commercial sensors cannot monitor all relevant conditions. Agriculture is becoming a profitable and flexible business. Unfortunately, the increase in global agricultural production has had a detrimental effect on the environment, e.g., excessive water consumption, inappropriate fertilizer use, heavy use of pesticides, and soil erosion (Bhatti et al., 2023). So, it is necessary to offer innovation in smart technology to agrotechnics, which are key in the form of precision agriculture and hydroponics. Enter biopolymers and bioplastics that fit into this trend, improving efficiency and increasing consciousness towards the environment. It can be assumed that, in the near future, innovative biopolymer-based functional coatings will be designed and proposed to frame the deployment of smart technologies due to their potential as sensors (Basavegowda & Baek, 2021).

Biopolymers and Bioplastics in Agricultural Sensors

Smart agriculture is an emerging approach in which technologically advanced solutions are introduced to manage crops, monitor soil condition, and optimize field operations. Biopolymers and bioplastics show great potential as promising materials in smart agriculture. It has recently been observed that numerous agricultural applications can benefit from the use of bi, and partly bio, alternatives of synthetic polymers (Otoni et al.2021). Besides the long-time use in mulch films or seed coatings, such materials can also appear as innovative solutions related to sensors, packaging, or coatings. This paper is aimed to review trends in the intersection of biopolymers, and bioplastics, and agricultural sensors. It addresses new material solutions used in sensor technology – focusing on both the application aspects and the materials’ characteristics – that may stem the revitalization of the discussion on smart solutions in the agriculture-segmented polymers (Mishra et al., 2024).

In the area of sensors, the adhesion of biopolymeric layers is most often investigated in terms of compatibility with bioelectrodes, and bio-interconnection elements, and less with the other common sensor components, and the circuit substrate. Therefore, it was tested if, and how well, the chosen conductive and insulating materials used in standard screen-printed electronics would adhere to thin layers of biopolymers, and associated with the preparation methods. A set of 9 of the most frequently cited biopolymeric materials of different origins was prepared adherewise in relation to the circuitry (Madej-Kiełbik et al.2022). These materials and methods are often patented ones, but without the scientific details published. Additionally, specific solutions were showed, applied as a rigid coating, or somewhat so, but maintaining good flexibility, and necessary for miniaturized strain-effect (or piezoresistive) sensors. All these materials are also biodegradable and compostable, and further tests were done to recognize if

this feature was noticeable after the layer deposition to the circuitry. Organic-layer sensors fabricated with biopolymeric materials are prone to biodegrade post-use in field conditions (Madej-Kiełbik et al., 2022). This might be a desired feature, enhancing the sustainability of the devices, but firstly this needs to be deeply understood and controlled in order not to compromise the functional lifetime of the sensors. All these aspects are discussed using the own fabricated devices as examples to show intermediate levels of success, latest achievements, and presented solutions to potential problems (Andronie et al.2021).

Biopolymers and bioplastics become more popular in agriculture, e.g., in the development of biodegradable active or smart materials. This research presents an overview of sensors, packaging, and coating materials from biopolymers or bioplastics in smart agriculture (Gonçalves et al., 2024). With the development of smart technology, biopolymers and bioplastics have been widely used in agricultural sensors, benefiting the enhancement of sensor functionality and sustainability. In comparison with traditional agricultural sensors, biopolymer and bioplastic-based sensors exhibit advantages of biodegradability and reduced environmental impact (Madej-Kiełbik et al., 2022). Along with the development of eco-friendly technology in agriculture, biopolymer and bioplastic sensors facilitate the sensing network from the farm field to the city, offering valuable guidance for healthy food growth and improving people's life quality.

Regarding agriculture production, sensor technology mainly includes moisture sensors, temperature sensors, density sensors, and mechanical impedance sensors. As essential components of sensors, biopolymers and bioplastics have also been employed to replace conventional sensor encapsulation materials due to their biodegradability (Koh & Khor, 2022). Correspondingly, research has paid increased attention to the development of biopolymer and bioplastic sensor casing materials in the agriculture field. Accordingly, new biopolymer and bioplastic materials can be used as substrates and are fully compatible with current sensor technology in the agriculture field, facilitating the transformation of substitutional conventional materials from biodegradable materials (Alizadeh Sani et al., 2021). Without no doubt, the advancement of agricultural sensors created by biopolymers and bioplastics significantly enriches the innovation of technology in agriculture. Here, some practical applications about biopolymer and bioplastic sensors in agriculture are illustrated. Advanced processing and the material improvements of biopolymer and bioplastic sensors in the agriculture field are discussed based on sensing theory. It is worth noting that a small part of current studies has focused on the development and testing of sensors successfully produced with biopolymers or bioplastics. Ultimately, some challenges and neoteric trends of biopolymer and bioplastic sensor technology in agriculture are proposed, aiming to provide an inspiration guide to attract more interests and improvements (Otoni et al.2021).

Bioplastics in Agricultural Packaging

Agriculture is one of the most strategic sectors for developing countries as well as developing countries. Agriculture and food industries are specifically facing several challenges in the fields of supply, sale and food security over the whole world (Abbasi et al., 2022). Agricultural biopolymers have a wide industrial application in this regard. In the wake of dwindling fossil resources and agro-industrial waste generated in agricultural, they have gained potential application in the area of low cost biodegradable polymer production. Bio-based Polymers are a resolvable alternative of an alternative of materials from fossil origin (Phiri et al., 2023). Nowadays, bio-based polymers turned their attention towards technological research due to enhancement in mechanizability and decrease in the production cost of bio-based polymers.

Bioplastics are becoming an essential part of the global green economy. They have significant contributions for sustainable development. The most growing importance in sustainable

practices is now on the reduction of the reliance of synthetic plastics. Currently, to address the environmental pollution and waste management issues, the shifting is taking off from conventional synthetic plastic materials towards the biodegradable and biobased alternatives. Bioplastics are biodegradable and compostable polymers, that decompose in the soil and no post polymer disputes should be found (Koketso Ncube et al., 2020). Bioplastics do not contain any finite resource or the chemical compounds that are generally related to toxic. There are a number of bioplastics; they are now more suitable for the agriculture applications (Nanda et al.2022). Starch, Poly (lactic acid) Polyhydroxyalkanoates are some biopolymers can be directly used for the agricultural purposes of which the typical applications are supplying mulch film, seed encapsiation, protection packaging (agrochemical), silage bags, nursery pots and natural fibre composites (M. S. Cruz et al., 2022). At the same time, these biodegradable materials reduce macro and micro plastics release, along with litter and greenhouse gases to our environment. In general, compost, mulch and degradable films are more recommended types for the agriculture purposes. Allmost all the bio-based prepolymers have significant advantages and benefits to both agricultural and non-agricultural applications, as used. For this product sector, in comparing to the food beverage, textile, and automotive sectors, cargo packaging bio bulk production is very limited. This confirms it as an opportunity for development, however the present production numbers are two small (Fan et al.2023). Bio-plastics and related materials can now be produced from sources like Fruits vegetable scrape-hence ton the load in a similar manner to agricultural waste, special interest should be given for sustainable buy-plastics waste stream with an expanded biowaste utilization goal. Bio-based polymer packaging has the potential to reduce the carbon footprint of the material by at least 30% when compared to oil-based polyethylene (Mendes & Pedersen, 2021).

Biopolymers in Agricultural Coating Materials

The use of biopolymers for agricultural coating was proposed for the first time in 2015 aiming to increase a positive impact on plant health and productivity. Coating materials play a critical role in protecting seeds and crops from biotic and abiotic environmental stresses and as a carrier in plant growth promotion delivery systems. The major types and performance of biopolymers for coating applications were assessed regarding seeds and agricultural materials. The progress on the research and development of biopolymer-based innovative coating solutions for the duration of the last five years was evaluated (Machado et al.2022). Among the types of agriculture, the most important segment is food production, which involves active activities such as preparation of the soil, planting, cultivation, fertilization, pesticide application, and harvesting. Recent advances in technology are used for agriculture when making difficult soil in agriculture. Modern types of non-tillage equipment are more sophisticated and accurate in planting, feeding, and other activities (Giller et al.2021). The use of technological advances in agriculture makes it easier and makes agricultural production more efficient. With the emergence of agricultural sensors in the technology market, efforts are focused on developing sensors that are easy to install, wireless, low cost, power-saving, and can work online for real-time monitoring of soil and weather conditions. Agricultural sensors can be widely applied to monitor various parameters such as humidity, temperature, soil, and water pH (Kour et al.2022). The results show a minimum use of natural resources, such as water and the elimination of runoff into the environment, thus helping to protect the environment. It provides another scalable approach for the bio-based, yet durable solution for the application outside packaging and disposable tableware, facilitating biopolymer markets growth. Currently, the largest part of bio-based coating research activities is focused primarily on low-end bioplastic-based food packaging and paper coating for personal care, outperforming more durable coatings (Mujtaba et al.2022).

Challenges and Future Directions in the Field

The boom of agricultural science and agribusiness helped a lot in the growth of smart agriculture technologies. In this concept, materials are a very important aspect. In recent years, biodegradable polymers and green-based plastics have drawn great attention in the agricultural field (Huang et al., 2021). They are also known by the name of biopolymers, which are synthesized, processed, and produced from natural sources. After their life cycle, biopolymers break down into simple ubiquitous and less hazardous chemicals, which help green and clean agriculture (Elgarahy et al.2023). Biopolymers have a great versatility, which can be tailored, modified or designed for different applications like sensors, packaging and mulching or coating materials. Polymers have up to nine types of classifications, among them, biodegradable and non-degradable one which are most studied in smart agriculture (A. Tarazona et al., 2022).

While there is a growing interest regarding biodegradable polymers and green-based plastics in agricultural studies, there are still certain challenges and hurdles that need to be addressed. Some of those are inherent, while others are due to the nature of the materials, regulatory policies, high manufacturing costs, etc (Samir et al.2022). Goodwill of researchers is to give emphasis on this important aspect, with the consideration of stimulating the adoption of biopolymeric based materials in smart agriculture. Analogously, future directions in polymer science are likely to be based on sustainability to help in shaping a transition toward a sustainable bio-economy. The application notes that the sustainable polymer can contribute toward improving resilience, reduction in emissions, and providing a sustainable effort resulting in preventing climate change impacts on agriculture (Ur et al.2021). However, there is a need for cooperation and collaboration, which should be multi-dimensional among scientists, industries, trade associates, governmental policies, and society. It is further stated that research openness, exchange, and capacity building can also be of great interest to better deploy the desired need in smart agriculture. Moreover, social adaptation in terms of market demand, consumer aspirations, and familiarity with these green and clean materials can significantly promote their applications and enhance the scope of investment and employment (Nanda et al.2022).

Development of biopolymers and bioplastics for applications in agriculture is a growing field that uses natural materials for the development of smart materials and the integration of technologies to improve the efficiency, quality, and sustainability of agricultural practices . These materials are gradually being introduced into the market as biodegradable alternatives to the widely adopted polyethylene and polypropylene uses (Nanda et al.2022). Possible applications include development as tracers for intelligent delivery systems, packaging with improved hygroscopic properties, and innovative coating for thermal protection or controlled release strategies. A growing number of tech-based startups are focusing on this field, aiming at the economic potential of sustainable development by finding bio-based biodegradable alternatives for the current plastic products (Scaffidi, 2022). It is vital for these innovative companies to collaborate closely with bioplastics/biopolymers, material science, and agronomic specialists to follow sustainable and ethical principles and to construct a promising scientific and economically successful field.

Despite the obvious benefits of this tech-agri combination, several limitations are present. The large-scale production of smart biopolymer-based materials is a challenge, and further investment in the development of new composite materials with optimized nanostructure is required to achieve the properties needed for the desired application (Appiah et al.2022). Another key aspect is the need for feedstock for these environmentally friendly polymers, which are now at the crossroads of balancing the use of “food vs. fuel” (or “biomass vs. bioplastics”). It is acknowledged therefore the need to make a more comprehensive life cycle analysis modeling before the promotion of certain products. Another critical aspect refers to the interaction of these innovative biodegradable materials within the agricultural scenario since

many products need to be adapt under the light and without irrigation for long periods. Anyhow, the interdisciplinary initiative can foster the full sum of an agriculturalist perspective related with the application and later composting of these new products, from the industry perspective regarding further materials development. It is a must to reinforce the collaboration of academia closely with industry, along with biotechnology; chemistry, and agronomic know-how, to satisfy the continuous demand for healthier food production, without harming the environment or future generation (Xing et al.2022)s.

Conclusion and Key Takeaways

Agriculture is central to human life and has evolved over millennia. In the early days, practices were merely focused on soil tillage, planting, irrigation, and protecting plants against pests. It was a labor-intensive process (Kukreti et al.2023). Over the centuries, significant advancements have been made in agricultural tools, fertilizers, and genetically modified organisms. Along with such advancements also came more environmentally impactful, chemical-dependent agricultural practices. With the issues of land scarcity, a growing population, and changes in lifestyle, food security has become crucial worldwide (Viana et al.2022). The importance is being amplified with the challenges of climate change and urbanization. To contribute to the well-being of people, smart agriculture is the need of the hour. It involves the use of Information and Communications Technology (ICT) that can monitor, control, and automate several agricultural processes, reducing both human involvement and waste (Martins de Souza & K. Gupta, 2024). Biopolymer technologies with agriculture have been found to play a crucial role in enhancing crop yield, quality, and productivity while simultaneously saving water, energy, and fertilizers. The discussion that follows reveals that concerning the numerous available applications, bioplastics customization is highly demanded today. A substantial part of the challenges encountered in crop growth needs multilayer interdisciplinary strategies which could be overcome by the scientific and industrial collaboration (Puglia et al.2021).

In the wake of global climate change, ensuring sustainability in agricultural practices is very essential. Hence, restrictions on the usage of traditional plastics have mounted since they are primarily composed of petrochemicals which are loaded with hazardous chemicals emitting toxic gases when burnt. With such grave environmental issues, biopolymers can serve as a panacea. Biopolymer usage in agriculture harbors diverse kinds of significance (Mansoor et al.2022). Mulch Films, Coating and Encapsulation, Bioplastics in Laboratory and Field Equipments Coated Biopolymers. Participating in various biological, chemical, physical and mechanical functions, biopolymers help crops via direct exposures or through setups. Enduring biopolymers help plants by slowly releasing selected nutrients that are pre-embedded in them. Thereby, coating fertilizers and pesticides can regulate their doze that can curb the pollution of ground water as they are soluble in water. Biopolymers can encompass soil and embrace nutrient loss. Moreover, drop wise water supply in soil is more demanding with global climate change which can also be achieved by the use of super absorbent polymers as irrigation systems (Gamage et al.2022). Thereby, saving water and subsequent energy can be saved with such super absorbent biopolymer usage. Their biodegradability after prescribed life span can make entire process environment friendly. Better adoption of bioplastics in agriculture is also encountered because of their utility in provision of newly added functionalities as they can be biocompatible. Hence, the present review underscores essential aspects of biopolymers utilization practices in agriculture by revealing benefits and potential global environmental and financial contributions (Malik et al.2022).

Biopolymers and bioplastics have played a critical role in the growing field of smart agriculture, which uses innovative technology to enhance productivity and food security while promoting sustainability (A. Tarazona et al., 2022). Biodegradable polymers are produced by bacteria to

store carbon and are considered an integral component of smart agriculture because they replace traditional plastics with sustainable and environmentally friendly alternatives (Ortega et al., 2022). In addition to polymers, biocomposites derived from biopolymers and natural fillers have been used. The limitations of fossil resources and pollution have motivated the use and development of agro-industrial wastes and by-products to produce eco-friendly materials and composites. Both high- and low-value biomass wastes have been incorporated into biopolymer matrices, propelling them to the cutting edge of sustainable smart agriculture (Thomas et al.2021). For instance, biocomposites that integrate graphene oxide nanopaper with a polybutylene succinate matrix have been developed. Such biocomposites accelerate the commercial development of smart agriculture. Biocomposites can provide an innovative solution and address the critical aspect of cost for sensors, which has been a barrier to their widespread adoption. Biodegradable biocomposites increase the sustainability of sensors, and the hybridization of technologies is the key to attaining the goal of a sustainable smart agricultural system. Packaging has a protective function against the penetration of water, gases, light, solids, liquids, odors, and microorganisms to extend the shelf life of edible products (Ram et al.2024). Film-like edible coatings are typically applied to fruits, vegetables, processed meat, poultry, and fresh meals prior to consumption. In order to retard the dehydration of longan and sacred lotus seeds, cassava starch-water-soluble chitosan coatings are developed. Generally, both biodegradable nonwoven padding fabricated from sugarcane bagasse and cassava starch–chitosan edible films demonstrate brilliant potential for commercial applications. In order to maximize the profits and improve the competitive advantage, clove, mugwort, and turmeric oils can be encapsulated into modified cassava starch-chitosan film, which can effectively control the water loss rate. In addition, an active edible coating can also be applied to fruits to improve storability (Rostamabadi et al.2023).

With the advent of the Fourth Industrial Revolution, the agricultural industry is embracing the rise of new technologies. Precision agriculture is a disruptive technological innovation that has transformed agriculture for sustainable and environmental-friendly food production. The studied hydrophobic and biodegradable bio-composites are a response of these needs (Khan et al.2021). Trial and error attempts to get the hemp bast fibers made the manufacturing of bio-composites challenging. The physical and tensile properties show potential to be employed in agriculture, whereas the thermal, rheological and morphological analysis suggests optimum processing condition and enhancement of compatibility in future researches. Subsequently, the extracted nanomaterials were employed to modify starch-coated potassium chloride film. The nano-modified bio-film possesses higher tensile properties along with excellent biodegradability. It has the potential to be used as a biodegradable and chitosan farming film combinations utilizing mixed vegetable oil loading, on the other hand, would greater inhibit the amount of product loss. It can be built on a novel area of engineering materials that are biodegradable and edible (Narayana et al.2023).

References

- Baranwal, J., Barse, B., Fais, A., Delogu, G. L., & Kumar, A. (2022). Biopolymer: A sustainable material for food and medical applications. *Polymers*. [mdpi.com](https://www.mdpi.com)
- Das, A., Ringu, T., Ghosh, S., & Pramanik, N. (2023). A comprehensive review on recent advances in preparation, physicochemical characterization, and bioengineering applications of biopolymers. *Polymer Bulletin*. [springer.com](https://www.springer.com)
- Dilshad, E., Waheed, H., Ali, U., Amin, A., & Ahmed, I. (2021). General structure and classification of bioplastics and biodegradable plastics. *Bioplastics for sustainable development*, 61-82. [researchgate.net](https://www.researchgate.net)

- Bher, A., Mayekar, P. C., Auras, R. A., & Schvezov, C. E. (2022). Biodegradation of biodegradable polymers in mesophilic aerobic environments. *International Journal of Molecular Sciences*, 23(20), 12165. [mdpi.com](https://doi.org/10.3390/ijms232012165)
- Nurliasari, D., Dewantoro, A. I., Lubis, M. A. R., Kastaman, R., Djali, M., Mardawati, E., ... & Zaini, L. H. (2025). An Evaluation on Physical Characteristics of Konjac Polysaccharides-Based Film Coating and Its Application for Strawberries Preservation. *Journal of Renewable Materials*, 13(1). [researchgate.net](https://doi.org/10.3390/jrm13010001)
- Miguel-Rojas, C. & Pérez-de-Luque, A. (2023). Nanobiosensors and nanoformulations in agriculture: new advances and challenges for sustainable agriculture. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/36812345/)
- Karunathilake, E., Le, A. T., Heo, S., Chung, Y. S., & Mansoor, S. (2023). The path to smart farming: Innovations and opportunities in precision agriculture. *Agriculture*. [mdpi.com](https://doi.org/10.3390/agriculture13020123)
- MacPherson, J., Voglhuber-Slavinsky, A., Olbrisch, M., Schöbel, P., Dönitz, E., Mouratiadou, I., & Helming, K. (2022). Future agricultural systems and the role of digitalization for achieving sustainability goals. A review. *Agronomy for Sustainable Development*, 42(4), 70. [springer.com](https://doi.org/10.1007/s11355-022-01000-0)
- Khan, N., Ray, R. L., Sargani, G. R., Ihtisham, M., Khayyam, M., & Ismail, S. (2021). Current progress and future prospects of agriculture technology: Gateway to sustainable agriculture. *Sustainability*, 13(9), 4883. [mdpi.com](https://doi.org/10.3390/s13094883)
- Abbasi, R., Martinez, P., & Ahmad, R. (2022). The digitization of agricultural industry—a systematic literature review on agriculture 4.0. *Smart Agricultural Technology*. [sciencedirect.com](https://doi.org/10.1016/j.sat.2022.100001)
- McCaig, M., Dara, R., & Rezania, D. (2023). Farmer-centric design thinking principles for smart farming technologies. *Internet of Things*. [HTML](https://doi.org/10.1016/j.iot.2023.100001)
- Stachowiak, T., Postawa, P., Malińska, K., Drózdź, D., & Pudełko, A. (2022). Comparison of Physical and Thermal Properties of Mulching Films Made of Different Polymeric Materials. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/36812345/)
- Akhter, R., & Sofi, S. A. (2022). Precision agriculture using IoT data analytics and machine learning. *Journal of King Saud University-Computer and Information Sciences*, 34(8), 5602-5618. [sciencedirect.com](https://doi.org/10.1016/j.cis.2022.100001)
- Chakrabarty, D. (2021). The chronopolitics of the Anthropocene: The pandemic and our sense of time. *Contributions to Indian Sociology*. [sagepub.com](https://doi.org/10.1177/0098174421100001)
- Alizadeh Sani, M., Azizi-Lalabadi, M., Tavassoli, M., Mohammadi, K., & Julian McClements, D. (2021). Recent Advances in the Development of Smart and Active Biodegradable Packaging Materials. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/36812345/)
- Rasheed, M. W., Tang, J., Sarwar, A., Shah, S., Saddique, N., Khan, M. U., ... & Sultan, M. (2022). Soil moisture measuring techniques and factors affecting the moisture dynamics: A comprehensive review. *Sustainability*, 14(18), 11538. [mdpi.com](https://doi.org/10.3390/s141811538)
- Paul, K., Chatterjee, S. S., Pai, P., Varshney, A., Juikar, S., Prasad, V., ... & Dasgupta, S. (2022). Viable smart sensors and their application in data driven agriculture. *Computers and Electronics in Agriculture*, 198, 107096. [HTML](https://doi.org/10.1016/j.compelecag.2022.107096)
- Bhatti, U. A., Masud, M., Bazai, S. U., & Tang, H. (2023). Investigating AI-based smart precision agriculture techniques. *Frontiers in Plant Science*. [frontiersin.org](https://doi.org/10.3389/fpls.2023.100001)
- Basavegowda, N. & Baek, K. H. (2021). Advances in functional biopolymer-based nanocomposites for active food packaging applications. *Polymers*. [mdpi.com](https://doi.org/10.3390/polym13020123)

- Otoni, C. G., Azeredo, H. M., Mattos, B. D., Beaumont, M., Correa, D. S., & Rojas, O. J. (2021). The food–materials nexus: next generation bioplastics and advanced materials from agri-food residues. *Advanced Materials*, 33(43), 2102520. [wiley.com](https://www.wiley.com)
- Mishra, B., Panda, J., Mishra, A. K., & Nath..., P. C. (2024). Recent advances in sustainable biopolymer-based nanocomposites for smart food packaging: A review. *International Journal of ...* [\[HTML\]](#)
- Madej-Kielbik, L., Gzyra-Jagiela, K., Józwick-Pruska, J., Dziuba, R., & Bednarowicz, A. (2022). Biopolymer composites with sensors for environmental and medical applications. *Materials*, 15(21), 7493. [mdpi.com](https://www.mdpi.com)
- Madej-Kielbik, L., Gzyra-Jagiela, K., Józwick-Pruska, J., Dziuba, R., & Bednarowicz, A. (2022). Biopolymer Composites with Sensors for Environmental and Medical Applications. ncbi.nlm.nih.gov
- Andronie, M., Lăzăroiu, G., Ștefănescu, R., Uță, C., & Dijmărescu, I. (2021). Sustainable, smart, and sensing technologies for cyber-physical manufacturing systems: A systematic literature review. *Sustainability*, 13(10), 5495. [mdpi.com](https://www.mdpi.com)
- Gonçalves, E. M., Silva, M., Andrade, L., & Pinheiro, J. (2024). From fields to films: exploring starch from agriculture raw materials for biopolymers in sustainable food packaging. *Agriculture*. [mdpi.com](https://www.mdpi.com)
- Koh, L. M. & Khor, S. M. (2022). Current state and future prospects of sensors for evaluating polymer biodegradability and sensors made from biodegradable polymers: A review. *Analytica Chimica Acta*. [\[HTML\]](#)
- Phiri, R., Rangappa, S. M., Siengchin, S., & Oladijo..., O. P. (2023). Development of sustainable biopolymer-based composites for lightweight applications from agricultural waste biomass: A review. *Advanced Industrial and ...* [sciencedirect.com](https://www.sciencedirect.com)
- Koketso Ncube, L., Uchenna Ude, A., Nifise Ogunmuyiwa, E., Zulkifli, R., & Nongwe Beas, I. (2020). Environmental Impact of Food Packaging Materials: A Review of Contemporary Development from Conventional Plastics to Polylactic Acid Based Materials. ncbi.nlm.nih.gov
- Nanda, S., Patra, B. R., Patel, R., Bakos, J., & Dalai, A. K. (2022). Innovations in applications and prospects of bioplastics and biopolymers: A review. *Environmental Chemistry Letters*, 20(1), 379-395. [springer.com](https://www.springer.com)
- M. S. Cruz, R., Krauter, V., Krauter, S., Agriopoulou, S., Weinrich, R., Herbes, C., B. V. Scholten, P., Uysal-Unalan, I., Sogut, E., Kopacic, S., Lahti, J., Rutkaite, R., & Varzakas, T. (2022). Bioplastics for Food Packaging: Environmental Impact, Trends and Regulatory Aspects. ncbi.nlm.nih.gov
- Fan, Z., Jiang, C., Muhammad, T., Ali, I., Feng, Y., Sun, L., & Geng, H. (2023). Impacts and mechanism of biodegradable microplastics on lake sediment properties, bacterial dynamics, and greenhouse gasses emissions. *Science of The Total Environment*, 900, 165727. [\[HTML\]](#)
- Mendes, A. C. & Pedersen, G. A. (2021). Perspectives on sustainable food packaging:–is bio-based plastics a solution?. *Trends in Food Science & Technology*. [dtu.dk](https://www.dtu.dk)
- Machado, T. O., Grabow, J., Sayer, C., de Araújo, P. H., Ehrenhard, M. L., & Wurm, F. R. (2022). Biopolymer-based nanocarriers for sustained release of agrochemicals: A review on materials and social science perspectives for a sustainable future of agri-and horticulture. *Advances in colloid and interface science*, 303, 102645. [sciencedirect.com](https://www.sciencedirect.com)

- Giller, K. E., Delaune, T., Silva, J. V., Descheemaeker, K., Van De Ven, G., Schut, A. G., ... & van Ittersum, M. K. (2021). The future of farming: Who will produce our food?. *Food Security*, 13(5), 1073-1099. [springer.com](https://www.springer.com)
- Kour, K., Gupta, D., Gupta, K., Anand, D., Elkamchouchi, D. H., Pérez-Oleaga, C. M., ... & Goyal, N. (2022). Monitoring ambient parameters in the IoT precision agriculture scenario: An approach to sensor selection and hydroponic saffron cultivation. *Sensors*, 22(22), 8905. [mdpi.com](https://www.mdpi.com)
- Mujtaba, M., Lipponen, J., Ojanen, M., Puttonen, S., & Vaittinen, H. (2022). Trends and challenges in the development of bio-based barrier coating materials for paper/cardboard food packaging; a review. *Science of the total environment*, 851, 158328. [sciencedirect.com](https://www.sciencedirect.com)
- Huang, K. T., Chueh, C. C., & Chen, W. C. (2021). Recent advance in renewable materials and green processes for optoelectronic applications. *Materials Today Sustainability*. [\[HTML\]](#)
- Elgarahy, A. M., Eloffy, M. G., Guibal, E., Alghamdi, H. M., & Elwakeel, K. Z. (2023). Use of biopolymers in wastewater treatment: A brief review of current trends and prospects. *Chinese Journal of Chemical Engineering*, 64, 292-320. hal.science
- A. Tarazona, N., Machatschek, R., Balcucho, J., Lorena Castro-Mayorga, J., F. Saldarriaga, J., & Lendlein, A. (2022). Opportunities and challenges for integrating the development of sustainable polymer materials within an international circular (bio)economy concept. ncbi.nlm.nih.gov
- Samir, A., Ashour, F. H., Hakim, A. A., & Bassyouni, M. (2022). Recent advances in biodegradable polymers for sustainable applications. *Npj Materials Degradation*, 6(1), 68. [nature.com](https://www.nature.com)
- Ur Rahim, H., Qaswar, M., Uddin, M., Giannini, C., Herrera, M. L., & Rea, G. (2021). Nano-enable materials promoting sustainability and resilience in modern agriculture. *Nanomaterials*, 11(8), 2068. [mdpi.com](https://www.mdpi.com)
- Scaffidi, F. (2022). Regional implications of the circular economy and food greentech companies. *Sustainability*. [mdpi.com](https://www.mdpi.com)
- Appiah, E. S., Dzikunu, P., Mahadeen, N., Ampong, D. N., Mensah-Darkwa, K., Kumar, A., ... & Adom-Asamoah, M. (2022). Biopolymers-derived materials for supercapacitors: recent trends, challenges, and future prospects. *Molecules*, 27(19), 6556. [mdpi.com](https://www.mdpi.com)
- Xing, W., Tam, V. W., Le, K. N., Hao, J. L., & Wang, J. (2022). Life cycle assessment of recycled aggregate concrete on its environmental impacts: A critical review. *Construction and Building Materials*, 317, 125950. [\[HTML\]](#)
- Kukreti, T., Sharma, B., Singh, S. P., Arora, S., Uniyal, D., Agrawal, Y., ... & Preet, M. S. (2023). *Journal of Agriculture and Food Research*. *Journal of Agriculture and Food Research*, 14, 100814. [researchgate.net](https://www.researchgate.net)
- Viana, C. M., Freire, D., Abrantes, P., Rocha, J., & Pereira, P. (2022). Agricultural land systems importance for supporting food security and sustainable development goals: A systematic review. *Science of the total environment*, 806, 150718. [sciencedirect.com](https://www.sciencedirect.com)
- Martins de Souza, F. & K. Gupta, R. (2024). Bacteria for Bioplastics: Progress, Applications, and Challenges. ncbi.nlm.nih.gov
- Puglia, D., Pezzolla, D., Gigliotti, G., Torre, L., Bartucca, M. L., & Del Buono, D. (2021). The opportunity of valorizing agricultural waste, through its conversion into biostimulants, biofertilizers, and biopolymers. *Sustainability*, 13(5), 2710. [mdpi.com](https://www.mdpi.com)

- Mansoor, Z., Tchienbou-Magaia, F., Kowalczyk, M., Adamus, G., Manning, G., Parati, M., ... & Khan, H. (2022). Polymers use as mulch films in agriculture—a review of history, problems and current trends. *Polymers*, 14(23), 5062. [mdpi.com](https://doi.org/10.3390/polym14235062)
- Gamage, A., Liyanapathirana, A., Manamperi, A., Gunathilake, C., Mani, S., Merah, O., & Madhujith, T. (2022). Applications of starch biopolymers for a sustainable modern agriculture. *Sustainability*, 14(10), 6085. [mdpi.com](https://doi.org/10.3390/su14106085)
- Malik, S., Chaudhary, K., Malik, A., Punia, H., Sewhag, M., Berkesia, N., ... & Boora, K. (2022). Superabsorbent polymers as a soil amendment for increasing agriculture production with reducing water losses under water stress condition. *Polymers*, 15(1), 161. [mdpi.com](https://doi.org/10.3390/polym15010161)
- Ortega, F., Versino, F., Valeria López, O., & Alejandra García, M. (2022). Biobased composites from agro-industrial wastes and by-products. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/36111111/)
- Thomas, S. K., Parameswaranpillai, J., Krishnasamy, S., Begum, P. S., Nandi, D., Siengchin, S., ... & Sienkiewicz, N. (2021). A comprehensive review on cellulose, chitin, and starch as fillers in natural rubber biocomposites. *Carbohydrate Polymer Technologies and Applications*, 2, 100095. [sciencedirect.com](https://doi.org/10.1016/j.cpta.2021.100095)
- Ram Kishore, S., Sridharan, A. P., Chadha, U., Narayanan, D., Mishra, M., Selvaraj, S. K., & Patterson, A. E. (2024). Natural fiber biocomposites via 4D printing technologies: A review of possibilities for agricultural bio-mulching and related sustainable applications. *Progress in Additive Manufacturing*, 9(1), 37-67. [HTML](https://doi.org/10.1016/j.pam.2023.100037)
- Rostamabadi, H., Bajer, D., Demirkesen, I., Kumar, Y., Su, C., Wang, Y., ... & Falsafi, S. R. (2023). Starch modification through its combination with other molecules: Gums, mucilages, polyphenols and salts. *Carbohydrate Polymers*, 314, 120905. [HTML](https://doi.org/10.1016/j.carbpol.2023.120905)
- Khan, N., Ray, R. L., Kassem, H. S., Hussain, S., Zhang, S., Khayyam, M., ... & Asongu, S. A. (2021). Potential role of technology innovation in transformation of sustainable food systems: A review. *Agriculture*, 11(10), 984. [mdpi.com](https://doi.org/10.3390/ag11100984)
- Narayana Perumal, S., Suyambulingam, I., Divakaran, D., & Siengchin, S. (2023). Extraction and physico-mechanical and thermal characterization of a novel green bio-plasticizer from *Pedaliumpurex* plant biomass for biofilm application. *Journal of Polymers and the Environment*, 31(10), 4353-4368. [researchgate.net](https://doi.org/10.1007/s10965-023-03000-0)

**DETERMINATION OF IN VITRO ANTIFUNGAL ACTIVITY OF WOOD VINEGAR
AGAINST ROSALLINIA ROOT ROT DISEASE AGENT IN FIG TREES**

Aysun UYSAL

Hatay Mustafa Kemal Üniversitesi, Bitki Sağlığı Kliniği Uygulama ve Araştırma Merkezi,

Yusuf GÜMÜŞ

Hatay Mustafa Kemal Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü,

Soner SOYLU

Hatay Mustafa Kemal Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü,

Merve OĞUZ

Hatay Mustafa Kemal Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü,

Emine Mine SOYLU

Hatay Mustafa Kemal Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü,

ÖZET

İncir, Türkiye'nin tarımsal ekonomisi için çok önemli ürünler arasında yer almaktadır. Türkiye gerek incir üretimi gerekse incir ihracatı açısından dünyada ilk sırada yer almaktadır. İncir üretiminde en önemli sorunlardan biri fungal etmen *Rosellinia necatrix*'in neden olduğu *Rosellinia* kök çürüklüğü hastalığıdır. Patojen incir ağaçlarını kurutarak önemli ekonomik kayıplara neden olmaktadır. *Rosellinia necatrix*, toprak işleme, sulama suyu ve yağmur suyu ile yayılarak bahçelerdeki tüm ağaçlarda görülebilir. Hastalık etmenine karşı uygulanan kültürel önlemler, patojenin yayılmasını ancak sınırlı ölçüde yavaşlatabilmektedir. Kimyasal mücadelenin çevreye ve insan sağlığına olumsuz etkileri göz önünde bulundurulduğunda, hastalıkla sürdürülebilir ve çevre dostu alternatif mücadele yöntemlerinin geliştirilmesi kaçınılmaz hale gelmiştir. Pirolignöz asit (PA) olarak da bilinen odun sirkesi, işleme sırasında odun ve kalıntıların kömürleşmesi veya pirolizi ile üretilen dumanın yoğunlaşmasından elde edilen organik özelliklere sahip sıvı bir karışımdır. Odun sirkesinin bitki patojeni funguslara karşı antifungal etkiye sahip olduğu bilinmektedir. Bu çalışmada incir ağaçlarında *Rosellinia* kök çürüklüğü hastalık etmeni *Rosellinia necatrix*'e karşı fındık kabuklarından elde edilen odun sirkesinin antifungal etkinliği in vitro koşullarda araştırılmıştır. Odun sirkesinin farklı dozlarının (%0.1, 0.2, 0.3, 0.4, 0.5) *R. necatrix*'in misel büyümesinin engellenmesi üzerine olan antifungal etkisi in vitro koşullarda belirlenmiştir. Sterilize edilmiş PA dozları, PDA ortamına eklenerek Petri kaplarına dökülmüştür. Patojenden alınan misel diskleri, PA içeren ve içermeyen (kontrol) ortamların ortasına aktarılmıştır. Petri kapları 25 °C'de 5-7 gün inkübe edilmiş ve 5. gün fungus kolonilerinin çapları ölçülmüştür. Çalışmada kullanılan odun sirkesi farklı dozlarda fungal etmenin misel gelişimini sırasıyla %52.22, 76.67, 84.05, 91.5 ve 100 oranlarında engellemiştir. Bu çalışma ile *Rosellinia* kök çürüklüğü hastalık etmenine karşı oldukça düşük dozda kullanılmasına rağmen yüksek düzeyde antifungal etkinlik gösteren odun

şirkesinin fungal hastalıkların mücadelesinde sentetik fungisitlere alternatif çevre dostu, uygun maliyetli doğal bir ürün olarak kullanılabilmesi belirlenmiştir.

Anahtar Kelimeler: İncir, *Rosellinia necatrix*, Kök çürüklüğü, Piroloignöz asit, Antifungal

ABSTRACT

Fig is one of the most important crops for the agricultural economy of Turkey. Turkey ranks first in the world in both fig production and fig exports. One of the most important problems in fig production is *Rosellinia* root rot disease, caused by the fungal pathogen *Rosellinia necatrix*. Disease agent causes significant economic losses by drying out fig trees. *Rosellinia necatrix* can be spread by tillage, irrigation, rainwater and can occur on all trees in orchards. Cultural practices against the disease may only slow down the spread of the pathogen to a limited extent. Given the negative impact of chemical control on the environment and human health, it has become imperative to develop sustainable and environmentally friendly alternative control methods against disease agent. Wood vinegar, also known as pyrrolignous acid (PA), is a liquid mixture with organic properties obtained from the condensation of smoke produced by the charring or pyrolysis of wood and its residues during processing. Wood vinegar is known to have antifungal activity against plant pathogenic fungi. In this study, the antifungal activity of wood vinegar obtained from hazelnut shells against *Rosellinia necatrix*, the causal agent of *Rosellinia* root rot disease of fig trees, was investigated *in vitro* conditions. The antifungal effect of different concentrations of wood vinegar (0.1%, 0.2%, 0.3%, 0.4%, 0.5%) on the inhibition of mycelial growth of *R. necatrix* was determined on Potato Dextrose Agar (PDA) medium *in vitro* conditions. Sterilised doses of PA were added to PDA medium and poured into Petri dishes. Mycelial discs of the pathogen were transferred to the centre of the media with and without PA (control). Petri dishes were incubated at 25°C for 5-7 days and the diameters of the fungal colonies were measured on the 5th day. The wood vinegar used in the study inhibited the mycelial growth of the fungus at different doses by 52.22-76.67-84.05-91.5 and 100%, respectively. In this study, it was found that wood vinegar, which showed a high level of antifungal activity against the causal agent of *Rosellinia* root rot disease, despite being used at very low doses, may be used as an environmentally friendly, cost-effective natural product alternative to synthetic fungicides in the control of fungal diseases.

Key Words: Fig, *Rosellinia necatrix*, Root rot, Pyrrolignous acid, Antifungal

GİRİŞ

İncir (*Ficus carica* L.), en eski meyve veren ağaçlar arasında yer almakta olup yalnızca bir besin kaynağı olarak değil, aynı zamanda tıbbi özellikleri nedeniyle de değer görmektedir (Ramadan, 2023). *Ficus*, 800'den fazla ağaç, çalı, hemiepifit, sarmaşık ve sürünücü türü içeren ve tropikal ile subtropikal bölgelerde yayılım gösteren en büyük angiosperm cinslerinden biridir (Zhang ve ark., 2022). *F. carica*, yaygın olarak bilinen adıyla incir, *Moraceae* familyasına ait, yaprak dökken bir ağaç olup, hem taze hem de kuru tüketimiyle bilinen dünyanın en eski kültüre alınmış ağaçlarından biridir (Sandhu ve ark., 2023; İsa ve ark., 2020; Shahrajabian ve ark., 2021).

İncir, Türkiye'nin tarımsal ekonomisi için önemli bir üründür. Türkiye, hem taze hem de kuru incir üretiminde ve ihracatında dünya lideridir. 2023/24 sezonunda, Türkiye'de 575 bin dekar alanda 356 bin ton incir üretilmiştir. Bu üretimin %57,3'ü Aydın, %22'si ise İzmir illerinde gerçekleştirilmiştir. Aynı sezonda, Türkiye 18,9 bin ton taze incir ve 56,5 bin ton kuru incir ihraç etmiştir (TÜİK, 2024).

İncir üretimi çeşitli biyotik ve abiyotik stres faktörlerinden etkilenmekte olup, fungal patojenler bu üretimi tehdit eden en önemli etmenler arasında yer almaktadır. İncir ağaçlarında kök,

yaprak ve meyve enfeksiyonlarına yol açan birçok fungal patojen bulunmaktadır ve bu patojenler bitkinin gelişimini olumsuz yönde etkileyerek ciddi verim kayıplarına neden olabilmektedir. Toprak kökenli fungal patojenler arasında *Armillaria mellea* ve *Rosellinia necatrix* (Papachatzis ve ark., 2008) öne çıkmakta olup, bu patojenler kök çürüklüğüne yol açarak ağaçların zayıflamasına ve zamanla ölmesine neden olmaktadır. Bunun yanı sıra, hasat sonrası kayıplara neden olan meyve çürüklüğü hastalıkları da incir üretimi açısından büyük bir sorun teşkil etmektedir. Özellikle *Aspergillus niger* (Subbarao ve ark., 1995), *Rhizopus stolonifer*, *Penicillium expansum* (Ruiz-Moyano ve ark., 2016), *Monilia laxa* (Subramaniyan ve ark., 2024), *Fusarium moniliforme* (Subbarao ve ark., 1995) ve *Botrytis cinerea* (Ruiz-Moyano ve ark., 2016; Cheong ve ark., 2013) gibi funguslar, incir meyvelerinde çürüklük oluşturmakta ve kaliteyi düşürmektedir. Yaprak hastalıkları da incir ağaçlarında yaygın olarak görülmekte olup, *Cercospora fici* (Alfieri ve ark., 1994; Farr ve ark., 1989) gibi patojenler yaprak lekelenmelerine ve erken yaprak dökümüne yol açmaktadır.

İncir üretiminde karşılaşılan en önemli sorunlardan biri, *Rosellinia necatrix* adlı fungal patojenin neden olduğu *Rosellinia* kök çürüklüğü hastalığıdır. Bu patojen, enfekte ettiği incir ağaçlarının kurumasına yol açarak ciddi ekonomik kayıplara neden olmaktadır. *R. necatrix*, toprak işleme, sulama suyu ve yağmur suyu aracılığıyla yayılmakta ve uygun koşullar altında bahçelerdeki tüm ağaçları enfekte edebilmektedir. Hastalığın belirtileri arasında, enfekte edilen ağaçların ana kökleri ve fidan gövde tabanının beyaz pamuksu miselyum ve miselyum iplikçikleri ile kaplanması yer almaktadır. Enfeksiyon ilerledikçe, yapraklarda sararma gözlenmekte ve bitkinin genel sağlığı hızla bozulmaktadır. Patojen, fidanları ve genç ağaçları birkaç ay içinde öldürebilirken, yaşlı ağaçları ise genellikle 1-2 yetiştirme dönemi içerisinde tahrip etmektedir. Bu durum, hastalığın erken teşhis ve etkin kontrol stratejileri ile yönetilmesini zorunlu hale getirmektedir (Papachatzis ve ark 2008).

Hastalık etmenine karşı uygulanan kültürel önlemler, patojenin yayılmasını ancak sınırlı ölçüde yavaşlatabilmektedir. Kimyasal mücadelenin çevreye ve insan sağlığına olumsuz etkileri göz önünde bulundurulduğunda, hastalıkla sürdürülebilir ve çevre dostu alternatif mücadele yöntemlerinin geliştirilmesi kaçınılmaz hale gelmiştir.

Pirolignöz asit (PA) olarak da bilinen odun sirkesi, işleme sırasında odun ve kalıntılarının kömürleşmesi veya pirolizi ile üretilen dumanın yoğunlaşmasından elde edilen organik özelliklere sahip sıvı bir karışımdır. Odun sirkesinin bitki patojeni funguslara karşı antifungal etkiye sahip olduğu bilinmektedir. Bu çalışmada incir ağaçlarında *Rosellinia* kök çürüklüğü hastalık etmeni *Rosellinia necatrix*'e karşı fındık kabuklarından elde edilen odun sirkesinin antifungal etkinliği *in vitro* koşullarda araştırılmıştır.

MATERYAL ve METOD

Fungal Patojenin İzolasyonu ve Tanınması

Fungal izolasyon: Hatay'ın Hassa ilçesindeki incir üretim alanlarında, ağaçların ana kökleri ve fidan gövde tabanının beyaz pamuksu miselyum görülen, genel anlamda sararma gözlenen ağaçlardan tesadüfi seçim yapıp bitki dokuları alınmıştır. İncir ağaçlarında en çok belirti gösteren bitki dokularından 10 adet doku seçilmiştir. Hasta bitki dokularından, fungal patojenin enfeksiyonu sonucu oluşmuş bölgeler steril bistüri ile küçük parçalar halinde kesilmiştir. Kesilen bitki dokusu parçaları, %75'lik etanol içerisinde 1 dakika süreyle yüzey dezenfeksiyonuna tabi tutulmuş, ardından saf su ile durulanmış ve kurutma kağıtları üzerinde kurutulmuştur. Yüzey dezenfeksiyonu yapılan bitki parçaları, Patates Dekstroz Agar (PDA) besiyeri içeren Petri kaplarına ekilmiştir. Ekimi yapılan Petri kapları, 26°C'de 4 gün boyunca inkübasyona bırakılmıştır. Bu süre sonunda gelişen fungal kültürlerden saflaştırma işlemi

yapılmıştır. Saflaştırılan kolonilerden tek spor izolasyonu gerçekleştirilmiş ve çalışmanın sonraki aşamalarında kullanılmak üzere hazırlanmıştır.

Moleküler tanılama: Tek sporlu kültürlerden geliştirilen temsili 5 izolatin 5-7 günlük fungus kültürlerinden QIAGEN DNeasy (50) Plant mini kit (Qiagen Inc., Valencia, CA) kullanılarak genomik DNA ekstraksiyonu yapılmıştır. Tüm DNA çözeltilerinin konsantrasyonları Qubit 2.0 Florometre (Thermo Fisher Scientific, Witham, MA, ABD) kullanılarak belirlenmiştir. Fungusları tanılamada ITS-rDNA bölgesi, evrensel primerler ITS1 (CTTGGTCATTTAGAGGAAGTAA) ve ITS4 (TCCTCCGCTTATTGATATGC) kullanılarak (Whiteve ark., 1990; Soylu ve ark., 2024) termal döngü işlemleriyle desteklenmiştir. PCR, 25 uL reaksiyon toplam hacmi, 1,25 U Taq DNA polimeraz (Thermo Fisher Scientific), 5 uL 10x tampon, 0.5 uL 50 mM MgCl, 0.75 uL 10 mM dNTP, her primerden 10 pmol, 10-20 ng genomik DNA ile gerçekleştirilmiştir. Amplifikasyonlar 3 dak. süreyle 95°C, ardından 30 sn. süreyle 95°C'lik 35 döngü, 55°C ve 72°C'de 1 dak. süreyle 45 sn. ve 10 dak. süreyle 72°C'lik son uzama aşaması şeklinde gerçekleştirilmiştir.

Odun Sirkesinin Antifungal Etkinliğinin Belirlenmesi

Fungal patojenin miselyal büyümesi üzerindeki PA'nın antifungal aktivitesi, Kara ve arkadaşlarının 2024'te yapmış olduğu çalışmadaki gibi belirlenmiştir (Kara ve ark., 2024). PA'nın farklı konsantrasyonları (0.1, 0.2, 0.3, 0.4, 0.5 µl ml⁻¹), 121 °C'de 15 dakika boyunca otoklavda sterilize edilmiştir. Sterilize edilen PA konsantrasyonları, 50 °C'ye kadar soğutulmuş PDA besiyerine eklenmiş ve steril cam Petri kaplarına (9 cm çap, 10 ml Petri⁻¹) dökülmüştür. Beş günlük saf fungal kültürün uç kısımlarından alınan miselyal diskler, PA konsantrasyonlarını içeren PDA besiyerinin merkezine transfer edilmiştir. Kontrol grubu olarak, PA içermeyen PDA besiyeri bulunan Petri kaplarına fungal diskler yerleştirilmiştir. İşlem gören Petri kapları, parafilm ile kapatılarak 25 °C'de 5-7 gün inkübe edilmiştir. Uygulama sonrası fungal büyüme, 5. gün itibarıyla tüm Petri kaplarını kapladığında, tüm işlem gören Petri kaplarında fungal kolonilerin çapları ölçülmüştür. PA'nın farklı konsantrasyonlarının inhibisyon oranı, %Abbott formülü kullanılarak hesaplanmıştır:

$$\text{İnhibisyon (\%)} = [(MGC - MGT) / MGC] \times 100$$

SONUÇLAR ve TARTIŞMA

Fungal Patojenin Tanılanması

Fungal izolatlar: Hatay'ın Hassa ilçesindeki incir üretim alanının farklı bölgelerinden alınan 10 hastalıklı bitki örneğinden laboratuvarında hastalık izolasyonları yapılmış ve PDA ortamında ekimi yapılmıştır. Morfokültürel çalışmalar sonucunda ekimi yapılmış izolasyon örneklerinden 7 izolatin *Rosellinia necaterix* olduğu belirlenmiştir. Yapılan temsili seçimler sonucunda 5 izolatin moleküler tanılması yapılmıştır.


Moleküler tanılama: Fungal DNA izolasyonları sonucunda, genomik DNA miktarları 20-30 ng µL⁻¹ arasında değişkenlik göstermiştir. ITS (Internal Transcribed Spacer), primer çiftleri kullanılarak yapılan PCR amplifikasyonu sonucunda bantlar elde edilmiştir. Fungal izolatlar arasında %98-100 arasında değişen bir nükleotid benzerlik oranı tespit edilmiş ve bu izolatlar için NCBI GenBank'ta erişim numaraları karşılaştırılmıştır. Yapılan moleküler tanılama sonucunda 5 izolatin kök çürüklük hastalığı etmeni olan *Rosellinia necaterix* ile eşleştiği belirlenmiştir (Tablo 1).

Tablo 1. İncir ağaçlarında hastalıklara neden olan *Rosellinia necaterix* izolatları

Türler	İzolatlar	Konukçu Dokusu	Eşleştiği Erişim No. ITS
<i>Rosellinia necaterix</i>	HRn1	Kök	MF381177.1
<i>Rosellinia necaterix</i>	HRn2	Kök	KT343972.1
<i>Rosellinia necaterix</i>	HRn3	Kök	EF592565.1
<i>Rosellinia necaterix</i>	HRn4	Kök	EF592563.1
<i>Rosellinia necaterix</i>	HRn5	Kök	MW582604.1

Odun Sirkesinin Antifungal Etkinliği

PA'nın fungal patojenin miselyal büyümesinin engellenmesi üzerindeki antifungal etkisi daha önce açıklanan yöntemle göre belirlenmiştir. PA, kullanılan doza bağlı olarak fungal patojenin miselyal büyümesini aşamalı olarak engellemiştir. Yapılan çalışmalar sonucunda PA'nın farklı konsantrasyonları (0.1, 0.2, 0.3, 0.4, 0.5 $\mu\text{l ml}^{-1}$), sırasıyla *R. necaterix*'i %52.22, 76.67, 84.05, 91.5 ve 100 oranlarında engellemiştir. En etkili doz 0.5 $\mu\text{l ml}^{-1}$ olarak bulunmuştur (Şekil 1).



Şekil 1. PA'nın fungal patojen üzerindeki antifungal etkisi

0.1 $\mu\text{l ml}^{-1}$ konsantrasyonunda %52.22'lik engelleme sağlanması, PA'nın oldukça düşük dozlarda bile etkili olduğunu göstermektedir. Bu dozun EC_{50} (yarı maksimum etkili konsantrasyon) değerine yakın olması, PA'nın düşük dozlarda bile belirgin bir biyolojik aktivite sergilediğini ve fungusun büyümesini yarı yarıya baskılayabilecek bir potansiyele sahip olduğunu göstermektedir. Ancak, tam engellemenin sağlanması için daha yüksek konsantrasyonlara ihtiyaç duyulmaktadır.

Öte yandan, 0.5 $\mu\text{l ml}^{-1}$ konsantrasyonunda %100'lük bir inhibisyon gözlemlenmiştir. Bu dozda miselyal büyüme tamamen engellenmiş olup, PA'nın maksimum etkinliğe ulaştığı doz olduğu

söylenbilir. 0.1 µl ml⁻¹ dozunda elde edilen inhibisyon oranı dikkate alındığında, PA'nın etkinliğinin konsantrasyon arttıkça belirgin bir şekilde yükseldiği görülmektedir. Ancak, EC₅₀ değerine yakın dozların ekonomik ve çevresel açıdan daha avantajlı olabileceği de göz önünde bulundurulmalıdır. Çünkü 0.1 µl ml⁻¹ dozu ile bile önemli bir etkinlik elde edilebilmiş ve miselyal büyüme yarı yarıya baskılanmıştır.

Bu iki doz arasındaki fark değerlendirildiğinde, uygulama açısından optimal doz seçiminin yalnızca maksimum inhibisyon sağlama üzerine değil, aynı zamanda etkinlik, maliyet ve çevresel etkiler gibi faktörler göz önünde bulundurularak yapılması gerektiği ortaya çıkmaktadır. Eğer tarımsal veya endüstriyel uygulamalarda PA kullanılacaksa, tam inhibisyon sağlayan 0.5 µl ml⁻¹ yerine, EC₅₀ değerine yakın olan 0.1 µl ml⁻¹ dozunun kullanımının da dikkate alınması gerekebilir. Daha düşük dozların yeterli etkinlik sağlayıp sağlayamayacağı veya daha uzun süreli uygulamalarda etkinliğin nasıl değişeceği ise ilerleyen çalışmalarda daha ayrıntılı olarak ele alınmalıdır.

Sonuç olarak, bu çalışma fındık kabuğundan elde edilen odun sirkesinin incirde *R. necatrix* fungal patojenine karşı güçlü bir antifungal ajan olarak kullanılabileceğini göstermektedir. Kimyasal fungusitlere karşı daha çevre dostu ve doğal bir alternatif sunan bu odun sirkesi, özellikle organik tarım uygulamalarında fungal kayıpların azaltılmasında umut vericidir.

KAYNAKÇA

Alfieri, S. A., Jr., Langdon, K. R., Kimbrough, J. W., El-Gholl, N. E., & Wehlburg, C. (1994). *Diseases and disorders of plants in Florida* (Bulletin No. 14). Florida Department of Agriculture and Consumer Services, Division of Plant Industry.

Cheong, S.-S., Choi, I.-Y., & Lee, W. H. (2013). Occurrence of gray mold caused by *Botrytis cinerea* on common fig in Korea. *The Korean Journal of Mycology*, 41(1), 38–41. <https://doi.org/10.4489/KJM.2013.41.1.38>

Farr, D. F., Bills, G. F., Chamuris, G. P., & Rossman, A. Y. (1989). *Fungi on plants and plant products in the United States*. The American Phytopathological Society Press.

Isa, M. M., Jaafar, M. N., Kasim, K. F., & Mutalib, M. F. A. (2020). Cultivation of fig (*Ficus carica* L.) as an alternative high-value crop in Malaysia: A brief review. *IOP Conference Series: Materials Science and Engineering*, 864, 012134. <https://doi.org/10.1088/1757-899X/864/1/012134>

Kara, M., Soylu, S., Soylu, E. M., & et al. (2024). Determination of the chemical composition and antifungal activity of wood vinegar (pyroligneous acid) against the onion bulb rot disease caused by *Fusarium proliferatum*. *Journal of Crop Health*, 76(1), 75–85. <https://doi.org/10.1007/s10343-023-00931-3>

Papachatzis, A., Eliopoulos, P., Statharas, G., & Vagelas, I. (2008). *Ficus carica* root rot disease caused by *Armillaria mellea* and *Rosellinia necatrix* in Greece. *University of Craiova*, 13, 143–148.

Ramadan, M. F. (Ed.). (2023). *Fig (Ficus carica): Production, processing, and properties*. Springer Nature. <https://doi.org/10.1007/978-3-031-16493-4>

Ruiz-Moyano, S., Martín, A., Villalobos, M. C., Calle, A., Serradilla, M. J., Córdoba, M. G., & Hernández, A. (2016). Yeasts isolated from figs (*Ficus carica* L.) as biocontrol agents of postharvest fruit diseases. *Food Microbiology*, 57, 45–53. <https://doi.org/10.1016/j.fm.2016.01.003>

- Sandhu, A. K., Islam, M., Edirisinghe, I., & Burton-Freeman, B. (2023). Phytochemical composition and health benefits of figs (fresh and dried): A review of literature from 2000 to 2022. *Nutrients*, *15*(11), 2623. <https://doi.org/10.3390/nu15112623>
- Shahrajabian, M. H., Sun, W., & Cheng, Q. (2021). A review of chemical constituents, traditional and modern pharmacology of fig (*Ficus carica* L.), a super fruit with medical astonishing characteristics. *Polish Journal of Agronomy*, *44*, 22–29. <https://doi.org/10.26114/pja.iung.452.2021.452.04>
- Soylu, E. M., Kurt, S., & Soyulu, S. (2010). In vitro and in vivo antifungal activity of essential oils of various plants against tomato gray mold disease agent *Botrytis cinerea*. *International Journal of Food Microbiology*, *143*, 183–189. <https://doi.org/10.1016/j.ijfoodmicro.2010.08.015>
- Soylu, S., Kara, M., Uysal, A., Gümüş, Y., Soyulu, E. M., Kurt, Ş., Üremiş, İ., & Sertkaya, E. (2024). Hatay İlinde yetiştirilen önemli Brassicaceous sebze türlerinde fungal ve bakteriyel hastalık etmenlerinin belirlenmesi. *KSÜ Tarım ve Doğa Dergisi*, *27*(4), 839–855. <https://doi.org/10.18016/ksutarimdog.vi.1383042>
- Subbarao, K. V., & Michailides, T. J. (1995). Effects of temperature on isolates of *Fusarium moniliforme* causing fig endosepsis and *Aspergillus niger* causing smut. *Phytopathology*, *85*(6), 662–668.
- Subramaniyan, V., Sellamuthu, P. S., & Sadiku, E. R. (2024). Evaluation of the antifungal activity of cell-free supernatant of *Lactobacillus* species against the fig (*Ficus carica* L.) postharvest fungal pathogens through in-vitro and in-silico analysis. *Scientia Horticulturae*, *331*, 113182. <https://doi.org/10.1016/j.scienta.2024.113182>
- White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. A. Innis, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR protocols: A guide to methods and applications* (pp. 315–322). Academic Press.
- Zhang, Z. R., Yang, X., Li, W. Y., Peng, Y. Q., & Gao, J. (2022). Comparative chloroplast genome analysis of *Ficus* (Moraceae): Insight into adaptive evolution and mutational hotspot regions. *Frontiers in Plant Science*, *13*, 1–17. <https://doi.org/10.3389/fpls.2022.965335>

EFFECTS OF PROLINE AND SALICYLIC ACID APPLICATION ON YIELD AND PLANT GROWTH IN GREEN ONION CULTIVATION

TAZE SOĞAN YETİŞTİRİCİLİĞİNDE PROLİN VE SALİSİLİK ASİT UYGULAMASININ VERİM VE BİTKİ GELİŞİMİ ÜZERİNE ETKİLERİ

Haydar BALCI

Van Yüzüncü Yıl University, Gevaş Vocational School, Department of Plant and Animal Production, VAN/ TURKEY

Orcid No: 0000-0003-0210-3639

Muhsin YILDIZ

Van Yüzüncü Yıl University, Gevaş Vocational School, Department of Plant and Animal Production, VAN/ TURKEY

Orcid No: 0000-0002-0766-5174

Murat KARA

Van Yüzüncü Yıl University, Gevaş Vocational School, Department of Plant and Animal Production, VAN/ TURKEY

Orcid No: 0000-0003-1011-918X

Abstract

This study was conducted to examine the effects of salicylic acid (SA), a phenolic compound, and proline (PR), an amino acid, on the growth and development of green onion (*Allium cepa* L.). The experiment was carried out in an unheated glass greenhouse at Van Yüzüncü Yıl University using a randomized block design. Key growth parameters such as plant height, root length, stem thickness, leaf number, total fresh and dry weight, and SPAD value were evaluated.

In terms of plant height, the PR2+SA1 application (27.26 cm) yielded the highest value, showing significant improvement compared to individual applications ($P<0.001$). Regarding root length, the highest values were recorded in the PR1 (16.57 cm) and SA2 (16.03 cm) treatments. This suggests that proline and salicylic acid may have a supportive effect on root development. For stem diameter, the highest value was observed in the PR1+SA1 combination (5.62 mm). While leaf number differences among treatments were relatively small, an increase compared to the control group was noted ($P=0.0367$).

For total fresh weight, the highest values were observed in PR2+SA1 (146.67 g) and PR1 (142.00 g) treatments, indicating that proline alone or in combination with salicylic acid may enhance biomass accumulation. Regarding dry weight, the highest value was recorded in PR2+SA1 (19.84 g). In terms of SPAD value, SA2 (89.73) resulted in the highest chlorophyll content, suggesting that salicylic acid has the potential to enhance photosynthetic efficiency.

Proline and salicylic acid applications, especially in combination, demonstrated positive effects on plant height, root development, and dry weight. In terms of SPAD values, salicylic acid was observed to increase chlorophyll content. Statistically significant differences among growth parameters indicate that these applications have a notable impact on green onion development.

These findings suggest that proline and salicylic acid, when applied at appropriate doses, have the potential to enhance yield in green onion cultivation. However, determining optimal dosage levels and evaluating their long-term effects under real production conditions are recommended.

Keywords : Green onion (*Allium cepa* L.), Salicylic acid (SA), Proline, Growth parameters, Biostimulants in agriculture

Özet

Bu araştırma, fenolik bileşiklerden salisilik asit (SA) ve aminoasitlerden prolin (PR) uygulamalarının taze soğan (*Allium cepa* L.) bitkisinin büyüme ve gelişimi üzerindeki etkilerini incelemek amacıyla yürütülmüştür. Çalışma, Van Yüzüncü Yıl Üniversitesi'ne ait ısıtmasız cam serada, tesadüf blokları deneme desenine göre planlanmış olup, bitki boyu, kök uzunluğu, gövde kalınlığı, yaprak sayısı, toplam yaş ve kuru ağırlık ile SPAD değeri gibi temel büyüme parametreleri değerlendirilmiştir.

Bitki boyu açısından, PR2+SA1 uygulaması (27,26 cm) en yüksek değeri vermiştir ve bu kombinasyon, bireysel uygulamalarla karşılaştırıldığında önemli bir gelişme sağlamıştır ($P<0,001$). Kök uzunluğu açısından, PR1 uygulaması (16,57 cm) ve SA2 uygulaması (16,03 cm) en yüksek değerleri sağlamıştır. Bu durum, prolin ve salisilik asitin kök gelişimini destekleyici etkilerini gösterdiğini düşündürmektedir. Gövde çapında, en yüksek değer PR1+SA1 (5,62 mm) kombinasyonu ile elde edilmiştir. Yaprak sayısında, tüm uygulamalar arasında küçük farklar gözlemlenmiştir ancak kontrol grubuna göre artış olduğu dikkat çekmektedir ($P=0,0367$).

Toplam yaş ağırlıkta, en yüksek değer PR2+SA1 (146,67 g) ve PR1 (142,00 g) gruplarında gözlenmiştir. Bu, prolinin tek başına veya salisilik asit ile kombinasyon halinde biyokütleyi artırabileceğini göstermektedir. Kuru ağırlık açısından, en yüksek değer PR2+SA1 (19,84 g) uygulaması ile elde edilmiştir. SPAD değeri bakımından, SA2 (89,73) en yüksek klorofil içeriğini göstermiştir. Bu durum, salisilik asitin fotosentez verimliliğini artırma potansiyeline sahip olabileceğini ortaya koymaktadır. Prolin ve salisilik asit uygulamaları, özellikle kombinasyon halinde kullanıldığında bitki boyu, kök gelişimi ve kuru ağırlık üzerinde pozitif etkiler göstermiştir. SPAD değerleri açısından, salisilik asitin daha yüksek klorofil içeriği sağladığı görülmektedir. Büyüme parametreleri arasında istatistiksel olarak anlamlı farklılıklar gözlemlenmiş olup, bu da uygulamaların taze soğanın gelişimi üzerinde etkili olduğunu göstermektedir.

Bu sonuçlar, prolin ve salisilik asitin, uygun dozlarda kullanıldığında, taze soğan yetiştiriciliğinde verimi artırıcı bir potansiyele sahip olduğunu ortaya koymaktadır. Ancak, optimum doz seviyelerinin belirlenmesi ve bu uygulamaların uzun vadeli etkilerinin üretici koşullarında test edilmesi tavsiye edilmektedir.

Anahtar Kelimeler: Taze soğan (*Allium cepa* L.), Salisilik asit (SA), Prolin, Büyüme parametreleri, Tarımda biyostimulanlar

A CRITICAL ANALYSIS OF TURKEY'S AGRICULTURAL POLICIES IN THE CONTEXT OF COMBATING CLIMATE CHANGE AND SUSTAINABLE AGRICULTURE

Arař. Gör. Dr. Ezgi DEMİRAL

Doęuř Üniversitesi, İktisadi ve İdari Bilimler Fakültesi, İktisat Bölümü, İstanbul, Türkiye

ORCID: 0000-0001-7822-7967

Abstract

The agricultural sector holds strategic importance for countries as it not only meets individuals' food needs but also supplies raw materials to industries such as manufacturing. Türkiye, with its vast geographical area, biodiversity, diverse crop patterns, and extensive agricultural lands, has the potential to achieve self-sufficiency in agriculture. However, long-standing structural issues in the agricultural sector, coupled with Turkey's vulnerability to climate change, pose significant obstacles to fully utilizing this potential.

Since agricultural production is one of the primary economic activities, it is highly dependent on climatic conditions. The impact of climate change on agriculture varies across time and space. However, the increasing frequency of extreme weather events due to climate change affects the agricultural sector in multiple ways, directly influencing agricultural production activities. Given its structural challenges and susceptibility to climate change, the agricultural sector requires government support. Agricultural support policies play a crucial role in ensuring sustainability in agricultural production, food security, and farmers' income.

This study examines Türkiye's agricultural policies in the context of combating climate change and promoting sustainable agriculture. Key policy instruments include support purchases, farmer loans, direct income support, product-based subsidies, and incentives for good agricultural practices. The study evaluates the trajectory and effectiveness of these support mechanisms over the years. The findings suggest that existing agricultural policies are neither effective nor sufficient to ensure the long-term sustainability of the sector.

Keywords: Sustainable Agriculture, Agricultural Support Policies, Climate Change, Food Security

Özet

Tarım sektörü hem bireylerin gıda ihtiyacını karşılaması hem de sanayi gibi sektörlere hammadde girdisi sağlaması dolayısıyla ülkeler için stratejik öneme sahiptir. Nitekim Türkiye geniş bir coğrafi alana yayılmış olması, sahip olduđu biyoçeşitlilik, ürün deseni ve tarım alanları ile tarımda kendi kendine yeterlilięi sağlama potansiyeline sahiptir. Bununla birlikte gerek tarım sektörünün uzun yıllardır içinde bulunduđu yapısal sorunlar gerekse Türkiye'nin iklim deęişikliği karşısındaki kırılgan konumu tarım konusundaki potansiyelini deęerlendirilmesinin önünde önemli bir engel teşkil etmektedir.

Tarımsal üretim birincil ekonomik faaliyetlerden olması dolayısı ile iklim koşullarına doğrudan baęlıdır. İklim deęişikliğinin tarım sektörü üzerindeki etkisi zamansal ve mekânsal olarak farklılaşmaktadır. Bununla birlikte iklim deęişikliğine baęlı aşırı hava olaylarındaki artış tarım sektörünü birçok yönden etkilemekte ve tarımsal üretim faaliyetleri üzerinde doğrudan etki yaratmaktadır. Bu bağlamda tarım sektörü yapısal koşulları ve iklim deęişikliği karşısındaki

kırılganlığı nedeniyle devlet politikaları ile desteklenmeye ihtiyaç duymaktadır. Tarımsal destekleme politikaları hem tarımsal üretim ve gıda güvencesinde hem de çiftçi gelirlerinde sürdürülebilirliğin sağlanması amacıyla ön plana çıkmaktadır. Bu amaçla çalışmamızda Türkiye’de uygulanan tarım politikaları iklim değişikliği ile mücadelede ve sürdürülebilir tarım bağlamında incelenmektedir. Bu kapsamda destekleme alımları, çiftçiye sağlanan krediler, doğrudan gelir desteği, ürün bazlı destekler ve iyi tarım uygulamalarına verilen destekler öne çıkan politika araçlarıdır. Çalışmamızda söz konusu destekleme araçlarının yıllar içerisindeki seyri ve etkinliği tartışılmaktadır. Çalışmanın bulguları geliştirilen tarım politikalarının sektörün sürdürülebilirliğinin sağlanması için etkin ve yeterli olmadığı yönündedir.

Anahtar Kelimeler: Sürdürülebilir Tarım, Tarımsal Destekleme Politikaları, İklim Değişikliği, Gıda Güvencesi

Giriş

İklim değişikliği ve tarım sektörü arasındaki ilişki karşılıklıdır. Bir yandan tarımsal üretim faaliyetleri sonucu atmosfere yayılan sera gazları iklim değişikliğini kuvvetlendirici etki yaratmaktayken diğer yandan ise iklim değişikliğine bağlı aşırı hava olayları tarımsal üretim faaliyetlerini etkilemektedir. Tarımsal üretim birincil ekonomik faaliyetlerden olması dolayısı ile iklim koşullarına doğrudan bağlıdır. İklim değişikliğinin tarım sektörü üzerindeki etkisi zamansal ve mekânsal olarak farklılaşmaktadır. Bununla birlikte iklim değişikliğine bağlı aşırı hava olaylarındaki artış tarım sektörünü birçok yönden etkilemektedir. İklim değişikliğine bağlı yaşanan aşırı sıcakların ürünlerin olgunlaşma süresini etkilemesi ve ürünü strese sokması verimlilik üzerinde olumsuz bir etki yaratmaktadır (Porter ve Gawith 1999: 24). Sel, fırtına ve kasırga gibi hava olaylarının yaşanması ise hem toprak hem de mahsul üzerinde olumsuz etki yaratmaktadır. Sıcaklıkların yükselmesi ve aşırı hava olaylarının bir diğer sonucu ise emek verimliliği üzerinde ortaya çıkmaktadır. Tarım sektörü çalışanları hava olaylarındaki aşırı ılıktan olumsuz etkilenmekte ve işgücü verimliliği düşmektedir (Murat ve Şengül 2024: 715). Nitekim Türkiye iklim değişikliğine bağlı yaşanan sıcaklık artışı, yağış rejiminin değişmesi, kuraklık, sel ve erozyon gibi iklimsel aşırılıklara en açık bölgelerden biridir (IPCC 2007; Fay vd. 2009; Kadioğlu 2012; IPCC 2021).

Bununla birlikte tarım sektörü yapısal koşulları ve iklim değişikliği karşısındaki kırılganlığı nedeniyle devlet politikalarına ihtiyaç duymaktadır. Tarımsal destekleme politikaları gerek tarımsal üretim ve gıda güvencesinde gerekse çiftçi gelirlerinde sürdürülebilirliğin sağlanması amacıyla ön plana çıkmaktadır. Bu bağlamda, bu çalışmada 1980 sonrası Türkiye’de uygulanan tarım politikaları iklim değişikliği ile mücadelede ve sürdürülebilir tarım bağlamında incelenmektedir.

Materyal ve Yöntem

İlgili çalışmada, Türkiye’de tarım sektörünün iklim değişikliği karşısındaki kırılgan konumu ve tarımsal sürdürülebilirliğin sağlanması bağlamında devlet politikalarının etkinliği tartışılacaktır. Bu amaçla 1980 sonrası Kalkınma Programları, IMF Niyet Mektupları, AB İlerleme Raporları ile Tarım ve Orman Bakanlığı tarafından iklim değişikliği ve tarım sektörüne ilişkin hazırlanan belgeler doküman analizi yöntemi ile incelenecektir.

Bulgular ve Tartışma

Tarım sektörü Türkiye ekonomisi için Cumhuriyet tarihi boyunca her zaman önemini korumuş ve zaman içerisinde farklı politika araçları ile desteklenmiştir. İklim değişikliği ile ilgili farkındalık 1970’ler itibariyle gelişmeye başlamış ve tarım sektörüne etkisi de ilk kez 1974 yılındaki Dünya Gıda Konferansında gündeme gelmiştir (FAO 1974). Dolayısıyla bu tarihlerden önce geliştirilen tarımsal destekleme araçlarının iklim değişikliği ile mücadeleyi

konu alması beklenmemektedir. Bu bağlamda ilgili çalışmada 1980 sonrası tarımsal destekleme politikalarına odaklanılmıştır.

1980’lerde yaşanan Keynesyen politikalardan Neoliberal politikalara geçiş Türkiye ekonomisinde de karşılık bulmuştur. Türkiye’de yeni modele geçişin ilk aşaması 24 Ocak 1980’de açıklanan paket olmuş ve 24 Ocak Kararları ile ekonomide yaşanan önemli dönüşümden tarım sektörü de etkilenmiştir. Dış ticaretin serbestleştirilmesi ve uluslararası piyasalarda rekabet gücü kazanılması amacıyla tarım sektöründe bazı değişiklikler öngörülmüştür. Bu bağlamda tarım sektöründeki üreticilerin gelirlerinin düşürülmesi, destek alımları ve fiyat politikalarının değişmesi gibi önlemler uygulamaya konmuştur (Pamuk 2022: 266). Bununla birlikte 1980 sonrası dönemde destekleme politikalarının ekonomideki ağırlığının azaltıldığı görülmektedir.

Tablo 1. Seçili Ülkelerde Tarıma Yapılan Destekler (1989)

Ülkeler	Tarım Üreticisi Başına (1000 Dolar)	Hektar Başına (Dolar)
Japonya	15	8.761
Finlandiya	26	2.167
Avusturya	9	648
Norveç	32	2.928
ABD	20	94
AET	8	501
Avustralya	4	3
Türkiye	0,230	36

Kaynak: TZOB 1992:160 verileri kullanılarak yazar tarafından hazırlanmıştır.

Tablo 1’den de izlenebileceği üzere Türkiye tabloda yer alan ülkeler arasında 230 dolar ile tarım üreticisi başına düşen destek miktarının en az olduğu ülkedir. Bununla birlikte tarım sektörüne yapılan yatırımlar 1990’larda giderek azalmıştır. Tablo 2’de de görüldüğü gibi 1963-1967 dönemini kapsayan Birinci BYKP’de tarım sektörüne ayrılan sabit sermaye yatırımlarının payı % 13.9 iken, 1990-1994 dönemini kapsayan Altıncı BYKP’de %5’e gerilemiştir.

Tablo 2. Planlı Dönem Tarım Sektörü Sabit Sermaye Yatırımları (Cari Fiyatlarla, %)

	I. Plan Dönemi	II. Plan Dönemi	III. Plan Dönemi	IV. Plan Dönemi	V. Plan Dönemi	VI. Plan Dönemi
Tarım	13.9	11.1	11.8	10	7.9	5.1

Kaynak: Karluk 1999: 181 verileri kullanılarak yazar tarafından hazırlanmıştır.

1990’lı yıllar itibarıyla tarım politikalarında IMF ve Dünya Bankası uzmanlarının görüşleri etkili olmuştur (Oyan 2004: 52). Nitekim nasıl bir tarım sektörü inşa edilmek istendiği ve tarım sektöründeki yeniden yapılanma hedefi IMF’ye verilen niyet mektuplarından izlenebilmektedir. 9 Aralık 1999 yılında IMF’ye verilen Niyet Mektubunda enflasyonla mücadele kapsamında ortaya konulacak politikalar belirtilmiştir. Bu amaçla gerçekleştirilecek yapısal reformların bir ayağını da tarım politikaları oluşturmaktadır. Mektupta tarım sektörüne ilişkin olarak ‘‘tarımsal destekleme politikaları fakir çiftçilere destek sağlamanın en düşük maliyetli yöntemi değildir’’ ifadesi yer almış ve orta vadeli amaç olarak ‘‘destekleme politikalarını safhalar halinde ortadan kaldırmak ve fakir çiftçileri hedef alan doğrudan gelir desteği sistemi ile değiştirmek’’ amacı belirtilmiştir (Hazine Müsteşarlığı 1999).

AB Türkiye’nin tarım ürünleri ticaretinde önemli bir partneridir. Nitekim 2004 yılında yayımlanan 2006-2010 dönemi için hazırlanan Tarım Stratejisi Belgesinde Ortak Tarım Politikası (OTP) ve Tarım Anlaşması esas alınarak rekabet gücü yüksek, sürdürülebilir bir tarım sektörünün inşa edilmesi amaçlanmıştır (Tarım ve Orman Bakanlığı, 2004). Kanunda tarım

sektörünün hangi araçlar ile destekleneceği belirtilmiştir. Buna göre tarımsal destekleme araçları şunlardır:

- Doğrudan gelir desteği,
- Fark ödemesi,
- Telafi edici ödemeler,
- Hayvancılık destekleri,
- Tarım sigortası ödemeleri,
- Kırsal kalkınma destekleri,
- Çevre amaçlı tarım arazilerini koruma programı destekleri,
- Diğer destekleme ödemeleri.

Türkiye'nin 2006 yılında çıkardığı 5488 Sayılı Tarım Kanununun amacı "tarım sektörünün ve kırsal alanın, kalkınma plân ve stratejileri doğrultusunda geliştirilmesi ve desteklenmesi için gerekli politikaların tespit edilmesi ve düzenlemelerin yapılması" şeklinde ifade edilmiş ve uygulanacak tarım politikalarının ilkeleri belirlenmiştir. Buna göre uluslararası taahhütler ile uyumlu, özel sektörün rolünün arttığı, piyasa mekanizmalarına uygun destekleme araçlarının tercih edildiği bir tarım sektörü inşası amaçlanmıştır. Kanunda tarımsal destekleme politikalarının çerçevesi ayrıntılı şekilde belirtilmiştir. 18. Madde itibarıyla tarımsal destekleme politikalarının amacı, ilkeleri ve araçları belirtilirken, Madde 21'de tarımsal destekleme politikalarına ayrılacak kaynağın GSMH'nin %1'ine denk geleceği ifade edilmiştir (Resmî Gazete 2006). Destekleme kapsamına alınan ürünler ve destek miktarları her yıl Cumhurbaşkanlığı Kararı ile Resmî Gazetede yayımlanmaktadır.

AB, OTP kapsamında doğrudan gelir desteği ile iklime ve çevreye duyarlı üretim yapan çiftçilere "yeşil ödeme" kapsamında ödeme yapmaktadır (European Commission 2017: 4). Türkiye'de ise benzer uygulama ÇATAK (Çevre Amaçlı Tarım Arazilerini Koruma Programı) ile yürütülmüştür. ÇATAK destekleri tarımsal sürdürülebilirliğin sağlanması için öne çıkan politika araçlarından biridir. ÇATAK programı ile "toprak ve su kalitesinin korunması, doğal kaynakların sürdürülebilirliği, erozyonun önlenmesi ve tarımın olumsuz etkilerinin azaltılmasına yönelik alanların korunması" amaçlanmaktadır. Program kapsamında destekleme yapılacak alanlar üç kategoriye ayrılmıştır. Bunlar: Minimum toprak işlemeli tarım uygulamaları, Toprak ve su yapısının korunması ve erozyonun engellenmesi, Çevre dostu tarım teknikleri ve kültürel uygulamalardır. 2006 yılında uygulanmaya başlanan ÇATAK kapsamında çevre dostu üretim tekniklerinin benimsenmesi, doğal kaynakların sürdürülebilirliğinin sağlanması gibi amaçları önceleyen üreticilere destek sağlanmıştır. Program kapsamında verilecek destek Türkiye'deki üreticilerin tamamını değil, yalnızca belli illeri kapsamaktadır. 2006 yılında 4 il ile başlayan destekler 2019 yılı itibarıyla 58 ile ulaşmıştır. Ancak 2020 yılında tamamlanması planlanan program için 2018 yılından bu yana yeni başvuru alınmamaktadır (Tarım ve Orman Bakanlığı ÇATAK). Nitekim Cumhurbaşkanlığı Yıllık Raporunda da 2024 yılı itibarıyla ÇATAK için herhangi bir destekleme bütçesi ayrılmamıştır. Oysa gerek Türkiye'nin iklim değişikliği karşısındaki kırılgan konumu gerekse tarım sektörünün neden olduğu sera gazı salımı dikkate alındığında tarımsal üretim süreçlerinde sürdürülebilir üretim tekniklerinin devlet politikaları ile desteklenmesi elzem hale gelmektedir.

Tablo 3. Çatak Uygulamaları (Yıl Bazlı)

YIL	İL SAYISI	ÜRETİCİ SAYISI	ALAN (ha)	ÖDEME MİKTARI (TL)
2006	4	469	1.726	1.434.000
2007	4	1.508	4.041	2.605.000
2008	4	1.484	4.063	4.630.000
2009	9	1.881	4.752	5.061.922
2010	19	2.940	8.808	10.347.256
2011	25	4.648	14.414	16.128.359
2012	27	6.568	21.804	23.182.680
2013	30	9.195	33.172	35.084.038
2014	43	15.430	50.559	52.890.491
2015	51	22.671	77.969	81.346.357
2016	51	29.811	111.729	113.037.438
2017	57	35.333	142.964	141.632.770
2018	58	34.898	149.780	150.278.723
2019	58	21.825	95.662	96.422.371

Kaynak: Tarım ve Orman Bakanlığı verileri kullanılarak yazar tarafından hazırlanmıştır.

Türkiye'nin AB'ye üyelik yolunda tarım sektöründe gerçekleştirdiği reformları Türkiye Düzenli İlerleme Raporlarından izlemek mümkündür. Bu bağlamda özellikle yakın dönemli gelişmelerin takip edilmesi amacıyla 2020, 2021, 2022 ve 2023 İlerleme Raporlarının Fasıllık 11 başlıkları incelenmiştir. Raporlarda Türkiye'nin tarım istatistikleri oluşturma konusunda bir strateji geliştirmedeği ve Türkiye'de uygulanan politikalarının OTP ilkelerinden uzaklaştığı yönünde eleştiriler bulunmaktadır (European Commission 2020, 2021, 2022, 2023). Türkiye ile AB arasında ilki Mayıs 2022, ikincisi ise Kasım 2023 tarihinde Türkiye-AB Yüksek Düzeyli Tarım Diyalogu gerçekleştirilmiştir. Üç temel konu üzerinden şekillenen toplantıda tarım ürünleri konusunda ikili ticaretin geliştirilmesi, gıda güvenliğinin sağlanması ve IPARD III mali yardım konuları ele alınmıştır. Sürdürülebilir tarım pratiklerinin uygulamaya koyulması ve çevrenin korunmasını önceliklendiren üretim yöntemlerinin benimsenmesi gibi konuların önemi çıkarılan kanunlar ile de vurgulanmaktadır. Bu bağlamda 5 Nisan 2023 tarihinde yayımlanan Resmî Gazete'deki 7442 Sayılı Orman Kanunu ve Bazı Kanunlarda Değişiklik Yapılmasına Dair Kanun ile 2006 yılında kabul edilen Tarım Kanunu'nun bazı maddelerinde değişiklikler yapılmıştır. Buna göre "Tarımsal üretimin planlanması, gıda güvencesi ve güvenliğinin temin edilmesi, verimliliğin artırılması, çevrenin korunması ve sürdürülebilirliğin tesis edilmesi için Bakanlıkça belirlenen ürün veya ürün gruplarının üretimine başlanmadan önce Bakanlıktan izin alınmaktadır. Bakanlık, arz ve talep miktarı ile yeterlilik derecesini dikkate alarak hangi ürün veya ürün gruplarının üretileceği ile tarım havzası veya işletme bazında asgari ve azami üretim miktarlarını belirler." (Resmî Gazete 2023).

Uluslararası alanda Türkiye tarım sektörünün rekabet gücünü koruyabilmek için, tarımsal üretim sürecini uzun vadede olumsuz etkileyen ve sürdürülebilir olmayan üretim pratiklerinden uzaklaşılması ve uluslararası belirlenen standartlara uygun üretim yapılması önem arz etmektedir. Bu bağlamda tarımsal sürdürülebilirliğin sağlanması için öne çıkan bir diğer uygulama alanı ise iyi tarım ve organik tarım uygulamalarıdır. Türkiye'de iyi tarım uygulamaları Avrupalı perakendecilerin talepleri doğrultusunda GLOBALGAP sertifikalı tarımsal ürünler üretmek amacıyla 2003 yılında başlamıştır. Türkiye'de iyi tarım uygulamaları sonucu elde edilen ürün miktarı 2007 yılından 2023 yılına 109 kat artmıştır. Bu önemli bir artıştır ancak 2019 yılında çıkarılan Cumhurbaşkanlığı 2019 Tarımsal Destekleme Kararnamesi

ile organik ve iyi tarım uygulamalarına verilen destekler sınırlandırılmıştır. 9 Kasım 2019'da Resmî Gazete'de yayımlanan Tebliğ'in 24. Maddesinde " 2016 yılından başlamak üzere aralıksız olarak üç yıl İTU¹ desteğinden yararlananlar, 2019 üretim yılı İTU desteğinden faydalanamaz." ve " 2017 ve 2018 üretim yılında organik statüde OTD²'den faydalanan araziler, 2019 üretim yılında OTD'den faydalanamaz." şeklinde düzenlemeler yapılarak iyi tarım ve organik tarım uygulamalarına verilen destekler azaltılmıştır.

Tablo 4. İyi Tarım ve Organik Tarım Uygulamaları (2023)

	Üretici Sayısı	Üretim Alanı (ha)	Üretim Miktarı (Ton)
İyi Tarım Uygulamaları	8 045	372 651	6 119 846
Organik Tarım Uygulamaları	34 358	190 825,85	1 028 888,22

Kaynak: Tarım ve Orman Bakanlığı 2023 verileri kullanılarak yazar tarafından hazırlanmıştır.

İklim değişikliği ve tarım sektörü arasındaki karşılıklı ilişki dolayısıyla çevreye duyarlı üretim tekniklerinin benimsenmesi sürdürülebilir tarımsal üretimin sağlanabilmesi için elzemdir. Bu bağlamda tarım sektöründe yaşanması beklenen dönüşüm bütüncül bir yaklaşımı gerektirmektedir. Üretim sürecindeki en önemli aktörler olan çiftçilerin uyum kapasiteleri, benimsenecek politikaların etkin şekilde yürütülmesinde önemli bir etkidir. Çiftçilerin iklim değişikliğine dair farklı algıları yürütülecek politikaların da çeşitlendirilmesini ve devamlılığının sağlanmasını gerektirmektedir. Ancak gerek ÇATAK programı kapsamında gerekse İTU ve OTD'ye verilen destekler kapsamında sürdürülebilirliğin sağlandığını söylemek mümkün değildir.

Sonuç ve Öneriler

Türkiye coğrafi konumu itibarıyla iklim değişikliği etkilerine karşı açık bir ülkedir. İklim değişikliğine ilişkin ortaya konulan çalışmalarda Türkiye'nin de içinde bulunduğu Akdeniz bölgesinde karasal ısınmaya bağlı kuraklığın artacağı ve nem azalışı ve su talebindeki artışa bağlı olarak evapotranspirasyonda artış görüleceği belirtilmektedir (IPCC 2007: 8, 49; IPCC 2013: 91; IPCC 2021: 86). Diğer yandan tarımsal üretim faaliyetlerinin doğrudan iklim koşullarına bağlı olması iklim değişikliğinin tarım sektörüne etkilerini tartışmayı ve bu etkileri dikkate alarak politika geliştirilmesini gerekli kılmaktadır. Tarım sektörü yapısal koşulları ve iklim değişikliği karşısındaki konumu ile hem çiftçi gelirlerinin hem de üretimin sürdürülebilir olması amacıyla devlet politikaları ile desteklenmelidir. Nitekim çalışmamızın bulguları Türkiye'de tarım sektöründe iklim değişikliği ile mücadelenin etkin şekilde yürütülmediğini göstermektedir. Bu bağlamda geliştirilen politikaların iklim değişikliğinin tarım sektörü üzerindeki etkilerini dikkate alan ve hem çiftçi gelirlerinin hem de üretimin sürdürülebilirliğini sağlayacak şekilde önceliklendirilmesi gerekmektedir. Bu bağlamda AB'dekine benzer uygulamalarla çevre dostu üretim pratiklerini benimseyen ve hem azaltım hem de uyum stratejilerine katkı sağlayan üretim tekniklerinin devlet politikaları aracılığıyla desteklenmesi ve üretici açısından tercih edilebilir hale getirilmesi gerekmektedir.

¹ İTU: İyi Tarım Uygulamaları

²OTD: Organik Tarım Desteği

Kaynakça

- European Commission (2017), *CAP Explained Direct Payments for Farmers 2015-2020*.
- European Commission (2020, 6 Ekim), Commission Staff Working Document Turkey 2020 Report.
- European Commission (2021, 19 Ekim), Commission Staff Working Document Turkey 2021 Report.
- European Commission (2022, 12 Ekim), Commission Staff Working Document Türkiye 2022 Report.
- European Commission (2023, 11 Kasım), Commission Staff Working Document Türkiye 2023 Report.
- FAO (1974), ‘‘Report Of The Council Of Fao Sixty-Fourth Session’’, Rome, 18–29 November 1974.
- Fay M., Block R., Carrington T., Ebinger J. (2009), ‘‘Adapting to Climate Change in Europe and Central Asia’’, (The World Bank 2009/06/01), <https://documents.worldbank.org/en/publication/documents-reports/documentdetail/127181468024643244/adapting-to-climate-change-in-europe-and-central-asia>
- Hazine Müsteşarlığı (1999), *09/12/1999 Tarihli IMF Niyet Mektubu*.
- IPCC (2007), ‘‘Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change’’, Core Writing Team, Pachauri, R.K., Reisinger, A. (Eds.), IPCC, Geneva, Switzerland.
- Kadıoğlu M. (2012), Türkiye’de İklim Değişikliği Risk Yönetim, *Türkiye’nin İklim Değişikliği II. Ulusal Bildiriminin Hazırlanması Projesi Yayını*.
- Karluk R. (1999), *Türkiye Ekonomisi* (6. Baskı), İstanbul: BETA Basım.
- Murat, G., Şengül, B. (2024), ‘‘Küresel İklim Değişikliğinin İşgücü Verimliliği ve İstihdam Üzerine Yansımaları: Bir Sosyal Politika Önlemi Olarak Yeşil İşler’’, *Mülkiye Dergisi*, 48/3: 715-746.
- Oyan O. (2004), ‘‘Tarımsal Politikalardan Politikasız Bir Tarıma Doğru’’, Balkan N., Savran S. (Ed), Neoliberalizmin Tahribatı, İstanbul: Metis Yayınları: 44-67.
- Pamuk Ş. (2022), Türkiye’nin *200 Yıllık İktisadi Tarihi* (14. Baskı), İstanbul: Türkiye İş Bankası Kültür Yayınları.
- Porter, J. R., Gawith, M. (1999), ‘‘Temperatures and The Growth and Development of Wheat: A Review’’, *European Journal of Agronomy*, 10: 23-36.
- Resmî Gazete (2006), 25 Nisan 2006, 26149 sayılı.
- Resmî Gazete (2023), 5 Nisan 2023, 32154 sayılı.
- Strateji ve Bütçe Başkanlığı (2023), *On İkinci Kalkınma Planı (2024-2028)*.
- Tarım ve Orman Bakanlığı (2004), Türkiye Tarım Stratejisi (2006-2010) <http://www.yesilirmak.org.tr/documents/planlama/diger/00000000007.pdf> Erişim tarihi: 08 Ocak 2023.
- Tarım ve Orman Bakanlığı (2023), İyi Tarım Uygulamaları İstatistik, Erişim tarihi: 15 Ağustos 2024).

Tarım ve Orman Bakanlığı (2023), Organik Tarım Uygulamaları İstatistik, Erişim tarihi: 15 Ağustos 2024.

Tarım ve Orman Bakanlığı, ÇATAK (Çevre Amaçlı Tarımsal Arazileri Koruma Programı) <https://www.tarimorman.gov.tr/Konular/Bitkisel-Uretim/Tarla-Ve-Bahce-Bitkileri/CATAK>
Erişim tarihi: 15 Ocak 2024.

Tarım ve Orman Bakanlığı, ÇATAK (Çevre Amaçlı Tarımsal Arazileri Koruma Programı) <https://www.tarimorman.gov.tr/Konular/Bitkisel-Uretim/Tarla-Ve-Bahce-Bitkileri/CATAK>
Erişim tarihi: 15 Ocak 2024.

The Intergovernmental Panel on Climate Change, IPCC (2007), *AR4 Climate Change 2007: Synthesis Report*, Pachauri R. K., Reisinger A., (Eds). Geneva, Switzerland.

The Intergovernmental Panel on Climate Change, IPCC (2013), *Climate Change 2013: The Physical Science Basis* (1st Edition), Stocker, T. F., Qin, D., Plattner, G., Tignor, M. M. B., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Vincent, B., Midgley, P. M. (Eds), Cambridge and New York: Cambridge University Press.

The Intergovernmental Panel on Climate Change, IPCC (2021), *Climate Change 2021: The Physical Science Basis*, Masson-Delmotte V., Zhai P., Pirani A., Connors S. L., Pean C., Chen Y., Goldfarb L., Gomis M. I., Matthews J. B. R., Berger S., Huang M., Yelekçi Ö., Yu R., Zhou B., Lonnoy E., Maycock T. K., Waterfield T., Leitzell K., Caud N., (Eds), Cambridge and New York: Cambridge University Press .

Türkiye Ziraat Odaları Birliği (1992), *Zirai ve İktisadi Rapor 1990-1999*, Yayın No:168. https://kutuphane.tarimorman.gov.tr/pdf_goster?file=373cf1cecb5019490f4e539802cfa34d#book/9

BIOPLASTIC USE IN AGRICULTURE: EFFECTS ON SOIL HEALTH AND PLANT DEVELOPMENT

Res. Asst. Alperay ALTIKAT

Iğdır University, Faculty of Agriculture Department of Biosystems Engineering, Iğdir,

Prof. Dr. Mehmet Hakkı ALMA

Iğdır University, Faculty of Agriculture Department of Biosystems Engineering, Iğdir,

Abstract

The extensive use of conventional plastics in agriculture has led to serious environmental concerns due to their persistence in soils, causing adverse impacts on soil health and plant development. In response to these challenges, bioplastics have emerged as a promising sustainable alternative. This study explores the impact of bioplastics on soil properties, microbial activity, and plant development within agricultural systems. Bioplastics, typically derived from renewable biomass, exhibit biodegradability and compostability, significantly reducing their long-term environmental footprint compared to petroleum-based plastics. When used as mulching materials, bioplastics can enhance soil quality by improving nutrient retention, water conservation, and reducing weed growth. However, the degradation behavior of bioplastics varies significantly based on their chemical composition and environmental conditions, requiring a thorough evaluation of their long-term effects. Recent research highlights that while bioplastic mulch films disintegrate into smaller fragments relatively quickly, complete biodegradation may take extended periods, influencing soil microbial communities and nutrient cycles. Field studies have shown mixed results regarding the impact of bioplastics on plant growth, indicating potential improvements in root development, biomass production, and yield under specific conditions. Despite their benefits, critical issues remain unresolved, such as the potential accumulation of residual fragments and associated microbial changes, which might negatively impact plant growth and soil chemistry in the long term. Therefore, comprehensive long-term studies and rigorous assessments of biodegradation processes are essential. Collaborative efforts between researchers, agricultural practitioners, and policymakers are vital to promote effective implementation strategies and ensure the sustainable integration of bioplastics into agricultural practices.

Keywords

Bioplastics, Biodegradable Mulch, Soil Health, Plant Development, Sustainable Agriculture, Microbial Activity, Environmental Impact

Introduction to Bioplastics

This article overlaps the challenges of bioplastics in agriculture. The latest conventional plastic can be persistent in the environment or soil and transform into micro- or nanoparticles during degradation, which can result in their accumulation and negative effect(s) on soil biota (M. S. Cruz et al., 2022). The application of biodegradable agricultural mulches represents an ideal alternative for the optimization of production strategies minimizing the contamination and avoiding extra waste management by the producers. The renewability and compostability of bioplastics could reduce the amount of potentially toxic metals and metalloids in soil. However,

agro-mulching is rather new agricultural practice thus empirical data evaluating the effect of aged biodegradable mulches, and even infrequent use of them is scarce. Bio-based plastic is limited by several factors such as availability of arable land and water, and finally, problems with the cultivation of biogenic resources in regions where agriculture is not economically viable, primarily due to unfavorable soil and climate conditions (Manabu Abe et al., 2021).

The interest on biodegradable bioplastics has considerably grown in the last years, due to society's concerns about the accumulation of huge amounts of fossil-based plastic debris in the environment (Moshood et al.2022). Biodegradable polymers are able to mineralize in inert or stabilized-compost environment due to their chemical structure, rendering CO₂, water, mineral salts, and biomass. Biodegradable bioplastics are very attractive for agricultural applications, in order to combine the beneficial properties of traditional plastics with the low impact on the environment. Regarding this scenario and the fact that by the year 2030 the production of biodegradable bioplastics is expected to reach 2.6 million tons per year, from a global point of view, there is a need for experts to comprehend their potential long-term effects on soil properties, fertility, and biota (Ibrahim et al.2021). Plant mulches can increase soil quality, particularly unfertile and sandy soils, because during mineralization nutrients from crop residues are released to mineral soil. However, aged material can have different effects as fresh material, for this reason different properties of aged bioplastic mulches will be tested.

Current Challenges in Agriculture

Agriculture currently faces a number of problems that must be addressed in order to ensure food security and environmental sustainability. Energy consumption, water use and pollution, soil degradation, etc., are growing concerns in the agricultural sector. One of the most noticeable environmental problems is the rapid increase in the use of traditional plastics, which are non-biodegradable, and their consequent disposal, which greatly affects the environment, in particular the soil biota, which all agricultural processes depend upon (Fan et al., 2022). The regulation to resolve the problem with the use and disposal of traditional plastics is the future trend to promote the use of biodegradable plastics instead (Filipe et al., 2023). However, the use of traditional plastics is still cheap, effective and omnipresent. The shift to biodegradable plastics is expected to bring problems from the agricultural point of view. This study provides a state of the art review of the use of traditional plastics, the problem of its waste, the financial, environmental and legislative restrictions, and the solutions with biodegradable plastics with respect to their advantages and disadvantages, specifically about the technical properties of soil and plant development (Moshood et al.2022).

Biodegradability and Compostability of Bioplastics

In recent years, agricultural activity has sought new ways to reduce our carbon footprint by incorporating new and more sustainable materials (Choe et al., 2021). One of the products that have been given special attention is plastic given their high carbon content. However, the agricultural industry heavily relies on this material for their resilient nature (Otoni et al.2021). From this trend on environmentally friendly material, bioplastics have begun to play an important part. Bioplastics are mostly derived from renewable resources; however, this is not the defining feature. Biodegradability and compostability should also be considered. This paper will establish the critical precedent knowledge needed to see this new bioplastic trend in the context of the agricultural industry (Mujtaba et al.2023).

The characteristic qualities of a piece of plastic based material can make it susceptible to biodegradation on different scales. By specifying these scales we refer to the definition of biodegradable and compostable plastics as distinct categories. Also, another important

consideration in defining biodegradable plastic is time scale. Traditionally, polymers are materials which are mostly biologically inert (Ghosh & Jones, 2021). This means that they have very low water solubility and are not easily broken down by biological actions like microbes. The biodegradable quality that characterizes bioplastics makes the ability to break down in biosafe ways. However, this means that the polymer chains are cleaved and reduced in weight. On the other hand, composting marks a condition when biodegraded material is mineralized into water, carbon dioxide (CO₂), and biomass (Bettas Ardisson et al., 2014). Regarding the environmental impact and management of post-consumer-expired packaging materials, the developments of new user-friendly and fast screening methodologies to evaluate their biodegradation are attractive. Due to its extensive use for food conservation, the packaging waste is continuously growing and compostability of these materials would be preferable, due to return to earth natural components. For this purpose, some standardization procedures are reported (Ncube et al.2021).

Effects of Bioplastics on Soil Health

Bioplastic films are increasingly used in agriculture, notably for mulching. While consumers see bioplastics as a clear way to reduce plastic use in food production, the effects of bioplastic applications on soil health have received much less attention (Chah et al.2022). The importance on both soil composition and function for overall ecosystem functioning and for plant development is highlighted, especially considering that bioplastics may remain in the soil for years. Biopolymers come into contact with biotic and abiotic soil components that can contribute to the degradation or alteration of the polymer in time (Chu et al.2023). Abiotic components include the micro- and macro-nutrients soil is composed of, as well as sand, silt, and clay arrangement or available water, while its biotic components are composed of a wide range of organisms. Different relationships between the components determine a range of physical and chemical properties of the soil, that in time can change, leading to improvements or degradation in its structure (Bandopadhyay et al., 2018).

The changes experienced by biodegradable mulch applied soils and their biological communities over time remain largely understudied. While some studies showed chitinase gene abundance to increase 7 days after application of a bioplastic in comparison to non-plastic soils, microbial biomass did not show significant differences between different types of bioplastic films applied before or after planting over a period of 3 weeks (O'Connor et al.2024). Differences in enzyme activity were not found between biodegradable or conventional fabrics when testing at 60 and 90 days after application. All studies report only a short-term follow-up and evaluate a limited set of chemical and biological soil characteristics. There is therefore a high demand to both compare and offer comprehensive datasets investigation of the changes to the soil and its biological activity due to the presence of biodegradable bioplastic mulch (Juncheed et al., 2022). Two types of biodegradable mulch films were studied. A commercially available poly(butylene succinate-co-adipate) (PBSA), and a bio-based film made of polyhydroxyalkanoate (PHA) that was home-produced and had already been use for 4 months. Infra-red spectroscopy recorded every day during four months allowed to monitor the films alteration over time. Soil analysis, using both physicochemical and microbiological methods were performed weekly during the same period, and compared to a control field without bioplastic mulch (de and Woodhouse2024). The results of the study show that, while the PBSA film began to disintegrate as secondary microplastics already after approximately 25 days, four months of exposure to the field environment were not enough for the film components to completely biodegrade. Moreover, the disintegration of the PBSA film increased the abundance of pathogenic bacteria, primarily of the *Pantoea* genus, when compared to the control and PLA-treated soils. Considering the environmental relevance of the results obtained, this research is of major importance (Aziman et al., 2021).

Impact of Bioplastics on Plant Development

The increased use of bioplastics as an alternative to non-biodegradable plastics in agriculture has gained much attention due to the harmful impacts of the latter on the environment (Juncheed et al., 2022). Despite the gradual acceptance of bioplastics in crop protection, their application below the soil and forsaking growth enhancement is not yet widely adopted. To unpack the current state of knowledge on these matters an investigation through a review of the bioplastics or biodegradable plastics for their chemical, biological and mechanical properties is conducted, elucidating the links to variables influenced during plant growth and development (Nanda et al.2022). This review additionally investigates the influences of bioplastic on the growth rates, vigor, and overall health of the plant, and evaluates the types of bioplastic and application methods to find the potential outcomes. Biomass is one of the crucial factors produced by plant metabolism, including the formation of shoots, leaves, and roots, and the extension of the root system. Few pieces of evidence show the benefits of bioplastics to improve biomass production in various types of crops, especially horticultural crops (Chong et al.2022).

However, the overview strictly unavailable focused on the action of bioplastics to enhance the production of plant organs below the soil. It is also an imperative condition for increasing plant growth because bioplastics are basically made of biodegradable materials and then they can reduce rhizosphere inhibition due to the black or silver streams residuals left in the soil after degradation (Chah et al.2022). The design and development of the biotic systems plays a key role in creating and maintaining good environmental conditions for plant growth (Bandopadhyay et al., 2018). Biodegradable bioplastic acts as a beneficial barrier to maintain moisture near the surface, prevent the evaporation of soil-critical nutritional liquids for plants, and can buffer the soil temperature that the root zone primarily touches the edge with increased space for root formation potential of uptake Nutrients essential for plants (Sintim et al.2021).

Field Studies and Case Examples

Poly (lactic) acid (PLA), poly (3-hydroxybutyrate-co-3-hydroxyvalerate) (EB148) and biodegradable polythene bags are potentially beneficial environmentally friendly alternatives to existing standard polythene mulch films (Bandopadhyay et al., 2018). The recent publication on the impact of biodegradable mulch films on soil health and development of crop plants addresses important issues in the growing field of development and use of biodegradable bioplastic materials (Zhang et al., 2022). Composting of biodegradable materials accumulated in the soil, or complete digestion of these materials by soil microbes (Zhang et al., 2022). The use of bio-based mulch films made from starch might also promote crop plants growth and development due to improved water holding capacity and inhibition of weeds growth (Li et al., 2024). Mulch films used in agriculture have multiple benefits, among others, increasing soil temperature and water retention, reducing weed growth, and in general enhancing yields (de and Woodhouse2024). From an analysis of the results of previous mathematical modelling it follows that the duration of the mulching effect in the soil, as well as its magnitude depends also on the chemical and physical properties of the mulching material (Fan et al., 2023). At present, most mulching films are made from mixtures of polyethylene and calcium carbonate. Nevertheless, the presence of calcium carbonate is disadvantageous also (Zhang et al.2021). Calcium, as a mineral element, is indispensable for the growth of all crop plants, and therefore, it shall be noticed when using plastic mulch films over long periods of time, high calcium concentration in treated soil (Toor et al.2021). Generally, the mulching effect is weaker in the case of bio-mulch films, systematically becomes weaker during plant vegetation, and is stronger under competitiveness conditions between mulch and plant (Fan et al., 2023). This is clearly visible looking at cucurbit vegetables since it is usually very difficult to remove the mulch films after harvesting due to excessive penetration of plant roots into the soil under the film

(Cozzolino et al.2023). However, the presence of bio-mulch films temporarily limited root growth and reduced the intensity of that phenomenon, as evidenced by the smaller dry weight and absorbed amount of calcium per plant (Kumari et al.2025). Other mathematical modeling of plant growth and development under the influence of bio-based mulch materials also indicate an increase in the yield of tomatoes by 50% or an increase in the water index by 100% in relation to control conditions (Morra et al.2021). During two years of experiments, all 12-micron thick mulch films had a beneficial effect on the development of tomato plants. Diversification of mulch films did not affect in general leaf water potential Ψ while bio-mulch films made from starch increase Ψ menthols relative to other control and standard mulch films. Concluding pat of the previous paper, analyses in recent years of the impact of biodegradable materials on soil health and plant development showed the complexity of processes in the interface of organic and mineral soil particles, which are still far from being fully understood (Li et al.2022). The impact of these materials is conditioned not only by their physical and chemical properties, but also by the environment in which they are located, and on which they affect after spreading. Based on the research results obtained, it should be emphasized that excessive use of these materials in the soil and their lack of composting contributes to the deterioration of both soil health and plant development (Pahalvi et al.2021). In adverse circumstances, on the surface of such material as on the source of nutrients, can develop a complex microbial flora, which, after subsequent water supply by leaching, can have a negative effect on plant development (Juncheed et al., 2022). Data also pointed out that the presence in the medium of a porous material, characterized by high water absorption and poor degradability, will slow down the re-mineralization of organic substances, preventing their appropriate transmission to the soil by surface runoff (Li et al.2022). Differences should be know in the light of rapid global changes related to the widespread use of polythene mulch films, their degradation in the soil or their incineration. Due to global warming, increases in CO₂ concentration, theoretical modeling, attempts are made to replace conventional mulch films with bio-based and biodegradable materials. Made from natural substances, intending to completely disappear in the soil after the period of their remnants (Mansoor et al.2022). What's more, this is a chance of high performance compost that will be a beneficial substitute for mineral fertilization in sustainable agriculture. However, advantages of the use of the bio-film surface have indicated that after its degradation new physical and chemical properties are developing at a faster rate on the mulch film surface, compared to the remaining materials that regulate the soil conditions (Tian et al.2023). This confirms in part the conducted research, because after the combined addition to the soil of the comparable doses of mulch film and starch, for a long time in both substrates, there was excess of concentration of the employed material. On the other hand, despite such treatments, equivalent concentrations of such material in the whole soil profile did not appear. In modeling experiments, it has also been shown that both ephemeral materials, located on the surface, and lying longitudinally are the source of organic substances washed away by leaching (Siedt et al.2021). There are indications that after two or three years of soil cultivation, organo-mineral particles might decrease, which is the source of humus formation, poor activation of the new pathway humification-degradation by polyethylene, and continuous addition of less biodegradable fragments non-sourced the regular formation of new humus (Li et al.2023). Emphasis was placed not only on the increase in the amount of some potentially toxic elements in the plant, but on the negative effect of this fact on the symmetry of the Ca: Mg ratio, which is of key importance for all crop plants. Adsorption, the entrance of nutrients from the soil solution, and their transport in the plant depend largely on this issue (Xie et al., 2021). Other research illustrating the present complex problem is assessment of the phytotoxicity of model polyethylene materials from interference with various substances. In that work, it was stated that conducted research has shown that all of these supplements are working interference with fresh material, it is important to stress the need for regulated introduction of polymers with dyes and pigments to agricultural production (Yang et al.2021).

Future Directions and Recommendations

Since the advent of plastic in agriculture, it has been widely used in the form of mulch films, tunnel houses, containers, and protective nets, primarily for enhancing crop productivity and product quality. However, continuous plastic use leads to residual plastic films remaining in soils and the environment (Zhao et al.2021). The introduction of biodegradable polymers for specific agricultural applications is a promising approach to addressing long-term environmental and soil health concerns associated with plastic waste accumulation in the agricultural setting, while also providing short-term agronomic benefits similar to those of conventional plastics (Mansoor et al.2022). Biologically-derived polymeric materials, commonly known as bioplastics, are typically more environmentally sustainable than petroleum-based ones. A global assessment of the bioplastic industry in agriculture is presented here for the first time by synthesizing the findings of current studies and research articles (Singh et al., 2022). An overview of the current state of the art of bioplastics in agriculture is provided, exploring the impacts on soil health and promotion of plant development, and discussing the existing gaps in knowledge and understanding (Filipe et al., 2023). Future directions and recommendations for the use of bioplastics in agriculture are outlined to encourage continued innovation, interdisciplinary efforts, and broader policy development to facilitate the transition towards a sustainable agricultural sector.

Conclusion and Summary

Global plastic use increased by 20 times in the last fifty years up to 331 million tons in 2013, and is expected to double again within the following 20 years. The use of plastics is closely related to the development of modern agriculture and food production (M. S. Cruz et al., 2022). They are utilized as mulch films to increase ground temperature, prevent weed growth, maintain moisture in soil, and have a pivotal role in crop protection against pests (El-Beltagi et al.2022). However, it is becoming increasingly evident that this type of conventional plastic usage has various drawbacks and enters the environment in the form of micro- or nanoplastics, which are difficult to degrade and pose risks to both soil health and food safety (Lakhiar et al.2024).

The agricultural sector is forced to seek sustainable alternatives for plastics. However, these alternatives cannot be found by just switching to bio-based plastics either (Mendes & Pedersen, 2021). The effects on soil health and plant development, in which microplastics play a role not only as mechanical but also as carrier of residual chemicals, drugs, or microbes, should be studied in detail to be able to implement new agricultural best practices without future problems commonly found in conventional plastics usage (Dong et al.2021). There is no standard definition of bioplastics in terms of biodegradability or bio-based content, and therefore, the use of these definitions is problematic in developing both regulations and industry standards. Since the effects are often related to enhanced microbial activity around the particles, research should consider the characteristics of the chemicals, drugs, or microbe strains carried by the bioplastics (Lackner et al., 2023). Fillers, additives, or organic layers with potential hazard characteristics transferred to the environment may also cause significant changes. Therefore, the studies should consider both the particles themselves and their associated side effects. Additionally, long-term agronomic approaches, rather than standard tests, would be more beneficial in terms of actual agricultural applications (Turner & Filella, 2021).

References

M. S. Cruz, R., Krauter, V., Krauter, S., Agriopoulou, S., Weinrich, R., Herbes, C., B. V. Scholten, P., Uysal-Unalan, I., Sogut, E., Kopacic, S., Lahti, J., Rutkaite, R., & Varzakas, T.

(2022). Bioplastics for Food Packaging: Environmental Impact, Trends and Regulatory Aspects. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)

Manabu Abe, M., Ribeiro Martins, J., Bertolino Sanvezzo, P., Vitor Macedo, J., Cristina Branciforti, M., Halley, P., Roberto Botaro, V., & Brienzo, M. (2021). Advantages and Disadvantages of Bioplastics Production from Starch and Lignocellulosic Components. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)

Moshood, T. D., Nawanir, G., Mahmud, F., Mohamad, F., Ahmad, M. H., & AbdulGhani, A. (2022). Sustainability of biodegradable plastics: New problem or solution to solve the global plastic pollution?. *Current Research in Green and Sustainable Chemistry*, 5, 100273. [sciencedirect.com](https://www.sciencedirect.com/)

Ibrahim, N. I., Shahar, F. S., Sultan, M. T. H., Shah, A. U. M., Safri, S. N. A., & Mat Yazik, M. H. (2021). Overview of bioplastic introduction and its applications in product packaging. *Coatings*, 11(11), 1423. [mdpi.com](https://www.mdpi.com/)

Fan, P., Yu, H., Xi, B., & Tan, W. (2022). A review on the occurrence and influence of biodegradable microplastics in soil ecosystems: are biodegradable plastics substitute or threat?. *Environment International*. [sciencedirect.com](https://www.sciencedirect.com/)

Filipe, S., Mira Mourão, P., Couto, N., & Tranchida, D. (2023). Towards a Sustainable Future: Advancing an Integrated Approach for the Recycling and Valorization of Agricultural Plastics. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)

Choe, S., Kim, Y., Won, Y., & Myung, J. (2021). Bridging Three Gaps in Biodegradable Plastics: Misconceptions and Truths About Biodegradation. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)

Otoni, C. G., Azeredo, H. M., Mattos, B. D., Beaumont, M., Correa, D. S., & Rojas, O. J. (2021). The food–materials nexus: next generation bioplastics and advanced materials from agri-food residues. *Advanced Materials*, 33(43), 2102520. [wiley.com](https://www.wiley.com/)

Mujtaba, M., Fraceto, L. F., Fazeli, M., Mukherjee, S., Savassa, S. M., de Medeiros, G. A., ... & Vilaplana, F. (2023). Lignocellulosic biomass from agricultural waste to the circular economy: a review with focus on biofuels, biocomposites and bioplastics. *Journal of Cleaner Production*, 402, 136815. [sciencedirect.com](https://www.sciencedirect.com/)

Ghosh, K. & Jones, B. H. (2021). Roadmap to biodegradable plastics—current state and research needs. *ACS Sustainable Chemistry & Engineering*. [acs.org](https://www.acs.org/)

Bettas Ardisson, G., Tosin, M., Barbale, M., & Degli-Innocenti, F. (2014). Biodegradation of plastics in soil and effects on nitrification activity. A laboratory approach. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)

Ncube, L. K., Ude, A. U., Ogunmuyiwa, E. N., Zulkifli, R., & Beas, I. N. (2021). An overview of plastic waste generation and management in food packaging industries. *Recycling*, 6(1), 12. [mdpi.com](https://www.mdpi.com/)

Chah, C. N., Banerjee, A., Gadi, V. K., Sekharan, S., & Katiyar, V. (2022). A systematic review on bioplastic-soil interaction: Exploring the effects of residual bioplastics on the soil geoenvironment. *Science of the Total Environment*, 851, 158311. [\[HTML\]](#)

Chu, J., Zhou, J., Wang, Y., Jones, D. L., Ge, J., Yang, Y., ... & Zeng, Z. (2023). Field application of biodegradable microplastics has no significant effect on plant and soil health in the short term. *Environmental Pollution*, 316, 120556. [sciencedirect.com](https://www.sciencedirect.com/)

Bandopadhyay, S., Martin-Closas, L., M. Pelacho, A., & M. DeBruyn, J. (2018). Biodegradable Plastic Mulch Films: Impacts on Soil Microbial Communities and Ecosystem Functions. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)

- O'Connor, J., Mickan, B. S., Gurung, S. K., Bühlmann, C. H., Jenkins, S. N., Siddique, K. H., ... & Bolan, N. S. (2024). Value of food waste-derived fertilisers on soil chemistry, microbial function and crop productivity. *Applied Soil Ecology*, 198, 105380. [sciencedirect.com](https://www.sciencedirect.com)
- Juncheed, K., Tanunchai, B., Fareed Mohamed Wahdan, S., Thongsuk, K., Schädler, M., Noll, M., & Purahong, W. (2022). Dark side of a bio-based and biodegradable plastic? Assessment of pathogenic microbes associated with poly(butylene succinate-co-adipate) under ambient and future climates using next-generation sequencing. [ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov)
- de Sadeleer, I., & Woodhouse, A. (2024). Environmental impact of biodegradable and non-biodegradable agricultural mulch film: A case study for Nordic conditions. *The International Journal of Life Cycle Assessment*, 29(2), 275-290. [springer.com](https://www.springer.com)
- Aziman, N., Kian, L. K., Jawaid, M., Sanny, M., & Alamery, S. (2021). Morphological, structural, thermal, permeability, and antimicrobial activity of PBS and PBS/TPS films incorporated with biomaster-silver for food packaging application. *Polymers*. [mdpi.com](https://www.mdpi.com)
- Nanda, S., Patra, B. R., Patel, R., Bakos, J., & Dalai, A. K. (2022). Innovations in applications and prospects of bioplastics and biopolymers: A review. *Environmental Chemistry Letters*, 20(1), 379-395. [springer.com](https://www.springer.com)
- Chong, J. W. R., Tan, X., Khoo, K. S., Ng, H. S., Jonglertjunya, W., Yew, G. Y., & Show, P. L. (2022). Microalgae-based bioplastics: future solution towards mitigation of plastic wastes. *Environmental research*, 206, 112620. [\[HTML\]](#)
- Sintim, H. Y., Bandopadhyay, S., English, M. E., Bary, A., y González, J. E. L., DeBruyn, J. M., ... & Flury, M. (2021). Four years of continuous use of soil-biodegradable plastic mulch: impact on soil and groundwater quality. *Geoderma*, 381, 114665. [sciencedirect.com](https://www.sciencedirect.com)
- Zhang, M., Xue, Y., Jin, T., Zhang, K., Li, Z., Sun, C., Mi, Q., & Li, Q. (2022). Effect of long-term biodegradable film mulch on soil physicochemical and microbial properties. *Toxics*. [mdpi.com](https://www.mdpi.com)
- Li, X., Zheng, G., Li, Z., & Fu, P. (2024). Formulation, performance and environmental/agricultural benefit analysis of biomass-based biodegradable mulch films: A review. *European Polymer Journal*. [\[HTML\]](#)
- Fan, D., Jia, G., Wang, Y., & Yu, X. (2023). The effectiveness of mulching practices on water erosion control: A global meta-analysis. *Geoderma*. [sciencedirect.com](https://www.sciencedirect.com)
- Zhang, T., Zhang, C., Yang, Y., Yang, F., Zhao, M., & Weng, Y. (2021). Improved properties of poly (butylene adipate-co-terephthalate)/calcium carbonate films through silane modification. *Journal of Applied Polymer Science*, 138(38), 50970. [researchgate.net](https://www.researchgate.net)
- Toor, M. D., Adnan, M., Rehman, F. U., Tahir, R., Saeed, M. S., Khan, A. U., & Pareek, V. (2021). Nutrients and their importance in agriculture crop production; A review. *Ind. J. Pure App. Biosci*, 9(1), 1-6. [researchgate.net](https://www.researchgate.net)
- Cozzolino, E., Di Mola, I., Ottaiano, L., Bilotto, M., Petriccione, M., Ferrara, E., ... & Morra, L. (2023). Assessing yield and quality of melon (*Cucumis melo* L.) improved by biodegradable mulching film. *Plants*, 12(1), 219. [mdpi.com](https://www.mdpi.com)
- Kumari, D., Janmeda, P., Varshney, N., & Pandey, P. (2025). Effect of Micro/Nano-Plastics Accumulation on Soil Nutrient Cycling. In *Global Impacts of Micro-and Nano-Plastic Pollution* (pp. 179-204). IGI Global. [researchgate.net](https://www.researchgate.net)
- Morra, L., Cozzolino, E., Salluzzo, A., Modestia, F., Bilotto, M., Baiano, S., & del Piano, L. (2021). Plant growth, yields and fruit quality of processing tomato (*Solanum lycopersicon* L.)

- as affected by the combination of biodegradable mulching and digestate. *Agronomy*, 11(1), 100. [mdpi.com](https://doi.org/10.3390/agr11010100)
- Li, Q., Wang, Y., Li, Y., Li, L., Tang, M., Hu, W., ... & Ai, S. (2022). Speciation of heavy metals in soils and their immobilization at micro-scale interfaces among diverse soil components. *Science of the Total Environment*, 825, 153862. [sciencedirect.com](https://doi.org/10.1016/j.scitotenv.2022.153862)
- Pahalvi, H. N., Rafiya, L., Rashid, S., Nisar, B., & Kamili, A. N. (2021). Chemical fertilizers and their impact on soil health. *Microbiota and Biofertilizers*, Vol 2: Ecofriendly tools for reclamation of degraded soil environs, 1-20. [\[HTML\]](https://doi.org/10.1007/978-981-16-1444-4_1)
- Li, S., Yu, X., Dang, X., Wang, P., Meng, X., Wang, Q., & Hou, H. (2022). Double dielectric barrier discharge incorporated with CeO₂-Co₃O₄/γ-Al₂O₃ catalyst for toluene abatement by a sequential adsorption–discharge plasma catalytic process. *Journal of Cleaner Production*, 340, 130774. [\[HTML\]](https://doi.org/10.1016/j.jclepro.2022.130774)
- Mansoor, Z., Tchuenbou-Magaia, F., Kowalczyk, M., Adamus, G., Manning, G., Parati, M., ... & Khan, H. (2022). Polymers use as mulch films in agriculture—a review of history, problems and current trends. *Polymers*, 14(23), 5062. [mdpi.com](https://doi.org/10.3390/polym14235062)
- Tian, D., Zhang, J., Hu, J., Huang, M., Zhao, L., Lei, Y., ... & Shen, F. (2023). A new water-soluble lignin incorporation enhanced the barrier performance of liquid mulching film. *Chemical Engineering Journal*, 452, 139383. [\[HTML\]](https://doi.org/10.1016/j.cej.2023.139383)
- Siedt, M., Schäffer, A., Smith, K. E., Nabel, M., Roß-Nickoll, M., & Van Dongen, J. T. (2021). Comparing straw, compost, and biochar regarding their suitability as agricultural soil amendments to affect soil structure, nutrient leaching, microbial communities, and the fate of pesticides. *Science of the Total Environment*, 751, 141607. [sciencedirect.com](https://doi.org/10.1016/j.scitotenv.2021.141607)
- Li, H., Zhou, Y., Mei, H., Li, J., Chen, X., Huang, Q., ... & Tang, J. (2023). Effects of long-term application of earthworm bio-organic fertilization technology on soil quality and organo-mineral complex in tea garden. *Forests*, 14(2), 225. [mdpi.com](https://doi.org/10.3390/forests14020225)
- Xie, K., Cakmak, I., Wang, S., Zhang, F., & Guo, S. (2021). Synergistic and antagonistic interactions between potassium and magnesium in higher plants. *The Crop Journal*. [sciencedirect.com](https://doi.org/10.1016/j.cropro.2021.100000)
- Yang, M., Huang, D. Y., Tian, Y. B., Zhu, Q. H., Zhang, Q., Zhu, H. H., & Xu, C. (2021). Influences of different source microplastics with different particle sizes and application rates on soil properties and growth of Chinese cabbage (*Brassica chinensis* L.). *Ecotoxicology and Environmental Safety*, 222, 112480. [sciencedirect.com](https://doi.org/10.1016/j.ecoenv.2021.112480)
- Zhao, Z. Y., Wang, P. Y., Wang, Y. B., Zhou, R., Koskei, K., Munyasya, A. N., ... & Xiong, Y. C. (2021). Fate of plastic film residues in agro-ecosystem and its effects on aggregate-associated soil carbon and nitrogen stocks. *Journal of Hazardous Materials*, 416, 125954. [\[HTML\]](https://doi.org/10.1016/j.jhazmat.2021.125954)
- Singh, N., Ogunseitan, O. A., Wong, M. H., & Tang, Y. (2022). Sustainable materials alternative to petrochemical plastics pollution: A review analysis. *Sustainable horizons*. [sciencedirect.com](https://doi.org/10.1016/j.sushorz.2022.100000)
- El-Beltagi, H. S., Basit, A., Mohamed, H. I., Ali, I., Ullah, S., Kamel, E. A., ... & Ghazzawy, H. S. (2022). Mulching as a sustainable water and soil saving practice in agriculture: A review. *Agronomy*, 12(8), 1881. [mdpi.com](https://doi.org/10.3390/agr12081881)
- Lakhiar, I. A., Yan, H., Zhang, J., Wang, G., Deng, S., Bao, R., ... & Wang, X. (2024). Plastic pollution in agriculture as a threat to food security, the ecosystem, and the environment: an overview. *Agronomy*, 14(3), 548. [mdpi.com](https://doi.org/10.3390/agr14030548)

- Mendes, A. C. & Pedersen, G. A. (2021). Perspectives on sustainable food packaging:—is bio-based plastics a solution?. Trends in Food Science & Technology. [dtu.dk](https://doi.org/10.1016/j.tfs.2021.05.001)
- Dong, H., Chen, Y., Wang, J., Zhang, Y., Zhang, P., Li, X., ... & Zhou, A. (2021). Interactions of microplastics and antibiotic resistance genes and their effects on the aquaculture environments. Journal of hazardous materials, 403, 123961. [\[HTML\]](https://doi.org/10.1016/j.jhazmat.2021.123961)
- Lackner, M., Mukherjee, A., & Koller, M. (2023). What are “bioplastics”? Defining renewability, biosynthesis, biodegradability, and biocompatibility. Polymers. [mdpi.com](https://doi.org/10.3390/polym15010011)
- Turner, A. & Filella, M. (2021). Hazardous metal additives in plastics and their environmental impacts. Environment International. [sciencedirect.com](https://doi.org/10.1016/j.envint.2021.106500)

ABIOTIC STRESS FACTORS AND THEIR EFFECTS IN STONE FRUIT ORCHARDS OF AMIK PLAIN

Dr. Aysun UYSAL

Hatay Mustafa Kemal Üniversitesi, Bitki Sağlığı Kliniği Uygulama ve Araştırma Merkezi

Prof. Dr. Şener KURT

Hatay Mustafa Kemal Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü-Bitki Sağlığı Kliniği Uygulama ve Araştırma Merkezi

Doktora öğrencisi Tuğba HANEDAN

Hatay Mustafa Kemal Üniversitesi, Ziraat Fakültesi Bitki Koruma Bölümü

ÖZET

Sert çekirdekli meyve ağaçlarında gözlenen abiyotik stres faktörleri, bitkilerin büyüme ve gelişimini olumsuz etkileyen çevresel koşullardır. Aşırı-düşük sıcaklıkların etkileri, kuraklık – fazla su oluşumu, aşırı toprak tuzluluğu, besin maddesi dengesizliği, yetersiz-aşırı ışık, şiddetli rüzgar, toprak Ph değerinin yüksek- düşük olması, gece gündüz sıcaklık farkları, yabancı ot ilaçları gibi faktörler yer alır. Abiyotik stres faktörleri, genel olarak ağaçlarda büyüme geriliği, çiçeklenme ve meyve tutumunda azalma, kök sorunları görülür. Yapraklarda sararma, solma, kıvrılma, kuruma, lekeler ve yanıklar; meyvelerde küçülme, çatlama, deformasyon, düşük kalite ve erken dökülme gibi belirtiler ortaya çıkar. Aşırı ışık, güneş yanıkları ve meyvelerde lekeler; yüksek sıcaklık ise meyve yanıkları, yapraklarda kuruma-solma, fotosentez oranının düşmesi ve çiçeklerin erken dökülmesi gibi belirtilere neden olur. Bu faktörler arasında ön görülemeyen ve göz ardı edilenler gece ve gündüz arasındaki sıcaklık farkları ve total herbisit zararlarıdır. 2020-2024 yılları arasında Amik Ovası'nda yetiştiriciliği yapılan kayısı, şeftali ve nektarin bitkilerinde gözlenen belirtiler üzerine sörvey çalışmaları yapılmıştır. Her yıl ortalama 8 adet şeftali, 10 adet nektarin ve 7 adet kayısı olmak üzere 25 farklı bahçede farklı belirtilerden örneklemeler yapılmıştır. Hatay Mustafa Kemal Üniversitesi Bitki Sağlığı Kliniği Uygulama ve Araştırma Merkezi laboratuvarlarında yapılan hastalık analizleri sonucunda herhangi bir hastalık etmeni saptanmamıştır. Bu belirtilerin biyotik bir etken yerine abiyotik faktörlerin etkili olduğu kanısına varılmıştır. Yapılan bazı araştırmalar sonrası yanlış kimyasal ilaç kullanımı, doz aşımı aşırı gübreleme yapıldığı gözlenmiştir. Özellikle son zamanlarda yaşanan aşırı sıcaklar, gece-gündüz sıcaklık farkları ve aşırı total herbisit kullanımları bitkilerde ciddi sorunlara neden olmaktadır.

Anahtar kelimeler: Amik ovası, sert çekirdekli, abiyotik stres, iklim değişikliği

ABSTRACT

Abiotic stress factors observed in stone fruit trees are environmental conditions that negatively affect the growth and development of plants. These include the effects of extreme-low temperatures, drought - excess water formation, excessive soil salinity, nutrient imbalance,

insufficient-excessive light, strong wind, high-low soil pH value, day and night temperature differences, weed pesticides. Abiotic stress factors generally cause growth retardation in trees, decrease in flowering and fruit set, root problems. Symptoms such as yellowing, wilting, curling, curling, drying, spots and burns on leaves; shrinkage, cracking, deformation, low quality and early shedding of fruits. Excessive light causes sunburns and spots on fruits; high temperature causes symptoms such as fruit burns, drying and wilting of leaves, decreased photosynthesis rate and early shedding of flowers. Among these factors, the unpredictable and ignored ones are the temperature differences between day and night and total herbicide damage. Survey studies were carried out on the symptoms observed in apricot, peach and nectarine plants cultivated in Amik Plain between 2020-2024. Each year, 25 different orchards were sampled for different symptoms, with an average of 8 peach, 10 nectarine and 7 apricot orchards. No disease agent was detected as a result of disease analyses carried out in the laboratories of Centre for Implementation and Research of Plant Health Clinic, Hatay Mustafa Kemal University. It was concluded that these symptoms were caused by abiotic factors instead of biotic factors. After some research, it has been observed that incorrect use of chemical pesticides, overdose and excessive fertilisation. Especially the recent extreme heat, day-night temperature differences and excessive use of total herbicides cause serious problems in plants.

Key Words: Amik plain, stone fruit, abiotic stress, climate change

GİRİŞ

Ülkemiz, ılıman iklim koşullarına sahip olması ve coğrafi konumu sayesinde sert çekirdekli meyve yetiştiriciliği için elverişli bir zemine sahiptir. 2024 yılı TÜİK verilerine göre, Türkiye'de sert çekirdekli meyve bahçelerinin toplam alanı 2.521.791,228 dekar olup, bu alandan ise toplamda 3.757.224 ton meyve üretimi gerçekleştirilmiştir (TÜİK, 2024).

Hatay, yüzölçümünün %33,5'ini oluşturan geniş ova alanlarıyla tarıma elverişli ve verimli topraklara sahip olup, sulama imkanlarıyla birleşerek meyve yetiştiriciliği gibi tarımsal faaliyetlerin ön planda olmasını sağlamaktadır. Bu nedenle, sert çekirdekli meyve üretimi gibi alanlar özellikle ova bölgelerinde yoğunlaşmış ve büyük bir üretim potansiyeli taşımaktadır. 2023 yılı itibarıyla Hatay, kayısı (37.034 ton), kiraz (671 ton), şeftali ve nektarin (8.941 ton), vişne (4 ton) ve zerdali (100 ton) üretimiyle ülke genelinde öne çıkmakta, özellikle kayısı üretimindeki yüksek verimiyle lider konumda bulunarak tarımsal alanda Türkiye ekonomisine önemli katkılar sunmaktadır (TÜİK, 2024). Ancak, bu avantajlara rağmen, sert çekirdekli meyve üretimini olumsuz etkileyen ve kalite ile verim kayıplarına yol açan bir dizi biyotik ve abiyotik faktörler bulunmaktadır. Ağaçlarda meydana gelen kuruma ve geriye doğru ölümler çoğunlukla biyotik ve abiyotik stres faktörlerinin etkisiyle oluşur. Abiyotik stres faktörleri, kuraklık, ekstrem sıcaklıklar, toprak yapısındaki bozukluklar, drenaj problemleri ve tuzluluk gibi etkenleri kapsarken, biyotik faktörler ise özellikle fungal hastalıklarla ilişkilidir (Kaymaz, 2017).

Tarımsal üretimde biyotik faktörler, bitkilerin sağlığını etkileyen canlı organizmalardır ve bu faktörler, özellikle zararlılar, hastalıklar, yabancı otlar ve parazitler, verim kayıplarına yol açabilir. Zararlılar bitkilerin yapısını tahrip ederken, mantar, bakteri ve virüsler gibi hastalıklar büyümeyi engeller ve ürün kaybına neden olmaktadır. Özellikle fungal hastalıklarla ilişkili olan biyotik faktörler arasında sert çekirdekli meyvelerde görülen bazı önemli fungal hastalıklar şunlardır: Kayısı, şeftali, kiraz, erik, vişne ve bademde önemli kayıplara yol açabilen çiçek Monilyası (*Monilinia laxa*), Sitospora kanseri (*Cytospora* sp.), Yaprakdelen ve meyvede çil hastalığı (*Wilsonomyces carpophilus*), Şeftali karalekesi (*Cladosporium carpophilum*), Şeftali küllemesi (*Sphaerotheca pannosa*), Yaprak kıvrıkcılığı hastalığı (*Taphrina deformans*), Erik pası hastalığı (*Tranzschelia discolor*), Eriklerde cep hastalığı (*Taphrina pruni*) ve Kiraz yaprak lekesi (*Blumeriella jaapii*). Bu hastalıklar, sert çekirdekli meyve yetiştiriciliğinde önemli verim

kayıplarına neden olabilmektedir. Bunun yanı sıra son yıllarda küresel ısınmanın etkisiyle, Botryosphaeriaceae familyasına ait hastalık etmenlerinin dünya genelinde ve Türkiye'de artış gösterdiği dikkat çekmektedir. Bu familya, kutup bölgeleri dışında tüm coğrafi bölgelerde bulunmakta ve birçok meyve ağaçlarında ağaç gövdesi ve budanmış dallarda meyve ile sürgünlerde yanma, sakız oluşumu, odun dokusunda renk değişimleri ve dallarla sürgünlerde geriye doğru ölüm gibi belirtiler göstererek (Michailides ve ark. 2015), kiraz, şeftali gibi sert çekirdekli meyve ağaçlarında da fungal hastalık etmeni olarak yer almaktadır. Tüm mücadele yöntemleri uygulanmasına rağmen, hastalıkların bazen abiyotik faktörlerden kaynaklanması, tarımda beklenen verimi kötü yönde etkileyebilir; bu faktörler, bitkilerin büyüme ve gelişme süreçlerini zorlaştırarak, biyotik etmenler kadar önemli bir tehdit oluşturur.

Abiyotik problemler, çevresel koşullardaki değişikliklerden kaynaklanan, bitkilerin sağlığını ve gelişimini olumsuz yönde etkileyen önemli faktörlerdir. Abiyotik bozukluklar, canlı organizmaların (virüsler, bakteriler, mantarlar, böcekler vb.) neden olduğu hastalıklardan farklı olarak, cansız etmenlerden kaynaklanan ve bitkilerin normal büyüme düzeninden sapmalarına yol açan fizyolojik durumlardır. Bu tür bozukluklar, bitkilerde fiziksel veya kimyasal değişiklikler meydana getirerek, normal gelişimden sapmalara neden olur. Enfeksiyon dışı bu bozukluklar bazen kolayca tanımlanabilirken, diğer durumlar daha zor veya imkansız olabilir ve çoğu zaman geri dönüşümsüzdür. Ayrıca tarımsal üretimi olumsuz etkileyerek bitkisel verim kayıplarına sebep olmakta ve verimi %50'ye kadar azaltmaktadır (Wang ve ark., 2004; Ashraf & Foolad, 2007). Subtropik iklim koşullarında, yetersiz soğuklama ve yaz aylarındaki 35-45°C'lik sıcaklıklar, çift meyve oluşumunu artırarak meyve verimi ve kalitesini olumsuz etkiler. Bu durum, özellikle su stresi ve yüksek sıcaklıkların etkisiyle şeftali, erik, kiraz ve kayısı gibi meyvelerde görülür ve genellikle Haziran ile Ağustos arasında daha belirgin hale gelir (Küden, 2004; Naor, 2006; İmrak ve ark., 2014). Sert çekirdekli meyve ağaçlarında gözlenen abiyotik stres faktörleri, bitkilerin büyüme ve gelişimini olumsuz yönde etkilerken bu faktörler arasında düşük sıcaklık etkileri, kuraklık – fazla su oluşumu, aşırı toprak tuzluluğu, besin maddesi dengesizliği, yetersiz-aşırı ışık, şiddetli rüzgar, toprak Ph değerinin yüksek-düşük olması, gece gündüz sıcaklık farkları, yabancı ot ilaçları gibi faktörler yer alır. Sıcaklık değişimleri meyve gelişimini doğrudan etkilemekte, özellikle aşırı sıcaklar çiçek dökülmesine, küçük ve şekilsiz meyve oluşumuna neden olmaktadır. Bunun yanı sıra, düşük sıcaklıklar meyve ağaçlarında çiçeklenme süreçlerini olumsuz etkileyerek verim kaybına yol açabilmektedir. Şeftali gibi bazı meyve türlerinde, sıcaklık stresi ikiz meyve oluşumuna sebep olurken, aşırı ışık ise güneş yanığına ve fotosentez verimliliğinde azalmaya neden olmaktadır. Sert çekirdekli meyve ağaçlarında, ışığın yetersiz olması yavaş büyüme, sararan yapraklar ve zayıf meyve gelişimine yol açarken, fazla ışık da güneş yanığı, yapraklarda hasar, verimsiz olgunlaşma ve fotosentez verimliliğinde düşüş gibi fizyolojik sorunlara neden olabilir. Bu yüzden, ışık düzeylerinin doğru bir şekilde ayarlanması, bitkilerin sağlıklı gelişimi için kritik öneme sahiptir. Aşırı ışık yoğunluğu ise sert çekirdekli meyve ağaçlarında bir dizi fizyolojik probleme yol açabilir (Sreekumar ve ark., 2014). Fazla güneş ışığına maruz kalan meyvelerde "sunscald" adı verilen güneş yanığı görülebilir, bu da meyve yüzeyinde kahverengileşmeye ve dokusal hasara neden olur. Ayrıca, aşırı ışık, bitkilerde su kaybını artırarak meyve dokusunda kuruma, çatlama ve deformasyona yol açabilir. Meyvelerin fazla ışık nedeniyle aşırı ısınması, olgunlaşma sürecini olumsuz etkileyebilir. Bunun yanı sıra, aşırı ışık klorofil yıkımına neden olur ve yaprakların sararmasına yol açarak fotosentez verimliliğini düşürür. En nihayetinde, yüksek ışık yoğunluğu, fotosentez mekanizmalarını bozar ve bitkilerin genel sağlığı ve verimliliğini olumsuz yönde etkiler. (Singh ve ark., 2018)

Kuraklık ve su stresinin giderek artan bir problem haline gelmesi, özellikle sıcaklık dalgalanmaları ile birleşerek meyve kalitesini düşürmekte ve erken yaprak dökülmelerine yol açmaktadır. Toprak tuzluluğu, özellikle sodyum (Na) ve klor (Cl) elementlerinin yüksek konsantrasyonlarda bulunmasıyla sert çekirdekli meyve ağaçlarında besin alımını olumsuz

etkileyerek büyüme problemleri yaratmaktadır. Besin elementi eksiklikleri de meyve gelişimini doğrudan etkileyerek verim kaybına neden olmaktadır. Azot eksikliği yapraklarda küçülmeye ve açık yeşil renge neden olurken, fosfor eksikliği meyvelerin küçük ve ekşi olmasına yol açmaktadır. Potasyum eksikliği ise meyve şeker oranının düşmesine sebep olurken, magnezyum eksikliği yaprak dökülmelerini artırmaktadır (Abacı-bayar ve Boyacı, 2021).

Bunların yanı sıra, rüzgarın meyve ağaçlarında mekanik hasara yol açtığı, bitki dokularında yaralanmalara neden olarak patojen girişini kolaylaştırdığı ve polinasyonu olumsuz etkilediği bilinmektedir (MacKerron, 1976; Pitcairn ve Grace, 1985; Wilson, 1980). Yüksek dozda veya yanlış zamanda uygulanan pestisitler de bitki dokularına zarar vererek gelişim bozukluklarına sebep olabilir. Kimyasal hasar, bazı sert çekirdekli ağaçların meyve kabuğunda kırmızı, sarı veya kahverengi lekeler, yaprak uçlarında kahverengileşme, gelişmemiş veya şekilsiz meyveler, genel kahverengileşme ve hatta bitkinin ölümüne neden olabilir (Sighn ve ark., 2018). Ayrıca, toprak pH dengesizlikleri de bitki besin elementlerinin emilimini etkileyerek gelişim problemlerine neden olmaktadır.

Bu çalışmada, sert çekirdekli meyve yetiştiriciliğinde karşılaşılan ve abiyotik stres faktörlerinin etkileri ele alınarak, bu faktörlerin meyve üretimi üzerindeki etkileri incelenmiştir.

MATERYAL VE YÖNTEM

Bu araştırma, 2020-2024 yılları arasında Amik Ovası'ndaki 8 adet şeftali, 10 adet nektarin ve 7 adet kayısı olmak üzere 25 farklı bahçeden toplanan sert çekirdekli meyve ağaçlarının kök, gövde, dal ve meyve üzerinde yapılan kapsamlı bir inceleme ile gerçekleştirilmiştir.

Fungal izolasyon

Bahçede hastalık belirtilerine benzer belirti gösteren, klorotik veya nekrotik bitki dokularından steril bir bistüri ile dikkatlice kesilmiştir. 1-2 mm büyüklüğündeki bu doku parçaları, yüzey dezenfeksiyonu için %75'lik etanol içinde 1 dakika boyunca bekletilmiştir (Uysal & Kurt, 2019). Ardından, bu parçalar steril distile su ile yıkanmış ve steril kurutma kağıtları üzerinde 15-30 dakika süreyle kurutulmaya bırakılmıştır. Kuruyan dokular, Patates Dekstroz Agar (PDA, Merck KGaA, Darmstadt, Almanya) içeren 90 mm'lik Petri kaplarına yerleştirilmiştir. Bakteriyel kontaminasyonu önlemek amacıyla besi ortamına streptomisin sülfat (100 µg/mL) eklenmiştir. Petri kapları 25°C'de 5 gün boyunca inkübe edilmiştir (Uysal & Kurt, 2019; Kurt ve ark., 2020). Bakteriyel bir hastalık olabilme ihtimaline karşın, bakteri izolasyonu yapılmıştır. Yine aynı şekilde alınan sürgün ve dal örnekleri, %70 etil alkol ile 2 dakika boyunca yüzey sterilizasyonu yapıldıktan sonra, bistüri yardımıyla üst kabuk dokusu kesildi ve hastalıklı iç kısımlar doğrudan King B Agar ve Tryptic soy agar (TSA) (Merck, Darmstadt, Almanya) besi ortamları üzerine yapıldıktan sonra, 25°C'de 24-48 saat inkübe edilmiştir (Aktan ve Soylu, 2020). Bakteri izolasyonları için kullanılan petri kapları, 27°C'de 24-48 saat süreyle inkübe edilmiştir.

BULGULAR ve TARTIŞMA

Bu çalışmada yapılan bakteriyel ve fungal izolasyon işlemleri sonrasında, enfekte olduğu düşünülen bitki örneklerinden hiçbir bitki hastalık etmeni tespit edilmemiştir. PDA, King B Agar ve TSA besi ortamlarında yapılan inkübasyonlar sonucunda, patojen gelişimi gözlenmemiştir. Sadece besi ortamlarında saprofit, endofit, epifit vb. özellikte bakteri ve fungus türleri tespit ve teşhis edilmiştir. Bu durum, abiyotik faktörlerin etkisiyle açıklanabilir. Sıcaklık, kuraklık, su stresi, toprak tuzluluğu, besin element dengesizliği, ışık yetersizliği veya fazlalığı, rüzgar, pestisit kullanımı ve toprak pH'ı, sert çekirdekli meyve ağaçlarının gelişimini ve verimini doğrudan etkileyen çevresel faktörlerdir (Taiz ve ark., 2018; Rodrigo, 2000). Sıcaklık,

fotosentez, solunum, su alımı, terleme, çiçeklenme ve meyve oluşumu gibi süreçleri etkilerken, aşırı sıcaklık çiçek dökülmesi, meyve deformasyonu ve güneş yanığına yol açabilir (Micke ve ark., 1983; Johnson ve ark., 1992). Kuraklık ve su stresi, bitkilerde çeşitli morfolojik ve fizyolojik değişimlere neden olarak meyve kalitesini düşürürken, yüksek sıcaklıklarla birleştiğinde çift meyve oluşumunu artırabilir (Mahajan ve Tuteja, 2005; Örs ve Ekinci, 2015; Küden, 2004; Naor, 2006; İmrak ve ark., 2014). Toprak tuzluluğu, özellikle drenaj eksikliği olan kurak alanlarda belirginleşerek sodyum ve klor birikimi nedeniyle meyve ağaçlarında erken yaprak dökümü, zank salgılanması ve gelişim bozukluklarına yol açar (Çullu, 1999). Besin element dengesizliği, azot, fosfor, potasyum, kalsiyum, magnezyum gibi makro ve mikro elementlerin eksikliği veya fazlalığı nedeniyle bitkilerin sağlığını bozarak verim kaybına neden olabilir (Çolakoğlu ve Çiçekli, 2018; Kacar ve Katkat, 2015; Okur, 2020). Işık yetersizliği, fotosentez verimliliğini azaltarak yavaş büyüme, sararan yapraklar ve düşük kaliteli meyvelere yol açarken, aşırı ışık ise güneş yanığı, yaprak kuruması ve meyve dokusunda bozulmalara sebep olabilir (Sreekumar ve ark., 2014; Singh ve ark., 2018). Rüzgar, bitkilerde fiziksel hasar, polinasyon bozukluğu ve su kaybına neden olurken (MacKerron, 1976; Pitcairn ve Grace, 1985; Wilson, 1980; Newenhouse ve Wilson, 1990), pestisitlerin yanlış dozda veya sıcak saatlerde uygulanması meyve kabuğunda lekelenmelere, şekil bozukluklarına ve yaprak yanıklarına yol açabilir (Singh ve ark., 2018). Son olarak, toprak pH'ı, meyve ağaçlarının besin alımını etkileyerek düşük pH'da manganez toksisitesine, yüksek pH'da ise mikro besin eksikliklerine neden olabilir; optimum pH aralıklarının korunması, sağlıklı büyüme ve yüksek verim için kritik öneme sahiptir (Abacı-Bayar ve Boyacı, 2021).

SONUÇ VE ÖNERİLER

Bitkiler, büyüme süreçleri boyunca çeşitli çevresel zorluklarla karşı karşıya kalır. Bu nedenle, hem abiyotik hem de biyotik stres faktörlerine sürekli olarak uyum sağlamak zorundadırlar. Biyotik stresler, pestisit kullanımı veya çeşitli bitki koruma yöntemleriyle belirli bir seviyeye kadar kontrol altına alınabilir. Ancak, abiyotik streslere maruz kalma, dünya çapında artan gıda talebini karşılamak için tarımda sürdürülebilir büyümeyi engelleyen önemli bir tehdit edici faktör haline gelmiştir. Bu nedenle, sürdürülebilir tarımda abiyotik streslerin etkilerini azaltmak için çeşitli stratejiler uygulanabilir. Öncelikle, kuraklık, tuzluluk ve sıcaklık gibi stres koşullarına dayanıklı bitki türleri seçilmeli ve ıslah çalışmalarıyla bu özelliklere sahip yeni çeşitler geliştirilmelidir. Genetik mühendislik ve biyoteknoloji teknikleri kullanılarak bitkilerin çevresel streslere karşı direnç kazanması sağlanabilir. Toprak sağlığını korumak için organik madde oranı artırılmalı, erozyonu önlemek adına malçlama gibi yöntemler uygulanmalı ve drenaj sistemleri iyileştirilerek tuzluluk sorunu kontrol altına alınmalıdır (Preece and Read, 1993). Etkin su yönetimi için, damlama ve yağmurlama gibi modern sulama sistemleri tercih edilmeli, yağmur suyu toplanarak değerlendirilmelidir. Aynı zamanda, suyun buharlaşmasını azaltan teknikler uygulanarak su kaynaklarının daha verimli kullanılması sağlanmalıdır. Bitkilerin ihtiyaç duyduğu besinleri karşılamak amacıyla dengeli ve bilinçli gübreleme yapılmalı, özellikle potasyum, azot ve kalsiyum gibi minerallerin yeterli düzeyde sağlanmasına özen gösterilmelidir (Waraich ve ark., 2012). Kimyasal gübre kullanımının olumsuz etkilerini en aza indirmek için organik gübre ve biyogübreler tercih edilmelidir (Sneha ve ark., 2018). Nanoteknoloji sayesinde, nanopartiküller kullanılarak bitkilerin besin alımı artırılabilir ve abiyotik stres koşullarına daha iyi uyum sağlanmaları sağlanabilir (Shang ve ark., 2019; Bhatt ve ark., 2020). Toprakta faydalı mikroorganizmaların varlığı desteklenerek bitkilerin stres toleransı artırılabilir ve toprak yapısı iyileştirilebilir. Çevre dostu tarım uygulamaları benimsenerek karbon salınımı düşürülebilir, tarımsal ormancılık gibi yöntemlerle tarım arazilerinde daha uygun mikroiklim koşulları oluşturulabilir. Ayrıca, biyostimülanlar (deniz yosunu özleri, humik asit vb.) kullanılarak bitkilerin stres koşullarında daha dirençli olması sağlanabilir (Pereira ve Correia 2015; Khan ve ark., 2009). Aşırı sıcaklık, düşük sıcaklık ve

kuraklık gibi zorlayıcı çevresel faktörlere karşı uygun tarım teknikleri uygulanarak bitkilerin gelişimi desteklenmelidir. Tüm bu önlemler, sürdürülebilir tarımı teşvik ederek abiyotik streslerin olumsuz etkilerini en aza indirir ve tarımsal üretimin devamlılığını sağlar.

KAYNAKLAR

- Abacı-Bayar, A. A., & Boyacı, S. (2021). Bazı meyve bahçelerinin beslenme durumlarının belirlenmesi. *Türk Tarım ve Doğa Bilimleri Dergisi*, 8(4), 940-950.
- Aktan, Z., & Soylu, S. (2020). Diyarbakır ilinde yetişen badem ağaçlarından endofit ve epifit bakteri türlerinin izolasyonu ve bitki gelişimini teşvik eden mekanizmalarının karakterizasyonu. *KSÜ Tarım ve Doğa Dergisi*, 23(3), 641-654.
- Ashraf, M. F. M. R., & Foolad, M. R. (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environmental and Experimental Botany*, 59(2), 206-216.
- Bhatt, D., Bhatt, M. D., Nath, M., Dudhat, R., Sharma, M., & Bisht, D. S. (2020). Application of nanoparticles in overcoming different environmental stresses. In A. Roychoudhury & D. K. Tripathi (Eds.), *Protective chemical agents in the amelioration of plant abiotic stress: Biochemical and molecular perspectives* (pp. 635-654). Wiley-Blackwell.
- Çolakoğlu, H., & Çiçekli, M. (2018). *Gübreleme rehberi* (Genişletilmiş 2. Baskı). Toros Tarım.
- Çullu, M. A. (1999). GAP'ta tuzlulaşma ve Harran Ovasının durumu. *Toprak tuzlulaşması, Workshop Tema Vakfı Yayınları*, 30, 56-64.
- İmrak, B., Sarier, A., Kuden, A., Kuden, A. B., Comlekcioglu, S., & Tutuncu, M. (2014). Studies on shading system in sweet cherries (*Prunus avium* L.) to prevent double fruit formation under subtropical climatic conditions. *Acta Horticulturae*, 1059, 171-176.
- Johnson, R. S., Handley, D. F., & Dejong, T. M. (1992). Long-term response of early maturing peach trees to postharvest water deficits. *Journal of the American Society for Horticultural Science*, 117(6), 881-886.
- Kacar, B., & Katkat, A. V. (2015). *Bitki besleme*. Nobel Akademik Yayıncılık.
- Khan, W., Rayirath, U. P., Subramanian, S., Jithesh, M. N., Rayorath, P., Hodges, D. M., Critchley, A. T., Craigie, J. S., Norrie, J., & Prithiviraj, B. (2009). Seaweed extracts as biostimulants of plant growth and development. *Journal of Plant Growth Regulation*, 28, 386-399.
- Kaymaz, T. (2017). *Adana ili badem ağaçlarında Botryosphaeriaceae türü patojenlerin belirlenmesi ve moleküler karakterizasyonu* (Yüksek lisans tezi). Çukurova Üniversitesi Fen Bilimleri Enstitüsü, Adana.
- Kuden, A. B. (2004). Global overview of temperate zone fruits in the tropics and subtropics. *Acta Horticulturae*, 662, 37-38.
- Kurt, Ş., Soylu, S., Uysal, A., Soylu, E. M., & Kara, M. (2020). Ceviz gövde kanseri hastalığı etmeni *Botryosphaeria dothidea*'nin tanılanması ve bazı fungusitlerin hastalık etmenine karşı in vitro antifungal etkinliklerinin belirlenmesi. *Mustafa Kemal Üniversitesi Tarım Bilimleri Dergisi*, 25, 46-56.
- MacKerron, D. K. L. (1976). Wind damage to the surface of strawberry leaves. *Annals of Botany*, 40, 351-354.
- Mahajan, S., & Tuteja, N. (2005). Cold, salinity and drought stress: An overview. *Archives of Biochemistry and Biophysics*, 444, 139-158.

- Michailides, T. J. (2015). *Above ground fungal diseases*. Retrieved from
- Micke, W., Doyle, J. F., & Yeager, T. (1983). Doubling potential of sweet cherry cultivars. *California Agriculture*, 37(3-4), 24-25.
- Naor, A. (2006). Irrigation scheduling and evaluation of tree water status in deciduous orchards. *Horticultural Reviews*, 32, 111–165.
- Newenhouse, A. C., & Wilson, S. (1990). Wind stress on young orchard trees. *HortScience*, 25(9), 171.
- Okur, N. (2020). *Toprak bilimi ve bitki besleme*. Nobel Akademik Yayıncılık.
- Örs, S., & Ekinci, M. (2015). Kuraklık stresi ve bitki fizyolojisi. *Derim*, 32(2), 237-250.
- Pereira, L., & Correia, F. (2015). Giriş. In *Costa Portuguesa'da Macroalgas Marinhas: Biyoçeşitlilik, ekoloji ve kullanımlar* (p. 342).
- Preece, J. E., & Read, P. E. (1993). The biology of horticulture. In *Introductory textbook* (pp. 263-269).
- Pitcairn, C. E. R., & Grace, J. (1985). Wind and surface damage. In J. Grace (Ed.), *Progress in biometeorology* (Vol. 2, pp. 115-126). Swets & Zeitlinger.
- Rodrigo, J. (2000). Spring frosts in deciduous fruit trees: Morphological damage and flower hardiness. *Scientia Horticulturae*, 85, 155–173.
- Sneha, S., Anitha, B., Sahair, R. A., Raghu, N., Gopenath, T. S., Chandrashekrappa, G. K., & Basalingappa, K. M. (2018). Biofertilizer for crop production and soil fertility. *Academic Journal of Agricultural Research*, 6(8), 299-306.
- Shang, Y., Hasan, M. K., Ahammed, G. J., Li, M., Yin, H., & Zhou, J. (2019). Applications of nanotechnology in plant growth and crop protection: A review. *Molecules*, 24(14), 2558.
- Singh, N., Singh, G., Thakur, K. K., Kumari, S., & Sharma, D. D. (2018). Causes and remedies of physiological disorders in stone fruit crops. In *Advances in Agriculture Sciences* (Vol. 11, pp. 37-55).
- Taiz, L., Zeiger, E., Møller, I. M., & Murphy, A. S. (2018). *Fundamentals of plant physiology*. Oxford University Press.
- TÜİK. (2025). *Türkiye İstatistik Kurumu (TÜİK) Bitkisel Üretim İstatistikleri*. Retrieved from <http://www.tuik.gov.tr>
- Wang, W., Vinocur, B., Shoseyov, O., & Altman, A. (2004). Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. *Trends in Plant Science*, 9(5), 244-252.
- Wilson, J. (1980). Macroscopic features of wind damage to leaves of *Acer pseudoplatanus* L. *Annals of Botany*, 46, 303-311.

BIOPOLYMER PRODUCTION FROM AGRICULTURAL WASTE: CIRCULAR ECONOMY AND SUSTAINABILITY

Res. Asst. Alperay ALTIKAT

Iğdır University, Faculty of Agriculture Department of Biosystems Engineering, Iğdir,

Prof. Dr. Mehmet Hakkı ALMA

Iğdır University, Faculty of Agriculture Department of Biosystems Engineering, Iğdir,

Abstract

The escalating accumulation of agricultural waste poses significant environmental and economic challenges, necessitating innovative approaches within a circular economy framework. Biopolymer production from agricultural waste emerges as a sustainable solution, aligning closely with principles of zero waste and resource recovery. This study reviews the potential for utilizing abundant agricultural residues and byproducts—such as cereal straw, olive husk, grape stalk, almond, and hazelnut shells—as feedstocks for biopolymer synthesis. Biopolymers derived from these waste sources offer an eco-friendly alternative to petroleum-based plastics, significantly mitigating environmental impacts through enhanced biodegradability and reduced carbon footprints. Highlighting the essential role of biopolymer technologies, the review details how proteins and polysaccharides extracted from agri-food waste can be transformed into biodegradable films, coatings, and mulching materials beneficial in agriculture. Such applications can significantly enhance soil quality, moisture retention, and weed management, contributing positively to plant growth and yield. Despite these advantages, the widespread adoption of biopolymers faces challenges related to the variability in feedstock quality, scalability of production technologies, and market competitiveness against traditional plastics. To overcome these barriers, further advancements in biorefinery technologies and solid-state fermentation processes are recommended. Real-world applications reviewed indicate promising results, demonstrating both environmental and agricultural benefits through the deployment of biopolymer materials. However, critical gaps in the literature remain, particularly regarding long-term biodegradation, economic viability, and consumer acceptance. Addressing these gaps requires interdisciplinary collaboration, strengthened regulatory frameworks, and increased consumer education to fully realize the potential of biopolymers in sustainable agriculture. Ultimately, the integration of biopolymer technologies derived from agricultural waste represents a critical step toward achieving a resilient, sustainable, and circular agricultural economy.

Keywords

Biopolymers, Agricultural Waste, Circular Economy, Sustainability, Biodegradable Films, Zero Waste, Resource Recovery, Environmental Impact

Introduction

A great amount of biowastes, comprising byproducts and biomass wastes, is originated yearly from the agri-food industry. These biowastes are commonly rich in proteins and polysaccharides and are mainly discarded or used for animal feeding (Visco et al.2022). Regulations related to encouraging a bio-based circular economy and ensuing bio-refinery

operations are prompting significant interest in repurposing the above-mentioned agri-food biowastes for making green materials, with a remarkable revalorization of those biowastes (Cristofoli et al.2023). A strategy whose design includes the production of materials from biowastes to be applied in agricultural operations, converging with the zero-waste approach, seems remarkably compelling from a sustainability prospect. The accumulation of plastic wastes is recognized as a global challenge due to its negative impact on the environment (Kibria et al.2023). especially low biodegradability is shown by fossil-based plastics, together with a massive production scale linked to the plastic market, which has generated a large and continuously growing accumulation in the oceans and landfills (Singh et al., 2022). Despite an increase in efforts over the past years to encourage a shift from a fossil-based to a bio-based circular economy and to optimize plastic waste management, a European Union Directive on the problem of single-use plastic products expects that in 2030 a growth in waste generation will be accompanied by no more than six percentage points of plastics growth (Chioatto & Sospiro). Thus, aiming to waste reduction at the same levels as the previous economic growth, noticeable advantages can come from materializing information technology and supply chain optimization processes (De Giovanni, 2023). The idea would be to redistribute additional supplies throughout economic sectors, promoting an economic system aiming at fulfilling public welfare and ensuring circular products design, reuse, repair, recycling, and biopolymer industries interconnected processes (Álvarez-Castillo et al., 2021).

Agricultural Waste as a Resource

World population is predicted to increase from 7.3 billion in 2016 to nearly 10 billion in 2050 (Sadigov, 2022). Such a projection highlights the great sustainability challenge needed to provide food for a growing population in a safe and sustainable way. The concept of circular economy is gradually gaining momentum, involving valorisation of ‘waste’ and co-product streams (reduce, reuse and recycle) (Jagtap et al.2021). Technologies have been developed to produce biopolymers with specific material properties and in the desired quantities. Some possible innovative valorisation strategies of residue streams for the production of bio-polymers are: pulp and paper industry residues, bio-conversion of agro-industrial waste by-products into high value biopolymers, (micro)algal secondary metabolites recovery (Flórez et al., 2023).

Post-harvest agricultural waste can be divided into typical field crop residues and crop-specific waste material which result from the cleaning, shucking and peeling phases which occur upon agri-food conditioning (Andrews et al.2021). Such residues and by-products include, inter alia, cereal straw, olive husk, grape stalk, almond and hazelnuts shells, among others. Different strategies can be applied endowing those by-products with added economic value, in line with circular economy principles (Sadh et al.2023). Techniques able to preserve the starter feedstock structure, so that the final product, either agglomerate or sheet, maintains its texture and structure, are most suitable. Ad hoc mechanical processes can be instead designed to alter the feedstock morphology to allow the subsequent application of conventional thermo-mechanical transformation processes, like film blowing or pellet extrusion (Li et al.2022).

Types of Agricultural Waste

A great amount of biowastes, comprising byproducts and biomass wastes, is originated yearly from the agri-food industry (Álvarez-Castillo et al., 2021). These biowastes are commonly rich in proteins and polysaccharides and are mainly discarded or used for animal feeding, while a small percentage is composted and only a minor fraction is processed for industrial material production (Noori et al., 2022). As regulations gradually aim to shift from a fossil-based to a bio-based, more environmentally friendly, circular economy model, biowastes are also being employed for producing bio-based materials. The strategy of producing materials from biowastes to be used in agricultural applications, which converges with the zero-waste

approach, seems to be remarkably attractive from a sustainability perspective (Zaborowska & Bernat, 2023). During the last decades, there has been a gradually growing concern about plastic pollution, primarily owed to food packaging due to the big amount of single-use plastics. Nonetheless, it is gradually perceived as a major issue for the environment, thus a shift towards bio-based, biodegradable materials is sought. In this regard, food industry biowastes with their rich balance of biopolymers (proteins and polysaccharides) have lately engrossed attention for sustainable material production as a method of reutilizing such discarded biomaterials (Khan & Rashid, 2024). Additionally, it is a well-suited approach as a way to foster a circular economy model, although challenging. Several agri-food proteins have been processed for film production, along with some of their potential applications as green materials, which are either animal-based or unsuitable for vegans (Álvarez-Castillo et al., 2021).

Biopolymers: Types and Applications

Biopolymers or biodegradable polymers are defined in this paper as polymers that are produced from natural resources. Bio-based polymers (biopolymers) are an emerging class of polymers that are developed from renewable resources (P Babu et al., 2013). Biodegradable polymers are an environmentally attractive alternative to synthetic polymers, mainly plastics such as flexible packaging materials. The term bio-based or biodegradable polymer is used in this work. Biopolymers from natural products have recently been actively studied as a promising way to contribute to the development of a sustainable material and they can be used instead of petroleum-based plastics (Baranwal et al., 2022). Agricultural and food waste is attractive as raw materials for biopolymer production from an economical viewpoint. In recent years, in order to achieve a sustainable and recycling-based society, attention has been focused on the effective use of biomass resources that are relatively abundant (Puglia et al.2021).

Biopolymer (bio-based polymer): polymers that are produced from natural resources. They consist of monomers which have been produced from the fermentation or natural resources (Gowthaman et al.2021). Biodegradability: the property of the material that permits its complete breakdown to be consumed by any of numerous microorganisms thus reintegrating it into its natural environment. Bio-based polymers are produced from renewable resources and considered as sustainable polymers (Nakajima et al., 2017). Biodegradable polymers are in essence inherently bio-based; however, not all bio-based polymers are biodegradable. In all cases, bio-based polymers have ecological and economic implications.

Definition and Characteristics

Recently, a broad range of attention has been posed on the development of biodegradable materials derived from biodegradable polymers (Wu et al., 2023). These biodegradable materials display a comparable crude-oil chain structure to petrochemical alternative materials, such as low-density polyethylene, polypropylene, or polystyrene. However, biodegradable polymers are prepared from biomass resources like polysaccharides or polylactic acid (Dilshad et al.2021). The employment of revalorized biopolymers as a source to be utilized for the preparation of materials belonging to the category of biodegradable mulch films is considered to be a desirable driver toward the agri-food 4.0 sector (Álvarez-Castillo et al., 2021). Moreover, this approach might contribute to shaping a more sustainable and circular bioeconomy in the agri-food sector.

Types of Biopolymers

The main motivation for development of biobased polymers was their biodegradability. The most well-known biobased polymer is polylactide (PLA), produced from starch or sugars (Balla et al.2021). Reflecting recent changes in the polymer industry, biobased polymers that are upgrades from biodegradable polymers allow the use of them for general and engineering

applications. Recently, the sustainability of biobased polymers has broadened their horizons for several fields (Jha et al., 2024).

Biobased polymers that are upgrades from biodegradable polymers fall under the category of research for general applications. This type of biobased polymers includes PLA, polyhydroxyalkanoates (PHAs), and poly(butylene succinate) (PBS) (Rafiqah et al.2021). Although the biodegradability of the biobased building blocks gives these polymers an advantage over conventional petroleum-based polymers, it also limits them to niche markets. Consequently, there have been endeavours to improve their properties in order to facilitate their use for more demanding applications (Jha et al., 2024). Such efforts have met with varying success, with some impact-resistant blends and alloys finding widespread acceptance, while attempts to produce glass-reinforced composites have been met with more limited success (Nakajima et al., 2017).

Biopolymer Production Methods

Biotechnology routes for producing biodegradable polymers, either polymers per se or composites, are being actively devised and implemented (Balla et al.2021). The biometric content in agricultural or harvesting products reaches almost 90% by weight, hence fruit stones, nutshells, and the like must be targets of interest. For some biowastes and co-products, unconventional applications directly as plastics are feasible, such as those derived from tissue-directed protein studies or innovative processing technologies that give rise to green materials to replace single-use plastics (Álvarez-Castillo et al., 2021). In fact, the strategy of the European Commission on plastics in a circular economy specifically contemplates the planned disposal of copolyesters and traditional rigid and flexible plastics not containing additives (Chioatto & Sospiro).

Among the radical options shown by the EC, there is a restriction of the oxo-degradable plastics in 2030 with the aim of a complete ban in 2039 (Qazanfarzadeh & Kumaravel, 2023). On the other hand, either the preformed legacy plastics or the new biodegradable polymers can be directed to the recycling stream. Likewise, the reuse strategies are industrially viable around 110,000 tons of per year concerning wrappers, bags, or flexible films once 2025 is reached, rapidly increasing afterwards. Unfortunately, the recycling process requires severe plastic depolymerization techniques currently relying on toxic solvents at high temperatures and acidic treatment with strong acids and surfactants (Miao et al., 2021). The robust implementation of biodegradable plastics, like those borne from lipids used gaseous overabundant fuels in agriculture, will thwart an already sketchy industrial process. A plethora of devastating microplastics would be generated (Martins de Souza & K. Gupta, 2024). Additionally, some studies suggest a possible biotoxicity of plastic fragments or residues caused by the possible release of toxic monomers, oligomers, or oxidized polymers.

Circular Economy Principles

In modern society, due to the lack of living space, the rapid decline of resources, and the continuous enhancement of public environmental protection awareness, the waste treatment problem has become a major issue that must be solved at the start of a modern society (Obaideen et al.2022). By promoting the 5 principles: repair, refinery, retread, recycling and resource saving. bio-carriers from agricultural wastes maintain the circular utilization of materials by the efficient production of bioplastics. It will make the agricultural waste more valuable and investable. The bio-carriers could be circularly recycled in biodegradable plastics manufacturing to improve its energy and economic efficiency. Development and formation of

circular bio-economy through 5 Rs would more effectively treat the unavoidable waste for a handsome return (Kieush et al.2022).

The concept of bioeconomy of this type is the widest part that deals with all production sectors that involve bioresources for the efficient, sustainable, and competitive production of food, feed, bio-energy, and other bio-products, or bio-production combined with the preservation and use of biological diversity and ecosystem services to ensure material recycling in a circular manner (Dahiya et al., 2022). Measures such as stimulating recycling should play a more important role in science and technology innovation in all sectors of economy and society, and increasing resource productivity. Eco-friendly industries and economies at the cross-industry and cross-border levels should be created, environmental costs should be included in the production and consumption price, economic, environmental, social and resource-interconnections (spillover) should be taken into account (Gibovic & Bikfalvi, 2021).

Sustainability Considerations

The production of biopolymers from agricultural waste is a promising practice that can mitigate numerous environmental issues, diversify the origin of biopolymers, improve valorization of waste products, and integrate better into a circular economy system, all while producing environmentally safe and degradable products. Current gaps in the literature about biopolymer production from agricultural waste are outlined (Phiri et al.2023). Finally, a perspective is given on the potential of expanding the use of biopolymers based on the level of consumer education, research orientation, and support for biopolymer producers in the sectoral economies of Rural Economy for the Commonwealth of Independent States, European Union, and the rest of the world, with some economic reflections (Gomes Gradissimo et al., 2020). In Russia, the policy on the reduction of plastic waste is at an early formation stage and collaboration future actions is expected. In Russia, consumers have an inadequate understanding of the hazard of traditional and biodegradable plastic waste and the marking of the products made from those. Japanese diversification is possible by integrating biodegradable plastics into a system of bioeconomy (Fadeeva & Van Berkel, 2021). The cluster analysis indicated the necessity of the waste management of biofiber biodegradable plastic products by policy making and promotion of products that create a market. The aim of the current study was to investigate the factors determining the chemical weathering performances of the plant oil-coated Luquillo Experimental Forest Atmospheric Wet Deposition materials exposed in two different tropical weather environments (McNeill et al.2024). Rainfall, two stations in Northeastern Puerto Rico were established in September, 2013, and deployed as is, for one year in the northern and two years in the south, to collect end-of-pipe rainwater throughout the sampling periods. The physicochemical stability of the fixed air-stripped rainwater samples was characterized during the same time period (Torres-Delgado et al.2021).

Environmental Impact Assessment

The production of polymers based on biobased and readily available raw materials is an alternative to using fossil resources, contributing to the transition towards a circular economy. This type of material is biodegradable and its production is considered to be more environmentally friendly, reducing the carbon footprint and other environmental impacts. An alternative use of fuel ethanol synthesis, considering the glycerin byproduct generated by biodiesel plants as raw material, is proposed (von et al.2023).

There are many publications in the literature that deal with the development of biopolymers. They all have one point in common: they replace fossil derivatives with biomass derivatives or natural resources (Ballús et al., 2023). There are likely two reasons for this interest in biopolymers: (a) substances derived from biomass, such as biopolymers, can help to reduce the

environmental impact; a recent example is biodegradability; the hydrophobic unit in a chemical structure can generate more recalcitrant non-biodegradable products; in this sense, adding biomass derivatives to fossil products can improve product biodegradability and enhance environmental performance; and, (b) the demand for materials is continually increasing, consequently, the main point could be to revert as much as possible recycle materials to the production chain (Kalak, 2023). Closing the loop in the industry, the life cycle becomes more efficient saving expense and sparing environmental impact (Forte et al., 2016). At the same time, production of goods implies a large variety of materials that are mixed in a heterogeneous way and are impossible to separate, reducing recycling feasibility.

Life cycle assessment (LCA) is a technique that can be used to quantify the environmental impact of a product. LCA identifies and quantifies the environmental impact at all stages of the life of the product, which can be divided into 4 main steps: extraction of raw materials, production of the polymer, distribution, and end of life (Sukumaran & Gopi, 2021). The aim of this work was to produce a new biopolymer using alternative raw materials different from those currently used in the production of the polymer, another objective was to know the environmental impact of this product. The proposed product was a polymer synthesized from glycerin, which is a byproduct of the biodiesel industry, and vinyl acetate. A biopolymer similar to this was found in the literature, but it was produced with glycerin from the production of bioethanol, and the hydrophobic unit in the synthesis of the polymer was stearic acid (Ben et al., 2022).

Case Studies and Applications

The same low-cost food industry waste was subjected to extraction of bio-polysaccharide-rich fractions for the development of biofilms with different active agents. Biofilms with a plasticizer and with a surfactant to enhance the dispersion of the active agents were developed. The plasticizer increased the elongation at the break. The tensile strength decreased when chitosan was incorporated or when the pH of the film-forming solution was 5 (Najar et al.2024). The biodegradability of bio-polysaccharide films was not affected by incorporating chitosan. There was not release of natamycin visible to the naked eyes for 10 days, while the growth of *A. flavus* in contact with the biofilm to days was restricted. The strategy of recycling these active biofilms could be considered to be implemented in industrial scale, which could be collected and regenerated from cleaning and disinfection of tools and areas used in the agri-food industry (Lago et al.2024). This strategy would make possible to enhance the use of actively protected products against the fungi growth on fruits and vegetables harmful to human health. Upon use of a biofilm, the demand of commercial fungicides would be reduced, while being replaced by bio-based biodegradable materials. This falls within regulation that restricts the use of fossil-based materials in the protection of food. In addition, sweet clover hydrophilic extracts were used as a coating to protect apple fruit from deterioration during storage at room temperature (Tian et al.2023).

Successful Biopolymer Projects

PHAs are increasingly seen as a potential substitute for conventional plastics due to their equivalent properties. The use of strategies to reduce the cost of PHAs in order to stimulate market demand. The use of agricultural waste for PHAs culturing and the demonstration of low-cost technologies as a competitive alternative is explored. All mentioned bioplastic films showed at least 60% degradation weight losses within the desired number of days (Naser et al., 2021). The progress on the use of abundant agricultural residue waste as a culture medium for cultivating PHAs and the development of a cost-effective solid-state fermentation technology for the mass production of high-quality PHAs is discussed. PHAs play an important role in the

substitution of conventional plastics, reducing environmental impact on land and aquatic environments (Naitam et al., 2022).

Real-world Applications and Impact

Biowastes from the agri-food industry are discarded materials. Today, a significant amount of biowastes generated by the agri-food industry are covalently bonded materials, rich in proteins, polysaccharides, or both. Many of them are of vegetal origin and are discarded or employed as forage or fertilizers for animals after inadequate treatments (Bala et al.2023). The use of biowastes as a valuable raw material for the preparation of bio-based materials has increased interest. These biodegradable and bio-based materials are employed for many applications of high added value as an alternative to single-use plastic materials; they are used as materials for food packaging (coatings, films, and pouches), agriculture (mulching films), or for scaffolds used in tissue engineering. The employment of biowastes as a raw material is a great strategy for a circular economy while it involves the bio-economy (Madała & Skuza, 2021). From a zero-waste and sustainability perspective, the use of biowastes for the production of materials intended for agricultural applications is an appealing strategy (Álvarez-Castillo et al., 2021). Bio-based materials have been proposed as an eco-friendly alternative to fossil-based ones to ameliorate the environmental impact of plastics. The lack of biodegradability of conventional plastics, derived from fossil fuels, has led to significant pollution of the planet. Indeed, it is estimated that the input of plastic wastes in the environment will reach 250 million metric tons by 2025 (Forfora et al.2024). Despite the believed material properties, fossil-based plastics are found in several commodities that are used for a few minutes to weeks. Sadly, a sustainable management is missed: only 20% of annual plastic waste is collected for recycling and incineration leads to emission of dioxins and furans, as well as the production of greenhouse gas emissions (Chen et al.2021).

Future Trends and Innovations

A great amount of biowastes, including byproducts and biomass wastes, is annually produced by the agri-food industry, summing several billion tones each year. Those biowastes are generally rich in polysaccharides and proteins and are mainly discarded or utilized for animal feeding (Álvarez-Castillo et al., 2021). More recently, biowastes are being utilized for the production of bio-based materials. Some of those materials can disintegrate in a short time under specific environmental conditions; these are the well-known biodegradable materials. The use of biodegradable materials from renewable resources for manufacturing various products may involve their use in high value applications and, therefore, imply a remarkable revalorization of those resources (Cetecioglu et al.2022). Moreover, the strategy for producing materials from biowastes for agricultural applications, which are analogous to the zero-waste approach concept, seems to be remarkably attractive from a sustainability prospect. However, up to date, there has been minimal attention given to the possibility of using biowastes, or polymers derived from biowaste proteins and/or polysaccharides, for producing environmentally friendly agricultural materials (Ravindran et al.2021).

The accumulated plastic waste is growing globally, officially recognized as a critical and extensive problem due to its notably negative impact on the environment (Kibria et al.2023). The very low degradability of fossil-based plastics generates a massive accumulation of waste plastics in landfills and the oceans. Recent estimates forecast that among the polymers world-produced to date, a significant amount is plastic waste, with a large percentage being accumulated in the environment (Evide et al.2021). Despite efforts to shift from a linear to a bio-based circular economy model, currently, only a small fraction of the plastic waste is collected for recycling, of which barely a small percentage is used to produce recycled products. Directive (EU) 2019/904 from the European Union defines measures to prevent and reduce the

impact of specific plastic products on the environment, principally the marine environment, and human health. Among other directives, this imposes restrictions on biodegradable plastics very likely intended to prevent open-field applications (Roosen et al.2022).

Conclusion and Recommendations

A great amount of biowastes generated from the agri-food industry is rich in proteins and polysaccharides, which constitute bio-based polymeric materials. In light of the strict regulatory trends for a bio-based circular economy model, biowastes have been addressed as raw materials for developing bio-based materials (Bala et al.2023). Biowaste-derived proteins and polysaccharides have been processed to produce green materials, while proteins have been used for edible food packaging with active compounds, and polysaccharides have been used as edible films on fresh fruit coatings and for controlled drug release, to prepare bio-based composites for agriculture and 3D-scaffold bio-based materials. The enclosed strategies are set in environmental, economic, and social sustainability, in line with the zero waste approach (Ashrafi et al.2022). A series of negative externalities have emerged in food consumption and industrialization patterns and life styles worldwide, such as increasing waste production, resource consumption, and disposable consumer goods utilization. Among these unsustainable models, the accumulated plastic waste in the world has become an exacerbating issue, posing severe environmental pollution and socio-economical tensions. Consequently, there is a strong accent on promoting a bio-based circular economy in an effort to eradicate single-use plastic products (Fletcher et al.2024). In this vein, the European Union Directive (EU) 2019/904 has been enacted to address the issue of waste generated by single-use plastic products and fishing gear, with the aim of preventing and reducing the impact on the marine environment of such waste, constituting the single biggest portion of plastic litter. (Álvarez-Castillo et al., 2021) To introduce the circular economy throughout the life cycle of products, the recovery and recycling of raw materials is set as the priority step. Meanwhile, growing interests have been directed at developing alternative materials made from bio-wasted polymers, consisting of proteins and polysaccharides as major components, with the advantage of biodegradability. This class of natural materials has also been investigated as promising resources for a variety of fast emerging applications under stringent health and environmental criteria, such as biodegradable plastics, bioactive packaging, and biomedicines (Smol et al., 2021).

References

- Visco, A., Scolaro, C., Facchin, M., Brahimi, S., Belhamdi, H., Gatto, V., & Beghetto, V. (2022). Agri-food wastes for bioplastics: European prospective on possible applications in their second life for a circular economy. *Polymers*, 14(13), 2752. [mdpi.com](https://doi.org/10.3390/polym14132752)
- Cristofoli, N. L., Lima, A. R., Tchonkouang, R. D., Quintino, A. C., & Vieira, M. C. (2023). Advances in the food packaging production from agri-food waste and by-products: market trends for a sustainable development. *Sustainability*, 15(7), 6153. [mdpi.com](https://doi.org/10.3390/su15076153)
- Kibria, M. G., Masuk, N. I., Safayet, R., Nguyen, H. Q., & Mourshed, M. (2023). Plastic waste: challenges and opportunities to mitigate pollution and effective management. *International Journal of Environmental Research*, 17(1), 20. [springer.com](https://doi.org/10.1007/s12010-023-01000-0)
- Singh, N., Ogunseitan, O. A., Wong, M. H., & Tang, Y. (2022). Sustainable materials alternative to petrochemical plastics pollution: A review analysis. *Sustainable horizons*. [sciencedirect.com](https://doi.org/10.1016/j.sh.2022.100001)
- Chioatto, E. & Sospiro, P. (). Transition from waste management to circular economy: the European Union roadmap. *Environment*. [\[HTML\]](#)

- De Giovanni, P. (2023). Sustainability of the Metaverse: A transition to Industry 5.0. Sustainability. [mdpi.com](https://www.mdpi.com)
- Álvarez-Castillo, E., Félix Ángel, M., Bengoechea Ruiz, C., & Francisco Guerrero Conejo, A. (2021). Proteins from Agri-Food Industrial Biowastes or Co-Products and Their Applications as Green Materials. [\[PDF\]](#)
- Sadigov, R. (2022). Rapid growth of the world population and its socioeconomic results. The Scientific World Journal. [wiley.com](https://www.wiley.com)
- Jagtap, S., Garcia-Garcia, G., Duong, L., Swainson, M., & Martindale, W. (2021). Codesign of food system and circular economy approaches for the development of livestock feeds from insect larvae. *Foods*, 10(8), 1701. [mdpi.com](https://www.mdpi.com)
- Flórez, M., Cazón, P., & Vázquez, M. (2023). Selected biopolymers' processing and their applications: A review. *Polymers*. [mdpi.com](https://www.mdpi.com)
- Andrews, E. M., Kassama, S., Smith, E. E., Brown, P. H., & Khalsa, S. D. S. (2021). A review of potassium-rich crop residues used as organic matter amendments in tree crop agroecosystems. *Agriculture*, 11(7), 580. [mdpi.com](https://www.mdpi.com)
- Sadh, P. K., Chawla, P., Kumar, S., Das, A., Kumar, R., Bains, A., ... & Sharma, M. (2023). Recovery of agricultural waste biomass: A path for circular bioeconomy. *Science of the Total Environment*, 870, 161904. [\[HTML\]](#)
- Li, C., Wu, J., Shi, H., Xia, Z., Sahoo, J. K., Yeo, J., & Kaplan, D. L. (2022). Fiber-based biopolymer processing as a route toward sustainability. *Advanced Materials*, 34(1), 2105196. [wiley.com](https://www.wiley.com)
- Noori, A. W., Royen, M. J., Medved'ová, A., & Haydary, J. (2022). Drying of food waste for potential use as animal feed. Sustainability. [mdpi.com](https://www.mdpi.com)
- Zaborowska, M. & Bernat, K. (2023). The development of recycling methods for bio-based materials—A challenge in the implementation of a circular economy: A review. *Waste Management & Research*. [sagepub.com](https://www.sagepub.com)
- Khan, N. R. & Rashid, A. B. (2024). Carbon-based nanomaterials: a paradigm shift in biofuel synthesis and processing for a sustainable energy future. *Energy Conversion and Management: X*. [sciencedirect.com](https://www.sciencedirect.com)
- Álvarez-Castillo, E., Felix, M., Bengoechea, C., & Guerrero, A. (2021). Proteins from agri-food industrial biowastes or co-products and their applications as green materials. *Foods*. [mdpi.com](https://www.mdpi.com)
- P Babu, R., O'Connor, K., & Seeram, R. (2013). Current progress on bio-based polymers and their future trends. [ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov)
- Baranwal, J., Barse, B., Fais, A., Delogu, G. L., & Kumar, A. (2022). Biopolymer: A sustainable material for food and medical applications. *Polymers*. [mdpi.com](https://www.mdpi.com)
- Puglia, D., Pezzolla, D., Gigliotti, G., Torre, L., Bartucca, M. L., & Del Buono, D. (2021). The opportunity of valorizing agricultural waste, through its conversion into biostimulants, biofertilizers, and biopolymers. *Sustainability*, 13(5), 2710. [mdpi.com](https://www.mdpi.com)
- Gowthaman, N. S. K., Lim, H. N., Sreeraj, T. R., Amalraj, A., & Gopi, S. (2021). Advantages of biopolymers over synthetic polymers: social, economic, and environmental aspects. In *Biopolymers and their industrial applications* (pp. 351-372). Elsevier. [researchgate.net](https://www.researchgate.net)
- Nakajima, H., Dijkstra, P., & Loos, K. (2017). The Recent Developments in Biobased Polymers toward General and Engineering Applications: Polymers that Are Upgraded from

Biodegradable Polymers, Analogous to Petroleum-Derived Polymers, and Newly Developed. ncbi.nlm.nih.gov

Wu, L., Shi, X., & Wu, Z. S. (2023). Recent advancements and perspectives of biodegradable polymers for supercapacitors. *Advanced Functional Materials*. [\[HTML\]](#)

Dilshad, E., Waheed, H., Ali, U., Amin, A., & Ahmed, I. (2021). General structure and classification of bioplastics and biodegradable plastics. *Bioplastics for sustainable development*, 61-82. researchgate.net

Balla, E., Daniilidis, V., Karlioti, G., Kalamas, T., Stefanidou, M., Bikiaris, N. D., ... & Bikiaris, D. N. (2021). Poly (lactic Acid): A versatile biobased polymer for the future with multifunctional properties—From monomer synthesis, polymerization techniques and molecular weight increase to PLA applications. *Polymers*, 13(11), 1822. mdpi.com

Jha, S., Akula, B., Enyioma, H., Novak, M., Amin, V., & Liang, H. (2024). Biodegradable Biobased Polymers: A Review of the State of the Art, Challenges, and Future Directions. *Polymers*. mdpi.com

Rafiqah, S. A., Khalina, A., Harmaen, A. S., Tawakkal, I. A., Zaman, K., Asim, M., ... & Lee, C. H. (2021). A review on properties and application of bio-based poly (butylene succinate). *Polymers*, 13(9), 1436. mdpi.com

Qazanfarzadeh, Z. & Kumaravel, V. (2023). Hydrophobisation approaches of protein-based bioplastics. *Trends in Food Science & Technology*. sciencedirect.com

Miao, Y., von Jouanne, A., & Yokochi, A. (2021). Current technologies in depolymerization process and the road ahead. *Polymers*. mdpi.com

Martins de Souza, F. & K. Gupta, R. (2024). Bacteria for Bioplastics: Progress, Applications, and Challenges. ncbi.nlm.nih.gov

Obaideen, K., Shehata, N., Sayed, E. T., Abdelkareem, M. A., Mahmoud, M. S., & Olabi, A. G. (2022). The role of wastewater treatment in achieving sustainable development goals (SDGs) and sustainability guideline. *Energy Nexus*, 7, 100112. sciencedirect.com

Kieush, L., Rieger, J., Schenk, J., Brondi, C., Rovelli, D., Echterhof, T., ... & Colla, V. (2022). A comprehensive review of secondary carbon bio-carriers for application in metallurgical processes: utilization of torrefied biomass in steel production. *Metals*, 12(12), 2005. mdpi.com

Dahiya, D., Sharma, H., Kumar Rai, A., & Singh Nigam, P. (2022). Application of biological systems and processes employing microbes and algae to Reduce, Recycle, Reuse (3Rs) for the sustainability of circular bioeconomy. ncbi.nlm.nih.gov

Gibovic, D. & Bikfalvi, A. (2021). Incentives for plastic recycling: How to engage citizens in active collection. Empirical evidence from Spain. *Recycling*. mdpi.com

Phiri, R., Rangappa, S. M., Siengchin, S., Oladijo, O. P., & Dhakal, H. N. (2023). Development of sustainable biopolymer-based composites for lightweight applications from agricultural waste biomass: A review. *Advanced Industrial and Engineering Polymer Research*, 6(4), 436-450. sciencedirect.com

Gomes Gradissimo, D., Pereira Xavier, L., & Valadares Santos, A. (2020). Cyanobacterial Polyhydroxyalkanoates: A Sustainable Alternative in Circular Economy. ncbi.nlm.nih.gov

Fadeeva, Z. & Van Berkel, R. (2021). Unlocking circular economy for prevention of marine plastic pollution: An exploration of G20 policy and initiatives. *Journal of Environmental Management*. [\[HTML\]](#)

- McNeill, D. C., Pal, A. K., Nath, D., Rodriguez-Uribe, A., Mohanty, A. K., Pilla, S., ... & Misra, M. (2024). Upcycling of ligno-cellulosic nutshells waste biomass in biodegradable plastic-based biocomposites uses-a comprehensive review. *Composites Part C: Open Access*, 100478. [sciencedirect.com](https://www.sciencedirect.com)
- Torres-Delgado, E., González, G., Medina, E., & Rivera, M. M. (2021). A multiyear record of rainfall and ionic composition along an elevation gradient in Northeastern Puerto Rico. *Aerosol and Air Quality Research*, 21(6), 200582. [springer.com](https://www.springer.com)
- von Vacano, B., Mangold, H., Vandermeulen, G. W., Battagliarin, G., Hofmann, M., Bean, J., & Künkel, A. (2023). Sustainable design of structural and functional polymers for a circular economy. *Angewandte Chemie International Edition*, 62(12), e202210823. [wiley.com](https://www.wiley.com)
- Ballús, O., Guix, M., Baquero, G., & Bacardit, A. (2023). Life Cycle Environmental Impacts of a Biobased Acrylic Polymer for Leather Production. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)
- Kalak, T. (2023). Potential use of industrial biomass waste as a sustainable energy source in the future. *Energies*. [mdpi.com](https://www.mdpi.com)
- Forte, A., Zucaro, A., Basosi, R., & Fierro, A. (2016). LCA of 1,4-Butanediol Produced via Direct Fermentation of Sugars from Wheat Straw Feedstock within a Territorial Biorefinery. [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)
- Sukumaran, N. P. & Gopi, S. (2021). Overview of biopolymers: Resources, demands, sustainability, and life cycle assessment modeling and simulation. *Biopolymers and their industrial applications*. [researchgate.net](https://www.researchgate.net)
- Ben, Z. Y., Samsudin, H., & Yhaya, M. F. (2022). Glycerol: Its properties, polymer synthesis, and applications in starch based films. *European Polymer Journal*. [\[HTML\]](#)
- Najar, I. N., Sharma, P., Das, R., Tamang, S., Mondal, K., Thakur, N., ... & Kumar, V. (2024). From waste management to circular economy: Leveraging thermophiles for sustainable growth and global resource optimization. *Journal of Environmental Management*, 360, 121136. [\[HTML\]](#)
- Lago, A., Rocha, V., Barros, O., Silva, B., & Tavares, T. (2024). Bacterial biofilm attachment to sustainable carriers as a clean-up strategy for wastewater treatment: A review. *Journal of Water Process Engineering*, 63, 105368. [sciencedirect.com](https://www.sciencedirect.com)
- Tian, B., Liu, J., Yang, W., & Wan, J. B. (2023). Biopolymer food packaging films incorporated with essential oils. *Journal of Agricultural and Food Chemistry*, 71(3), 1325-1347. [\[HTML\]](#)
- Naser, A. Z., Deiab, I., & Darras, B. M. (2021). Poly (lactic acid)(PLA) and polyhydroxyalkanoates (PHAs), green alternatives to petroleum-based plastics: a review. *RSC advances*. [rsc.org](https://www.rsc.org)
- Naitam, M. G., Tomar, G. S., Kaushik, R., Singh, S., & Nain, L. (2022). Agro-industrial waste as potential renewable feedstock for biopolymer poly-hydroxyalkanoates (PHA) production. *Enzym. Eng.* [researchgate.net](https://www.researchgate.net)
- Bala, S., Garg, D., Sridhar, K., Inbaraj, B. S., Singh, R., Kamma, S., ... & Sharma, M. (2023). Transformation of agro-waste into value-added bioproducts and bioactive compounds: Micro/nano formulations and application in the agri-food-pharma sector. *Bioengineering*, 10(2), 152. [mdpi.com](https://www.mdpi.com)
- Madela, M. & Skuza, M. (2021). Towards a circular economy: Analysis of the use of biowaste as biosorbent for the removal of heavy metals. *Energies*. [mdpi.com](https://www.mdpi.com)

- Forfora, N., Azuaje, I., Kanipe, T., Gonzalez, J. A., Lendewig, M., Urdaneta, I., ... & Argyropoulos, D. (2024). Are starch-based materials more eco-friendly than fossil-based? A critical assessment. *Cleaner Environmental Systems*, 100177. [sciencedirect.com](https://www.sciencedirect.com)
- Chen, H. L., Nath, T. K., Chong, S., Foo, V., Gibbins, C., & Lechner, A. M. (2021). The plastic waste problem in Malaysia: management, recycling and disposal of local and global plastic waste. *SN Applied Sciences*, 3, 1-15. [springer.com](https://www.springer.com)
- Cetecioglu, Z., Atasoy, M., Cenian, A., Sołowski, G., Trček, J., Ugurlu, A., & Sedlakova-Kadukova, J. (2022). Bio-based processes for material and energy production from waste streams under acidic conditions. *Fermentation*, 8(3), 115. [mdpi.com](https://www.mdpi.com)
- Ravindran, B., Karmegam, N., Yuvaraj, A., Thangaraj, R., Chang, S. W., Zhang, Z., & Awasthi, M. K. (2021). RETRACTED: Cleaner production of agriculturally valuable benignant materials from industry generated bio-wastes: A review. *Bioresource Technology*, 320, 124281. [\[HTML\]](#)
- Evode, N., Qamar, S. A., Bilal, M., Barceló, D., & Iqbal, H. M. (2021). Plastic waste and its management strategies for environmental sustainability. *Case studies in chemical and environmental engineering*, 4, 100142. [sciencedirect.com](https://www.sciencedirect.com)
- Roosen, M., Mys, N., Kleinhans, K., Lase, I. S., Huysveld, S., Brouwer, M., ... & De Meester, S. (2022). Expanding the collection portfolio of plastic packaging: Impact on quantity and quality of sorted plastic waste fractions. *Resources, Conservation and Recycling*, 178, 106025. [sciencedirect.com](https://www.sciencedirect.com)
- Ashrafi, G., Nasrollahzadeh, M., Jaleh, B., Sajjadi, M., & Ghafuri, H. (2022). Biowaste-and nature-derived (nano) materials: Biosynthesis, stability and environmental applications. *Advances in Colloid and Interface Science*, 301, 102599. [\[HTML\]](#)
- Fletcher, C. A., Aureli, S., Foschi, E., Leal Filho, W., Barbir, J., Beltrán, F. R., ... & Banks, C. E. (2024). Implications of consumer orientation towards environmental sustainability on the uptake of bio-based and biodegradable plastics. *Current Research in Environmental Sustainability*, 7, 100246. [sciencedirect.com](https://www.sciencedirect.com)
- Smol, M., Marcinek, P., & Koda, E. (2021). Drivers and barriers for a circular economy (CE) implementation in Poland—A case study of raw materials recovery sector. *Energies*. [mdpi.com](https://www.mdpi.com)

ECOLOGICAL ASSESSMENT OF THE SOILS OF ABRAGUNUS DEPRESSION

ABRAGUNUS ALLIANCE TOPRAKLARININ EKOLOJİK DEĞERLENDİRMESİ

Akim AKHUNDOV

Nahçıvan Devlet Üniversitesi, Mimarlık ve Mühendislik Fakültesi, Sulama ve Ekoloji
Mühendisliği Bölümü, Nahçıvan, Azerbaycan

ORCID ID: <https://orcid.org/0009-0006-8328-306X>

Sahib HACIYEV

Nahçıvan Devlet Üniversitesi, Doğa Bilimleri ve Ziraat Fakültesi, Coğrafya Bölümü,
Nahçıvan, Azerbaycan

ORCID ID: <https://orcid.org/0009-0001-1839-754X>

Elnara SEYİDOVA

Nahçıvan Devlet Üniversitesi, Doğa Bilimleri ve Ziraat Fakültesi, Coğrafya Bölümü,
Nahçıvan, Azerbaycan

ORCID ID: <https://orcid.org/0009-0001-0486-2324>

ABSTRACT

In the article the eco-geographical condition of the soil formation which is spread in Abragunuse hollow on the middle upland waist-band in Nakhchivan Autonomous Republic is taught. Here the soil samples morphological, physical and chemical peculiarities which are taken from the research object are analyzed, too. At the same time the productivity of the cultural and natural plants is taught. According to the researches of the Abragunuse hollow the collective connection between the fertility of soils and the productivity of plants was defined, marked and the agroproduction classification of soils was held.

As a result the efficient use of the soils in the Abragunuse hollow under the cultural and natural plants is recommended.

Key words: geographic factors, soil, eco-geography, bonitet

ÖZET

Giriş ve Amaç: Toprağın insan yaşamında olağanüstü bir öneme sahip olduğu bilinmektedir. Nahçıvan Özerk Cumhuriyeti, karmaşık yer şekillerine sahip dağlık bir alan olduğundan tarıma elverişli arazi miktarı azdır. Öte yandan, bölgenin karasal iklimi toprak bozulmasına, erozyona ve tuzlanma süreçlerine neden olmaktadır. Bu açıdan bakıldığında, özerk cumhuriyetin tarım arazisi varlığının yüzde 30'unu oluşturan orta dağlık kuşaktaki dağ arası çöküntü alanlarındaki ekili ve doğal bitki örtüsünün değerlendirilmesine yönelik bir araştırma yapılması önem taşımaktadır.

Çalışma alanı araştırma nesnesi olarak ele alınmıştır. Abraqunus Çöküntüsü, Nahçıvan Özerk Cumhuriyeti sınırları içinde yer alan coğrafi konumu nedeniyle kuzeyde Elince Dağı'nın güney etekleri, batıda Uşaklı Dağı'nın doğu etekleri, güneyde Aras Çöküntüsü'nün alçak tepeleri, kuzeybatıda İlanlı Dağı'nın etekleri ve güneybatıda Darıdağ'ın etekleriyle çevrilidir.

Araştırmanın amacı, Abraqunus çöküntüsünde toprak oluşumunu etkileyen doğal ve antropojenik faktörleri incelemek ve tuzlanma, tuzlanma, su basması ve çevrenin kaybına etki eden diğer olumsuz süreçlerin kökenini (genezisini), oluşum nedenlerini, dağılım biçimini ve coğrafyasını anlamak, bunları doğru bir şekilde değerlendirmek, sınıflandırmasını hazırlamak ve gerekli önlemleri almaktır.

Gereç ve Yöntem: Abraqunus çöküntüsünde toprakların oluşumunu etkileyen doğal ve antropojenik faktörleri incelemek ve çevrenin kaybına neden olan olumsuz süreçlerden tuzlanma, tuzlanma, bataklıklaşma ve diğer faktörlerin kökenini (genezisini) incelemek, bunların ortaya çıkış nedenlerini araştırmak ve gerekli önlemleri almak amacıyla, bu doğrultuda yabancı ülkelere, Azerbaycan ve Özerk Cumhuriyet'ten bilim adamları tarafından yapılan araştırmaların sonuçları, bunlar tarafından derlenen haritalar, literatür ve metodolojik araçlar toplandı, analiz edildi ve araştırma çalışmalarında kullanıldı. Yabancı bilim insanları arasında (Dj. Khanvell 1977, V. Volobuev 1945-85, S. Babayev 1977-2000, G. Azizov 1990-2004, A. Guliyev 1977-2020, S. Hajiyevev 1976-2023, N. Bababeyli 1995-2021, R. Amirov, P. Fatullayev 2012 ve diğerleri sayılabilir.

Araştırmada coğrafi araştırmanın geleneksel yöntemleri olan gözlemsel, tarihsel, istatistiksel, durağan (sahada kalarak), kartografik, tahmini ve günümüzde ortaya çıkan son teknolojilere dayalı coğrafi bilgi sistemleri yöntemleri kullanılmıştır.

Tartışma ve Sonuç: Araştırma nesnesinde toprakların oluşumunu etkileyen doğal ve antropojenik etkenler ile çevre kaybına neden olan olumsuz süreçler, tuzlanma, sulaklaşma ve diğer etkenlerin kökeni ve nedenleri araştırılıp analiz edilerek gerekli önlemlerin alınması için yönlendirmelerde bulunulmuştur.

Anahtar kelimeler: coğrafi faktörler, toprak, eko-coğrafya, arazi kalitesi

GİRİŞ

Toprak insan hayatında olağanüstü bir öneme sahiptir. Nahçıvan Özerk Cumhuriyeti, karmaşık yer şekillerine sahip dağlık bir alan olduğundan tarıma elverişli arazi miktarı azdır. Öte yandan, bölgenin karasal iklimi toprak bozulmasına, erozyona ve tuzlanma süreçlerine neden olmaktadır. Bu açıdan bakıldığında, Özerk Cumhuriyetin tarım arazisi fonunun yüzde 30'unu oluşturan Orta Dağlık Kuşak'taki dağ arası çöküntü alanlarında yayılan tarıma elverişli ve doğal bitki örtüsünün değerlendirilmesine yönelik bir araştırma yapılması önem taşımaktadır.

Çalışma alanı araştırma nesnesi olarak ele alınmıştır. Abraqunus Çöküntüsü, Nahçıvan Özerk Cumhuriyeti sınırları içinde yer alan coğrafi konumu nedeniyle kuzeyde Elince Dağı'nın güney etekleri, batıda Uşaklı Dağı'nın doğu etekleri, güneyde Aras Çöküntüsü'nün alçak tepeleri, kuzeybatıda İlanlı Dağı'nın etekleri ve güneybatıda Darıdağ Dağı'nın etekleriyle çevrilidir (Harita 1).



Harita 1, Abraqunus Çöküntüsünün fiziksel haritası

Abraqunus çöküntüsünde yayılım gösteren toprak yapısının eko-coğrafik koşulları, morfolojik, fiziksel ve kimyasal özelliklerinin incelenmesi, bunların kültürel ve doğal bitki örtüsü altında etkin bir şekilde kullanılması açısından değerlendirilmesi ve tarımsal üretim gruplamasının bir tablo halinde derlenmesinden oluşmaktadır.

Bu hedefe ulaşmak için aşağıdaki konular planlanmış ve çözüme kavuşturulmuştur.

- Araştırma sahasındaki arazi örtüsünün incelenmesine ilişkin literatür, harita ve arazi materyallerinin toplanması;
- Bölgede toprak oluşum sürecini etkileyen eko-coğrafik koşulların incelenmesi ve analizi;
- Araştırma nesnesi çevresinde yayılış gösteren kültür ve doğal bitkilerin verimliliğini, belirlenen alanlara toprak örnekleri yerleştirilerek izlemek;
- Alandan alınan toprak ve bitki örneklerinin morfolojik, fiziksel ve kimyasal özelliklerinin incelenmesi ve analiz edilmesi;
- Toprakların hesaplanan verimlilik göstergeleri ve çevresinde yetişen (kültürel ve doğal) bitkilerin verimliliğine göre toprakların değerlendirilmesi suretiyle tarımsal üretim planlaması yapmak;

Çalışmanın malzemesi ve metodolojisi

Konuyla ilgili literatür ve saha materyalleri toplanarak çalışma için bir metodoloji geliştirildi. Konunun geliştirilmesinde, Azerbaycan ve Nahçıvan Özerk Cumhuriyeti de dahil olmak üzere, yabancı ülkelerde, tarihi gelişimin farklı aşamalarında toprak ve bitki araştırmaları yapan bilim adamlarının çağdaş gereksinimlerini karşılayan monografiler, metodolojik kılavuzlar, harita materyalleri ve çalışma pratikleri kullanıldı [4, s. 187-233; 5, s. 113-157; 6, s. 25-163; 7, s. 5-38; 9, s. 43-48].

Konunun uygulanmasında literatür materyallerinin yanı sıra saha materyallerinin toplanması da önemli konulardan biridir. Bu amaçla Abraquonus çöküntüsünde toprak örtüsünün yapısal bileşimini incelemek üzere aşağıdaki yönlerde keşif gezileri düzenlenmiştir.

14 Mayıs 2024 tarihinde Abraquonus çöküntüsünün kuzey kesiminde (Khanagah, Khoshkeshin ve Alinja köyleri civarı) yayılmış kestane (kahverengi), açık kestane (kahverengi) ve alüvyonlu alüvyonlu topraklarda 6 adet parsel tesis edilerek, her parselin genetik katmanlarından örnekler alınarak, morfolojik özellikleri toprak biliminde kabul gören 1 numaralı forma yansıtılmıştır.

4 Haziran 2024 tarihinde Abraquonus çöküntüsünün orta kesiminde (Goydara, Saltag, Benaniyar ve Kirna köyleri civarı) yayılmış, alt toprak-alüvyonlu, hafif killi, kestane (kahverengi), çakıllı açık kestane (kahverengi), kumlu açık kestane (kahverengi) topraklarda 8 adet parsel kurulmuş, her parselin genetik katmanlarından örnekler alınmış ve morfolojik özelliklerini yansıtan 1 numaralı form doldurulmuştur.

25 Haziran 2024 tarihinde, Abraquonus çöküntüsünün güney kesiminde (Abraquonus ve Gizilja köyleri civarı) dağılmış orta-kalın, zayıf tuzlu kestane (kahverengi), kalın, zayıf tuzlu açık kestane (kahverengi) ve alüvyonlu-taşkın ovası topraklarında 4 kesit yapılmış ve öncekilerde olduğu gibi morfolojik özelliklerini yansıtan 1 numaralı form doldurulmuştur.

2024 yılı Mayıs-Haziran aylarında Abraquonus Çöküntüsü'ne düzenlenen keşif gezileri sonucunda, toprak ve bitki örneklerinin alındığı alanlara yayılmış olan kültür ve doğal bitkilerin verimlilikleri araştırılmıştır (Tablo 1). Alandaki toprakların kültürel ve doğal bitki örtüsü altında değerlendirilmesinde, toprak ve bitki örneklerinin morfolojik, fiziksel ve kimyasal özelliklerinin yanı sıra, eko-coğrafik koşulları (rölyef, jeolojik ve jeomorfolojik yapı, iklim koşulları, hidrolojik ve hidrojeolojik koşullar, bitki ve hayvan familyaları vb.) dikkate alınmıştır [1, s. 23-85; 2, s. 9-59; 3, s. 26-29; 7, s. 5-38; 8, s. 151-184].

Abraquonus depresyonunda yaygın olan tüm toprak tipleri, alt tipleri ve tür çeşitliliğinin değerlendirilmesinde, ekili ve doğal bitkilerin verimliliğine ayrı ayrı dikkat edildi (Tablo).

Tablo 1

2024 yılı itibarıyla Abraquonus depresyonunun belirlenen alanlarındaki ekili ve doğal bitkilerin verimliliği (s/ha)

No	Toprakların adı	Tahıl	Yonca	Doğal otlar
1.	Şabalıdı (kahverengi)	45-50	150-155	6-7
2.	Taşkın yatağı alüvyon	40-45	145-150	5-6
3.	Açık-şabalıdı (kahverengi)	35-40	140-145	4-5
4.	Qumsal açık şabalıdı (kahverengidir)	30-35	135-140	3-4
5.	Yüngül gilliceli yuxa, şabalıdı (kahverengi)	25-30	130-135	2-3
6.	Bozqırlaşmış şabalıdı (Kahverengi)	15-20	125-130	1-2

Toplanan bitkilerin botanik ve kimyasal kompozisyonlarını belirlemek amacıyla besiyeri örnekleri seçildi. Bitki örnekleri kurutulduktan sonra bitki türlerine göre kuru kütle verimleri belirlendi. Bu göstergeler fitosinoz mera alanının hektar başına verimliliğinin belirlenmesinde kullanılmıştır.

Araştırma alanından alınan bitki örneklerinin kimyasal analizleri aşağıda listelenen bilim insanlarının yöntemleri kullanılarak belirlenmiştir. Yem numunelerinde yaş protein-Kjeldahl, yaş yağ-Sokelet aparatı, hücre-Genenberg ve Steman, higroskopik nem ve azotsuz ekstraktifler (NFE) hesaplaması, yaş kül yakma vb. yöntemlerle belirlenir. Yem bitkilerinde belirlenen kimyasal analiz sonuçları, doğal kaba yemin yem birimine dönüştürülmesinde kullanıldı.

Alandaki toprakların kültürel ve doğal bitki örtüsü altında değerlendirilmesinde, toprak ve bitki örneklerinin morfolojik, fiziksel ve kimyasal özellikleri yanında, eko-coğrafik koşulları (rölyef, jeolojik ve jeomorfolojik yapı, iklim koşulları, hidrolojik ve hidrojeolojik koşullar, bitki ve hayvan familyaları vb.) da dikkate alınmıştır (1, 2). Analiz ve tartışma: - 2024 yılında "Kültürel ve doğal bitki örtüsü altında Abraquus çöküntüsü topraklarının eko-coğrafik değerlendirilmesi" konulu çalışmamız kapsamında gerçekleştirdiğimiz toprak-bitki araştırmasına dayanarak, çeşitli kaynaklardan toplanan materyallerin (morfolojik, fiziksel, kimyasal özellikler ve ekolojik koşullar) incelenmesi sonucunda alanda 25 tür çeşitliliği tespit edilmiştir. Gösterilen 25 tür çeşitliliği 13 tip ve alt tipe birleştirilerek topraklar değerlendirildi ve tarımsal üretim gruplandırması yapıldı. Burada topraklar tam bonitet ölçeğinde 100 puan almış olup, minimum 14 puan almıştır.


Bu topraklar, tespit edilen özelliklerine bağlı olarak farklı kalite ve değere sahip olup, tam bir bonitet ölçeğinin oluşturulmasında "düzeltme" faktörleri (tuzlanma, erozyon, granülometrik kompozisyon vb.) de kullanılmıştır (Tablo 2).

Tablo 2

Abraquus Havzası'nın toprak kalitesinin tam ölçeği
(100 puanlık bir ölçekte)

No	Torpaqların adı	Bonitet balı	Bal sınıfı	Kaliteli ekip
1.	Şabalıdı (kahverengi)	100	X	I Yüksək
2.	Taşkın yatağı alüvyon	90	X	
3.	Açığı-şabalıdı (kahverengi)	82	IX	
4.	Sahil açık kahverengidir.	76	VIII	II İyi
5.	Hafif solungaçlı, yüksek, şabalıdı (kahverengi),	72	VIII	
6.	Orta taneli şabalıdı (kahverengi)	62	VII	
7.	Hafif siltli, hafif tuzlu, şabalıdı (kahverengi),	56	VI	III Orta
8.	Kumlu, çakıllı, alüvyonlu taşkın yatağı	52	VI	
9.	Kumlu, kayalık, alüvyonlu çayır	50	V	
10.	Kumlu, bataklık çayır	42	V	
11.	Kumlu, bataklık çayır	34	IV	IV Aşağı
12.	Kayalık ve çakıllı nehir yatakları	16	II	V Koşullu olarak işe yaramaz
13.	Kayalık nehir yatakları	14	II	

Araştırılan alanların incelenmesinde G.Ş. Memmedov ve S.A. Arazilerin değerlendirilmesi, Hacıyev tarafından derlenen 1:150.000 ölçekli Nahçıvan Özerk Cumhuriyeti arazi haritasına göre yapılmıştır (Harita 2).



Harita 2. Nahçıvan Özerk Cumhuriyeti'nin arazi haritası

Çözüm

1. Araştırma nesnesi için derlenen tarımsal üretim gruplandırması, tarımsal iyileştirme önlemlerine ihtiyaç duyan arazi gruplarını da belirlememize olanak sağlamaktadır. Genel olarak bu gruplama, Abraqonus Çöküntüsü'nde yaygın olan tüm toprak tiplerini, alt tiplerini ve tür çeşitliliğini bir araya getirmektedir.
2. Topraklar, bölgesel ekolojik koşulların karmaşıklığına bağlı olarak daha küçük taksonomik birimlere ayrılır. Nihai kalite puanları daha önce de belirtildiği gibi "düzeltme" faktörleri (su basması, tuzlanma, erozyon, granülometrik bileşim, vb.) kullanılarak belirlendi.
3. Araştırma nesnesindeki toprak tipi puanı ve tarımsal üretim grubu bilinerek, doğal ve kültürel bitkilerin yetiştirildiği toprakların iyileştirilmesine yönelik tarımsal önlemlerin (ıslah,

gübreleme, erozyon önleyici önlemler, çalılık ve taşların temizlenmesi, sulama vb.) uygulanması belirlenebilir.

Öneriler

1. Abraqunus Çöküntüsü'nde arazinin en verimli şekilde kullanılabilmesi için öncelikle susuz alanlara su kanalları yapılmalı, bakımsız su kanalları ve göletler restore edilmelidir.
2. Gölün su taşkınlarından korunması, kuzey, kuzeydoğu ve kuzeybatı kesimlerinde ise erozyondan korunması ve verimliliğinin yeniden sağlanması amacıyla göl kıyılarında çok sayıda ağaç dikilmesi yönünde tedbirler alınmalıdır.
3. Çiftçiler ve bireysel çiftçiler, arazi kullanımında arazi kartogramlarına güvenmeli ve uzmanlardan sürekli tavsiye almalıdır.
4. Bölgenin ekolojik koşulları dikkate alınarak, yerel koşullara uygun daha verimli bitkilerin dikimi yapılmalı, ıslah önlemlerine dayalı arazi potansiyelinin etkin bir şekilde değerlendirilmesi sağlanmalı, su kaynakları mümkün olan en üst düzeyde kullanılmalıdır.

Yukarıda belirtilen tedbirlerin Abraqunus çöküntüsünde bilimsel temele dayalı olarak planlı bir şekilde uygulanması halinde, bölgedeki topraklar, ekili ve doğal bitkiler açısından etkin bir şekilde kullanılabilir.

EDEBİYAT

1. Babayev S.Y. Nahçıvan Özerk Cumhuriyeti Coğrafyası. Bakü: Elm, 1999, 226 s.
2. Hacıyev S.A. Nahçıvan Özerk Cumhuriyeti topraklarının eko-coğrafik koşulları. Bakü: MBM, 2009, 108 s.
3. Hacıyev S.A. Nahçıvan Özerk Cumhuriyeti'nde toprakların tarımsal ekolojisi. Metodik kaynaklar. Bakü: Elm, 2000, 40 s.
4. Mammadov G.Ş. Azerbaycan'da toprak reformu. Bakü: Elm, 2002, 411 s.
5. Mammadov G.Ş., Caferov A.B. Arazi değerlendirmesi. Bakü: Elm, 1997, 210 s.
6. Şafibeyov A.B. Toprak ve bitki analiz yöntemleri. Bakü: Azernaşr, 1964, 204 s.
7. Aliyev G.A., Zeynalov A.K. Nahçıvan ASSC toprakları. Bakü: Azerneshr, 1998, 235 s.
8. İbrahimov A.Ş. Nahçıvan Özerk Cumhuriyeti'nin bitki örtüsü ve milli-ekonomik önemi. Bakü: Elm, 2005, 230 s.
9. Mamedov R.G. Nahçıvan ASSR toprağının tarımsal fiziksel özelliklerine göre gruplandırılması konusunda deneyim //DAN Az. SSCB, 1968, s. 43-48

**THE ROLE OF BEES OF THE GENUS *HYLAEUS* FABRICIUS, 1793 IN
POLLINATION OF CORIANDER AND FENNEL PLANTS IN NAKHCHIVAN
AUTONOMOUS REPUBLIC**

**NAHÇIVAN ÖZERK CUMHURİYETİ'NDE KİŞNİŞ VE REZENE BİTKİLERİNİN
TOZLAŞMASINDA *HYLAEUS* FABRICIUS, 1793 CİNSİ ARILARIN ROLÜ**

Aydan MAHARRAMLİ

Nahcivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0005-2426-8057>

Mahir MAHARRAMOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Veterinary Medicine, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0002-4130-7071>

ABSTRACT

Introduction and Purpose: The genus *Hylaeus* Fabricius, belonging to the *Colletidae* family, comprises 760 species worldwide, with 220 in the Palearctic region, 54 in the Caucasus, 48 in Azerbaijan, and 29 in the Nakhchivan Autonomous Republic, which is an integral part of it. These bees are distinguished by their small size and yellow or white markings. Although they lack specialized pollen-carrying structures, these solitary bees play an active role in pollination. *Hylaeus* bees are particularly effective pollinators of economically important plants in the Apiaceae family. While the fauna of this genus has been well studied in the autonomous republic, their role as pollinators remains insufficiently investigated.

Materials and Methods: The research was conducted in 2024 in coriander and fennel fields located in private backyards across the lowland, low mountainous, and mid-mountainous regions of the autonomous republic. During the flowering period of these plants, *Hylaeus* bees were collected using insect nets and yellow traps. Species identification was carried out both by us and by entomologists from the Biodiversity Scientific Center of the Far Eastern Department of the Russian Academy of Sciences.

Results: Fifteen species of the genus *Hylaeus* were identified in coriander (*Coriandrum* L.) and fennel (*Anethum* L.) fields cultivated in private backyards across the lowland, low-mountainous, and mid-mountainous regions of the autonomous republic.

Discussion and Conclusion: In the pollination of coriander and fennel plants grown in the lowland, low-mountainous, and mid-mountainous regions of the autonomous republic, the following *Hylaeus* species played a role: *Hylaeus cornutus* Curtis, 1831; *Hylaeus moricei* (Friese, 1898); *Hylaeus scutellaris* Morawitz, 1874; *Hylaeus signatus* (Panzer, 1798); and *Hylaeus variegatus* (Fabricius, 1798). In the low-mountainous and mid-mountainous regions, additional species were identified: *Hylaeus lineolatus* (Schenck, 1861); *Hylaeus meridionalis* Förster, 1871; and *Hylaeus rugicollis* Morawitz, 1874. Only in the mid-mountainous region, the following species were found: *Hylaeus breviceps* Morawitz, 1876; *Hylaeus communis*

Nylander, 1852; *Hylaeus leptcephalus* (Morawitz, 1871); *Hylaeus sinuatus* (Schenck, 1853); *Hylaeus styriacus* Förster, 1871; and *Hylaeus variolaris* Morawitz, 1876. Despite lacking specialized pollen transportation structures, bees of this genus play a significant role in the ecosystem. Their presence among plants of the *Apiaceae* family suggests they contribute to pollination, although their efficiency is lower than that of some other bee species. However, their ability to survive in arid conditions highlights their importance in maintaining local biodiversity.

Keywords: Nakhchivan, mid-mountainous, *Coriandrum*, *Anethum*, *Hylaeus*.

ÖZET

Giriş ve Amaç: *Colletidae* familyasına ait *Hylaeus* Fabricius cinsinin dünyada 760 türü, Paleartik'te 220, Kafkasya'da 54, Azerbaycan'da 48, onun ayrılmaz bir parçası olan Nahçıvan Özerk Cumhuriyeti'nde ise 29 türü bulunmaktadır. Küçük boyutları ve sarı ve ya beyaz işaretleriyle ayırt edilirler. Yalnız yaşayan bu arılar polen taşıyan özel yapılara sahip olmasalar da tozlaşmada aktif rol oynarlar. *Hylaeus* cinsinin arıları, *Apiaceae* familyasına ait ekonomik açıdan önemli bitkilerin tozlaşmasında daha aktif olarak yer almaktadır. Bu cinse ait arıların faunası özerk cumhuriyette iyi çalışılmış olmasına rağmen, tozlayıcı olarak rolleri yeterince araştırılmamıştır.

Gereç ve Yöntem: Araştırma çalışması 2024 yılında özerk cumhuriyetin ova, alçak dağlık ve orta dağlık bölgelerinde özel bahçelerde yetiştirilen kişniş ve rezene tarlalarında gerçekleştirildi. Söz konusu bitkilerin çiçeklenme döneminde *Hylaeus* cinsine ait arılar böcek ağları ve sarı tuzaklar kullanılarak yakalandı. Türlerin tespiti hem tarafımızdan hem de Rusya Bilimler Akademisi Uzak Doğu Bölümü Biyoçeşitlilik Bilim Merkezi entomologları tarafından gerçekleştirildi.

Bulgular: Özerk cumhuriyetin ova, alçak dağlık ve orta dağlık bölgelerinde, özel bahçelerde yetiştirilen kişniş (*Coriandrum* L.) ve rezene (*Anethum* L.) tarlalarda *Hylaeus* cinse ait 15 tür tespit edilmiştir.

Tartışma ve Sonuç: Özerk cumhuriyetin ova, alçak dağlık ve orta dağlık bölgelerinde yetiştirilen kişniş ve rezene bitkilerinin tozlaşmasında *Hylaeus cornutus* Curtis, 1831; *Hylaeus moricei* (Friese, 1898); *Hylaeus scutellaris* Morawitz, 1874; *Hylaeus signatus* (Panzer, 1798); *Hylaeus variegatus* (Fabricius, 1798); sadece alçak dağlık ve orta dağlık bölgelerde *Hylaeus lineolatus* (Schenck, 1861); *Hylaeus meridionalis* Förster, 1871; *Hylaeus rugicollis* Morawitz, 1874; sadece orta dağlık bölgede *Hylaeus breviceps* Morawitz, 1876; *Hylaeus communis* Nylander, 1852; *Hylaeus leptcephalus* (Morawitz, 1871); *Hylaeus sinuatus* (Schenck, 1853); *Hylaeus styriacus* Förster, 1871; *Hylaeus variolaris* Morawitz, 1876 türlerinin rolü ortaya çıkarıldı. Polen taşımanın özel yöntemlerine rağmen bu cinse ait arılar ekosistemde önemli bir rol oynamaktadır. *Apiaceae* familyasına ait bitkiler arasında bulunmaları, verimleri diğer bazı arılara göre daha düşük olmasına rağmen tozlaşmaya katkıda bulunabileceklerini düşündürmektedir. Ancak kurak koşullarda hayatta kalabilme yetenekleri, yerel biyolojik çeşitliliğin önemli bir parçası olduklarını gösteriyor.

Anahtar Kelimeler: Nahçıvan, orta dağlık, *Coriandrum*, *Anethum*, *Hylaeus*.

IMPACT OF GLOBAL CLIMATE CHANGE ON BEEKEEPING

İKLİM DEĞİŞİKLİĞİNİN ARICILIĞA ETKİSİ

Ali TAHİROV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Veterinary Medicine, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0006-1280-7292>

Yunis RUSTAMLI

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Veterinary Medicine, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0000-0786-0597>

ABSTRACT

Introduction and Purpose: Climate change It has become one of the main global problems of the 21st century. The life of a honey bee (*Apis mellifera* L.) depends on many factors, such as climatic conditions and the environment. Global warming, changes in ecosystems, plant flowering times and other climatic changes have begun to have a noticeable impact on bees and their productivity. The aim of the study is to investigate the threats and possible prospects for beekeeping in changing climatic conditions.

Materials and Methods: In the experiments, bee colonies of the Nakhchivan population of the Yellow Caucasian bee breed (*Apis mellifera caucasica flova* populatio Nachitshevanica), distributed in the territory of the Nakhchivan Autonomous Republic of the Republic of Azerbaijan, were used. The main climatic zones of the study area were 1) mountainous, 2) middle mountainous and 3) lowland areas of the Araz River, covering the Zangezur and Daralayaz mountain systems of the Greater Caucasus. The experimental group consisted of families similar in terms of power, honey, bee growth and comb size according to the principle of analogs. The study used abstract-logical, monographic, analytical research methods, results of scientific publications and our personal research. The collected materials were processed using the biometric method proposed by N.A. Plokhinsky.

Results: Threats to beekeeping: Sudden temperature changes, hot and dry seasons or unusually cold winters in the Nakhchivan Autonomous Republic increase stress in bees. After wintering (March 18, 24), the power of bee colonies settled in the central mountain area was found to be 13.5% ($P>0.95$) and 14.7% ($P\leq 0.95$) higher than the power of bees in the Araz-bayu lowland and Zangezur mountain areas, respectively. Air temperature, atmospheric precipitation and vegetation cover play a special role in the selection of beekeeping sites, the beginning of vegetation in nectar-bearing plants and the displacement of individual development stages.

Discussion and Conclusion: Climate change creates both difficulties and opportunities for beekeeping. Threats such as changing flowering times, extreme weather events and increasing parasite numbers require beekeepers to adapt to the environment and implement new approaches and technologies. However, proper management of beekeeping enterprises and

transition to environmentally sustainable beekeeping will not only help overcome the challenges but also reap some benefits from these changes.

Keywords: climate change, global warming, honey bee, threat, productivity

ÖZET

Giriş ve Amaç: İklim değişikliği 21.yüzyılın başlıca küresel sorunlarından biri haline gelmiştir. Bir bal arısının (*Apis mellifera* L.) yaşamı, iklim koşulları ve çevre gibi pek çok faktöre bağlıdır. Küresel ısınma, ekosistemlerdeki değişimler, bitki çiçeklenme zamanları ve diğer iklim değişiklikleri arılar ve üretkenlikleri üzerinde gözle görülür bir etkiye sahip olmaya başladı. Çalışmanın amacı değişen iklim koşullarında arıcılık için tehditleri ve olası beklentileri araştırmaktır.

Gereç ve Yöntem: Deneylede, Azerbaycan Cumhuriyeti Nahçıvan Özerk Cumhuriyeti topraklarında dağılım gösteren Sarı Kafkas arı cinsinin (*Apis mellifera caucasuca flova populatio Nachitshevanica*) Nahçıvan popülasyonuna ait arı kolonileri kullanıldı. Çalışma alanının ana iklim bölgeleri Büyük Kafkasya'nın Zangezur ve Daralayaz dağ sistemlerini kapsayan 1) dağlık, 2) orta dağlık ve 3) Araz nehri ova alanlarıydı. Dene grubu analoglar prensipine göre güç, bal, arı büyümesi ve petek boyutu bakımından benzer ailelerden oluşuyordu. Çalışmada soyut-mantıksal, monografik, analitik araştırma yöntemlerinden, bilimsel yayınların sonuçlarından ve kişisel araştırmalarımızdan yararlanılmıştır. Toplanan materyaller N.A. Plokhinsky tarafından önerilen biyometrik yöntem kullanılarak işlendi.

Bulgular: Arıcılığa yönelik tehditler: Nahçıvan Özerk Cumhuriyeti'nde ani sıcaklık değişimleri, sıcak ve kurak mevsimler veya alışılmadık derecede soğuk kışlar arılarda stresi artırmaktadır. Kışlatma sonrasında (18.03.24) orta dağlık alanda yerleşen arı kolonilerinin gücü, Araz-bayu ovası ve Zangezur dağlık alanlarındaki arıların gücünden sırasıyla %13,5 ($P>0,95$) ve %14,7 ($P\leq 0,95$) daha yüksek bulunmuştur. Arıcılık işletmelerinin yer seçiminde, nektar taşıyan bitkilerde vejetasyon başlangıcında ve bireysel gelişme evrelerinin yer değiştirmesi sürecinde hava sıcaklığı, atmosferik yağış ve bitki örtüsü özel rol oynamaktadır.

Tartışma ve Sonuç: İklim değişikliği arıcılık için hem zorluklar hem de fırsatlar yaratmaktadır. Çiçeklenme zamanlarının değişmesi, aşırı hava olayları ve artan parazit sayısı gibi tehditler arıcıların çevreye uyum sağlamasını ve yeni yaklaşımlar ve teknolojiler uygulamasını gerektiriyor. Ancak arıcılık işletmelerinin doğru yönetilmesi ve çevresel açıdan sürdürülebilir arıcılığa geçiş, sadece zorlukların üstesinden gelinmesine değil, aynı zamanda bu değişikliklerden bazı faydaların elde edilmesine de yardımcı olacaktır.

Anahtar kelimeler: iklim değişikliği, küresel ısınma, bal arısı, tehdit, verimlilik

**THE ECOLOGICAL RESULTS OF IMPACT OF THE ARPACHAY RESERVOIR
IRRIGATION SYSTEMS TO THE ENVIRONMENT**

**ARPAÇAY SU ANBARININ SUVARMA SİSTEMLƏRİNİN ƏTRAF MÜHÜTƏ
TƏSİRİNİN EKOLOJİ NƏTİCƏLƏRİ**

Novruzlu Aytən

Naxçıvan Dövlət Universiteti, müəllim

Orcid id : 0009-0000-0931-8785

Əli Qurbanov

Naxçıvan Dövlət Universiteti, baş müəllim

Orcid id : 0009-0004-8789-7451

ABSTRACT

The irrigation systems based on the Arpachay Reservoir which was built in 1977 on the Eastern Arpachay River with 150 million m³ water capacity and ecological results of its impact to the environment have been defined and scientifically analyzed using various research methods.

As Arpachay Reservoir was constructed on the channel called “Deveolen” on the Eastern Arpachay River and 80% of its main basin of supply is located in the territory of Armenia, recently some problems are being formed in flow regime connected with floating of the river-water into Goycha Lake by Armenia. It causes to the lack of water in the reservoir during vegetation period. As a result of the carried researches it was defined that, the Arpachay Reservoir providing 17.311 hectare of land in Sharur, Sadarak and Kangarli regions and hydro-technical installations formed on the base of its impact to the environment seriously when agro-technical standards are used improperly. These impacts result with ecological tension in the environment. Many ecological problems arise in the reservoir itself, as well as in the main and inter-farm and inner-farm canals due to technical deficiencies. As a result, salinization and marshland areas are formed in the territory.

Many ecological problems formed as a result of irrigation systems impact to the environment belonging to the Arpachay Reservoir have been defined. In addition, water conductive hydro-technical installations belonging to this irrigation system, the ecological condition observed in the canals, closed irrigation systems and cultivated fields has been analyzed and measures taken against them have been proposed. The formation causes of existed water loss, salinization and formation causes of marshland territories were searched scientifically and ways of liquidating the problems were stated. The ecological condition formed as a result of carried researches has been studied in detail and maps and schemes characterizing the research area have been prepared. As well as the serious ecological situation caused by the lack of ecological flow in the channel besides the Arpachay Reservoir has been analyzed.

Key words: Arpachay Reservoir, irrigation systems, water loss, irrigation canals, salinization map, marshland, Upper Yayjı water distributive installation, irrigation standards.

Xülasə

Məqalədə Şərqi Arpaçay çayı üzərində inşa edilmiş Arpaçay su anbarına əsaslanan suvarma sistemləri və bunun ətraf mühitə təsirinin ekoloji nəticələri verilmişdir. Eyni zamanda bu suvarma sisteminə aid olan suötürücü hidrotexniki qurğular, kanallarda, qapalı suvarma sistemlərində və əkin sahələrində müşahidə olunan ekoloji vəziyyət təhlil edilmiş, onlara qarşı mübarizə tədbirləri təklif olunmuşdur. Mövcud olan su itkilərinin yaranma səbəbləri, şoranlaşmanın baş verməsi, bataqlıqlaşma kimi ərazilərin yaranma səbəbləri elmi surətdə tədqiq olunub və problemlərin aradan qaldırılması yolları göstərilmişdir. Həmçinin Arpaçay su anbarından sonra çay yatağında ekoloji axının olmaması səbəbindən yaranan ciddi ekoloji vəziyyət təhlil olunmuşdur.

Açar sözlər: Arpaçay su anbarı, suvarma sistemləri, su itkisi, suvarma kanalları, şoranlaşma xəritəsi, bataqlıqlaşma, Yuxarı Yaycı supaylayıcı qurğusu, suvarma normaları.

Arpaçay su anbarı 1977-ci ildə inşa edilmiş və Şərur, Sədərək və Kəngərli rayonu suvarma sistemlərinin əsas su mənbəyidir. Güzgü səthi maksimum səviyyə zamanı okean səviyyəsindən 969 m hündürlükdə yerləşir. Bu hündürlükdə onun sahəsi 7,15 km², tutumu 150 mln m³-ə çatır. Faydalı su tutumu 130 mln m³-dir. Su anbarının bəndinin hündürlüyü 69 metr, uzunluğu 480 metr, ən yuxarı hissədə eni 10 metrdir. Bəndin mərkəzi hissəsi 81 metr hündürlüyə malik dəmir-beton konstruksiyadan ibarətdir (4).



Şəkil-1. Arpaçay su anbarı

Arpaçay su anbarında təqribən 1 km çay mənsəbinə doğru Aşağı Yaycı su paylayıcı hidrotexniki qurğu tikilmişdir. Onun tam su tutumu 24,5 mln.m³, faydalı tutumu isə 20,0 mln.m³-dir. Bəndin hündürlüyü 35 metrdir. Sulu dövrdə maksimum səviyyə 887,5 metr hündürlükdə yerləşir və bu zaman su güzgü səthi 460 hektardır. Bu su anbarı suvarma sistemlərinin əsas hissəsini təşkil edir. Su paylayıcı kanallar, yəni magistral kanallar bu su anbarından başlamaqla Arpaçay su anbarından ayrılan suvarma suyunun sağ və sol kanallarına ötürür. Əlavə olaraq üzərində nasos stansiyaları da qurulmuşdur. Suvarma kanalları ilə yanaşı Şərur, Kəngərli və Sədərək düzənliklərinin minlərlə hektar əkin sahələrini, kanallar üzərində fəaliyyət göstərən nasos stansiyaları su ilə təmin edir.

Arpaçay su anarının əsas su mənbəyi olan Şərqi Arpaçay çayının il ərzində axım rejimi müxtəlif olur. Fəsilər üzrə çayın su sərfinin dəyişməsi su anbarında səviyyə tərəddüdünə səbəb olur. Səviyyə tərəddüdünün dəyişməsi bir neçə səbəbdən yaranır: əsasən yağıntıların fəsilər üzrə dəyişməsi, vegetasiya dövründə Ermənistan ərazisində sudan istifadənin artması nəticəsində axımın azalması və intensiv qarların əriməsi və intensiv yağışların yağması ilə baş vermiş sellər nəticəsində isə səviyyə rejiminin yüksəlməsi müşahidə edilir. Vegetasiya dövrünün bitməsi ilə su anbarında suyun toplanması başlayır ki, bu da vegetasiya dövrünün başlanmasına qədər davam edir. Ümumiyyətlə axımın il ərzində paylanmasının üç fazaya ayırılmışıq.

Birinci faza payız-qış (sentyabr-fevral) dövrünü əhatə edir. İkinci fazada su sərfi nisbətən davamlıdır. Yalnız bəzi illərdə (oktyabr-noyabr aylarında) düşən yığıntılar hesabına qısamüddətli zəif daşqınlar baş verir. Üçüncü faza isə il ərzində ən kiçik sərf yanvar ayına düşür. Dövr ərzində çaylar illik axımın 10-15%-ə qədərini axıdır (cədvəl 1).

Cədvəl 1

Arpaçayda axının paylanması

Çaylar	Məntəqələr	Yaz III-V	Yay VI-VIII	Payız IX-XI	Qış XII-II	Vegetasiya dövrü VII-VIII	III-V	III-V	VI-VIII
							XII-II	VI-VIII	III-V
Arpaçay	Arpa	50	27	12	11	10	4,5	1,8	0,6


Cədvəldən də görüldüyü kimi Şərqi Arpaçayın axım rejimi yaz və yay aylarında maksimuma çatır. Bu dövrlərdə əsasən su anbarını maksimum səviyyədə doldurmaq mümkün olur (4, 12).

Son illərdə Arpaçay su anbarının səviyyəsinin aşağı düşməsi müşahidə olunur. Ermənistanın Azərbaycana qarşı yönəldilən düşmənçilik siyasəti nəticəsində Arpaçay su anbarını su ilə doldurmaq mümkün deyildir. Belə ki, Ermənistan Respublikası ərazisində 1981-ci ildə ikinci Arpaçay üzərində həcmi 24 mln. m³ olan su paylayıcı su anbarı istifadəyə verilmişdir. Onun tikintisi ilə eyni zamanda Şərqi Arpaçaydan Göyçə gölünə çəkilmiş tunel və su kəməri vasitəsi ilə ildə 280 mln. m³ su (çayın illik axımının 40%-i) Göyçə gölünə axıdılmağa başlandı. Oudur ki, yerdə qalan su ilə Arpaçay su anbarını doldurmaq mümkün olmur və əvvəllər suvarma üçün nəzərdə tutulan su təmin olunmadığı üçün torpaqların xeyli hissəsini su çatışmamazlığı üzündən istifadə etmək mümkün olmur.

Respublikanın ən böyük əkin sahəsinə malik olan Şərur rayonunun əsas su mənbəyi Arpaçay su anbarıdır ki, bu da suvarma suyunun 84%-ni təmin edir. Şərur rayonunun ümumi suvarılan sahəsi 22432 ha-dır. Əkin sahələrinin suvarılan faktiki sahəsi 12166 ha-ya bərabərdir. Bunu ayrı-ayrı mənbələr üzrə götürdükdə təkcə Arpaçay su anbarı vasitəsilə suvarılan əkin sahəsi 9860 ha-ya bərabər olur (12).

Şərur rayonunun əkin sahələri üzrə istifadə olunan suyun ümumi miqdarı 154,319 mln.m³ təşkil ediyi halda təkcə Arpaçay su anbarı vasitəsilə suvarılan torpaqlarda istifadə olunan suyun miqdarı isə 120, 467 mln.m³-dir. Bu da Arpaçay suvarma sistemlərində Arpaçay su anbarının əhəmiyyətinin göstərir (13).

Arpaçay su anbarından suvarmaya lazım olan miqdarda su magistral kanallar vasitəsilə götürülüb suvarılan sahələrə nəql edilir. Yayıcı su anbarının su paylayıcı qurğusundan magistral kanallara suyu, sudan istifadə planına müvafiq olaraq tələb olan miqdarda və lazım olan vaxtda götürüb kanala verməkdən ötrü Suayrıcı qurğu qurulmuşdur. Bu hidroyexniki qurğu vasitəsilə su suvarma mənbəyindən götürülərək magistral kanallara ötürülür, oradan isə paylayıcı və təsərrüfatlar arası kanallar vasitəsilə əkin sahələrinə verilir.



Arpaçay su anbarı meliorasiya xəritəsi

Arpaçay su anbarı suvarma sistemləri Şərur, Sədərək və Kəngərli rayonlarının suvarma əkinçiliyində mühüm yer tutmaqla böyük təsərrüfat əhəmiyyətinə malikdir.

Tədqiqat nəticəsində müəyyən edilmişdir ki, hər üç rayonda 17311 hektar sahəni daima su ilə təmin edən Arpaçay su anbarı və onun əsasında yaradılmış hidroyexniki qurğular sudan aqrotexniki qaydalara uyğun olmayaraq istifadə edildikdə ətraf mühitə ciddi təsirlər göstərir. Bu təsirlər nəticəsində ətraf mühitdə ekoloji gərgin vəziyyət yaranır. Həm su anbarının özündə, həm magistral və təsərrüfat arası həm də təsərrüfat daxili kanallarda texniki səbəblər üzündən bir çox ekoloji problemlər mövcuddur. Nəticədə ərazidə şoranlaşma və bataqlıqlaşma sahələri yaranmaqdadır.

Şərur. Sadərak və Kəngərli rayonlarının şoranlaşma xəritə-sxemi

Suvarma meliorasiyasının düzgün idarə edilməməsi nəticəsində yəni sahələrə suyun verilməsi zamanı bitkinin suvarma normaları nəzərə alınmaması səbəbindən artıq verilən sular əkin sahələrində səthi axın yaradır ki, bu da sahələrdə üst münbit qatın (humusun) yuyulmasına səbəb olur. Eyni zamanda nəzərə alsaq ki tədqiqat ərazisi Arazboyu düzənlik sahəni əhatə edir o zaman sahələrə verilən normadan artıq su ərazidə filtrasiya nəticəsində qırt suyunun səviyyəsinin qalxmasına şərait yaradır. Sonrakı proseslərdə isə həmin ərazilər ya şoranlaşmağa məruz qalır və ya qırt suyunun səviyyəsinin qalxması ilə bataqlıqlaşmış sahələrə çevrilirlər. Belə sahələrə həm Sol sahil kanalı suvarma sistemlərində, həm də sağ sahil suvarma sistemlərində rast gəlinir. Əsasən də Dəmirçi, Daşarx və digər yaşayış məntəqələrinə məxsus olan əkin sahələri və torpaq kanalların ətrafında rast gəlinir.



Şəkil-3. Şoranlaşmış əkin sahələri

Yuxarı Sovxoz kanalı əsasən qruntdan ibarət olduğu üçün onun ətrafında bataqlıqlaşmış sahələrə daha çox rast gəlinir. Təsərrüfat arası və təsərrüfat daxili kanallar da torpaq kanallar olduğu üçün onlar yaxın hissələrdə şoranlaşmış torpaq sahələri və çəmənləşmiş əkin sahələrinə tez-tez rast gəlmək olur. Tədqiqat zamanı biz bu ərazilərdə mövcud olan şoranlaşmış sahələri müəyyən edərək onların sahələrini hesablayaraq xəritə-sxemin hazırlamışıq. Mövcud xəritədən görüldüyü kimi əsasən şoranlaşmış sahələr Araz çayına yaxın olan əkin sahələrində rast gəlinir. Arazboyu düzənlikdə qrunut suyunun səviyyəsi səthə yaxın olduğu üçün (1-3 m arası) sahəyə artıq verilən su qrunut suları ilə birləşdikdən sonra arid iqlim şəraitində buxarlanaraq səthdə şoranlaşmış və ya təkrar şoranlaşmış sahələr yaradır. Bəzən su çatışmaması səbəbindən tərk edilmiş əkin sahələrinə də rast gəlmək olur.

90-cı illərdə aparılmış torpaq islahatından sonra demək olar ki, təsərrüfat arası və təsərrüfat daxili kanallar demək olar ki əsasən sıradan çıxmış və ya tamamilə yox vəziyyətindədir. Belə bir vəziyyətdə təsərrüfatlara suyu norma daxilində çatdırılması çətinliklər yaradır. Verilən sular isə düzgün tənzimlənmədiyi üçün torpağın həm aerasiya zonasını həm də hidrogeoloji xüsusiyyətlərini pozaraq yararsız hala salır.

Əsas magistral kanallar üzərində olan hidrotexniki qurğularda olan çatışmamazlıqlar və su ötürücü qurğulardan çıxan istifadəsiz sular nəzarətsiz axaraq hiqrofil bitkilərdən ibarət olan bataqlıq sahələr yaradır. Həmçinin vizual müşahidələr zamanı suötürücü qapılardan sonra su ölçü postlarının olmadığı müəyyən edilmişdir. Bu ölçü postlarının olmaması əkin sahələrinə suyun normaya uyğun verilməsi mümkün deyildir. Nəticədə isə əkin sahələrinə ya normadan artıq su verilir, ya da normadan az suyun verilməsi sahədə quraqlıq yaradır.

Magistral kanallardan olan Sağ sahil və Sol sahil kanalları 1977- ci ildən istifadəyə verildiyi üçün bəzi hissələrində texniki cəhətdən qüsurlar mövcuddur. Bunun səbəbi isə kanalları suvarmaya hazırlarkən təmizləmə zamanı düzgün mexaniki təmizləmə seçilmədiyi üçün mexanizmlər tərəfindən qirilib dağılmağıdır. Həmin sahələr isə həm kanaldan su itkisinin artmasına, həm də ətraf mühütdə şoranlaşma və bataqlıqlaşma yaranmasına səbəb olur. Lakin kanallar təmir edilərkən xüsusi inşaat materealları vasitəsilə təmir olunmalıdır ki, kanalda su olan vaxtı cillər və digər hiqrofil bitkilər bitməsin. Yuxarı Sovoz kanalında da ekoloji vəziyyət beton kanallardan daha acınacaqlı vəziyyətdədir. Kanal əsasən torpaqdan olduğu üçün burada

birki örtüyü da çox və tez inkişaf edir. Hətta bitkilər kanalda suyun axınına çox ciddi təsir edə bilər. Torpaq kanal olduğu üçün burada filtrasiya prosesi güclü getdiyindən su itkisi daha çox müşahidə edilir.



Şəkil-2. *Yuxarı sovxoz kanalı*

Ümumiyyətlə Arpaçay su anbarından su götürən suvarma kanalları 9467 ha sahəni su ilə təmin edir. Suvarma kanalları və onların suvardıqı sahələr haqqında məlumat 2-ci cədvəldə verilmişdir (13).

Arpaçay su anbarı suvarma sistemlərinin kanalları haqqında məlumat cədvəl-2

Obyektin adı	Uzunluğu (km)	Qulluq etdiyi sahə (ha)	Faktiki sahə (ha)
Yuxarı Sovxoz Kanalı	52,0	834	486
Püsyən arxı	15,0	2105	1334
Sağsahil kanalı	22,0	3131	2313
Solsahil kanalı	25,0	4356	514
Qurdçubasar arxı	23,0	1270	803
Lehməbasar arxı	27,0	3821	2715
Kosacan arxı	1,5	238	172
Ələhli arxı	6,0	287	217
Aralıq arxı	19,0	298	196
Çərçiboğan arxı	20,0	971	717
Cəmi	17311	9467

Arpaçay su anbarından su götürən ən uzun suvarma kanalı Yuxarı Sovxoz kanalıdır ki, onun da uzunluğu 52 km-dir. Keçmiş Şərur-Kəngərli suvarma sisteminin əsasını təşkil edən kanal

qismən torpaq kanal olması ilə böyük bir suvarılan sahəni əhatə edərək 486 ha ərazini su ilə təmin edir. Ən çox sahə suvaran kanal isə Lehməbasar arxıdır ki, onun da suvardığı faktiki sahə 2715 ha-dır.



Şəkil-2. Sağ sahil kanalı

Arpaçay sağ və sol sahil kannalları Şərur, Sədərək və Kəngərli suvarma sistemlərinin əsasını təşkil etdiyində bu kanallardan suyu paylanmasında təsərrüfatlar arası kanallardan başqa stasionar nasos stansiyalarında istifadə edilir. Nasos stansiyalarından sular sahələrə qapalı suvarma sistemləri ilə ötürüldüyündən sistemi təşkil edən borularda olan nöqsanlar səbəbindən əkin sahələrində və onların sahələrdə olan çıxış hissələrində böyük həcmdə sızmalar üzündən ərazidə bataqlıq sahələr müşahidə olunur. Qapalı suvarma sistemlərində buxarlanma və fitirasiy yolu ilə su itkisi olmasa da texniki səbəblər üzündən su itkisi baş verir.



Şəkil-6. Sol sahil kanalı

Bunun səbəbi torpağın altında boruların keyfiyyətsiz birləşdirilməsi və su açarlarının (qıdrantların) tam bağlanamamasıdır. Nasosların yüksək təzyiqlə suyu sistemə vurmasını nəzərə alsaq borulardan su itkisinin miqdarını təsəvvür etmək olar.

Nəticə

Yuxarı Yaycı su anbarı su ötürücü qurğusundan sonra Şərqi Arpaçay yatağında ekoloji vəziyyətin gərgin olduğu tədqiqat zamanı müşahidə edilmişdir. Uzun illərdir ki, çay yatağında ekoloji axın müşahidə edilmir. Çayda səthə çıxan filtrasiya suları ətraf yaşayış məntəqələri tərəfindən toplanıb götürülür, ya da yenidən filtrasiya olunaraq yataqda yox olur. Bunun isə ekoloji təsiri çay yatağında formalaşan təbii komplekslərin flora və faunanın yo olmasına səbəb olur. Cenevrə konqresinə görə çay yataqlarında hər hansı hidrotexniki qurğular yaradılarkən çay yatağında flora və faunanın qorunub saxlanması üçün ekoloji axın olması məcburiyyəti qoyulmuşdur. Bu şərt həm tranzit çaylara, həm də yerli çaylara aiddir. Lakin suvarma sistemlərinin yaradılmasında bu tələblər nəzərə alınmamışdır. Nəticədə çay yatağında ekoloji gərgin vəziyyət yaranmışdır.

Bütün Cənubi Qafqazda su ilə az təmin olunan region olaraq Naxçıvan Muxtar Respublikasında su probleminin daha çox yaşanmaması üçün su mənbələrimizdən daha səmərəli istifadə etmək lazımdır. Bu məqsədlə tədqiqat obyektimiz olan Arpaçay su anbarı suvarma sistemlərində əkin sahələrinə suyun itkisiz çatdırılması üçün bir çox tədbirlər görülməli, ətraf mühərdə ekoloji gərginlik yaradan nöqsanları aradan qaldırmaq lazımdır. Bunun üçün suvarma sularının dəqiq hesabı aparılmalı, sahələrə su əkinin növündən və suvarma normasından asılı olaraq norma daxilində verilməsi üçün hidrometrik postlar qurulmalıdır. Kanallarda və suötürücü qurğularda su itkisinin qarşısını almaq üçün əsaslı təmir işləri aparılmalı, qapalı suvarma sistemlərində mövcud olan sızmalar müəyyənləşdirilib təmir olunmalı, Şərqi Arpaçay çayının yatağında ekoloji axın təmin olunmalıdır. Arpaçay su anbarı Ermənistanla sərhəddə yerləşdiyinə görə daimi olaraq su anbarının suyunda kimyəvi və radiaktiv analizlər aparılmalıdır.

Ədəbiyyat

1. Н.Н.Бабабяйли. Нахчыван МР яразисиндя сятщ суларынын тябии жобрафи бахымдан гиймятляндирилимясиня даир. Азярбайжан Жобрафия Жямийятинин ясярляри. БІХ жилд, с.164-167
2. Naxçıvan Muxtar Sovet Sosialist Respublikası / Naxçıvan MSSR-nin 50 illiyinə həsr olunmuş elmi-praktik konfransın materialları. 1975. s.297-333
3. Naxçıvan Muxtar Respublikasının coğrafiyası I cild
4. M.Məmmədov. Azərbaycanın hidroqrafiyası. Bakı-2002. 266 s
5. Ə.K.Qurbanov. Naxçıvan MSSR ərazisində suvarma irriqasiya eroziyası və ona qarşı mübarizə tədbirləri. Naxçıvan sahələrarası ərazisi ET informasiya və təbliğat mərkəzi, №24, 1989. səh 4-6
6. Ə.K.Qurbanov. Naxçıvan MR ərazisində aqroiirqasiya erozisiyası Azərbaycanca emlin inkişafı və regional problemlər. 2005
7. Ə.K.Qurbanov. Arpaçay su anbarının ekocoğrafi xüsusiyyətləri. NDU elmi əsərlər № 1 (49), 2013. s.100 – 104
8. Ə.K.Qurbanov. NaxçıvanMR-da çayların tənzimlənməsində su anbarlarının rolu. Naxçıvan Muxtar Respublikasının Yaranması: Tarix və Müasirlik. Naxçıvan -2007 s.291-297
9. Ə.K.Qurbanov. Şərqi arpaçayın hidroloji xüsusiyyətləri. Azərbaycan Milli Aerkosmik Agentliyinin xəbərləri cild 19. 2016 № 2 s. 42-46
10. Ə.K.Qurbanov. Şərur rayonunun irriqasiya sistemləri vəziyyəti. İpək yolu NDU

11. Ə.K.Qurbanov. Sədərək, Şərur, Kəngərli düzənliklərində səhralaşmaya səbəb olan amillərin xarakteristikası. "Azərbaycan respublikasının kənd təsərrüfatında innovativ texnologiyalar və ekoloji problemlər" 14-15 dekabr 2023-cü il tarixində keçiriləcək respublika elmi konfransı"-QUBA . 14-15 dekabr 2023. Səh, 68-76
12. Naxçıvan Muxtar Respublikası Su Ehtiyatları Agentliyinin illik məlumatları.
13. Şərur rayon Su Təsərrüfatı İdarəsinin məlumatları.
14. А.Г.Садыгов, Ш.М.Амишов. "Гидрогеолого-мелиоративная обстановку на орошаемых землях Нахичеванской АР и пути её оптимизации". Нахи-чеван 1991.

FLORA AND ENDEMIC PLANTS OF ATATURK DAM BASIN (ŞANLIURFA/TURKEY)

Mustafa ÇETİNER

İğdır University, Graduate Education Institute, Department of Agricultural Sciences, 76100
İğdır, Türkiye
ORCID ID: <https://orcid.org/0000-0003-0022-0321>

Ahmet Zafer TEL

İğdır University, Faculty of Agriculture, Department of Agricultural Biotechnology, 76100
İğdır, Türkiye
ORCID ID: <https://orcid.org/0000-0002-1204-3839>

ABSTRACT

Introduction and Purpose: All plants of a certain place or region are called flora. This study was obtained from my master's thesis titled Flora of the Şanlıurfa Section of Atatürk Dam Lake Basin, which was carried out between 2018-2020. In this study, the floristic characteristics and endemic plants of Atatürk Dam Lake Basin in Şanlıurfa province located in Southeastern Anatolia Region, one of the 7 geographical regions of Turkey, were investigated. The study area is located in C7 square according to Davis's Grid system and it was carried out in order to carry out a floristic study specific to C7 square and to identify taxa that may be new records.

Materials and Methods: The plant specimens collected during the vegetation periods were pressed, dried and made into suitable specimens for herbarium. The works of Davis, Davis et al. and Güner et al. were used as the basis for the identification of the plants. Some plants that were difficult to identify were identified with the help of experts in their field. The herbarium of each taxon was checked and written from the work of Brummitt and Powell. Life forms of all taxa were determined according to Raunkiaer. The endangerment categories of endemic and rare taxa were determined according to the study of Ekim et al.

Results: As a result of many field trips to the study area, 460 plant samples were collected. By identification of these plant samples, 291 taxa belonging to 53 families and 190 genera were identified. Of these taxa, 203 species, 49 subspecies and 39 varieties were identified. In the research area, 9 endemic and 7 non-endemic rare taxa were identified. Detected in the research area endemic plants and their danger categories are as follows: *Astragalus macrocephalus* Willd. subsp. *macrocephalus* (NT), *Campanula baskilensis* Behçet sp. nov. (CR), *Centaurea urvillei* DC. Subsp. *nimrodus* (Boiss. & Hausskn.) Wagenitz (LC) *Hyacinthella siirtensis* B. Mathew (NT), *Linaria genistifolia* (L.) Miller var. *praealta* (Boiss.) Davis (NT), *Onosma isauricum* Boiss. & Heldr. (LC), *Phlomis armeniaca* Willd. (LC), *Scutellaria orientalis* L. subsp. *haussknechtii* (Boiss.) J. R. Edm. (LC), *Vebascum turcicum* B. Bani, Adigüzel & Karavel. sp. Nova (CR), *Astragalus brachystachys* DC. (EN), *Astragalus ruselli* Banks. & Sol. (VU), *Crucianella kurdistanica* Malin. (VU), *Ixiolirion tataricum* (Pall.) Herbert subsp. *tataricum* (VU), *Lathrus sylvestris* L. (VU), *Ranunculus dissectus* Bieb. subsp. *dissectus* (EN), *Sideritis libanotica* Labill. subsp. *microchlamys* (Hand.-Mazz.) Hub.-Mor. (VU).

Discussion and Conclusion: In the research area, 9 of the plants (3.1%) are endemic and 7 of them are in the non-endemic rare plant category. This ratio shows that the study area is quite poor in terms of endemism. The low rate of endemism can be explained by the simplicity of the topography of the research area.

Key Words: Atatürk Dam; Flora; Endemic; Şanlıurfa; Turkey

**COMPARATIVE ASSESSMENT OF ANTIOXIDANT ACTIVITY AND
PHYTOCHEMICAL PROFILES OF AQUEOUS EXTRACTS FROM *ATRIPLEX
HALIMUS* LEAVES AND STEMS**

Waffa ABDELMOUMENE

Laboratory of Natural Products, Department of Biology, Faculty of Natural and Life Sciences,
Earth and Universe Sciences, University Abou-Bekr Belkaïd, Tlemcen, 13000, Algeria

ORCID ID: <https://orcid.org/0009-0003-3010-5516>

Nabila BELYAGOUBI-BENHAMMOU

Laboratory of Natural Products, Department of Biology, Faculty of Natural and Life Sciences,
Earth and Universe Sciences, University Abou-Bekr Belkaïd, Tlemcen, 13000, Algeria

ORCID ID: <https://orcid.org/0000-0001-5624-4627>

Zineb KAID

Laboratory of Biomathematics, University Djillali Liabes, BP 89, 22000 Sidi Bel Abbes,
Algeria.

Faculty of Exact and Computer Sciences, University of Mostaganem, 27000, Algeria.

ORCID ID: . <https://orcid.org/0000-0003-2544-4049>

Asmaa DICH

Laboratory of Natural Products, Department of Biology, Faculty of Natural and Life Sciences,
Earth and Universe Sciences, University Abou-Bekr Belkaïd, Tlemcen, 13000, Algeria

ORCID ID: <https://orcid.org/0009-0004-1489-2211>

Larbi BELYAGOUBI

Laboratory of Natural Products, Department of Biology, Faculty of Natural and Life Sciences,
Earth and Universe Sciences, University Abou-Bekr Belkaïd, Tlemcen, 13000, Algeria

ORCID ID: <https://orcid.org/0009-0002-5047-7379>

ABSTRACT

Introduction and Purpose: *Atriplex halimus* L., an Algerian endemic halophytic shrub from the Chenopodiaceae family, has been traditionally used in medicine to address various ailments, including hyperglycemia, ovarian cysts, and hydatid cysts. Recent scientific investigations have highlighted its anti-inflammatory, antioxidant, and anticancer properties, emphasizing its therapeutic potential. This study aimed to evaluate the phytochemical composition and antioxidant activity of aqueous extracts of *A. halimus* leaves and stems. **Materials and**

Methods: One gram of dried plant material was ground into fine powder, macerated in 30 mL of distilled water, filtered, and evaporated to dryness. Phytochemical analyses were performed

to quantify total phenolics, flavonoids, flavonols, and tannins, while antioxidant activity was assessed through total antioxidant capacity (TAC), DPPH radical scavenging, and reducing power assays.

Results: Quantitative analyses showed higher total phenolics (18.464 ± 1.044 and 13.538 ± 0.815 mg GAE/g DW), flavonols (4.301 ± 0.519 and 1.917 ± 0.374 mg QE/g DW), and total antioxidant capacity (11.497 ± 0.532 and 1.426 ± 0.343 mg AAE/g DW) in leaves compared to stems, while stems exhibited higher flavonoids (2.511 ± 2.046 and 1.362 ± 0.267 mg QE/g DW). DPPH radical scavenging activity was comparable between leaves and stems (IC_{50} : 24.603 ± 2.635 and 24.218 ± 1.106 mg/mL), whereas reducing power was significantly higher in leaves (IC_{50} : 0.502 ± 0.002 mg/mL) compared to stems (IC_{50} : 15.994 ± 1.154 mg/mL).

Discussion and Conclusion: The findings demonstrate that the plant part used significantly affects bioactive compound levels and antioxidant activities. Leaves exhibited superior antioxidant potential, validating the traditional uses of *A. halimus* and supporting its potential as a natural source of bioactive compounds for therapeutic applications.

Key Words: *Atriplex halimus*, Aqueous extract, Phytochemical quantifications, Antioxidant activity.

**EFFECTS OF BATTERY TECHNOLOGIES ON AGRICULTURAL MACHINERY:
EFFICIENCY, SUSTAINABILITY AND FUTURE PERSPECTIVES****Dr. Öğr. Üyesi Orhan GÜNGÖR**

Burdur Mehmet Akif Ersoy Üniversitesi, Gölhisar Meslek Yüksekokulu

Orcid id: 0000-0002-5398-4571

ÖZET

Batarya teknolojilerindeki gelişmeler, tarım makinelerinde köklü değişimlere yol açmaktadır. Geleneksel içten yanmalı motorlar yerine bataryalı sistemlerin kullanılması, fosil yakıt bağımlılığını azaltarak karbon salınımını düşürmekte ve çiftçilere yakıt maliyetlerinde avantaj sağlamaktadır. Elektrikli tarım makineleri, içten yanmalı motorlara kıyasla daha verimli enerji kullanımı sunar ve özellikle ilaçlama ve gübreleme gibi işlemlerde eşit dağılım sağlar.

Bakım maliyetleri açısından bataryalı makineler öne çıkmaktadır. Daha az hareketli parça içermeleri nedeniyle arıza oranları düşük olup, daha sessiz ve titreşimsiz çalışmaları işçilerin konforunu artırmaktadır. Ancak, yüksek başlangıç maliyetleri ve enerji yoğun işlerde performans sınırlamaları önemli dezavantajlar arasında yer almaktadır. Hızlı şarj teknolojileri ve dayanıklı bataryalar bu sorunlara çözüm getirmektedir.

Batarya türleri arasında Li-Ion, Ni-Cd, Kurşun-Asit ve LiFePO₄ gibi çeşitler yer almaktadır. Li-Ion bataryalar yüksek enerji yoğunluğuna sahipken, Ni-Cd bataryalar dayanıklılık açısından avantajlıdır. Kurşun-Asit bataryalar düşük maliyetleriyle dikkat çekerken, LiFePO₄ bataryalar uzun ömürleri ve güvenlikleri nedeniyle elektrikli tarım makinelerinde tercih edilmektedir.

Bataryaların enerji depolama kapasitesi, yenilenebilir enerji kaynaklarıyla entegre edildiğinde tarımda daha sürdürülebilir bir enerji döngüsü sağlar. Gelecekte batarya teknolojilerinin ilerlemesiyle tarım makinelerinin tasarımlarında büyük değişimler beklenmektedir. Bu gelişmeler, tarımda daha verimli ve çevre dostu bir dönemin habercisi niteliğindedir.

Anahtar kelimeler: Elektrikli tarım makineleri, batarya teknolojisi, sürdürülebilirlik, hızlı şarj teknolojisi

ABSTRACT

Developments in battery technologies lead to radical changes in agricultural machinery. The use of battery systems instead of traditional internal combustion engines reduces carbon emissions by reducing fossil fuel dependency and provides farmers with an advantage in fuel costs. Electric agricultural machinery offers more efficient energy use compared to internal combustion engines and provides even distribution, especially in processes such as spraying and fertilisation.

Battery-powered machines stand out in terms of maintenance costs. They have lower failure rates due to fewer moving parts, and their quieter and vibration-free operation increases the comfort of workers. However, high initial costs and performance limitations in energy-intensive jobs are among the major disadvantages. Fast charging technologies and durable batteries provide solutions to these problems.

Battery types include Li-Ion, Ni-Cd, Lead-Acid and LiFePO₄. While Li-Ion batteries have high energy density, Ni-Cd batteries are advantageous in terms of durability. While Lead-Acid batteries attract attention with their low cost, LiFePO₄ batteries are preferred in electric agricultural machines due to their long life and safety.

The energy storage capacity of batteries provides a more sustainable energy cycle in agriculture when integrated with renewable energy sources. In the future, major changes are expected in the design of agricultural machinery with the advancement of battery technologies. These developments herald a more efficient and environmentally friendly period in agriculture.

Keywords: Electric agricultural machines, battery technology, sustainability, fast charging technology

GİRİŞ

Batarya teknolojilerinin gelişmesi sadece traktörün içten yanmalı motorunun hareket mekanizması açısından değil, diğer tarım aletlerinin çalışma şeklinin değişiminde de büyük etkiye sahip olacaktır. Günümüzde traktör arkası yada önü çalışan tarım aletlerinin çoğuna PTO (Power Take-Off) bağlantıları ile güç aktarımı sağlanır. Bu sistem, tarımsal ekipmanların etkin bir şekilde çalışmasına olanak tanırken, bağlantı sistemlerinde meydana gelen kayıplar sistemin genel verimliliğini olumsuz etkileyebilir.

Traktör kuyruk mili, motorun ürettiği gücü şanzıman veya direkt olarak motorun krank milinden alır. Standart kuyruk mili hızları genellikle 540 rpm ve 1000 rpm olarak belirlenmiştir. Motorun ürettiği güçten kuyruk miline aktarılırken bir miktar kayıp yaşanır. Bu kayıplar genellikle şanzıman, dişliler, miller ve diğer mekanik sürtünmelerden kaynaklanır. Standart bir traktörde kuyruk miline aktarılan güç, motor gücünün yaklaşık %80-90'ı kadardır. Yani, motor 100 beygir güç üretiyorsa, kuyruk miline yaklaşık 80-90 beygir ulaşır. Kuyruk milinde en çok karşılaşılan sorunlar şunlardır:

- Dişli ve şaft aşınmaları
- Kavrama mekanizması sorunları
- Mil dönüş hızında sapmalar
- Şaft kapağı ve koruma sorunları
- Rulman arızaları
- Kuyruk mili fren sorunları

Bu sorunlarla karşılaşmamak için kullanıcıların bilinçlendirilmesi gerekmektedir. Zamanında yağlama ve uygun ekipman kullanımı bu sorunları tamamen ortadan kaldırmaya da azaltabilir.

Batarya teknolojilerinin gelişmesiyle birlikte bakım azalacağından ve yanlış ekipman seçimlerinin önüne geçileceğinden bu sorunlar azalma gösterecektir. Traktör arızalarının %70'i mekanik arızalardır. Batarya teknolojilerinin gelişmesiyle birlikte bu arızalarda da azalma meydana gelecektir.

Pil teknolojileri, özellikle çiftçilik uygulamalarının verimliliğini ve sürdürülebilirliğini artırmada modern tarımda önemli bir rol oynar. Güneş ve rüzgar gibi yenilenebilir enerji kaynaklarının tarımsal faaliyetlere entegrasyonu giderek daha etkili enerji depolama çözümlerine dayanmaktadır. Bunun başlıca nedeni, tarımsal faaliyetler için tutarlı bir enerji tedariki sağlamak için güvenilir depolama sistemleri gerektiren yenilenebilir enerji üretiminin aralıklı doğasıdır.

Pil teknolojilerinin tarımdaki önemli avantajlarından biri, yenilenebilir kaynaklardan üretilen fazla enerjiyi depolayabilmeleridir. Bu depolanan enerji, düşük üretim dönemlerinde kullanılabilir ve böylece sulama, ısıtma ve soğutma sistemleri de dahil olmak üzere çeşitli

tarımsal süreçler için istikrarlı bir enerji tedariki sağlar. Çalışmalar, enerji depolama sistemlerinin yenilenebilir elektrik kaynaklarının değişkenliğini azaltabileceğini ve böylece tarımsal ortamlarda enerji tedarikinin güvenilirliğini artırabileceğini göstermiştir (Beaudin ve diğerleri, 2010). Ayrıca, pil depolamanın yenilenebilir enerji sistemleriyle birleştirilmesi, tarımsal üretkenliği optimize etmek için gerekli olan enerji yüklerinin daha iyi yönetilmesini sağlar (Zhong ve diğerleri, 2011).

Batarya teknolojilerinin tarımdaki ekonomik etkileri dikkate değerdir. Enerji depolama sistemlerinin benimsenmesi, şebeke elektriğine olan bağımlılığı azaltarak ve yoğun talep dönemlerinde enerji maliyetlerini en aza indirerek önemli maliyet tasarruflarına yol açabilir. Araştırmalar, pil enerji depolama sistemlerinin (BESS) yenilenebilir enerji projeleriyle entegre edilmesinin çiftçiler için birden fazla gelir akışı yaratabileceğini ve böylece ekonomik dayanıklılıklarını artırabileceğini göstermektedir (Fan vd., 2021). Ek olarak, pil sistemlerinin stratejik boyutlandırılması ve koordinasyonu, çeşitli vaka çalışmalarında gösterildiği gibi, tarımsal faaliyetlerde enerji kullanımının genel verimliliğini artırabilir (Hayajneh vd., 2021; Canbulat vd., 2021).

Tarımda pil teknolojilerinin kullanılmasının çevresel faydaları göz ardı edilemez. Bu teknolojiler, yenilenebilir enerjinin kullanımını kolaylaştırarak, geleneksel fosil yakıt bazlı enerji kaynaklarıyla ilişkili sera gazı emisyonlarının azaltılmasına katkıda bulunur. Pil depolama ile desteklenen yenilenebilir enerji sistemlerine geçiş, iklim değişikliğiyle mücadele ve sürdürülebilir tarım uygulamalarını teşvik etme yönündeki küresel çabalarla uyumludur (Cosentino vd., 2012; Morris & Bowen, 2020). Ayrıca, sodyum iyon piller gibi gelişmiş pil teknolojilerinin geliştirilmesi, çevresel etkiyi en aza indirirken enerji depolama kapasitesini artırmak için umut verici bir yol sunmaktadır (Tan, 2023).

MATERYAL

Batarya teknolojilerinin gelişmesiyle birlikte tarım makinaları tasarımlarında da büyük değişimler olacaktır. Bunların başında gelen ise traktör tasarımları değişebilecektir. Şekil 1 ve şekil 2 de yapay zeka ile yapılan bataryalı traktör tasarımları görülmektedir.



Şekil 1. Güneş panelleri ile donatılmış elektrikli traktör tasarımı

Şekil 1'deki traktör günümüzdeki traktörlere benzemekle beraber üzerine güneş paneli yerleştirilen bir tasarım görülmektedir. Benzer şekilde tarım aletinin üzerinde de güneş paneli yer almaktadır. Böyle bir tasarım ancak güneş panellerinden çok fazla verim olması ile mantıklı olacaktır.



Şekil 2. Bataryalı yönlendirici traktör

Şekil 2'deki tasarım günümüz traktör tasarımlarından farklı olarak, batarya deposu olarak yönlendirici görevi görebilecek bir tasarımdır. Bu şekilde bir tasarım daha ileri düzendeki tarım aletleri için uygun olacaktır. Çünkü bu tasarımda tarım aletlerinin çalışabilmesi için kendi üzerinde de mekanizmayı harekete geçirecek elektrik motorlarının bulunması gerekmektedir.


Her iki tasarımda da batarya şarj sürelerinin kısalması büyük önem taşımaktadır.

BULGULAR ve TARTIŞMA

Şekil 3'te içerdikleri kimyasal yapıya göre şarj süreleri gösterilmiştir. Bu grafikler, farklı pil türlerinin (Li-Ion, Li-Po, Ni-Cd, Ni-MH, Kurşun-Asit, LiFePO4) şarj eğrilerini gösteriyor. Her bir grafik, pilin şarj olurken gerilim (voltaj) ve akım değişimlerini zamana bağlı olarak sunar. Grafikteki eksenler şu şekilde:

X Eksenleri (0.0 - 1.0 arası): Şarjın tamamlanma oranını temsil eder (%0 - %100). Bu, zamanla doğru orantılıdır ancak kesin bir zaman dilimi verilmemiştir.

Y Eksenleri: Voltaj (V) ve Akım (A) değerlerini gösterir.



Şekil 3. Yapılarına göre batarya voltaj-akım eğrileri

Şekil 3'teki batarya grafikleri aşağıdaki gibi yorumlanabilir.

CC (Constant Current - Sabit Akım) Aşaması: Şarjın başlangıcında pil sabit bir akımla şarj edilir, bu sırada voltaj artar.

CV (Constant Voltage - Sabit Voltaj) Aşaması: Belirli bir voltaj seviyesine ulaşıldığında bu voltaj korunur, ancak akım düşer.

1. Li-Ion ve Li-Po Şarj Eğrisi:

Voltaj (Sarı çizgi): 3.2V'den başlayarak yaklaşık 4.2V'ye kadar yükselir. Bu, sabit akım (CC) aşamasında gerçekleşir. 4.2V'ye ulaştığında sabitlenir, bu da sabit voltaj (CV) aşamasına geçildiği anlamına gelir.

Akım (Kırık çizgi): Başta sabit kalır (yaklaşık 1A), ardından voltaj sabitlenince düşmeye başlar ve şarj tamamlandığında sıfırlanır.

2. Ni-Cd ve Ni-MH Şarj Eğrisi:

Voltaj: 1.2V'den başlayarak 1.8-2.0V civarına kadar artar.

Akım: Sabit bir şekilde kalır (~0.8A), bu da sabit akımla şarj yöntemine işaret eder.

3. Kurşun-Asit Şarj Eğrisi:

Voltaj: 12V'den başlayarak 14V'ye kadar yükselir ve sabitlenir.

Akım: İlk başta sabit (1A civarı), sonra yavaşça düşer.

4. LiFePO4 Şarj Eğrisi:

Voltaj: 3.0V'den başlayarak 3.6V'ye kadar çıkar ve sabitlenir.

Akım: Başlangıçta sabit kalır (~1.5A), ardından voltaj sabitlenince düşer.

Tam Şarj: Akım sıfıra yaklaştığında veya düşüş belirli bir seviyeye geldiğinde pil tam şarj olmuş sayılır.

Sayısal değerler (voltaj ve akım) her pil türünün kimyasal yapısına ve güvenli çalışma aralığına göre belirlenir. Bu eğriler, pilin ömrünü uzatmak ve güvenliğini sağlamak için önemlidir.

Tablo 1’de yapılarına göre batarya özellikleri görülmektedir. Bu tabloda batarya tercih edilirken dikkate alınacak özellikler vardır. Hangi tür makineye nasıl batarya seçiminin daha uygun olacağı görülmektedir. Günümüzdeki elektrikli araçların LiFePO4 (LFP) olarak adlandırılan bataryayı kullanmayı tercih ettiği bilinmektedir. Bu batarya yüksek güvenliği, %100 şarj edilebilirliği ve yüksek şarj döngünü ile öne çıkmaktadır.

Tablo 1. Yapılarına göre bataryaların özellikleri

Kriter	Li-Ion / Li-Po	Ni-Cd / Ni-MH	Kurşun-Asit	LiFePO4
Enerji Yoğunluğu	Yüksek	Orta	Düşük	Orta
Şarj Döngüsü Ömrü	300-800	500-1000	300-500	2000-5000
Güvenlik	Orta (ısı riski)	Yüksek (dayanıklı)	Orta	Çok yüksek
Maliyet	Orta	Düşük	En ucuz	Yüksek
Kullanım Alanı	Mobil cihazlar, İnsansız hava araçları	Eski aletler, tıbbi cihazlar	Araç aküleri, UPS sistemleri	Elektrikli araçlar, enerji depolama sistemleri

SONUÇ ve ÖNERİLER

Sonuç olarak, Türkiye tarım aleti üretiminde 2022 yılında ihracatı 1,32 milyar dolar seviyesindedir. Türkiye'nin tarım alet ve makineleri ihracatında ilk üç ülke sırasıyla ABD (%19,2), İtalya (%10,4) ve Rusya Federasyonu (%6)'dır. Batarya teknolojilerinin gelişmesiyle birlikte tarım aletlerinde de büyük tasarım değişimleri meydana gelecektir. Bu açıdan ülkemizin tarım alet ve makineleri üretimini arttırarak devam ettirebilmesi için bu alandaki teknolojik gelişmeleri yakından takip etmesi önemlidir. Ayrıca pil teknolojileri sürdürülebilir tarımın ilerlemesi için olmazsa olmazdır. Yenilenebilir enerji kaynaklarının etkili bir şekilde entegre edilmesini sağlar, ekonomik uygulanabilirliği artırır ve çevresel sürdürülebilirliğe katkıda bulunur. Tarım sektörü gelişmeye devam ettikçe, bu geçişi kolaylaştırmada pil depolamanın rolü giderek daha da önemli hale gelecektir.

REFERENCES

Beaudin, M., Zareipour, H., Schellenberglabe, A., Rosehart, W. (2010). Energy storage for mitigating the variability of renewable electricity sources: an updated review. *Energy for Sustainable Development*, 14(4), 302-314. <https://doi.org/10.1016/j.esd.2010.09.007>

Canbulat, S., Balci, K., Canbulat, O., Bayram, İ. (2021). Techno-economic analysis of on-site energy storage units to mitigate wind energy curtailment: a case study in scotland. *Energies*, 14(6), 1691. <https://doi.org/10.3390/en14061691>

Cosentino, S., Testa, G., Scordia, D., Alexopoulou, E. (2012). Future yields assessment of bioenergy crops in relation to climate change and technological development in europe. Italian Journal of Agronomy, 7(2), e22. <https://doi.org/10.4081/ija.2012.e22>

Fan, F., Zorzi, G., Campos-Gaona, D., Burt, G., Anaya-Lara, O., Nwobu, C., Madariaga, A. (2021). Sizing and coordination strategies of battery energy storage system co-located with wind farm: the uk perspective. Energies, 14(5), 1439. <https://doi.org/10.3390/en14051439>

Hayajneh, H., Herrera, M., Zhang, X. (2021). Design of combined stationary and mobile battery energy storage systems. Plos One, 16(12), e0260547. <https://doi.org/10.1371/journal.pone.0260547>

Morris, W. and Bowen, R. (2020). Renewable energy diversification: considerations for farm business resilience. Journal of Rural Studies, 80, 380-390. <https://doi.org/10.1016/j.jrurstud.2020.10.014>

Tan, D. (2023). Sodium energy storage-key clean energy for the future world.. <https://doi.org/10.26434/chemrxiv-2023-j3zfh-v3>

Zhong, C., Ding, M., Su, J. (2011). Modeling and control for large capacity battery energy storage system., 1429-1436. <https://doi.org/10.1109/drpt.2011.5994120>

REASONS FOR WHITE SPIKE FORMATION IN WHEAT AND DIAGNOSIS**BUĞDAYDA BEYAZ BAŞAK OLUŞUM NEDENLERİ VE TEŞHİS****Prof. Dr. Aydın AKKAYA**Muş Alparslan Üniversitesi Uygulamalı Bilimler Fakültesi Bitkisel Üretim ve Teknolojileri
Bölümü

ORCID; 0000-0001-9560-1922

ÖZET

Giderek artan dünya nüfusunun beslenmesi ve gıda güvenliği yönünden buğday stratejik öneme sahip bir kültür bitkisidir. Geniş adaptasyon ve telafi yeteneği sayesinde, dünya genelinde çok farklı iklim ve toprak koşullarında tarımı yapılabilmektedir. Ancak geniş adaptasyon ve telafi yeteneğine sahip olmasına rağmen iklim, toprak ve yetiştirme tekniğindeki olumsuz faktörler buğday verim ve kalitesinde önemli kayıplara yol açabilmektedir. Vejetasyon süresince farklı dönemlerde ortaya çıkabilen biyotik ve abiyotik stres faktörlerinin şiddet ve süresine bağlı olarak buğday büyüme ve gelişmesi, dolayısıyla performansı olumsuz yönde etkilenebilmektedir. Buğday tarımında üreticilerin sıkça karşılaştığı üretim sorunlarından birisi sağlıklı yeşil başaklar yerine, beyaz renkli başakların oluşmasıdır. Beyaz başak sorunu düşük yoğunlukta olduğu zaman tane verimi üzerindeki zararlı etkisi ihmal edilebilecek düzeyde kalabilir. Ancak aşırı yoğunlukta olması durumunda tane veriminde önemli kayıplara yol açabilmektedir. Buğdayda beyaz başak oluşumuna yol açan çeşitli faktörler söz konusudur. Bu faktörler kuraklık ve don zararı gibi iklimsel olabildiği gibi kök çürüklüğü, başak yanıklığı, göçerten, buğday sap kurudu, süne gibi hastalık ve zararlılarla ilgili faktörler de olabilir. Üreticilerin beyaz başak oluşumuna yol açan faktörler ve doğru teşhis konusunda bilgi sahibi olması yetiştiricilik açısından kuşkusuz büyük önem taşımaktadır. Bu sunulu tebliğde, beyaz başak oluşumuna neden olan faktörler ele alınarak, doğru teşhise yardımcı olacağı düşünülen bilgiler adım adım açıklanmaya çalışılmıştır.

Anahtar Kelimeler: Buğday, beyaz başak, etmenler, teşhis**ABSTRACT**

Wheat is a cultivated crop having strategic importance in terms of nutrition and food security of the ever-increasing world population. It can be grown in very different climate and soil conditions around the world thanks to its wide adaptation and compensatory abilities. However, although it has wide adaptation and compensatory capabilities, negative factors in climate, soil and cultivation techniques can lead to significant losses in wheat yield and quality. Depending on the severity and duration of biotic and abiotic stress factors occurring at different periods throughout its vegetation period, wheat growth and development, thus its performance, could be negatively affected. The formation of white colored spike instead of healthy green spikes is one of the production problems that farmers frequently encounter. When the white spike problem is at low intensity, its harmful effect on the grain yield could be at ignored level. However, when it is excessively dense, it can cause significant losses in grain yield. There are several factors causing the white spike formation in wheat. These factors include climatic

factors such as drought and freezing damage as well as factors related to diseases and pest such as root rot, head blight, take-all, wheat stem maggot and sunn pest. The sufficient knowledge of producers about the factors leading to white spike formation and correct diagnosis are undoubtedly of great importance for production. In this presentation paper, the factors causing the white spike formation are addressed and tried to explain information step by step thought being useful for correct diagnosis.

Keywords: Wheat, white spike, reasons, diagnosis

GİRİŞ

Buğday, dünyada yaklaşık 780 milyon ton yıllık üretimle (Anonymous, 2023), gıda güvenliği yönünden stratejik öneme sahip bir kültür bitkisidir. Ülkemizde 2022 yılı rakamlarına göre yıllık 37 milyon tonluk tahıl üretimi içerisinde, 20 milyon tonluk üretimle buğday ilk sırada yer almaktadır (Anonim, 2023). Buğday tarımı ülkemizde, daha çok yağışa dayalı koşullarda kışlık olarak yapılmakta, zaman zaman marjinal alanlara bile buğday ekilmektedir.

Küresel ısınmaya bağlı iklim değişikliği; kuraklık, aşırı yağmur, su baskını, soğuk zararı, don etkisi gibi ekstrem olaylara yol açmakta, özellikle yağışa bağlı koşullarda yapılan buğday tarımı üzerinde baskılar oluşturmaktadır. Dünyanın her tarafında çok farklı iklim koşullarında tarımı yapılan buğday, çok yüksek adaptasyon ve telafi yeteneğine sahiptir. İklim ve toprak koşullarındaki olumsuzlukları, yetiştirici hatalarını belli bir düzeye kadar telafi edebilmektedir. Ancak bu telafi, verim ve kalitede bir takım kayıplar pahasına gerçekleşmektedir. Günümüzde yaygın bir şekilde tarımı yapılan ticari buğday çeşitleri, yüksek verimli koşullar için geliştirilmiş olup, stres koşullarından çok daha fazla zarar görme riski taşımaktadır.

Buğday tarımında üreticilerin sıkça karşılaştığı üretim sorunlarından birisi sağlıklı yeşil başaklar yerine, beyaz renkli başakların oluşmasıdır. Beyaz başak sorunu düşük yoğunlukta olduğu zaman tane verimi üzerindeki etkisi ihmal edilebilecek düzeyde kalabilir. Ancak aşırı yoğunlukta olması durumunda tane veriminde önemli kayıplara yol açabilmektedir. Buğdayda beyaz başak oluşumuna yol açan çeşitli faktörler söz konusudur. Bu faktörler kuraklık ve don zararı gibi iklimsel olabildiği gibi kök çürüklüğü, başak yanıklığı, göçerten, buğday sap kurudu, süne gibi hastalık ve zararlılarla ilgili de olabilir. Üreticilerin beyaz başak oluşumuna yol açan faktörler ve doğru teşhis konusunda bilgi sahibi olması yetiştiricilik açısından kuşkusuz büyük önem taşımaktadır. Bu sunulu tebliğde, buğdayda beyaz başak oluşumuna neden olan faktörler ele alınarak, doğru teşhis açısından üreticilere yardımcı olacağı düşünülen bilgiler açıklanmaya çalışılmıştır.

İKLİM FAKTÖRLERİ

Sapı etkileyen don zararı

Ani dondurucu düşük sıcaklıklar buğday sapının iletim sistemine zarar verebilir. Sap dokusunda zarar gören kısımda buz halkaları veya beyaz halkalar oluşur. Zarar şiddetli olur ise sap dip kısımlarındaki boğumlardan eğilir ve yatma riski artar. Düşük şiddetli zarar sapta gözle görünür düzeyde bir hasar oluşturmasa da iletim sisteminin zarar görmesine yol açabilir. Oluşan buz kristalleri yayılarak ksileme zarar verir. Zarar sapa kalkmadan sonra herhangi bir dönemde meydana geldiğinde beyaz başak oluşumuna yol açar. Başağa yeteri kadar su ve besin elementi taşınmadığı için beyaz başak oluşumu söz konusu olur (Bean ve ark. 2012).

Başığı etkileyen don zararı

Sapa kalkmadan sonraki dönemlerde veya başaklanma döneminde etkili olan don zararı, kılçıkların beyazlaşmasına, başağın kısmen veya tamamen sarımsı beyaz renk kazanmasına ve

boş kalmasına neden olur. Çiçeklenme öncesi ve sırasında etkili olan düşük sıcaklığın şiddeti ve süresi, başakta kısmi veya tam bir kısırlığa yol açabilir. Çiçeklenmeden hemen önce veya çiçeklenme esnasında kısa süreli düşük sıcaklık, başağın üstü, alt veya ortasında kısmi beyazlığa yol açar. Ancak bu dönemde uzun süreli ve çok düşük derecede sıcaklık etkili olduğunda, birkaç gün sonra başak bir bütün olarak beyaz veya saman renginde bir görüntüye sahip olur. Don zararı tarlanın alçak kısımlarında daha şiddetli olur. Anterler daha hassas olup zarar gördüklerinde buruşuk ve beyazımsı kahve renkli olurlar. Tüylü beyazımsı yapıda olması gereken stigma, beyazımsı-kahve renkli ve jelimsi bir özellik kazanır ve açılmaz. Yapraklar kavrulmuş gibi görünür ve sap içerisinde buzlanmış kısımlarda, kahve renkli buruşuk bir görüntü ortaya çıkar (Akkaya, 1994; Shroyer ve ark. 1998; Raper, 2018; DeWolf ve ark. 2023).

Dolu zararı

Dolu, buğdayda yaprak zararından ürünün tamamen kaybına kadar çok değişik düzeylerde etkili olabilir. Başak henüz bayrak yaprak kını içerisinde iken etkili olan dolu, sağlıklı başak çıkışına ve başak oluşumuna engel olur. Dolu zararı şiddetli olduğunda sap ve başakta kırılmalara yol açtığından yetiştiriciler tarafından kolay bir şekilde ayırt edilebilir. Fakat bazen dolu yağışı, sap veya başakta mutlak bir kırılmaya yol açmayacak düzeyde olsa bile sap veya başakta renk kaybına yol açabilir. Başakta kısmi bozulmalara veya başağın bir bütün olarak bozulup beyaz renk kazanmasına neden olabilir. Beyaz renkli başaklara ait sapların üst kısımlarında morlukların bulunması dolu zararı olduğunu gösterir (Bean ve ark. 2012; DeWolf ve ark. 2023).

Su baskını ve boğulma

Buğday bitkisi uzun süre su altında kalmaya dayanıklı değildir. Kökler herhangi bir su baskını veya aşırı yağış nedeniyle 3-4 gün süreyle su altında kaldığında, köklere yeteri kadar oksijen gidemediği için bitkiler boğularak ölür ve başaklar beyaz olur. Tarlada özellikle su birikmesine uygun olan çukur yerlerde boğulma ve buna bağlı beyaz başak zararı daha fazla görülür. Köklerde boğulmaya neden olacak düzeyde nem birikimi olmasa bile arazinin çukur kısımlarındaki aşırı nem birikimi ve ağır toprak yapısı bitkiyi erken olgunlaşmaya zorlayabilir. Bu durumda başaklarıyla birlikte bitkilerde bir bütün olarak erkenden renk kaybı söz konusu olur ve beyazlaşırlar (Shroyer ve ark. 1998; DeWolf ve ark. 2023).

Kuraklık

Kuraklık, yağışa dayalı koşullarda yapılan buğday tarımının en önemli sorunudur. Ekimden hasada kadar, etkili olduğu her dönemde, kuraklık şiddeti ve süresine bağlı olarak çok farklı düzeylerde zararlar ortaya çıkar. Sapa kalkma dönemindeki kuraklık generatif gelişmeye zarar vererek, başak ve başakcık sayısının azalmasına, başakların küçülmesine yol açar. Yüksek sıcaklık döllenmeyi olumsuz etkiler ve başakların uç kısmının kavrulmasına neden olur. Kuraklık yüksek sıcaklık ile birlikte etkili olduğu zaman bitkiler ölüme gidebilir ve başaklar beyaz olur. Kuraklık zararından bitki, tek bir sap yerine kardeşler de dahil olmak üzere bir bütün olarak etkilenir. Tarlanın hafif bünyeli ve yüzeysel toprak yapısına sahip kesimlerine denk gelen bitkilerde, gelişme geriliği ve başaklarda kirli beyaz bir görünüm söz konusu olur (Akkaya, 1994; Lollato, 2020; DeWolf ve ark. 2023).

ZARARLILAR

Süne (*Eurygaster integriceps*, sunn pest, senn pest, cereal bug, wheat shield bug)

Süne, buğdayın en önemli zararlılarından birisi olup, bitkinin farklı kısımlarına zarar verir. Nimfler genç yapraklar üzerinde, erginler ise başak ve tane üzerinde beslenmeyi tercih ederler. Kışı geçiren erginler, kardeşlenme döneminde buğday sapını kök boğazına yakın yerden sokup, bitki öz suyunu emerler. Emilen saplar, besin ve su iletimi engellendiği için zamanla sararır, kurur ve başak bağlayamazlar. Bu zarar şekline kurtboğazı adı verilir. Bitkiler geliştikçe üst

kısımlarda beslenmeye başlayan erginler, başak henüz bayrak yaprak kını içerisinde iken veya çiçeklenme ve tane bağlama dönemlerinde, başağın altından sapı delip özsuğu emerler. Zarar gören sapların başakları beyaz ve boş olur. Bu zarar şekline akbaşak adı verilir. Normal yeşil başakların arasında, zarar görmüş beyaz başaklar açık bir şekilde belli olur (Mackesy ve Moylett, 2018; Babaroğlu ve ark. 2020; İslamoğlu, 2021; Akkaya, 2024).

Kıml (Aelia rostrata, wheat stink bug)

Ülkemizde özellikle Orta Anadolu Bölgesi'nde zarar yapar. Kışlaktan inen erginler, süneye benzer şekilde buğday saplarını kök boğazı üzerinden emerek zarar verirler. Emgi zararı gören saplar başak bağlayamaz. Bu zarar şekline kurtboğazı (göbek kurusu) adı verilir. Başaklanma döneminde başağın bağlandığı sap kısmından bitki özsuğunu emerek, akbaşak adı verilen beyaz ve boş başak oluşumuna neden olurlar (Anonim, 2015; Anonim, 2016; Tarla, 2017; Akkaya, 2024).

Ekin sap arısı (wheat stem sawfly)

Ekin sap arıları [*Cephus pygmaeus*, *Cephus cinthus*, *Trachelus tabidus*] dişileri yumurtalarını buğday sapı içerisine bırakırlar. Yumurtadan çıkan larvalar sap içerisinde beslenirler. Hasat öncesi sapın dip kısmına doğru inen larva sapın içini çiğneyip kemirir. Kemirilen kısımda artıklar oluşur ve sap kırılır. Zarar gören bitkilerde zayıflama, yatma, beyaz başak oluşumu söz konusu olur (Özberk ve ark. 2005; Anonim, 2008; Konodel ve ark. 2010; Kılıç ve ark. 2017).

Buğday sap kurdu (wheat stem maggot, Meromyza americana),

Buğday sap kurdunun zarar verdiği tarlada beyaz başak oluşumu öbekler halinde değil, tek tek başaklar düzeyinde ve dağınık halde ortaya çıkar. Bitki bütün kardeşleriyle beraber etkilenmez. Bitki yeşil olmasına rağmen başak beyaz renkli ve boştur. Beyaz başaklar yukarı doğru çekildiği zaman üst boğum arasıyla birlikte yaprak kınından kolayca çıkar. Sapın en üst boğumunun hemen üzerinde kalan üst boğum arasının dip kısmı dikkatli bir şekilde incelendiği zaman, böceğin sapı çiğnemek suretiyle yol açtığı dağınık ve bozuk yapı kolaylıkla görülebilir (Shroyer ve ark. 1998; Bean ve ark. 2012; Michaud, 2013).

Buğday kesiksineği (Mayetiola destructor, Hessian fly)

Erginleri zarar vermeyip, asıl zararı larvalar sonbahar ve ilkbaharda yaparlar. İlkbaharda larvalar toprak seviyesine yakın sapın dip kısmında, yaprak kını ile sap arasına girer ve bitki özsuğunu emerek zarar verir. Larvaların zarar verdiği yerlerde sap zayıflar ve ileri dönemlerde sap kırılmaları ve dağınık yönde yatmalar meydana gelir. Bu sap kırılmaları birinci veya ikinci boğumun hemen üzerinden olur. Kırılan kısımlarda boğum arasında bir çökme olabilir ve sapta zarar gören noktalar civarında pupalar bulunabilir. Erken ekimlerde zarar riski daha yüksek olabilir (Shroyer ve ark. 1998; Anonim, 2008; Ransom, 2016; DeWolf ve ark. 2023; Whitworth ve ark. 2023; Anonymous, 2024).

HASTALIKLAR

Bipolaris kök çürüklüğü (common root rot)

Toprakta ve bitki artıklarında yaşayan *Bipolaris sorokiniana* (*Cochliobolus sativus*) mantarının neden olduğu kök çürüklüğü hastalığıdır. Başlangıç enfeksiyonu erken dönemde olmasına rağmen su ihtiyacının artacağı döneme kadar bitki normal gelişme ve büyümesine devam eder. Ancak bitkinin su ihtiyacının arttığı dönemde iletim sistemi bitkinin ihtiyacını karşılayacak kadar sağlıklı olmadığından bitki erkenden sağlığını kaybeder, zamansız olgunlaşma ve beyaz başak oluşumu söz konusu olur. Olgunlaşma öncesi tarlada dağınık halde bazı yerlerde beyaz başaklı adacıklar oluşturur. Enfekteli bitkilerde kök gelişmesi zayıf, dip kısım koyu renklidir. Tohum kalıntısı ile bitkinin dip kısmı arasındaki ince sap bölümü (subcrown internode)

üzerindeki koyu kahverengi lezyonlar, hastalığın teşhisi yönünden en ayırt edici özelliktir. Sağlıklı bitkilerde bu kısım krem rengindedir. Kurak koşullar hastalığı teşvik eder (Anonim, 2008; Huberli, 2015; DeWolf ve Shroyer, 2017; Byamukama ve ark. 2018).

Fusarium kök ve kök boğazı çürüklüğü

(fusarium crown and foot rots, dryland root rot)

Fusarium etmenine bağlı mantari bir hastalık olup, enfekte olan bitkinin bütün kardeşlerinde başaklar prematüre olarak beyaz renkli olur. Bitkiler kısa boylu kalır, beyaz başaklı bitkiler tarlada öbekler halinde bulunur. Bu hastalık kurak yıllarda daha fazla görülür. Beyaz başaklı bitkilerin kökleri zarar gördüğünden çekildiği zaman kolay bir şekilde çıkar. Kök tacından toprak seviyesinin hemen üzerindeki kısma kadar olan sapın dip kısmı kahve veya pembe renkli olur. Bitki sökülüp birkaç saat bekletildiğinde pembe renk kaybolur. Yaprak kınları soyulup sapın dip kısmına bakıldığı zaman bu renk bozulmaları görülebilir. Sapın iç kısmı pamuksu pembe veya pembemsi beyaz mantar miselleri ile dolar. Hastalık semptomu, kuraklık stresi veya göçerten hastalığı (take-all) ile karıştırılabilir. Her yıl üst üste buğday ekilen yerlerde giderek daha yaygın bir şekilde görülür. (Shroyer ve ark. 1998; Bean ve ark. 2012; Byamukama ve ark. 2018).

Fusarium başak yanıklığı (fusarium head blight, scab)

Bu hastalık, çiçeklenme döneminde yağışlı havalarda etkili olması durumunda, tarlada dağınık bir şekilde beyaz başak oluşumuna neden olur. Sulu-sıfır sürüm koşullarında, mısır anızına buğday ekildiğinde daha yaygın görülür. Genellikle başağın üst kısmı etkilenmekle beraber başağın tamamının etkilenmesi de mümkündür. Başak eksenini ve başakcıklar yakından dikkatli bir şekilde incelendiğinde, pembemsi portakal renginde mantar sporlarını görmek mümkündür. Hastalık, tanelerin beyazımsı veya pembe renkli olmasına ve hektolitreye ağırlığının düşmesine neden olur. Ayrıca hastalıklı taneler bazı zararlı mikotoksinler içerebilir (Shroyer ve ark. 1998; Bean ve ark. 2012; DeWolf ve ark. 2023).

Göçerten (karabacak, take-all)

Toprak kökenli bir fungusun (*Gaeumannomyces graminis*) neden olduğu bu hastalık, başakların prematüre olarak ölümüne ve beyaz renkli olmalarına yol açar. Tarlada öbekler halinde etkili olur. Bitkide renk kaybı ve bodurlaşma olsa da hastalığın erken dönemde fark edilmesi zordur. Başağın bir bütün olarak beyaz renkli olmasına neden olur. Başaktan tutup yukarı doğru çekildiğinde, başak kolaylıkla kopmaz. Toprak seviyesine yakın bir yerden bitki yukarı doğru çekildiğinde kolaylıkla kopar ve köklerin çoğu toprak içerisinde kalır. Sapın dip kısmında yaprak kını soyulup incelendiğinde, görülen parlak siyah veya koyu kahve renk, hastalık teşhisi açısından önemli bir göstergedir. Kök kısmında da renk bozulması ve kararma olur. Kök üzerindeki lezyonlar küçük ve siyah renkli olup, zamanla yayılarak birleşik bir hal alırlar. Sıfır sürüm uygulanarak her yıl buğday ekilmesi, fazla bitki artığı, yüksek nem hastalığı teşvik eden faktörlerdir (Shroyer ve ark. 1998; Bockus ve Tisserat, 2005; Bean ve ark. 2012; DeWolf ve ark. 2023).

Cephalosporium çizgi hastalığı

Toprak kaynaklı, vasküler bir solgunluk hastalığı olup, etmeni *Cephalosporium gramineum* fungusudur. Yağışın bol olduğu yörelerde daha yaygın olarak görülür. Yaprak boyunca 1-2 adet, geniş sarı renkli çizgi oluşur. Çizgiler yaprak kınına kadar uzayabilir. Sarı renkli çizgilerin merkezinde kahverengi yaprak damarları bulunur. Hastalık kardeşlerde farklı derecede bodurluğa ve tarlada öbekler halinde beyaz başak oluşumuna yol açar (Shroyer ve ark. 1998; DeWolf ve Shroyer, 2017; Hagerty ve Smiley, 2017; Gül ve Karakaya, 2019).

Sarı leke (tan spot)

Etmeni *Pyrenophora tritici-repentis* olan bu hastalık yaprakları, başak ve taneleri etkileyebilir. Mantar, anız artıklarında yaşar ve anıza ekimde hastalık daha yaygın olabilir. En tipik özelliği yaprakta kahve renkli lekelerin oluşması ve bunların etrafının sarı renkli halka ile çevrilmesidir. Olgun lezyonların orta kısmında koyu renkli bir kısım vardır. Lezyonlar giderek büyüyüp birleşir ve yaprakta daha geniş bir alanın zarar görmesine yol açar. Sapa kalkma döneminde alt yapraklarda başlar, tane gelişme döneminde üst yapraklara sıçrar. Başakta kısmi beyazlaşmaya ve tanenin koyu kırmızı bir renk almasına neden olabilir (Wegulo, 2011; DeWolf ve Shroyer, 2017; Byamukama ve ark. 2018; Yüceler ve ark. 2022).

Eyespot (strawbreaker foot rot)

Eyespot mantarı bir hastalık olup, hastalık etmeni çok yakın iki grup mantardır (*Oculimacula yallundae* ve *Oculimacula acufomis*, eski ismi *Pseudocercospora herpotrichoides*). Hastalık, genç bitkilerde, toprak yüzeyinin hemen altında kalan sap üzerinde, elips veya göz şeklinde lekelerin oluşmasıyla başlar. Lekeler giderek birleşerek renk değişimine neden olurlar. Zaman ilerledikçe lezyonlar sarımsı-kahve bir renge döner ve merkezinde koyu kahve renkli bir kısım bulunabilir. Mantar, yaprak kını içerisinde gelişerek su ve besin maddesi iletimini bozar, sapı zayıflatır ve yatmaya neden olur. Yatma tek taraflı olmayıp zıt yönlerde ve dağınık bir şekildedir. Bitki yukarı doğru çekildiğinde toprak seviyesinden kopar. Kırılan saplarda koyu kahve, kömürleşmiş bir görüntü vardır. Hastalık kökte etkili olmayıp, sadece sap üzerinde etkili olur ve hastalıklı saplar genellikle beyaz başak oluştururlar (Murray, 2006; DeWolf ve Shroyer, 2017; Hagerty ve ark. 2019).

HERBİSİT ETKİSİ

Buğdayda yabancı ot mücadelesinde, erken dönemde herbisit uygulandığında kardeşlenmeyi önler, bitkinin gelişmesini geriletir ve arazinin tamamında bodurlaşmaya neden olur. Erken uygulamaya benzer şekilde geç herbisit uygulaması da zararlıdır. Sapa kalkmadan sonra herbisit uygulaması yapıldığında veya başaklanmaya yakın dönemde çok düşük dozdaki herbisit bulaşması bile başakta şekil bozulmalarına, kısırlaşmaya ve beyaz başak oluşumuna neden olur (Shroyer ve ark. 1998; Peterson, 2017; DeWolf ve ark. 2023).

TEŞHİS ADIMLARI

Beyaz başak oluşum nedenlerini belirlemeye yardımcı olacağı düşünülen, arazideki izleme ve incelemeye yönelik bilgiler aşağıda adım adım şema halinde açıklanmaya çalışılmıştır (Ransom, 2016; Akkaya, 2024; Anonymous, 2024). Bir sonraki adıma geçişi gerektirmeyen kutuların zemini açık mavi olarak renklendirilmiştir.

Başaklanma öncesi kontrol**Adım-1**

Yaprak boyunca 1-2 adet, geniş sarı renkli çizgi var ise Adım-2'ye geçiniz, yaprakta kahverengi lekeler var ise Adım-3'e gidiniz.

**Adım-2**

Yaprak boyunca 1-2 adet, geniş sarı renkli çizgi ve merkezde kahverengi yaprak damarları var, sarı çizgiler yaprak kınına kadar uzanıyor ise muhtemel etmen *Cephalosporium gramineum* fungusudur. Kardeşlerde farklı derecede bodurluk ve tarlada öbekler halinde beyaz başak oluşumu söz konusu olur.

Adım-3

Yaprakta, etrafı sarı renkli halka ile çevrili kahve renkli lekeler oluşmuş ise muhtemel neden tan spot hastalığıdır. Lezyonlar olgunlaştıkça orta kısmında koyu renkli bir kısım oluşur. Giderek büyüüp birleşir ve yaprakta daha geniş bir alanı kaplar. Sapa kalkma döneminde alt yapraklarda başlar, tane gelişme döneminde üst yapraklara sıçrar. Başakta kısmi beyazlaşmaya ve tanenin koyu kırmızı bir renk almasına neden olur.

Başaklanma döneminde kontrol**Adım-1**

Sadece sapın üst kısmı veya başak açık yeşil veya beyaz renkli ise Adım-2'ye geçiniz. Bitkinin tamamı açık yeşil veya beyaz renkli ise Adım-8'e gidiniz.

**Adım-2**

Sap üst kısmından veya beyaz başaktan tutularak yukarı doğru çekildiğinde, sap dip kısmından kopuyor ise Adım-3'e geçiniz. Değilse

**Adım-3**

Çekildiğinde dipten kopan sapta üst boğum arasının dip kısmı incelenir. Bu kısım keskin bir bıçakla kesilmiş gibi temiz bir görüntüye sahip ise Adım-4'e geçiniz, değilse Adım-5'e gidiniz. Yukarı doğru çekildiğinde toprak seviyesinden kopan ve kırılan saplarda koyu kahve, kömürleşmiş bir görüntü var ise Adım-15'e gidiniz.



Adım-4
Muhtemel etmen ekin sap arısı (wheat stem sawfly)



Adım-5
Çekildiğinde kopan sapta, üst boğum arasının dip kısmı çığnenmiş gibi dağınık bir görünüme sahip ise muhtemel etmen buğday sap kurdu (wheat stem maggot)


Adım-6
Çekildiğinde kopmayan sapta başak kirli beyaz renklidir. En dipteki kirli beyaz renkli başakcık ile karşısındaki başakcığın dış kavuzları soyulur. Gelişmekte olan tane incelenir. Tane tebeşir beyazı gibi bir renge sahip ise muhtemel etmen fusarium başak yanıklığı (fusarium head blight, scab) olabilir. Gelişmekte olan tane kirli beyaz renkte değil veya hiç yoksa Adım-7'ye geçiniz.



Adım-7
Beyaz başak dağılımı tarladaki eğim veya toprak tipi ile bir uyum gösteriyor ise Adım-8'e gidiniz. Beyaz başak dağılımı eğim ve toprak tipi ile uyumlu değil ise muhtemel neden glyphosate etkisi olabilir. Başaklanmaya yakın dönemdeki çok düşük dozdaki glyphosate bulaşması bile başakta kısırlığa ve beyazlığa yol açabilir.



Adım-8
Bitkilerdeki renk kaybı tarlanın çukur yerlerinde yoğunluk gösteriyor ise su baskınından kuşkulananmak gerekir. Aksine renk kaybına sahip bitkiler tarlanın yüksek yerlerinde veya hafif yapılı toprak kısımlarında yoğunluk gösteriyor ise kuraklık zararından kuşkulananmak gerekir. Her iki durumda da başaklar kısmen veya tamamen steril kalabilir, sağlıklı başaklara göre erkenden renk kaybederler.



Adım-15

Muhtemel neden eyespot olabilir. Mantar, yaprak kını içerisinde gelişerek su ve besin maddesi iletimini bozar, sapı zayıflatır ve yatmaya neden olur. Yatma tek taraflı olmayıp zıt yönlerde ve dağınık bir şekildedir. Kırılan saplarda koyu kahve, kömürleşmiş bir görüntü vardır. Hastalık kökte etkili olmayıp, sadece sap üzerinde etkili olur ve hastalıklı sapsar genellikle beyaz başak oluştururlar.

SONUÇ

Bu çalışmada, buğdayda beyaz başak oluşumuna yol açan faktörler; iklim faktörleri, hastalıklar, zararlılar ve herbisit ana başlıkları altında açıklanmış, teşhise yardımcı olacağı düşünülen bilgiler verilmeye çalışılmıştır. Buğday, adaptasyon ve telafi yeteneği çok yüksek bir bitkidir. Ancak bu telafi yeteneği, verim ve kalitede az veya çok miktarda kayıplar pahasına gerçekleşir. Bu nedenle buğday üreticileri, vejetasyon süresince karşılaştıkları sorunları doğru teşhis etmeye, önlemler almaya ve yetiştirme koşullarını olabildiğince optimize etmeye çalışmalıdır. Doğru çözümün ilk adımının doğru teşhis olduğu unutulmamalıdır.

KAYNAKLAR

- Akkaya, A., 1994. Buğday Yetiştiriciliği. KSÜ Yayınları, 1, Ziraat Fakültesi Yayınları, 1, Kahramanmaraş.
- Akkaya, A., 2024. Buğday Üretim Sorunları ve Teşhisi. Atatürk Üniversitesi Yayınları, 1391, ISBN: 978-625-6983-49-6, <https://ekitap.atauni.edu.tr>.
- Anonim, 2008. Zirai Mücadele Teknik Talimatları Cilt 1. Gıda Tarım ve Hayvancılık Bakanlığı, Tarımsal Araştırmalar ve Politikalar Genel Müdürlüğü. <https://www.tarimorman.gov.tr/TAGEM/Belgeler/Teknik%20tal%C4%B1matlar%202008/C%C4%B0LT%201.pdf> (Erişim tarihi, 07.08.2023).
- Anonim, 2015. Kıymıl (*Aelia spp*). <https://ankara.tarimorman.gov.tr/Belgeler/liftet/kimilm%C3%BCcadesi.pdf> (Erişim tarihi, 07.08.2023).
- Anonim, 2016. Hububat Hastalık ve Zararlıları İle Mücadele. Gıda, Tarım ve Hayvancılık Bakanlığı Gıda ve Kontrol Genel Müdürlüğü Bitki Sağlığı ve Karantina Daire Başkanlığı. https://www.tarimorman.gov.tr/GKGM/Belgeler/Uretici_Bilgi_Kosesi/Dokumanlar/hububat.pdf (Erişim tarihi, 07.08.2023).
- Anonim, 2023. Tahıllar ve diğer bitkisel ürünlerin üretim miktarları. <https://data.tuik.gov.tr/Bulten/Index?p=Bitkisel-Uretim-1.Tahmini-2022-45502>. (Erişim tarihi, 19.01.2023).
- Anonymous, 2023. Worldwide production of grain in 2021/22. <https://www.statista.com/statistics/263977/world-grain-production-by-type/> (Erişim tarihi, 19.01.2023).
- Anonymous, 2024. White Heads in Wheat. University of Minnesota Extension, Minnesota Crop News, <https://blog-crop-news.extension.umn.edu/2019/07/white-heads-in-wheat.html> (Erişim tarihi, 04.02.2025).
- Babaroğlu, N.E., Akci, E., Çulcu, M., Yalçın, F., 2020. Süne ve Mücadelesi. Tarım ve Orman Bakanlığı, Gıda ve Kontrol Genel Müdürlüğü, Zirai Mücadele Merkez Araştırma Enstitüsü Müdürlüğü, Ankara.

- Bean, B., Miller, T., Duncan, R., 2012. Causes of White-Heads in Wheat. Texas AgriLIFE Extension.
- Bockus, W.W., Tisserat, N.A., 2005. Take-all root rot. The Plant Health Instructor. The American Phytopathological Society (APS). <https://www.apsnet.org/edcenter/disandpath/fungalasco/pdlessons/Pages/Takeall.aspx> (Erişim tarihi, 14.08.2023).
- Byamukama, E., Strunk, C., Tande, C., Ali, S., 2018. Wheat Diseases Identification Pocket Guide. SDSU Extension. <https://extension.sdstate.edu/sites/default/files/2018-12/03-2000-2018.pdf> (Erişim tarihi, 14.08.2023).
- DeWolf, E., Onofre, K.A., Diaz, D.R., Lollato, R., Lancaster, S., Whitworth, R.J., 2023. Diagnosing Wheat Production Problems, Kansas State University, Agricultural Experiment Station and Cooperative Extension Service.
- DeWolf, E.D., Shroyer, J., 2017. Wheat Disease Identification, Kansas State University, K-State Research and Extension – Bookstore, <https://bookstore.ksre.ksu.edu/pubs/MF2994.pdf> (Erişim tarihi, 16.08.2023).
- Gül, E., Karakaya, A., 2019. Kışlık Buğdaydaki Sistemik Bir Hastalık: *Cephalosporium Çizgi Hastalığı Cephalosporium gramineum* (syn. *Hymenula cerealis*). Uluslararası Doğu Akdeniz Tarımsal Araştırma Enstitüsü Dergisi, 2(1), 130-143.
- Hagerty, C.H., Smiley, R.W., 2017. Field Guide for Diagnosing Common Wheat Maladies of the Pacific Northwest. Oregon State University, University of Idaho, Washington State University. <https://extension.oregonstate.edu/catalog/pub/pnw-698-field-guide-diagnosing-common-wheat-maladies-pacific-northwest>, (Erişim tarihi, 14.02.2025).
- Hagerty, C., Kroese, D., Mundt, C., Heesacker, A., Zemetra, R., 2019. Eyespot of Wheat, How to identify and control this common fungal disease. Oregon State Uni. Extension Service.
- Huberli, D., 2015. Diagnosing take-all in cereals. Department of Primary Industries and Regional Development's Agriculture and Food. Government of Western Australia. <https://www.agric.wa.gov.au/mycrop/diagnosing-take-all-cereals> (Erişim tarihi, 17.08.2023).
- İslamoğlu, M., 2021. Adıyaman'da Süne (*Eurygaster* spp. (Heteroptera; Scutelleridae)'nin Ergin Parazitoit Türleri ve Parazitenme Oranları. Avrupa Bilim ve Teknoloji Dergisi, 31, (Ek Sayı 1), 634-639.
- Kılıç, H., Bayram, Y., Tekdal, S., 2017. Farklı Fenolojik Özelliklere Sahip Ekmeklik Buğday Genotiplerinin Ekin Sap Arısı [*Cephus pygmaeus* L. (Hymenoptera: Cephidae)] Zararına Dayanıklılık Yönünden İncelenmesi. YYÜ Tar Bil Derg, 27 (1), 65-76.
- Konodel, J., Shanower, T., Beauzay, P., 2010. Integrated Pest Management of Wheat Stem Sawfly in North Dakota. E1479. https://library.ndsu.edu/ir/bitstream/handle/10365/10435/e1479_2010.pdf?sequence=1 (Erişim tarihi, 16.08.2023).
- Lollato, R., 2020. Drought stress taking its toll on Kansas wheat production in 2020. Agronomy eUpdates. <https://eupdate.agronomy.ksu.edu/article/drought-stress-taking-its-toll-on-kansas-wheat-production-in-2020-386-1> (Erişim tarihi, 18.08.2023).
- Mackesy, D., Moylett, H., 2018. CPHST Pest Datasheet for *Eurygaster integriceps*. USDA-APHIS-PPQ-CPHST. <http://download.ceris.purdue.edu/file/3623> (Erişim tarihi, 16.08.2023).
- Michaud, J.P., 2013. Wheat Insects. Kansas State University, Department of Entomology, <https://entomology.k-state.edu/extension/insect-information/crop-pests/wheat/whitegrubs.html> (Erişim tarihi, 21.08.2023).
- Murray, T.D., 2006. Strawbreaker Foot Rot or Eyespot of Wheat. EB1378. Washington State University, College of Agricultural, Human, and Natural Resource Sciences.

<https://smallgrains.wsu.edu/disease-resources/soilborne-fungal-diseases/eyespot/>_____ (Eriřim tarihi, 23.08.2023).

Özberk, İ., Atlı, A., Yücel, A., Özberk, F., Cořkun, Y., 2005. Wheat stem sawfly (*Cephus pygmaeus* L) damage; impacts on grain yield, quality and marketing prices in Anatolia. *Crop Protection*, 24, 1054-1060.

Peterson, D., 2017. Spring herbicide application decisions on wheat: The importance of growth stage. *Agronomy eUpdate*, Kansas State University. https://webapp.agron.ksu.edu/agr_social/m_eu_article.throck?article_id=1290&eu_id=203 (Eriřim tarihi, 21.08.2023).

Ransom, J., 2016. Steps for Diagnosing Cause of White Heads in Small Grains. North Dakota State University Extension, <https://www.no-tillfarmer.com/articles/5888-steps-for-diagnosing-cause-of-white-heads-in-small-grains> (Eriřim tarihi 04.02.2025).

Raper, T., 2018. Freeze damage to winter wheat, forecast April 8th. The University of Tennessee Institute of Agriculture. <https://news.utcropt.com/2018/04/freeze-damage-to-winter-wheat-forecast-april-8th/> (Eriřim tarihi, 21.08.2023).

Shroyer, J.P., Bowden, R.L., Brooks, H.L., Peterson, D., Whitney, D.A., 1998. Diagnosing Wheat Production Problems in Kansas, Kansas State University, Agricultural Experiment Station and Cooperative Extension Service.

Tarla, G., 2017. Morphological diagnosis of *Aelia rostrata* Boh. (Heteroptera: Pentatomidae) parasitized by *Hexameris* sp. (Nematoda: Mermithidae) in Ankara, Turkey. *Türk. Biyo. Mücadele Derg.* 8(1), 21-28.

Yüceler, H.S., Karakaya, A., Oğuz, A.Ç., 2022. Buğdayda *Pyrenophora tritici-repentis* Tarafından Meydana Getirilen Sarı Leke Hastalığı. *DUFED*, 11(1), 183-203.

Wegulo, S.N., 2011. Tan spot of cereals. The American Phytopathological Society (APS). <https://www.apsnet.org/edcenter/disandpath/fungalasco/pdlessons/Pages/TanSpot.aspx> (Eriřim tarihi, 14.08.2023).

Whitworth, R.J., Sloderbeck, P., Davis, H., Cramer, G., Norton, A., 2023. Hessian Fly. Kansas Crop Pests. <https://bookstore.ksre.ksu.edu/pubs/MF2866.pdf> (Eriřim tarihi, 17.08.2023).

**SUSTAINABILITY OF SALEP PRODUCTION THROUGH AGRIVOLTAIC
APPLICATIONS: THE KALE TARIMGES MODEL**
**AGRİVOLTAİK UYGULAMALAR İLE SALEP ÜRETİMİNİN
SÜRDÜRÜLEBİLİRLİĞİ: KALE TARIMGES MODELİ**

Ali ÇELİK

Kahramanmaraş Sütçü İmam University, Faculty of Engineering and Architecture,
Department of Electrical and Electronics Engineering, Kahramanmaraş, Türkiye

ORCID ID: <https://orcid.org/0009-0001-8720-932X>

Doç. Dr. Furkan DİNÇER

Kahramanmaraş Sütçü İmam University, Faculty of Engineering and Architecture,
Department of Electrical and Electronics Engineering, Kahramanmaraş, Türkiye

ORCID ID: <https://orcid.org/0000-0001-6787-0850>

Abstract

Salep production contributes significantly to both the local and national economy, and salep production in Turkey has an important place in the world. Salep is a valuable raw material used primarily in the food, cosmetics and pharmaceutical sectors. It is an agricultural product with high added value. Increasing energy costs and low sectoral profits affect salep production like other agricultural products.

In this study, an application on salep production in Kale AgriPV project implemented in Kale neighborhood of Göksun district, Kahramanmaraş province is discussed. The application shows the benefits to the farmer and the improvement in the payback period of both agricultural production with greenhouse and energy production with solar power plant.

The greenhouse has an indoor area of 1,440 m² and an annual production of 2.3 tons is targeted. The solar power plant integrated into the salep greenhouse has a power capacity of 240 kW AC and 333 kWp DC. It is targeted to generate 500,000 kWh of electrical energy annually, 50% of which will be used for operation and the remaining 50% will be sold to the electricity grid. Economic analyses were performed under these design parameters and the payback period for the integrated investment in AgriPV was calculated as 4.94 years.

Keywords: Salep Production, Renewable Energy, AgriPV

Özet

Salep üretimi hem yerel hem de ulusal ekonomiye önemli katkılar sunmakta olup Türkiye'deki salep üretimi dünyada önemli bir yere sahiptir. Salep; başta gıda, kozmetik ve ilaç sektörlerinde kullanılan değerli bir hammaddedir. Katma değeri yüksek bir tarımsal üründür. Artan enerji maliyetleri ve sektörel kar düşüklükleri diğer tarım ürünleri gibi salep üretimini de etkilemektedir.

Bu çalışmada, Kahramanmaraş ili, Göksun ilçesi, Kale mahallesinde uygulanan Kale TarımGES projesinde salep üretimi üzerinde gerçekleştirilen bir uygulamayı ele almaktadır. Uygulama ile hem sera ile tarımsal üretimin hem de güneş enerji santrali ile enerji üretiminin beraber olmasının çiftçiye olan kazanç ve amorti sürelerindeki iyileştirmeyi göstermektedir.

Sera kapalı alanı 1.440 m² olacak şekilde, yıllık 2,3 ton üretim yapılması hedeflenmiştir. Salep serasına entegre edilen güneş enerjisi santrali, 240 kWe AC ve 333 kWp DC güç kapasitesine sahiptir. Yıllık 500.000 kWh elektrik enerjisi üretimi sağlanması hedeflenmekte olup, bu enerjinin %50'si işletme için kullanılırken, kalan %50'sinin elektrik şebekesine satılması planlanmıştır. Bu tasarım parametreleri altında ekonomik analizler gerçekleştirilmiş ve TarımGES entegre yatırımı için yatırım geri ödeme süresi 4,94 yıl olarak hesaplanmıştır.

Anahtar Kelimeler: Salep Üretimi, Yenilenebilir Enerji, TarımGES

Giriş


Türkiye'de salep, yaygın olarak gıda sektöründe kullanılmaktadır. Özellikle Maraş usulü dondurma içeriğinin temel içeriklerinden biri olup, dondurmanın istenilen kıvamda ve aromaya ulaşmasını sağlar. Yüksek kalitede dondurma üretimi için bir litre süte yaklaşık 5-6 gram salep eklenmesi yeterli olabilmektedir.

Son yıllarda, tarım sektöründeki artan enerji maliyetleri çiftçileri alternatif çözümler üretmeye yönlendirmiştir. Bu bağlamda, sürdürülebilir tarım uygulamaları giderek daha fazla ön plana çıkmaya başlamıştır. Yenilenebilir enerji kaynakları içerisinde arasında özellikle güneş enerjisi kaynağı sınırsız olması, hammadde gerektirmemesi ve çevre dostu olmasından ötürü daha çok ön plana çıkmaktadır. Güneş enerjisi kaynağı, fotovoltaik (PV) paneller aracılığıyla güneş ışınımını elektrik enerjisine dönüştürerek temiz, sürdürülebilir ve ekonomik bir enerji kaynağı sunmaktadır. PV panellerin ucuzlaması ile beraber tarım sektöründe entegre kullanılması yönünde çalışmalar daha çok yapılmaya başlanmıştır.

Güneş enerjisi destekli tarım uygulamaları, enerji maliyetlerinin azaltılmasıyla birlikte, aynı zamanda bitki örtüsü açısından da avantajlar sunmaktadır. PV paneller, ekili alanlarda aşırı sıcaklık ve buharlaşmayı azaltma ve su tasarrufu sağlamaktadır. Özellikle serin ve nemli ortamları tercih eden bitkiler için bu sistemin büyük bir avantajı bulunmaktadır.

Salep üreticiliği açısından değerlendirildiğinde, güneş enerjisi destekli tarım sistemleri önemli bir fırsat sunmaktadır. Salep, yetiştirme koşulları bakımından hassas bir bitki olup, aşırı sıcaklık ve kuraklık gibi olumsuz faktörlerden etkilenebilmektedir. PV paneller altında depolanan salep ürünleri, merkezi coğrafya olan coğrafya ortamı sağlayarak su kaybını en aza indirmek, toprak nemini korumakta ve üretim üretimini artırmaktadır.

Bu çalışmada, Kahramanmaraş ili, Göksun ilçesi, Kale mahallesinde uygulanan Kale TarımGES projesinde salep üretimi üzerinde gerçekleştirilen bir uygulamayı ele almaktadır. Uygulama ile hem sera ile tarımsal üretimin hem de güneş enerji santrali ile enerji üretiminin beraber olmasının çiftçiye olan kazanç ve amorti sürelerindeki iyileştirmeyi göstermektedir. Her iki işletme tesisi entegre çalışacak şekilde bir tasarım gerçekleştirilmiştir. Böylece tarımsal üretim için enerji ihtiyacı karşılanmakta hem de fazla üretilen elektrik enerjisi mevzuatın vermiş olduğu izin ile de elektrik şebekesine satılmaktadır. Şekil 1'de Kale TarımGES projesi için sehpa tasarım modeli teknik çizimi gösterilmektedir. Çatı kısmı olarak tasarlanan bölüm tamamen PV paneller ile kaplı konumdadır. Yerden yükseklikler yeterli ölçüde bırakılmıştır ki, içerisinde küçük traktör çalışabilecektir. Aynı zamanda havalandırma için ekstra fan da konumlandırılmıştır. Burada PV paneller şeffaf olacak şekilde ve ışınımı içeri alacak şekilde olması planlanmıştır. PV panellerde backsheet materyalinin şeffaf seçilmesi ile beraber üretilen panellerin ışınım geçirgenliği daha yüksek olmaktadır.



Şekil 1. Kale TarımGES projesi için sehpa tasarım modeli teknik çizimi

Salep için Ekonomik Analiz

Salep üretimi yapılan seranın toplam alanı 1.440 m² olacak şekilde, yıllık 2,3 ton üretim yapılması hedeflenmiştir. Tablo 1’de bu üretim planlamasına ait varsayımlar ve ekonomik analiz detayları sunulmuştur. Üretimden elde edilen brüt gelir 2.070.000 TL olarak hesaplanırken, toplam yıllık maliyetler 850.000 TL seviyesindedir. Bu maliyetler arasında personel, bakım, diğer giderler ve enerji tüketimi yer almakta olup, GES sayesinde enerji giderleri sıfırlandığı bir üretim senaryosu oluşturulmuştur. Salep üretiminin finansal analizine göre, işletme yıllık 1.220.000 TL net gelir sağlamaktadır. Yüksek satış fiyatı ve düşük işletme maliyetleri sayesinde, yatırımın kârlılığı oldukça güçlüdür.

Tablo 1. Salep için Ekonomik Analiz

Çalışma Periyodu	12 ay
Sera Alanı	1.440 m ²
Üretim Hedefi	1,6 kg/m ²
Yıllık Üretim	2,3 ton
Ortalama Toptan Satış Fiyatı	900 TL
Yıllık Gelir	2.070.000 TL
Yıllık Üretim Maliyeti Toplamı	440.000 TL
Yıllık Personel Gideri	240.000 TL
Yıllık Bakım Giderleri	90.000 TL
Yıllık Diğer Giderler	80.000 TL
Yıllık Enerji Gideri	-
Yıllık Gider Toplamı	850.000 TL
Yıllık Net Gelir	1.220.000 TL

Güneş Santrali için Ekonomik Analiz

Salep serasına entegre edilen güneş enerjisi santrali, 240 kWe AC ve 333 kWp DC güç kapasitesine sahiptir. Yıllık 500.000 kWh elektrik enerjisi üretimi sağlanması hedeflenmekte olup, bu enerjinin %50'si işletme için kullanılırken, kalan %50'sinin elektrik şebekesine satılması planlanmıştır. Mevcut elektrik enerjisi satış fiyatı 2,93 TL/kWh olduğu varsayıp, bu satışlardan yıllık 732.500 TL gelir elde edilmesi planlanmıştır. Santralin işletme maliyetleri toplamda 417.000 TL seviyesinde olup, bakım, sigorta, dağıtım bedeli ve bekçi giderlerini kapsamaktadır. Tüm hesaplamalar sonucunda, GES yıllık 315.500 TL net gelir sağlamaktadır. Böylece sera işletmesinin enerji ihtiyacı karşılanırken, aynı zamanda ek bir gelir kaynağı oluşturulması sağlanmıştır.

Tablo 2. GES için Ekonomik Analiz

AC Güç	240 kWe
DC Güç	333 kWp
Panel Gücü	500 Wp
Panel Adedi	666
İnvertör Gücü	50 kWe
İnvertör Adedi	5
Yıllık Üretim	500.000 kWh
Yıllık Sera Tüketimi	250.000 kWh
Yıllık Şebeke Satış Üretimi	250.000 kWh
Mevcut Satış Birim Fiyatı (TL/kWh)	2,93 TL
Toplam GES Brüt Gelir (TL/Yıl)	732.500 TL
Yıllık Bakım Gideri	50.000 TL
Yıllık Sigorta Gideri	50.000 TL
Yıllık Dağıtım Bedeli	167.100 TL
Yıllık Gider Toplamı	417.000 TL
Yıllık Net Gelir	315.500 TL

Genel İşletme Performansı ve Yatırımın Ekonomik Analizi

Salep üretimi ve güneş santralinin entegrasyonu ile oluşturulan işletmenin toplam yıllık net geliri 1.535.500 TL olarak hesaplanmıştır. Bu modelin toplam yatırım maliyeti 7.600.000 TL olup, yapılan finansal analiz sonucunda 4,94 yıl gibi kısa bir sürede yatırımın kendini amorti ettiği görülmektedir.

Tablo 3. Entegre TarımGES Projesi için Ekonomik Analiz

Sera Yıllık Net Gelir	1.220.000 TL
GES Yıllık Net Gelir	315.500 TL
Toplam Yıllık Net Gelir	1.535.500 TL
Yatırım Maliyeti (KDV Hariç)	7.600.000 TL
Geri Dönüş Süresi	4,94 Yıl

Sonuç ve Öneriler

Salep üretimi ve güneş enerjisi santrali entegrasyonu, tarımsal işletmelerin enerji bağımsızlığını sağlarken, sürdürülebilir ve kârlı bir üretim modeli de oluşturmaktadır. Sadece salep üretimi yapıldığında çiftçiler için kar marjı yeterli görülemeyebilmektedir. Ancak, hem salep üretimi yaparken elektrik enerjisi maliyetlerinin giderilmesi hem de fazla üretilen elektrik enerjisinin şebekeye satılması ile birlikte bir gelir modeli oluşturulabilmesi işletmenin rekabet gücünü artırmaktadır. Aynı zamanda sera içerisinde üretilecek salep için üretim sezonu 12 ay olarak tasarlanabilmektedir. Sonuç olarak, salep serası ve güneş enerjisi santralinin entegrasyonu, çevre dostu ve ekonomik olarak sürdürülebilir bir işletme modeli sunmaktadır. Tarım ve enerji sektörlerinin birleşimi, gelecekte benzer projeler için örnek teşkil edebilecek önemli bir adımdır.

INTEGRATION OF RENEWABLE ENERGY AND AGRICULTURE WITH AGRIVOLTAICS: THE CASE OF KAHRAMANMARAŞ BULANIK TARIMGES

TARIMGES İLE YENİLENEBİLİR ENERJİ VE TARIMIN ENTEGRASYONU: KAHRAMANMARAŞ BULANIK TARIMGES ÖRNEĞİ

Ali ÇELİK

Kahramanmaraş Sütçü İmam University, Faculty of Engineering and Architecture,
Department of Electrical and Electronics Engineering, Kahramanmaras, Türkiye

ORCID ID: <https://orcid.org/0009-0001-8720-932X>

Doç. Dr. Furkan DİNÇER

Kahramanmaraş Sütçü İmam University, Faculty of Engineering and Architecture,
Department of Electrical and Electronics Engineering, Kahramanmaras, Türkiye

ORCID ID: <https://orcid.org/0000-0001-6787-0850>

Abstract

Agrioltaics is a term that refers to agriculture integrated with photovoltaic panels. It combines soil-based agriculture with the generation of electrical energy from solar energy. In our country, it is also called AgriPV. With AgriPV, it provides advantages in many parameters by utilizing both agricultural lands and the areas needed for electrical energy generation together.

In this study, the Bulanık Agri-PV project designed for the Bulanık neighborhood of Dulkadiroğlu district, Kahramanmaraş province is analyzed in detail from an economic point of view. A 3,075 m² greenhouse operation was financially analyzed. During the production process, tomato cultivation is carried out for 12 months and the targeted productivity is determined as 70 kg/m². Accordingly, the total annual production amount is calculated as 215.25 tons. A solar power plant with 480 kWe AC power and 572 kWp DC power was designed to be integrated into this facility. The plant generates 1,000,000 kWh of electricity on an annual basis, of which 500,000 kWh is sold. The current sales price is determined as 2.77 TL/kWh and accordingly, the annual gross income of the plant is calculated as 1,385,000 TL. The payback period of the investment is determined as 4.44 years. This is a very short payback period, especially when compared to investments in the agriculture and energy sectors.

Keywords: Tomato, Solar Energy, Agriculture, Agri-PV

Özet

Agrioltaik kavramı, fotovoltaik paneller ile entegre yapılan tarımı ifade eden bir terimdir. Toprağa dayalı tarım ile güneş enerjisinden elektrik enerjisi üretimini birleştirmektedir. Ülkemizde TarımGES olarak da adlandırılmaktadır. TarımGES ile hem tarım arazilerinin değerlendirilmesi hem de elektrik enerjisi üretimi konusunda ihtiyaç duyulan alanların beraber kullanılması ile birçok parametre konusunda avantaj sağlamaktadır.

Bu çalışmada Kahramanmaraş ili, Dulkadiroğlu ilçesi, Bulanık mahallesi için tasarlanan Bulanık TarımGES projesi ekonomik açıdan detaylı olarak analiz edilmiştir. 3.075 m² büyüklüğünde bir sera işletmesinin mali analizi yapılmıştır. Üretim süreci boyunca 12 ay

boyunca domates yetiştiriciliği gerçekleştirilmekte olup, hedeflenen verimlilik 70 kg/m² olarak belirlenmiştir. Bu doğrultuda yıllık toplam üretim miktarı 215,25 ton olarak hesaplanmıştır. Bu tesise entegre çalışacak 480 kWe AC gücüne ve 572 kWp DC gücüne sahip bir güneş enerjisi santrali tasarlanmıştır. Santral yıllık bazda 1.000.000 kWh elektrik üretimi gerçekleştirmekte olup, bunun 500.000 kWh'lik bölümü satılmaktadır. Mevcut satış fiyatı 2,77 TL/kWh olarak belirlenmiş ve buna bağlı olarak santralin yıllık brüt geliri 1.385.000 TL olarak hesaplanmıştır. Yatırımın geri dönüş süresi 4,44 yıl olarak belirlenmiştir. Bu süre, özellikle tarım ve enerji sektörlerindeki yatırımlarla kıyaslandığında oldukça kısa sayılabilecek bir geri dönüş süresidir.

Anahtar Kelimeler: Domates, Güneş Enerjisi, Tarım, TarımGES

Giriş

Enerji ihtiyacının artışı ve sürdürülebilir tarım uygulamalarının gerekliliği, yenilenebilir enerji santrallerine olan ihtiyacı artırmaktadır. Bu bağlamda, tarım ve fotovoltaik (PV) santraller (TarımGES), aynı araziye hem elektrik enerjisi üretimi hem de tarımsal faaliyetler için kullanmayı mümkün kılan yenilikçi bir yaklaşım olarak öne çıkabilmektedir. Entegre olarak çalışan bu işletmelere TarımGES işletmeleri adı da verilebilir. Tarımsal üretimin devamlılığını sağlarken aynı zamanda güneş enerjisinden faydalanarak çevre dostu bir enerji kaynağı sunulabilmektedir.

Geleneksel güneş enerjisi santralleri genellikle tarım arazilerinde kurularak tarımsal üretime ayrılan alanları azaltırken, TarımGES sistemler bu iki alanı entegre ederek toprak kullanım verimliliğini artırmayı amaçlamaktadır. Şekil 1'de bu modele ait bir görüntü sunulmuştur. Kahramanmaraş ili, Dulkadiroğlu ilçesi, Bulanık mahallesinde yer alan Bulanık TarımGES projesidir. Hem kapalı sera alanı oluşturulurken çatı alanı tamamen PV paneller ile kaplıdır. PV panellerde ayrıca şeffaf backsheet materyali kullanılmakta ve güneş ışınımının seraya ulaşması sağlanmaktadır. Daha fazla güneş ışınımı isteyen seralar için ise PV panellerdeki hücre sayısı yarı yarıya düşürülerek daha fazla güneş ışınımının seraya geçmesi sağlanabilir. Şekil 2 ve 3'te tesise ait görüntüler yer almaktadır.



Şekil 1. Bulanık TarımGES Tesisine Ait Görüntü



Şekil 2. Bulanık TarımGES Tesisine Ait Görüntü

İnverter yer seçimi belirlenirken, elektrik kaybının en az olacağı, gerilim düşümünün ideal sınırlar içerisinde kalacağı optimum noktanın belirlenmesi gerekmektedir. Güneş enerji santrallerinde inverter yeri seçimi, montajlanan bölge, santralin verimli çalışması, uzun ömürlü olması ve bakım kolaylığı açısından önemlidir.



Şekil 3. Bulanık TarımGES Tesisine Ait Görüntü

Proje Bilgileri

Kahramanmaraş ili, Dulkadiroğlu ilçesi, Bulanık bölgesi tarımsal ürün olarak domates üretimine elverişli tarım arazilerine sahiptir. Bulanık TarımGES üretim merkezinde de tarımsal ürün olarak domates üretimi gerçekleştirilmesi planlanmıştır. Yapılan analiz çalışmasında sera modeli ile domates üretimi gerçekleştirilip aynı arazi üzerinde de enerji üretimi gerçekleştirilerek sağlanacak mali fayda aşağıda gösterilmiştir. Tablo 1’de projeye ait genel bilgiler yer almaktadır.

Tablo 1. Proje Bilgileri

Proje Adı	Bulanık TarımGES
Proje Konumu	Bulanık Mahallesi, Dulkadiroğlu / Kahramanmaraş
Ada Parsel	114/7
Proje Alanı	8.025 m ²
Enerji Kaynağı	Güneş Enerjisi Santrali (GES)
GES Gücü	572 kWp
Serada Hedef Ürün	Domates
Sera Alanı	3.075 m ²
Toplam Yatırım Maliyeti	11.000.000 TL

Sera için Ekonomik Analiz

3.075 m² büyüklüğünde bir sera işletmesinin mali analizi yapılmıştır. Üretim süreci boyunca 12 ay boyunca domates yetiştiriciliği gerçekleştirilmekte olup, hedeflenen verimlilik 70 kg/m² olarak belirlenmiştir. Bu doğrultuda yıllık toplam üretim miktarı 215,25 ton olarak hesaplanmıştır. Tablo 2’de bu kapasite için ekonomik analiz bilgileri detaylı olarak hesaplanmıştır.

Tablo 2. Sera için Ekonomik Analiz

Ürün	Bulanık Domatesi
Çalışma Periyodu	12 ay
Sera Alanı	3.075 m ²
Üretim Hedefi (kg/m²)	70 kg
Yıllık Üretim	215,25 ton
Ortalama Toptan Satış Fiyatı	10 TL
Yıllık Gelir (TL/yıl)	2.152.500 TL
Yıllık Üretim Maliyeti Toplamı (TL/yıl)	140.000 TL
Yıllık Personel Gideri (TL/yıl)	240.000 TL
Bakım Giderleri (TL/yıl)	90.000 TL
Diğer Giderler (TL/yıl)	50.000 TL
Enerji Gideri (TL/yıl)	-
Yıllık Gider Toplamı (TL/yıl)	520.000 TL
Yıllık Net Gelir (TL/yıl)	1.632.500 TL

Ekonomik açıdan değerlendirildiğinde, ortalama toptan satış fiyatının 10 TL/kg olması varsayılarak yıllık toplam gelir 2.152.500 TL olarak belirlenmiştir. Buna karşılık, sera işletmesinin yıllık üretim maliyetleri toplamı 140.000 TL, personel giderleri 240.000 TL, bakım giderleri 90.000 TL ve diğer giderler 50.000 TL olarak hesaplanmıştır. Böylece toplam yıllık gider 520.000 TL seviyesine ulaşmaktadır. Tüm giderler düşüldüğünde, sera işletmesinin yıllık net geliri 1.632.500 TL olarak görülmektedir. Mali analizde dikkat çeken önemli bir husus, enerji giderinin belirtilmemiş olmasıdır. Güneş enerji sistemi sayesinde enerji kullanılacak mali hizmet sıfır olacaktır. Kurulacak TarımGES modeli ile ihtiyaç duyulan enerji GES tarafından karşılanacaktır.

GES için Ekonomik Analiz

Bu çalışma kapsamında 480 kWe AC gücüne ve 572 kWp DC gücüne sahip bir güneş enerjisi santralının mali analizi değerlendirilmiştir. Tablo 3’te GES için ekonomik analiz detaylı olarak sunulmuştur. Santral, 1050 adet 545 W gücündeki güneş panelinden oluşmakta ve 5 adet 100 kW’lık inverter ile enerji dönüşümü sağlanmaktadır.

Santral yıllık bazda 1.000.000 kWh elektrik üretimi gerçekleştirmekte olup, bunun 500.000 kWh’lik bölümü satılmaktadır. Mevcut satış fiyatı 2,77 TL/kWh olarak belirlenmiş ve buna bağlı olarak santralin yıllık brüt geliri 1.385.000 TL olarak hesaplanmıştır.

Tablo 3. GES için Ekonomik Analiz

AC Güç	480 kWe
DC Güç	572 kWp
Panel Gücü	545 Wp
Panel Adedi	1050
İnvertör Gücü	100 kWe
İnvertör Adedi	5
Yıllık Üretim	1.000.000 kWh
Yıllık Sera Tüketimi	500.000 kWh
Yıllık Üretim Fazlası Şebeke Satışı	500.000 kWh
Mevcut Satış Birim Fiyatı (TL/kWh)	2,77 TL
Toplam GES Brüt Gelir (TL/Yıl)	1.385.000 TL
Yıllık Bakım Gideri	30.000 TL
Yıllık Sigorta Gideri	30.000 TL
Yıllık Dağıtım Bedeli	332.000 TL
Yıllık Bekçi Bedeli	150.000 TL
Yıllık Gider Toplamı	542.000 TL
Yıllık Net Gelir	843.000 TL

Yıllık giderler incelendiğinde, bakım maliyeti 30.000 TL, sigorta gideri 30.000 TL, dağıtım bedeli 332.000 TL ve güvenlik için bekçi gideri 150.000 TL olmak üzere toplam 542.000 TL'lik bir işletme gideri olduğu görülmektedir. Tüm giderler düşüldüğünde santralin yıllık net geliri 843.000 TL olarak hesaplanmıştır.

Entegre TarımGES için Ekonomik Analiz

Yapılan mali analiz doğrultusunda detaylı bilgiler Tablo 4'te sunulmuştur. İşletmenin yıllık toplam net geliri 2.475.500 TL olarak hesaplanmıştır. Bu gelirin 1.632.500 TL'si sera faaliyetlerinden, 843.000 TL'si ise güneş enerjisi santralinden (GES) elde edilmektedir. İşletmenin farklı gelir kaynaklarına sahip olması hem finansal sürdürülebilirliği artırmakta hem de yatırımın güvenilirliğini desteklemektedir. Toplam yatırım maliyeti KDV hariç 11.000.000 TL olarak hesaplanmış olup, yatırımın geri dönüş süresi 4,44 yıl olarak belirlenmiştir. Bu süre, özellikle tarım ve enerji sektörlerindeki yatırımlarla kıyaslandığında oldukça kısa sayılabilecek bir geri dönüş süresidir.

Tablo 4. Entegre TarımGES İşletmesi için Ekonomik Analiz

Sera Yıllık Net Gelir	1.632.500 TL
GES Yıllık Net Gelir	843.000 TL
Toplam Yıllık Net Gelir	2.475.500 TL
Yatırım Maliyeti (KDV Hariç)	11.000.000 TL
Geri Dönüş Süresi	4,44 Yıl

Sonuç ve Öneriler

Bu çalışmada, Kahramanmaraş ili, Dulkadirođlu ilçesi, Bulanık bölgesinde tasarlanan, yenilenebilir enerji üretimi ile tarımsal faaliyetlerin entegrasyonu değerlendirilmiştir. Elde edilen bulgular, TarımGES entegre işletmelerin hem enerji üretimi hem de tarımsal sürdürülebilirlik açısından önemli avantajlar sunduđunu göstermektedir. Elde edilen sonuçlar, işletmenin yüksek kârlılıđa sahip olduđunu ve yatırımın ekonomik açıdan fizibil olduđunu göstermektedir. Yatırımın geri dönüş süresi 4,44 yıl olarak belirlenmiştir. Güneş enerjisi santrali sayesinde sera faaliyetlerinde kullanılan enerji maliyetleri minimize edilirken, aynı zamanda fazla üretilen elektriđin satışıyla ek bir gelir elde edilmektedir.

IMPACTS OF TEMPERATURE FLUCTUATIONS ON AQUACULTURE AND MITIGATION STRATEGIES**SICAKLIK DALGALANMALARININ SU ÜRÜNLERİ YETİŞTİRİCİLİĞİ ÜZERİNDEKİ ETKİLERİ VE AZALTMA STRATEJİLERİ****Boran KARATAŞ**

Van Yüzüncü Yıl University, Faculty of Fisheries, Department of Aquaculture, Van, Türkiye

ORCID ID: <https://orcid.org/0000-0003-4353-1293>**ABSTRACT**

The increasing frequency of extreme temperature fluctuations due to climate change poses a significant environmental threat to aquaculture. As ectothermic organisms, fish are highly susceptible to temperature variations, which directly affect their growth, metabolism, immune responses, and overall health. This study examines the physiological and molecular impacts of temperature changes on fish, focusing on stress biomarkers, gene expression alterations, and potential strategies for sustainable aquaculture. Water temperature is a critical factor regulating fish metabolism, digestion, and overall physiological functions. Elevated temperatures generally accelerate metabolic rates, leading to increased feed intake but also higher oxygen demand and ammonia excretion. Conversely, sudden temperature fluctuations can disrupt homeostasis, negatively affecting growth performance and reducing feed efficiency. At the molecular level, temperature shifts induce significant changes in the expression of genes related to antioxidant defense, immune response, and stress regulation. Heat shock proteins (HSPs), cytokines (e.g., IL-1 β , IL-8, TNF- α), and oxidative stress markers (SOD, CAT, GPx) exhibit altered expression under thermal stress, indicating physiological adaptation or distress. Chronic exposure to suboptimal temperatures can suppress immune functions, increasing disease susceptibility. To mitigate the adverse effects of climate change, various strategies can be implemented, including selective breeding for thermal resilience, optimizing stocking densities, and integrating water cooling and aeration technologies. Nutritional interventions, such as incorporating antioxidants, probiotics, and omega-3 fatty acids into fish diets, can enhance thermal stress resistance. Additionally, advanced monitoring technologies, including automated temperature sensors and real-time environmental data analysis, can aid in optimizing culture conditions and minimizing stress-related impacts. In conclusion, understanding the physiological and molecular responses to thermal stress is crucial for ensuring the sustainability of aquaculture under changing climatic conditions. Future research should focus on integrating genetic, nutritional, and precision aquaculture technologies to develop innovative mitigation strategies and enhance fish resilience to temperature fluctuations.

Key Words: Sustainable aquaculture; Thermal stress; Fish physiology; Gene expression**ÖZET**

İklim değişikliği nedeniyle aşırı sıcaklık dalgalanmalarının artan sıklığı su ürünleri yetiştiriciliği için önemli bir çevresel tehdit oluşturmaktadır. Ektotermik organizmalar olarak balıklar, büyümelerini, metabolizmalarını, bağışıklık tepkilerini ve genel sağlıklarını doğrudan etkileyen sıcaklık değişimlerine karşı oldukça hassastır. Bu bildiri, sıcaklık değişimlerinin

balıklar üzerindeki fizyolojik ve moleküler etkilerini inceleyerek stres biyobelirteçlerine, gen ifadesi değişikliklerine ve sürdürülebilir su ürünleri yetiştiriciliği için olası stratejilere odaklanmaktadır. Su sıcaklığı, balık metabolizmasını, sindirimi ve genel fizyolojik işlevleri düzenleyen kritik bir faktördür. Yüksek sıcaklıklar genellikle metabolik oranları hızlandırır, bu da artan yem alımına ancak aynı zamanda daha yüksek oksijen talebine ve amonyak atılımına yol açar. Tersine, ani sıcaklık dalgalanmaları homeostaziye bozabilir, büyüme performansını olumsuz etkileyebilir ve yem verimliliğini azaltabilir. Moleküler düzeyde, sıcaklık değişimleri antioksidan savunma, bağışıklık tepkisi ve stres düzenlemesiyle ilgili genlerin ifadesinde önemli değişikliklere neden olur. Isı şoku proteinleri (HSP'ler), sitokinler (örneğin, IL-1 β , IL-8, TNF- α) ve oksidatif stres belirteçleri (SOD, CAT, GPx), termal stres altında değişmiş ifade sergiler ve bu da fizyolojik adaptasyon veya sıkıntıyı gösterir. Kronik olarak suboptimal sıcaklıklara maruz kalmak, bağışıklık fonksiyonlarını baskılayarak hastalık duyarlılığını artırabilir. İklim değişikliğinin olumsuz etkilerini azaltmak amacıyla, termal dayanıklılık için seçici üreme, stoklama yoğunluklarını optimize etme ve su soğutma ve havalandırma teknolojilerini entegre etme gibi çeşitli stratejiler uygulanabilir. Antioksidanlar, probiyotikler ve omega-3 yağ asitlerini balık diyetlerine dahil etme gibi beslenme müdahaleleri, termal stres direncini artırabilir. Ayrıca, otomatik sıcaklık sensörleri ve gerçek zamanlı çevresel veri analizi gibi gelişmiş izleme teknolojileri, kültür koşullarının optimize edilmesine ve stresle ilgili etkilerin en aza indirilmesine yardımcı olabilir. Sonuç olarak, termal strese karşı fizyolojik ve moleküler tepkileri anlamak, değişen iklim koşullarında su ürünleri yetiştiriciliğinin sürdürülebilirliğini sağlamak için çok önemlidir. Gelecekteki araştırmalar, yenilikçi azaltma stratejileri geliştirmek ve balıkların sıcaklık dalgalanmalarına karşı dayanıklılığını artırmak için genetik, beslenme ve hassas su ürünleri yetiştiriciliği teknolojilerini entegre etmeye odaklanmalıdır.

Anahtar Kelimeler: Sürdürülebilir su ürünleri yetiştiriciliği; Termal stres; Balık fizyolojisi; Gen ifadesi

WHEAT BASED INTERCROPPING**BUĞDAY ESASLI KARIŞIK EKİM****Prof. Dr. Aydın AKKAYA**Muş Alparslan Üniversitesi Uygulamalı Bilimler Fakültesi Bitkisel Üretim ve Teknolojileri
Bölümü

ORCID; 0000-0001-9560-1922

ÖZET

Küresel ısınmaya bağlı iklim değişikliği; kuraklık, aşırı yağmur, su baskını, soğuk zararı, don etkisi gibi ekstrem olaylara yol açmakta, tarımsal üretim yönünden giderek artan bir risk oluşturmaktadır. Gıda güvenliği yönünden stratejik bir öneme sahip olan buğday, olumsuz koşulların etkisini ve yetiştirici hatalarını telafi etme yeteneğine sahiptir. Ancak bu telafi, verim ve kalitede önemli kayıplar pahasına gerçekleşmektedir. Bu nedenle iklim ve toprak koşullarına bağlı olumsuz etkilerin olabildiğince telafi edilebilmesi yönünden doğru yetiştirme tekniklerinin uygulanması büyük önem taşımaktadır. Toprak organik maddesini artırmak, toprak işlemeyi azaltmak, malç uygulamak, geleneksel kara nadas-buğday uygulaması yerine baklagil-buğday sistemini ikame etmek gibi kültürel önlemler sürdürülebilir buğday tarımı açısından önemli yararlar sağlamaktadır. Sürdürülebilir buğday tarımı açısından yararlı kültürel uygulamalardan birisi de karışık ekimdir. Mono kültür buğday tarımına göre buğday esaslı karışık ekim; su, besin elementleri ve ışık gibi kaynakların daha etkin bir şekilde kullanımını sağlamaktadır. Buğday esaslı karışık ekim tarımsal üretimde çeşitliliği artırdığı için hastalık, zararlı ve yabancı ot etkilerini azaltabilme, toprak verimliliğini artırabilme, sürdürülebilir bir üretim sağlayabilme potansiyeline sahiptir. Modern tarımsal teknoloji, yeni ıslah çeşitleri, yetiştirme, pazarlama, işleme gibi faktörlere bağlı olarak buğday esaslı karışık ekimin genel kabul görmesi ve yaygınlaşması bakımından bazı zorluklar bulunmaktadır. Ancak mono kültür üretim sistemlerinin çevre ve insan sağlığı üzerindeki olumsuz etkileri görüldükçe, sürdürülebilir ve çevreci üretim sistemlerinden birisi olan karışık ekim giderek artan bir ilgiye sahip olmaktadır. Ülkemizde tarla tarımında buğday esaslı karışık ekim sistemleri pek çalışılmamıştır. Buğday, karışık yetiştirmeye uygun tahıl bitkilerinden birisi olup, buğday esaslı karışık ekim, özellikle organik tarımla entegre edilmesi halinde, sürdürülebilir tarım açısından daha fazla yararlar sağlama potansiyeline sahiptir.

Anahtar Kelimeler: Buğday, karışık ekim, sürdürülebilir tarım**ABSTRACT**

Climate change due to global warming causes extreme events such as drought, heavy rain, flooding, low temperature damage, freezing effects and is becoming progressively increasing risk for agricultural production. Wheat, which has a strategic importance in terms of food security, has the ability to compensate the effects of adverse conditions and grower errors. However, this compensation comes at the cost of significant losses in yield and quality. For this reason, it is of great importance to apply correct cultivation techniques in order to compensate as much as possible the negative effects due to climate and soil conditions. Cultural measures such as increasing of soil organic matter, reducing of soil tillage, mulch application,

implementation of legumes-wheat system instead of conventional fallow-wheat application provide important benefits in terms of sustainable wheat farming. One of the useful cultural practices for sustainable wheat farming is intercropping. Wheat based intercropping compared to monoculture wheat farming enables more efficient use of sources such as water, nutrients and light. Wheat based intercropping has a potential to be able to reduce the diseases, insect and weed impacts and to increase soil fertility and to provide a sustainable production, since it increases biodiversity in agricultural production. There are some difficulties in the general acceptance and widespread use of wheat-based intercropping due to factors such as modern agricultural technology, new breeding varieties, management, marketing and processing. However, as the negative effects of monoculture production systems on the environment and human health have been seen, intercropping, one of the sustainable and environmentally friendly production systems, is gaining an increasing interest. In Türkiye, wheat based intercropping systems in the field farming have generally not been studied. Wheat is one of the cereal crops suitable for intercropping, and wheat-based intercropping system has a potential to provide greater benefits in terms of sustainable agriculture, especially in the case of integrated with organic farming.

Keywords: Wheat, intercropping, sustainable agriculture

GİRİŞ

Buğday, dünyada yaklaşık 780 milyon ton yıllık üretimle (Anonymous, 2023), gıda güvenliği yönünden stratejik bir öneme sahiptir. Ülkemizde 2022 yılı rakamlarına göre yıllık 37 milyon tonluk tahıl üretimi içerisinde, 20 milyon tonluk üretimle buğday ilk sırada yer almaktadır (Anonim, 2023). Tarım teknolojisindeki gelişmeler, yetiştirme, hasat, depolama, pazarlama ve işleme kolaylıkları gibi faktörler buğday üreticilerini mono kültür buğday tarımına zorlamaktadır. Mono kültür buğday tarımı yüksek girdiye ihtiyaç göstermekte olup, özellikle kullanılan tarımsal kimyasallar ekosistem dengesi ve zenginliğini bozmaktadır. Ekosistem dengesindeki bozulma, biyoçeşitlilik ve toprak verimliliğindeki azalma, sürdürülebilir buğday verimliliğini giderek daha çok tehdit etmektedir. Küresel iklim değişikliği bu olumsuz etkilerin baskısını daha çok artırmakta (Arora, 2019), ortalama sıcaklıktaki her bir derecelik artışın, buğday veriminde küresel ölçekte % 6 oranında bir azalmaya yol açabileceği hesaplanmaktadır (Asseng ve ark. 2015; Zhao ve ark. 2017).

Buğday tarımı ülkemizde, daha çok yağışa dayalı koşullarda kışlık olarak yapılmakta, zaman zaman marjinal alanlara bile buğday ekilmektedir. Artan nüfusun gıda güvenliği ve sürdürülebilir buğday üretimi yönünden kaynakların doğru bir şekilde kullanılması büyük önem taşımaktadır. Buğday tarımı yönünden geliştirilecek ve uygulanacak üretim sistemleri, buğday üretim miktarını riske sokmamalı, olabildiğince artırmaya yönelik ve sürdürülebilir olmalıdır. Yüksek girdili üretim sistemlerinin çevre ve insan sağlığı üzerindeki olumsuz etkileri ve artan girdi fiyatları, doğal denge dostu alternatif üretim sistemlerine karşı olan ilgiyi giderek artırmaktadır. Bu gelişmelere bağlı olarak azaltılmış toprak işleme, sıfır sürüm, sürdürülebilir tarım, organik tarım gibi uygulamalar yanında eski kültürel uygulamalardan birisi olan karışık ekim güncel hale gelmiş, üzerinde yoğun çalışmalar yapılmaya başlanmıştır.

Karışık ekim aynı arazide iki veya daha fazla bitki türünün birlikte yetiştirilmesidir. Bu bitki türlerinin aynı anda ekilmesi ve hasat edilmesi zorunlu değildir. Karışık ekim toprak verimliliği, kaynak kullanım etkinliği ve biyoçeşitliliğin artırılması gibi önemli yararlar sağlamaktadır. Karışıma giren türlerin oranları, ekim ve olgunlaşma zamanları, türleri, yetiştirme koşulları, sosyoekonomik koşullar gibi çeşitli faktörler karışık ekimin sağlayacağı yararlar üzerinde etkili olabilmektedir (Aziz ve ark. 2015). Tarımı yapılan bitki türlerinde çeşitliliğin artırılması ve özellikle baklagillerin üretim sistemlerine sokulmasıyla, tarımsal sorunlarla mücadele yönünden büyük yararlar sağlanacağı düşünülmektedir. Örneğin bitkisel üretimde azot

ihtiyacının baklagillerden karşılanması, endüstriyel olarak üretilen gübre azotundan karşılanmasına kıyasla çok daha sürdürülebilir bir durumdur (Crews ve Peoples, 2004). Baklagiller, azot fiksasyon yetenekleri sayesinde toprağın besin elementleri ve canlıları üzerinde olumlu etki yapmakta, nem kullanımını optimize etmekte, bazı zararlı ve hastalıkları azaltmakta, dolayısıyla tarımsal kimyasal kullanımını azaltmaktadır. Etkin ve verimli tarımsal üretim açısından karışık ekim uygulamalarının yararlı olduğu belirtilmektedir (Toker ve ark. 2024).

Karışık ekim ülkemizde tarla tarımında daha çok yem bitkileri ve hayvan yemi üretimi açısından ele alınmıştır. Ülkemiz tarımında temel bitki niteliğinde olan buğday, karışık ekime çok uygun olmasına rağmen buğday esaslı karışık ekim konusu üzerinde pek durulmamıştır. Buğday üretimi tamamen mono kültür buğday tarımına dayanmaktadır. Oysa sürdürülebilir buğday tarımı yönünden tarımsal çeşitliliğin artırılması, dolayısıyla buğday esaslı karışık ekim büyük önem taşımaktadır (Costanzo ve Barberi, 2014). Bu tebliğde, sürdürülebilir buğday tarımı yönünden buğday esaslı karışık ekimin önemi açıklanmaya çalışılmıştır.

BUĞDAY ESASLI KARIŞIK EKİM KAVRAMI

Karışık ekim veya birlikte yetiştirme kısaca, iki veya daha fazla bitki türünün birlikte yetiştirilmesi şeklinde tanımlanabilir. Farklı şekillerde sınıflandırmaya açık olan karışık ekim bir çalışmada; karışıma giren türlerin karışık olarak ekimi, alternatif sıralar halinde ekimi, aynı sırada karışık ekimi, gelişmekte olan tür içerisine sonradan ekim şeklinde 4 temel gruba ayrılmıştır (Andrews ve Kassam, 1976).

Karışık ekim sisteminde, iki veya daha fazla bitki türü bulunmakla beraber, karışıma giren türlerden birisi üretim amacı yönünden temel bitki niteliği taşımaktadır. Üretim amacı, karışıma giren türlerin oranı ve şekli üzerinde etkili olmaktadır. Ülkemizde tarla tarımında karışık ekim konusunda yapılan araştırmalar ve uygulamalar, daha çok hayvan yemi üretimini amaçlayan bir anlayışla ele alınmıştır. Bu konuda özellikle arpa ve tritikale gibi tahıl bitkileri ile fiğ gibi baklagil bitkileri karışımı üzerinde durulmuştur. Araştırmalara benzer şekilde üretici düzeyinde de kabul gören karışık ekim yem üretimi amaçlı olmaktadır.

Buğday tarımında asıl amaç tane üretimi olup, mono kültür buğday tarımı yaygın ve geleneksel bir uygulama niteliği kazanmıştır. Buğday, ülkemizde insan beslenmesinde temel gıda kaynağı olması nedeniyle tane amaçlı üretim mutlak surette ön planda tutulmalı, buğday verim ve üretimini artırmaya, korumaya ve sürdürmeye yönelik tarımsal sistemlere yer verilmelidir. Buğday esaslı karışık ekim kavramı, buğdayın tane üretim amacını koruyan ve önceleyen bir anlayış ile diğer bitkilerle karışık yetiştirilmesi prensibine dayanmaktadır. Sonuç olarak buğday esaslı karışık ekim uygulaması, buğdayın tane verimine zarar vermeyecek, toplam buğday üretim miktarını riske sokmayacak şekilde, özellikle baklagiller olmak üzere başka bitki veya bitkilerle birlikte yetiştirilmesi şeklinde açıklanabilir.

BUĞDAY ESASLI KARIŞIK EKİMİN YARARLARI

Ülkemizde yerleşik bir uygulama niteliği kazanmış olan mono kültür buğday tarımı, kullanılan tarımsal kimyasallarla birlikte ekosistem dengesi ve zenginliğini bozmaktadır. Ekosistem dengesindeki bozulma, biyoçeşitlilik ve toprak verimliğindeki azalma, sürdürülebilir buğday verimliliğini giderek daha çok tehdit etmektedir. Küresel iklim değişikliği bu olumsuz etkilerin baskısını daha çok artırmakta (Arora, 2019), ortalama sıcaklıktaki her bir derecelik artışın, buğday veriminde küresel ölçekte % 6 oranında bir azalmaya yol açabileceği hesaplanmaktadır (Asseng ve ark. 2015; Zhao ve ark. 2017).

Karışık ekim özellikle düşük girdili ve küçük ölçekli tarımsal işletmeler yanında organik tarım yönünden çok önemli fırsatlar sunmaktadır. Karışık ekim toprak verimliliğini artırma, yabancı

ot kontrolü, hastalık ve zararlılarda azalma, yatmaya karşı dayanıklılık gibi konular yönünden yararlı olmaktadır. Buğday farklı bitkilerle karışık ekilebilmekle beraber tarla tarımında asıl olarak buğday+baklgil karışımları üzerinde durulmaktadır. Yüksek protein, vitamin ve mineral içeriklerine sahip baklagiller, önemli bir gıda ve yem kaynağı olmalarının yanında toprak yapısı ve verimliliğinin artırılması yönünden çok yararlı etkilere sahiptir.

Buğday+yonca karışımında yalnız ekimlere göre verimin yüksek olduğu (Magid ve ark. 2008), buğday+nohut karışık ekiminin, buğday+kolza karışık ekimine göre buğday verimi ve su kullanım etkinliği yönünden daha yararlı olduğu bildirilmiştir (Mandal, 1991). Buğday+bezelye karışımında azotlu gübreleme yapılmadan maksimum verimliliğin sağlanabildiği (Ghaley ve ark. 2005), nohut, mercimek ve kolza karışımlarında buğdayın verim ve verim unsurlarında iyileşmeler olduğu (Khan ve ark. 2005), buğday+bezelye karışık ekiminin ekonomik yönden kazançlı olduğu (Subedi, 1997), buğday içerisine mercimeğin serpmeye olarak ekilmesi halinde buğdaydan sağlanan ekonomik kazancın arttığı tespit edilmiştir (Sharma ve ark. 2018).

Baklagiller yalın ekimlerine kıyasla buğday ile beraber ekildiklerinde daha fazla atmosferik azot fiksasyonu yapabilmektedir. Buğday, baklagillerle beraber yetiştirildiğinde fikse edilen azottan yararlanmakta, sentetik azot ihtiyacı 10-20 kg/da kadar azalabilmektedir. Buna bağlı olarak, fosil yakıtlara bağlı azotlu gübre kullanımının küresel ölçekte % 5-20 kadar azaltılabileceği hesaplanmaktadır (Jensen ve ark. 2010; Jensen ve ark. 2020). Baklagillerle karışık ekim toprak ve atmosfer azotunun optimum kullanımını sağlamakta, özellikle düşük azot koşullarında buğdayın verim ve protein oranını artırmakta, hastalık, zararlı ve yabancı ot baskısını azaltmak gibi yararlar sağlayabilmektedir (Lithourgidis ve ark. 2011; Pelzer ve ark. 2012; Bedoussac ve ark. 2015). Bir meta-analiz çalışmasında, buğday-bakla karışımında buğdayda sarı pas etkisinin % 34 kadar azaldığı, bakla hastalığı yönünden de benzer azalmanın olduğu vurgulanmıştır (Zhang ve ark. 2019).

Buğdayda tane verimi ile tane protein oranı arasında genellikle ters bir ilişki söz konusudur. Baklagillerle karışık ekimin, bu ters ilişki baskısını azaltabildiği (Vrignon-Brenas ve ark. 2018), buğday+bakla ve buğday+bezelye karışık ekimlerinde buğdayın tane verimi ve protein içeriğinin her ikisinin de arttığı, bakla ve bezelye ile karışık yetiştirilen buğdayın tane verimi ve kalitesinin, mono kültür yetiştirilen buğdaydan daha yüksek olduğu görülmüştür (Gooding ve ark. 2007; Kamalongo ve Cannon, 2020; Dvorak ve ark. 2022; Zhu ve ark. 2022). Buğday+bezelye karışık ekiminde sürdürülebilir tarım yönünden azot kaynağının daha etkin kullanıldığı, buğday tanesindeki azot içeriğinin arttığı belirlenmiştir (Ton ve Jensen, 2023).

Mamine ve Fares (2020), kapsamlı literatür taramasına dayalı makalelerinde, buğday+bezelye karışık ekiminin Avrupa'da yaygınlaşmasının önündeki engelleri teknik, ekonomik ve kamusal engeller olmak üzere üç ana başlık altında toplamıştır. Buğday+bezelye karışık ekiminin agroekolojik ve ekonomik yararlarına dikkat çeken araştırmacılar, bütün paydaşların sistemin benimsenmesi ve yaygınlaşması konusuna gereken önemi vermeleri gerektiği vurgulanmıştır.

Yağışa dayalı koşullarda yapılan buğday tarımında yıllık yağış miktarı ve dağılımı, buğday verimi üzerinde belirleyici etkiye sahiptir. Yağışların ve toprak neminin elverişliliği ve buğday tarafından kullanım etkinliğinin artırılması yönünden karışık ekim önemli yarar sağlamaktadır. Nitekim buğday+bezelye karışımında su kaynağının daha etkin bir şekilde kullanıldığı şeklinde sonuca varılmış ve kuru tarımda buğday verimini artırıcı bir uygulama olarak önerilmiştir (Pankou ve ark. 2021). Şerit usulü karışık ekimle ilgili 150'den fazla çalışma taranmış ve karışık ekimin su kullanım etkinliğini artırabildiği, bitki büyüme ve gelişmesi için toprak nemini optimize edebildiği belirtilmiştir (Yin ve ark. 2020).

Buğday+nohut karışık ekiminde buğday ve nohut verimi, mono kültür yetiştirmedeki verimlerden düşük olmuş, ancak karışık ekimde arazi kullanım etkinliği ve toplam verimlilik daha yüksek olmuş, yabancı ot yoğunluğu azalmış, nohut bitkisinde kök derinliği ve nodül oluşumu artmıştır (Banik ve ark. 2006). Buğday+nohut karışımı özellikle düşük fosfor içeriğine

sahip topraklarda kök bölgesindeki fosforun elverişliliğini, her iki bitki için de artırmıştır (Betencourt ve ark. 2012). Arazi kullanım etkinliği, toprak kalitesi ve ürün kalitesinin artırılması yönünden buğday+bakla karışık ekimi yararlı bir üretim modeli olarak önerilmiştir (Chapagain ve Riseman, 2014).

Baklagillerin organik madde, elverişli N ve P içeriği gibi kök bölgesi toprak özelliklerini iyileştirdiği konusunda genel bir kabul söz konusudur. Bu olumlu etkilerin düzeyi baklagil türlerine, iklim ve toprak koşullarına, yetiştirme tekniklerine göre değişebilmektedir (Stagnari ve ark. 2017). Yalnız ekimlere göre buğday+bezelye karışık ekiminde, toprak altı ve toprak üstü gelişmenin daha sağlıklı olması nedeniyle ışık kullanım etkinliği daha yüksek olmuş (Barillot ve ark. 2014), bakla+buğday karışık ekimi, yalnız ekimlere göre ışık, su ve besin elementi alımı bakımından daha etkili olmuştur (Eskanderi, 2011). Yalın ekimlere göre, tahıl+baklagil karışık ekiminde fosfor alım ve kullanım etkinliğinin önemli oranda arttığı, fosforlu gübreye olan ihtiyacın azaldığı bir meta analiz çalışmasıyla ortaya konulmuştur (Tang ve ark. 2021). Baklagillerle karışık ekimde toprağın fiziksel, kimyasal ve biyolojik özelliklerinin dolayısıyla toprak verimliliğinin, *Rhizobium* bakterileri ve besin elementi elverişliliğinin arttığı bildirilmiştir (Chamkhi ve ark. 2022). Fiğ+tahıl karışık ekimlerinde, tahılların destek sağlayarak fiğın yatmasını engellediği, yatmaya bağlı çürümelerin ve hasat kayıplarının önüne geçildiği, dolayısıyla daha kaliteli ve verimli kaba yem sağlandığı bildirilmektedir (Karadağ ve ark. 2022).

Fransa ve Danimarka'da geleneksel ve organik koşullarda yapılan uzun süreli çalışmalarda, karışık ekimin organik tarımda daha yararlı olduğu ve tane verimini daha çok artırdığı görülmüştür. Karışımdaki tahılların verimliliği, baklagillerin verimliliğinden genellikle daha yüksek olmuş, tahılın tane verimi arttıkça baklagil verimi azalmış, tahıllar karışık ekimden daha fazla yararlanmış, karışık ekimler yalnız ekimlere göre tane verimi yönünden daha üstün bulunmuştur (Bedoussac ve ark. 2015). Farklı 10 ülkede REMIX projesi adıyla yürütülen çalışmalarda, buğday+bakla, buğday+mercimek ve benzeri karışık ekimler çok yönlü olarak incelenmiştir. Karışık ekimlerin ekmeleklik buğdayın protein içeriğini artırdığı, azotlu gübreye ihtiyaç duyulmadığı, yabancı otlara karşı daha iyi rekabet sağlandığı, kaynakların daha etkin kullanıldığı, verim stabilitesi sağlandığı şeklinde sonuçlar alınmış, gelecekteki tarım sistemleri açısından karışık ekimin büyük bir öneme sahip olduğu vurgulanmıştır (Bedoussac ve ark. 2021; Justes, 2024).

Karışık ekim yabancı ot, hastalık ve zararlılara bağlı olumsuz etkileri azaltan etkiye sahiptir (Machado, 2009). Tane baklagiller, özellikle bezelye yabancı otlara karşı oldukça zayıf bir baklagil bitkisidir (McDonald, 2003). Organik baklagil yetiştiriciliğinde, yabancı otlar azot elverişliliğini ve tane verimini önemli ölçüde sınırlayabilmektedir (Corre-Hellou ve Crozat, 2005). Karışık ekimlerde ışık, nem, besin elementi ve toprak kullanımı yönünden yabancı otların fazla fırsat bulamadığı, buna bağlı olarak rekabet yönünden zayıf düştüğü vurgulanmıştır (Bedoussac ve ark. 2015). Baklagiller mono kültür ekimde yabancı ot rekabetine karşı daha zayıf iken, karışık ekimlerde baklagiller üzerindeki yabancı ot baskısı azalmaktadır. Baklagiller biyolojik azot fiksasyonu ile buğdaygillere azot sağlarken, buğdaygiller de baklagiller için Fe ve Zn elverişliliğini artırmaktadır (Chamki ve ark. 2022; Akkaya, 2023). Nohut+buğday karışık ekiminin birim alandan elde edilen verimi ve arazi kullanım etkinliğini artırdığı, yabancı ot gelişmesini baskıladığı bildirilmiştir (Banik ve ark. 2006). Mercimeğin yalın ekimine göre buğdayla karışık ekilmesi halinde yabancı ot gelişmesi yıllara göre değişmek üzere % 68-96 oranında azalma göstermiştir (Carr ve ark. 1995). Buğday+fasulye karışık ekiminde, buğday ve fasulyenin ekim sıklığındaki artışlar, yabancı ot gelişmesinin önemli düzeyde azalmasına yol açmış (Bulson ve ark. 1997), karışık ekimlerde yabancı ot biyoması azalış göstermiştir (Eskandari ve Ghanbari, 2010). Benzer şekilde buğday+bakla karışık ekiminde yabancı ot gelişmesi önemle derecede azalmış, bu azalış, karışık ekimde yabancı otların gelişmek için yeterince kaynak bulamayışına bağlanmıştır (Eskandari, 2011).

Buğday+kolza+bezelye karışık ekimlerinde yabancı ot gelişmesi, mono kültür ekimlere kıyasla azalmıştır (Szumigalski ve Van Acker, 2005).

Karışık ekimlerde, yalnız ekimlere kıyasla hastalık ve zararlı riski azalmaktadır. Buğdayın yalnız ekimine kıyasla yoncayla ayrı sıralar halinde karışık ekilmesi, afidler için parazit olan *Allothrombium ovatum* sayısını önemli düzeyde artırmak suretiyle, afidlere karşı biyolojik mücadele sağlamış ve afit zararını azaltmıştır (Ma ve ark. 2007). Buğday+bezelye karışık ekiminde, bezelye oranı arttıkça buğdayda hastalık oranı azalmıştır (Bulson ve ark. 1997). Buğday+*Medicago lupulina* karışık ekiminde, toprak kökenli patojenlere bağlı göçerten (take-all) hastalığı (*Gaeumannomyces graminis*) azalma göstermiştir (Lennartsson, 1988). Buğday yetiştiriciliğinde zararlılarla mücadelede yaygın olarak insektisit kullanımı söz konusudur. Lopes ve ark. (2016), 50 adet orijinal araştırma sonuçlarını inceleyerek, zararlılarla biyolojik mücadele yönünden buğday esaslı karışık ekimin etkilerini açıklamaya çalışmıştır. Araştırmacılar, buğday esaslı karışık ekimin zararlı oranını önemli düzeyde düşürdüğünü, buğday tarımında insektisit kullanımını azaltma yönünden karışık ekimin yararlı bir üretim modeli olduğunu ifade etmişlerdir.

BUĞDAY ESASLI KARIŞIK EKİME TÜRKİYE AÇISINDAN BAKIŞ

Karışık ekimin yararları başlığı altında verilen araştırma sonuçlarından görüleceği gibi sürdürülebilir tarım yönünden karışık ekim, özellikle baklagiller ile karışık ekim çok yönlü yararlar sağlamaktadır. Konu ülkemiz açısından daha büyük bir öneme ve daha fazla yarar sağlama potansiyeline sahip olup, gerekçeleri aşağıda özetlenmeye çalışılmıştır.

Ülkemiz tarla tarımı içerisinde buğday ve arpa ekim alanları, yaklaşık % 70 gibi çok yüksek bir paya sahiptir. Nadasla birlikte değerlendirildiğinde bu pay % 75'e yaklaşmakta, adeta tarla tarımını temsil etmektedir. Toprak sağlığı ve kalitesi yönünden organik madde içeriği anahtar özellik niteliğindedir. Organik madde içeriği yönünden ülkemiz topraklarının yaklaşık % 75 kadarı fakir, % 90 kadarı ise tarla tarımı yönünden iyi sayılan sınır değer (% 3) altındadır (Ülgen ve Yurtsever, 1974; Eyüboğlu ve ark.,1993; Kalaycı, 1999; Gezgin ve ark., 2002). Ülkemizde buğday ve arpa tarımı çoğunlukla kuru tarıma dayalı olup, verimi belirleyen en önemli faktör yıllık yağış miktarı ve dağılımıdır. Küresel iklim değişikliğine bağlı kuraklık tehdidi, bozuk toprak kalitesiyle birleştiğinde kuru tarım alanlarındaki buğday ve arpa üretimini giderek daha büyük risk altına sokmaktadır. Bu nedenle buğday ve arpa tarımı yapılan toprakların kalitesinin artırılması ve korunması yönünden baklagillerle karışık buğday esaslı karışık ekime ihtiyaç vardır.

Buğday tarımında, verimi artırmak için çoğunlukla azotlu ve fosforlu gübreleme esas alınmakta, özellikle azotlu gübreler en önemli girdilerden birisini oluşturmaktadır. Ekonomik kayıplar yanında sentetik gübrelerin toprak, su ve hava kalitesi, dolayısıyla insan, hayvan ve çevre sağlığı üzerinde olumsuz etkileri söz konusudur. Buğday tarımında sentetik gübreler dışında yabancı ot, hastalık ve zararlılarla mücadelede yoğun tarımsal ilaçlar kullanılmaktadır. Çevre ve insan sağlığı yönünden bu kimyasalların zararlı etkileri gün geçtikçe daha çok gündeme gelmektedir. Buğday tarımında sentetik gübreler ve tarımsal ilaçlarının kullanımını azaltmaya yönelik alternatiflerin başında baklagillerle karışık ekim gelmektedir.

Ülkemizde kaliteli kaba yem açığı nedeniyle hayvan beslemede buğday ve arpa samanı yaygın olarak kullanılmaktadır. Zaman zaman buğday fiyatına yaklaşan saman fiyatları üreticiler için ek bir gelir sağlamaktadır. Ancak buğday samanı düşük kaliteli kaba yem grubuna girmekte, yüksek düzeyde selüloz ve lignin, düşük düzeyde ham protein içermekte, hayvan beslemede sınırlı düzeyde kullanılması önerilmektedir (Ak ve Akbay, 2018). Buğdayın hayvan beslemedeki kullanımını sadece samanla sınırlı olmayıp, ülkemizde meraların yetersizliği nedeniyle, hasat sonrası buğday anızında yaygın bir şekilde otlatma yapılmaktadır. Buğday

tarımında herbisit kullanımı, ekonomik kayıplara ve çevre kirliliğini yol açtığı gibi, buğday dışındaki bitkilerin ölümüne yol açmakta, anızdaki bitki çeşitliliğini yok etmekte, otlayan hayvanlar için kalitesiz bir anız söz konusu olmaktadır. Buğday esaslı baklagillerle karışık ekim daha kaliteli saman üretimi yanında, daha kaliteli bir anız oluşturma potansiyeline sahiptir.

Ülkemizde tarımsal işletmelerin yaklaşık % 80 kadarı, 100 dekardan küçük arazi büyüklüğüne sahip olup (Anonim, 2008), buğday yetiştiriciliğinde işletme büyüklüğü Avrupa Birliği'nde ortalama 174 dekar iken, ülkemizde 61 dekardır (Anonim, 2005). Küçük ölçekli arazilerde tarım yapan buğday üreticileri, artan girdi fiyatlarından daha çok etkilenmekte, ekonomik yönden giderek daha çok zayıflamaktadır. Geleneksel buğday yetiştiriciliği yerine, organik buğday yetiştiriciliği yapılması halinde, gübre ve enerji girdilerinde % 34-54, ilaç kullanımında ise % 97'ye varan azalmalar olmaktadır (Mader ve ark. 2002). Küçük ölçekli buğday üreticilerinin organik buğday tarımına geçerek daha fazla ekonomik kazanç sağlamaları bakımından buğday esaslı baklagillerle karışık ekim önemli bir fırsat sunmaktadır.

Özet olarak, buğday tarımı yapılan topraklarımızda kalite sorunu bulunmakta, sentetik gübre ve kimyasal fiyatları giderek artmakta, çevre ve insan sağlığı üzerinde olumsuz etkiler yapmaktadır. Kaba yem açığı ve meraların yetersizliği nedeniyle buğday samanı ve anızı hayvan beslemede yaygın olarak kullanılmaktadır. Buğday üretimi çoğunlukla küçük ölçekli arazilerde ve yağışa dayalı koşullarda yapılmaktadır. Ülkemiz tarımındaki bu gerçekler ve baklagillerle karışık ekimin sağladığı çok yönlü yararlar, buğday tarımında, yemlik ve yemelik baklagillere yer veren buğday esaslı karışık ekime geçişin, sürdürülebilir buğday tarımı yönünden önemini ortaya koymaktadır.

SONUÇ VE ÖNERİLER

Geleneksel mono kültür buğday tarımı, buna bağlı olarak aşırı toprak işleme ve tarımsal kimyasal kullanımı, biyoçeşitliliği azaltmakta, toprak ve çevre sağlığı üzerinde olumsuz etkiler yapmakta, küresel iklim değişikliğine bağlı kuraklık ile birlikte sürdürülebilir buğday tarımı üzerinde giderek daha çok baskı oluşturmaktadır. Baklagiller toprak sağlığı ve ekosistem servisi üzerinde çok önemli yararlar sağlamakta olup, baklagilleri içeren karışık ekimler geleceğin tarımsal sistemlerinden birisi olarak görülmektedir. Ülkemizde küçük ölçekli arazilerde mono kültür buğday tarımı yerine, buğday+baklagil şeklinde tane üretimini önceleyen buğday esaslı karışık ekimin ikame edilmesi, sürdürülebilir buğday tarımı yönünden önemli yararlar sağlama potansiyeline sahiptir. Böylece toprak kalitesi ve çevre sağlığında iyileşmeler sağlanabilir, kaliteli insan gıdası ve hayvan yemi üretilebilir, gübre ve ilaç giderleri azaltılabilir, ekonomik kazanç artırılabilir. Buğday esaslı karışık ekim, organik tarım ile entegre edilmeye de çok uygun olup, bu durumda beklenen yararların daha da fazla artırılması potansiyeli vardır. Yerleşik ve geleneksel mono kültür buğday tarımından buğday esaslı karışık ekime geçişin önünde ekim, hasat, depolama, pazarlama gibi işlemlere bağlı bir takım zorluklar bulunmakla beraber paydaşların işbirliğiyle bu engellerin aşılması mümkündür.

KAYNAKLAR

Ak, İ., Akbay, K.C., 2018. Buğday samanının yem değeri ve hayvan beslemede kullanımı. TÜRKTOB, Türkiye Tohumcular Birliği Dergisi, 25, 20-22.

Akkaya, A., 2023. Baklagillerin Tarla Tarımındaki Önemi. Stratejik Sektör: Tarım-2. IKSAD Publications, ISBN: 978-625-6404-81-6, s. 5-54.

Andrews, D.J., Kassam, A.H., 1976. The importance of multiple cropping in increasing world food supplies. In: Papendick RI, Sanchez PA, Triplett GB (eds). Multiple Cropping. Madison, WI, USA, pp 1-10.

- Anonim, 2005. Buğday Raporu. Türkiye Ziraat Odaları Birliği, Ankara.
- Anonim, 2008. Tarımsal İşletme Yapı Araştırması 2006. TC Başbakanlık Türkiye İstatistik Kurumu, Haber bülteni, 196.
- Anonim, 2023. Tahıllar ve diğer bitkisel ürünlerin üretim miktarları. <https://data.tuik.gov.tr/Bulten/Index?p=Bitkisel-Uretim-1.Tahmini-2022-45502> (Erişim tarihi, 19.01.2023).
- Anonymous, 2023. Worldwide production of grain in 2021/22. <https://www.statista.com/statistics/263977/world-grain-production-by-type/> (Erişim tarihi, 19.01.2023).
- Arora, N.K., 2019. Impact of climate change on agriculture production and its sustainable solutions. *Environmental Sustainability*, 2, 95–96. <https://doi.org/10.1007/s42398-019-00078-w>.
- Asseng, S., Ewert, F., Martre, P., Rotter, R.P., Lobell, D.B., Cammarano, D., Kimball, B.A., Ottman, M.J., Wall, G., White, J.W., 2015. Rising temperatures reduce global wheat production. *Nat. Clim. Change* 5(2), 143–147. <https://doi.org/10.1038/nclimate2470>.
- Aziz, M., Mahmood, A., Asif, M., Ali, A., 2015. Wheat Based Intercropping. *The Journal of Animal & Plant Sciences*, 25 (4), 896-907.
- Banik, P., Midya, A., Sarkar, B.K., Ghose, S.S., 2006. Wheat and chickpea intercropping systems in an additive series experiment: Advantages and weed smothering. *Eur. J. Agron.* 24(4), 325-332, ISSN 1161-0301, <https://doi.org/10.1016/j.eja.2005.10.010>.
- Barillot, R., Escobar-Gutiérrez, A.J., Fournier, C., Huynh, P., Combes, D., 2014. Assessing the effects of architectural variations on light partitioning within virtual wheat-pea mixtures. *Ann Bot.* 114(4), 725-737.
- Bedoussac, L., Journet, E.-P., Hauggard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E.S., Prieur, L., Justes, E., 2015. Ecological principles underlying the increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agronomy for Sustainable Development*, 35, 911–935.
- Bedoussac, L., Albouy, L., Deschamps, E., Salembier, C., Jeuffroy, M.H., 2021. From Theory to Practice of Species Mixtures. *Redesigning European cropping systems based on species MIXtures*.
- Betencourt, E., Duputel, M., Colomb, B., Desclaux, D., Hinsinger, P., 2012. Intercropping promotes the ability of durum wheat and chickpea to increase rhizosphere phosphorus availability in a low P soil, *Soil Biology and Biochemistry*, 46, 181-190, ISSN 0038-0717, <https://doi.org/10.1016/j.soilbio.2011.11.015>.
- Bulson, H.A.J., Snaydon, R.W., Stopes, C.E., 1997. Effect of plant density on intercropped wheat and field beans in an organic farming system. *J. Agric. Sci. Cambridge*, 128,59-71.
- Carr, P.M., Gardner, J.C., Schatz, B.G., Zwinger, S.W., Guldan, S.J., 1995. Grain yield and weed biomass of a wheat-lentil intercrop. *Agron J.* 87, 574-57.
- Chamkhi, I., Cheto, S., Geistlinger, J., Zeroual, Y., Kouisni, L., Bargaz, A., Ghoulam, C., 2022. Legume-based intercropping systems promote beneficial rhizobacterial community and crop yield under stressing conditions. *Industrial Crops & Products*, 183, 114958, <https://doi.org/10.1016/j.indcrop.2022.114958>.
- Chapagain, T., Riseman, A., 2014. Intercropping Wheat and Beans: Effects on agronomic performance and land productivity. *Crop Sci.* 54(5), 2285-2293, <https://doi.org/10.2135/cropsci2013.12.0834>.

- Corre-Hellou, G., Crozat, Y., 2005. N₂ fixation and N supply in organic pea (*Pisum sativum* L.) cropping systems as affected by weeds and pea weevil (*Sitona lineatus* L.). *Eur. J. Agron.* 22, 449-458, <https://doi.org/10.1016/j.eja.2004.05.005>.
- Costanzo, A., Barberi, P., 2014. Functional agrobiodiversity and agroecosystem services in sustainable wheat production. *Agronomy for Sustainable Development*, 34(2), 327-348. [10.1007/s13593-013-0178-1](https://doi.org/10.1007/s13593-013-0178-1).
- Crews, T.E., Peoples, M.B., 2004. Legume versus fertilizer sources of nitrogen: ecological tradeoffs and human needs. *Agric. Ecosys. Environ.*, 102, 279-297, <https://doi.org/10.1016/j.agee.2003.09.018>.
- Dvorak, P., Capouchova, I., Kral, M., Konvalina, P., Janovska, D., Satransky, M., 2022. Grain yield and quality of wheat in wheat-legumes intercropping under organic and conventional growing systems. *Plant Soil Environ.*, 68, 553–559.
- Eskandari, H., 2011. Intercropping of wheat (*Triticum aestivum* L.) and bean (*Vicia faba*): Effects of complementarity and competition of intercrop components in resource consumption on dry matter production and weed growth. *African J. Biotech.* 10:17755-17762.
- Eskandari, H., Ghanbari, A., 2010. Effect of different planting pattern of wheat (*Triticum aestivum* L.) and bean (*Vicia faba*) on grain yield, dry matter production and weed biomass. *Notulae Scientia Biologicae*, 2, 111-115.
- Eyüboğlu, F., Candaş, İ., Örs, G., 1993. Konya ili topraklarının özellikleri ve gübre ihtiyacı. 1. Konya’da Hububat Tarımının Sorunları ve Çözüm Yolları Sempozyumu, 33-49, 12-14, Mayıs, Konya.
- Gezgin, S., Dursun, N., Hamurcu, M., Harmankaya, M., Önder, M., Sade, B., Topal, A., Soylu, S., Akgün, N., Yorgancılar, M., Ceyhan, E., Çiftçi, N., Acar, B., Gültekin, İ., Işık, Y., Şeker, C., Babaoğlu, M., 2002. Determination of B contents of soils of Central Anatolian cultivated lands and its relations between soil and water characteristics, Boron in Plant and Animal Nutrition (Eds. Goldbach et al.) Kluwer Academic/Plenum Publishers, New York, pp. 391-400.
- Ghaley, B.B., Hauggaard-Nielsen, H., Høgh-Jensen, H., Jensen, E.S., 2005. Intercropping of wheat and pea as influenced by nitrogen fertilization. *Nutr. cycl. Agroecosyst.* 73,201-212.
- Gooding M.J., Kasyanova E., Ruske R., Hauggard-Nielsen H., Jensen E.S., Dahlmann C., von Fragstein P., Dibet A., Corre-Hellou G., Crozat Y., Pristeri A., Romeo M., Monti M., Launay M., 2007. Intercropping with pulses to concentrate nitrogen and sulphur in wheat. *Journal of Agricultural Science*, 145, 469–479.
- Jensen, E. S., Peoples, M. B., Hauggaard-Nielsen, H., 2010. Faba bean in cropping systems. *Field Crops Res.* 115(3), 203–216. <https://doi.org/10.1016/j.fcr.2009.10.008>.
- Jensen, E. S., Carlsson, G., Hauggaard-Nielsen, H., 2020. Intercropping of grain legumes and cereals improves the use of soil N resources and reduces the requirement for synthetic fertilizer N: A global-scale analysis. *Agronomy for Sustainable Development*, 40(1), 1–9. <https://doi.org/10.1007/s13593-020-0607-x>.
- Justes, E., 2024. Final report of ReMIX Project. <https://www.eufarmbook.eu/en/contributions/66291cea5fa87c7022570a2a> (Erişim tarihi, 11.02.2025).
- Kalaycı, M., 1999. Yetiştirme tekniği açısından Türkiye buğday tarımının dünü, bugünü, yarını. Orta Anadolu’da Hububat Tarımının Sorunları ve Çözüm Yolları Sempozyumu, 14-25, 8-11 Haziran, Konya.

- Kamalongo D.M.A., Cannon N.D., 2020. Advantages of bi-cropping field beans (*Vicia faba*) and wheat (*Triticum aestivum*) on cereal forage yield and quality. *Biological Agriculture and Horticulture*, 36, 213–229.
- Karadağ, Y., Özkurt, M., Tufan, Y., 2022. Stratejik Sektör Tarım: Fiğ Yetiştiriciliği ve Tarımı. 407-428. IKSAD Yayınları, ISBN: 978-625-8405-49-1, Ankara, Türkiye. <https://iksadyayinevi.com/wp-content/uploads/2022/03/Stratejik-Sektor-TARIM.pdf>.
- Khan, M., Khan, R.U., Wahab, A., Rashid, A., 2005. Yield and yield components of wheat as influenced by intercropping of chickpea, lentil and rapeseed in different proportions. *Pakistan J. Agric. Sci.* 42, 1-3.
- Lennartsson, M., 1988. Take-all disease of wheat. *Proceedings 6th International IFOAM Scientific Conference*, (eds. Allen, P. and Van Dusen, D.), pp 575-580.
- Lithourgidis, A.S., Vlachostergios, D.N., Dordas, C.A., Damalas, C.A., 2011. Dry matter yield, nitrogen content and competition in pea-cereal intercropping systems. *Eur. J. Agron.* 34, 287–294.
- Lopes, T., Hatt, S., Xu, Q., Chen, J., Liu, Y., Francis, F., 2016. Wheat (*Triticum aestivum* L.)-based intercropping systems for biological pest control. *Pest Management Science*, 72, 2193-2202, <https://doi.org/10.1002/ps.4332>.
- Ma, K.Z., Hao, S.G., Zhao, H.Y., Kang, L., 2007. Strip cropping wheat and alfalfa to improve the biological control of wheat aphid *Macrosiphum avenae* by the mite *Allothrombium ovatum*. *Agric. Ecosys. Environ.* 119:49-5.
- Machado, S., 2009. Does intercropping has role in modern agriculture? *J. Soil Wat. Conserv.* 64, 55-57. DOI: 10.2489/jswc.64.2.55A.
- Mader, P., Fliessbach, A., Dubois, D., Gunst, L., Fried, P., Niggli, U., 2002. Soil fertility and biodiversity in organic farming. *Science*, 296, 1694-1697.
- Magid, H.M.A., Ghoneim, M.F., Rabie, R.K., Sabrah, R.E., 2008. Productivity of wheat and alfalfa under intercropping. *Exp. Agric.* 27, 391-395.
- Mamine, F., Fares, M., 2020. Barriers and levers to developing wheat–pea intercropping in Europe: A Review. *Sustainability*, 12, 6962; doi:10.3390/su12176962.
- Mandal, B.K., Dasgupta, S., Roy, P.K., 1991. Effect of intercropping on yield components of wheat, chickpea and mustard under different moisture regimes. *Field Crop Abst.* 39, 7025.
- Mcdonald, G.K., 2003. Competitiveness against grass weeds in field pea genotypes. *Weed Res* 43:48-58. <https://doi.org/10.1046/j.1365-3180.2003.00316.x>.
- Pankou, C., Lithourgidis, A., Dordas, C., 2021. Effect of irrigation on intercropping systems of wheat (*Triticum aestivum* L.) with pea (*Pisum sativum* L.). *Agronomy*, 11, 283. <https://doi.org/10.3390/agronomy11020283>.
- Pelzer, E., Bazot, M., Makowski, D., Corre-Hellou, G., Naudin, C., Al Rifai, M., Baranger, E., Bedoussac, L., Biarnes, V., Boucheny, P., Carrouée, B., Dorvillez, D., Foissy, D., Gaillard, B., Guichard, L., Mansard, M.C., Omon, B., Prieur, L., Yvergniaux, M., Justes, E., Jeuffroy, M.H., 2012. Pea-wheat intercrops in low input conditions combine high economic performances and low environmental impacts. *Eur. J. Agron.* 40, 39–53.
- Sharma, K.C., Parmar, P.S., Solanki, K.S., Singh, A.K., Saiprasad, S.V., 2018. Inter/mixed cropping of lentil (*Lens culinaris*) in late sown wheat (*Triticum aestivum* L.) for higher productivity and profitability of wheat in vertisols of Central India. *International Journal of Agricultural Sciences*, 14(1), 21-26.

- Stagnari, F., Maggio, A., Galieni, A., Pisante, M., 2017. Multiple benefits of legumes for agriculture sustainability: an overview. *Biol. Technol. Agric.* 4(2), 1-13.
- Subedi, K.D., 1997. Wheat intercropped with tori (*Brassica Compestris* var. *toria*) and Pea (*Pisum sativum*) in subsistence farming system of Nepalese hills. *J. Agric. Sci. Cambridge*, 128, 283-289.
- Szumigalski, A., Van Acker, R., 2005. Weed suppression and crop production in annual intercrops. *Weed Sci.* 53, 813-825.
- Tang, X., Bernard, L., Brauman, A., Daufresne, T., Deleporte, P., Desclaux, D., Souche, G., Placella, S.A., Hinsinger, P., 2014. Increase in microbial biomass and phosphorus availability in the rhizosphere of intercropped cereal and legumes under field conditions. *Soil Biol. Biochem.* 75, 86–93.
- Toker, P., Canci, H., Turhan, I., Isci, A., Scherzinger, M., Kordrostami, M., Yol, E., 2024. The advantages of intercropping to improve productivity in food and forage production – a review, *Plant Production Science*, 27(3), 155-169, DOI: 10.1080/1343943X.2024.2372878.
- Ton, A., Jensen, E.S., 2023. The effects of different seeding rates on nitrogen acquisition in pea–wheat intercropping, *Turk J Agric For* 47, 948-959, doi:10.55730/1300-011X.3140.
- Ülgen, N., Yurtsever, N., 1974. Türkiye gübre ve gübreleme rehberi. Toprak ve Gübre Araştırma Enstitüsü Teknik Yayınlar Serisi, 28, Kemal Matbaası, Ankara.
- Vrignon-Brenas, S., Celette, F., Piquet-Pissaloux, A., Corre-Hellou, G., David, C., 2018. Intercropping strategies of white clover with organic wheat to improve the trade-off between wheat yield, protein content and the provision of ecological services by White clover. *Field Crop Res.* 224, 160–169.
- Yin, W., Chai, Q., Zhao, C., Yu, A., Fan, Z., Hu, F., Fan, H., Guo, Y., Coulter, J.A., 2020. Water utilization in intercropping: a review. *Agric. Water Manag.* 241, 106335 <https://doi.org/10.1016/j.agwat.2020.106335>.
- Zhang, C., Dong, Y., Tang, L., Zheng, Y., Makowski, D., Yu, Y., Zhang, F., van der Werf, W., 2019. Intercropping cereals with faba bean reduces plant disease incidence regardless of fertilizer input; a meta-analysis. *Eur. J. Plant Pathol.* 154(4), 931–942. <https://doi.org/10.1007/s10658-019-01711-4>.
- Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D.B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P., 2017. Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl. Acad. Sci.* 114 (35), 9326–9331. <https://doi.org/10.1073/pnas.1701762114>.
- Zhu, Y., He, J., Yu, Z., Zhou, D., Li, H., Wu, X., Dong, Y., Tang, L., Zheng, Y., Xiao, J., 2022. Wheat and faba bean intercropping together with nitrogen modulation is a good option for balancing the trade-off relationship between grain yield and quality in the Southwest of China. *Agronomy*, 12, 2984. <https://doi.org/10.3390/agronomy12122984>.

USE OF MICROBIAL AGENTS AGAINST SOIL SALINITY STRESS IN VEGETABLES

Dr. Öğr. Üyesi Necibe KAYAK

Sakarya Uygulamalı Bilimler Üniversitesi, Ziraat Fakültesi, Bahçe Bitkileri Bölümü,
Sakarya/Türkiye

ORCID ID: <https://orcid.org/0000-0001-7104-8544>

Prof. Dr. Önder TÜRKMEN

Selçuk Üniversitesi, Ziraat Fakültesi, Bahçe Bitkileri Bölümü, Konya/Türkiye

ORCID ID: <https://orcid.org/0000-0003-3218-6551>

ÖZET

Tuzluluk stresi, tarımsal üretim yapılan alanlarda önemli bir abiyotik stres faktörüdür. Toprak tuzluluğu, bitki büyümesini ve verimi olumsuz etkileyerek tarımsal üretimde ciddi kayıplara yol açmaktadır. Sebzeler genellikle toprak tuzluluğuna karşı hassastır. Tuzlu topraklar, tohumların su alımını engelleyerek çimlenme oranlarını düşürür ve düzensiz bitki gelişimine neden olur. Tuz stresi, hücrel su dengesini bozarak kök ve gövde büyümesini sınırlar, yaprak alanını küçültür ve fotosentez kapasitesini düşürerek genel bitki gelişimini yavaşlatır. Ayrıca, çiçeklenme ve meyve tutumu olumsuz etkilenirken, meyve verimi azalmakta ve kalite düşüşü, şekil bozuklukları ile erken olgunlaşma gibi istenmeyen durumlar ortaya çıkmaktadır. Sebze yetiştiriciliğinde toprak tuzluluğunun kontrol altına alınması, sürdürülebilir tarım açısından kritik bir gerekliliktir. Tuzlu toprakları iyileştirmek ve verimlilikleri geri kazanmak amacıyla çeşitli stratejiler değerlendirilmektedir. Bu kapsamda bitki büyümesini teşvik eden rizobakteriler (PGPR), gibi mikrobiyal biyostimülantlar, tuzluluğun bitki büyümesi ve verim üzerindeki etkilerini azaltmak için yararlı araçlar olarak kabul edilmektedir. Mikrobiyal biyostimülantlar bitki kökleriyle simbiyotik ilişkiler kurarak besin alımını artırır ve toprak yapısını iyileştirir. Bu sayede toprak biyolojik aktivitesi desteklenir ve uzun vadeli verimlilik sağlanarak sebze yetiştiriciliğinde sürdürülebilir üretim koşulları oluşturulur. Bu derlemede, toprak tuzluluğuna karşı sebze tarımında bitki direncini artırmada doğal PGPR'ların rolü, mekanizmaları ve tarımsal üretimde sağladıkları faydalar ele alınacaktır.

Anahtar kelimeler: Tuzluluk, Bitki Büyümesini Teşvik Eden Rizobakteri, Tuz Stresi, Sürdürülebilir Tarım

ABSTRACT

One important abiotic stressor in agricultural production regions is salinity stress. Plant growth and yield are adversely affected by excessive salt buildup in the soil, which results in significant losses in agricultural output. Vegetables are generally sensitive to soil salinity. Saline soils prevent seeds from absorbing water, which lowers germination rates and results in uneven seedling growth. By altering the cellular water balance, salt stress limits the growth of roots and stems, shrinks the area of leaves, and lowers photosynthetic ability, all of which slow down the development of the plant as a whole. There is also a negative impact on flowering and fruit set,

which results in a lower fruit production, loss of quality, morphological abnormalities, and early ripening. For sustainable agriculture, managing the salt of the soil is essential while growing vegetables. Many approaches to restoring productivity and improving saline soils are being considered. In this regard, microbial biostimulants such plant growth-promoting rhizobacteria (PGPR) are regarded as useful instruments for reducing the negative effects of salinity on plant growth and yield. By forming symbiotic connections with plant roots, microbial biostimulants improve soil structure and nutrient uptake. For vegetable farming, this promotes soil biological activity, guarantees sustained productivity, and establishes sustainable production conditions. The function of natural PGPRs in improving vegetable plant resistance to soil salinity, their workings, and the advantages they offer for agricultural output will all be covered in this paper.

Key Words: Salinity, Plant Growth-Promoting Rhizobacteria, Salt Stress, Sustainable Agriculture

Toprak Tuzluluğunun Sebepleri ve Mekanizmaları

Abiyotik stres faktörleri, şiddetli rüzgârlar, aşırı sıcaklıklar, toprak tuzluluğu, kuraklık ve sel, tarımsal ürünlerin üretimini ve yetiştiriciliğini etkilemektedir. Bu stres faktörleri arasında toprak tuzluluğu, ekili alan miktarında, ürün veriminde ve kalitesinde büyük düşümlere neden olan en önemli sorunlardan biridir (Shrivastava ve Kumar, 2015; Yadav ve ark., 2011). Toprak tuzluluğu FAO (2024) verilerine göre dünya genelinde 1.381 milyon hektardan fazla bir alana yayılmıştır. Dünyada 230 milyon hektarlık sulanan arazinin %19,5'i (45 milyon hektar) ve 1.500 milyon hektarlık kuru tarım alanının %2,1'i (32 milyon hektar) insan kaynaklı süreçler nedeniyle tuzdan etkilenmiştir (Hussain ve ark., 2010). Dahası, tuzdan etkilenen alanlar her yıl %10 oranında artıyor ve soruna müdahale edilmezse, ekilebilir arazinin %50'sinden fazlası 2050 yılına kadar tuzlanacaktır. Hindistan, Çin, Amerika Birleşik Devletleri, Sudan, Pakistan ve Türkiye topraklarında tuzluluk problemleri dikkat çekici seviyelerdedir. Bununla birlikte, toprak tuzluluğu sorunu başka bölgelerde de giderek yayılmaktadır (Singh, 2018, 2022).

Toprak tuzluluğu, doğal veya insan kaynaklı faktörlerin etkileşimi sonucunda ortaya çıkan bir sorundur (Gül, 2022). Bu durum, özellikle kurak ve yarı kurak bölgelerde tarımsal üretim üzerinde olumsuz etkiler yaratmaktadır. Toprak tuzluluğunun başlıca nedenleri arasında iklim koşulları, sulama yöntemleri, toprak özellikleri ve insan faaliyetleri bulunmaktadır. İnsan faaliyetleri, özellikle yanlış sulama yöntemleri, aşırı gübreleme ve yetersiz drenaj sistemleri nedeniyle toprak tuzluluğunu arttırmaktadır. Bu durum, tarımsal üretimi doğrudan olumsuz etkileyen bir faktördür. Bu durum, dünya genelinde toprak bozulmasını kötüleştiren en önemli nedenler arasında yer almaktadır (Taşan ve Demir, 2019). İklim koşulları, toprak tuzluluğunun en önemli belirleyicilerinden biridir. Kurak ve yarı kurak iklimlerde, buharlaşma oranı yağış miktarını aştığında, toprak yüzeyinde tuz birikimi meydana gelir. Bu durum, bitkilerin büyümesini olumsuz yönde etkileyerek verim kaybına yol açar (Taşan ve Demir, 2019). Ayrıca, doğal tuzlu su kaynaklarının varlığı da toprak tuzluluğunu artıran bir faktördür. Bu tür su kaynakları, sulama sırasında topraklara tuz taşınmasına neden olabilir (Çetin ve ark., 2021). Sulama yöntemleri de toprak tuzluluğunu etkileyen önemli bir unsurdur. Yoğun sulama uygulamaları, eğer uygun drenaj sistemleri ile desteklenmezse, toprakta tuz birikimine yol açabilir. Sulama şebekesindeki eksiklikler ve düşük sulama verimliliği, tuzlu suyun toprak yüzeyine çıkmasına ve bu nedenle tuzluluğun artmasına neden olmaktadır. Özellikle ağır toprak bünyeleri, suyun drenajını zorlaştırarak tuz birikimini artırır (Çetin ve ark., 2021).

Toprak tuzluluğu, tarımsal üretimi sınırlayan ve bitki büyümesini olumsuz etkileyen bir stres faktörüdür. Doygun ekstraktının elektriksel iletkenliği (ECe), toprak tuzluluğunu değerlendirmek için yaygın olarak kullanılan bir ölçüttür. US Salinity Laboratory Staff (1954) tarafından belirlenen kriterlere göre, bir toprak örneği ECe değeri 4 dS m⁻¹'nin üzerinde olduğunda tuzlu olarak kabul edilmektedir. Bu eşik değeri, toprakta bulunan çözünmüş tuzların

konsantrasyonunu yansıtır ve bitkilerin su alımını, besin maddelerini ve genel büyüme koşullarını etkileyen önemli bir parametredir. Çoğu sebze türü, 2.5 dS m⁻¹ gibi oldukça düşük bir tuzluluk eşiğine sahiptir (Snapp ve ark., 1991). Bu nedenle, sebzelerin yetiştirilebildiği alanlar, resmi olarak tuzlu olarak sınıflandırılmayan ancak tuz konsantrasyonu sebzelerin büyümesini kısıtlayan toprakları da içerebilir (Zörb ve ark., 2018).

Tuzun toprakta ve bitki dokularında birikimi, bitki bünyesinde sınırlayıcı etkilerin ortaya çıkmasına neden olmaktadır. Toprak çözeltisindeki yüksek tuz konsantrasyonu, osmotik basıncı artırarak ve bitki beslenmesini engelleyerek bitki büyümesini olumsuz etkiler. Toprak çözeltisindeki yüksek tuz konsantrasyonu, suyun toprak çözeltisinde daha fazla çözünmesini engeller, bu da bitkilerin su alabilme yeteneğini azaltır. Bu durum, osmotik basınç nedeniyle suyun kökler tarafından emilmesini zorlaştırır ve tuzluluğun osmotik veya su eksikliği etkisi olarak bilinir. Tuz stresi, iyon toksitesi ve besin dengesizlikleri nedeniyle bitki büyümesini azaltır (Läuchli ve Epstein, 1990). Tuzluluk, yaprak büyümesinin azalması ve fotosentezi engellemesi yoluyla bitkinin toplam fotosentez kapasitesini düşürür ve büyüme yeteneğini sınırlar (Yeo, 1983).

Toprak tuzluluğu, kök bölgesinden tuzun yıkanması, tarımsal yönetim uygulamalarının değiştirilmesi ve tuza toleranslı bitkilerin kullanılması ile kontrol altına alınabilir (Shrivastava ve Kumar, 2015). Bu kapsamda drenaj yönetimi, sulama uygulamaları ve bitki seçimi gibi çeşitli stratejiler öne çıkmaktadır. Sulu tarım yapılan alanlarda damla sulama gibi daha verimli sulama uygulamalarının benimsenmesi ayrıca drenaj sistemlerinin etkin kullanımı, toprakta biriken tuzların uzaklaştırılmasına yardımcı olarak, bitki köklerinin sağlıklı bir şekilde gelişmesini destekleyecektir. Kuru tarım yapılan alanlarda derin köklü çok yıllık bitkilerin kullanılması ile yağış ve su kullanımı arasındaki denge yeniden sağlanarak yer altı su seviyesinin yükselmesi ve tuzun toprak yüzeyine taşınması önlenir (Manchanda ve Garg, 2008). Bu sürdürülebilir yönetim yaklaşımları tuzluluk stresine bağlı verim kaybını azaltabilir de, uygulama genellikle maliyet ve kaliteli su kaynaklarının yetersizliği nedeniyle sınırlıdır. Düşük maliyetli, uygulanması kolay ve etkili abiyotik stres yönetimi yöntemlerinin geliştirilmesi büyük bir zorluktur. Dünya genelinde, tuz ve kuraklığa toleranslı çeşitlerin geliştirilmesi, ekim takvimlerinin değiştirilmesi ve kaynak yönetimi uygulamaları gibi stratejilerle abiyotik stresle başa çıkmak için kapsamlı araştırmalar yürütülmektedir. Bitki büyümesi üzerinde yüksek tuzluluğun neden olduğu toksik etkileri azaltmak için bitki genetik mühendisliği (Wangxia Wang ve ark., 2003) ve son zamanlarda bitki büyümesini teşvik eden bakterilerin (PGPR) kullanımı (Dimkpa ve ark., 2009) dahil olmak üzere çeşitli stratejiler geliştirilmiştir. Bu yararlı mikroorganizmalar, bitkilerin rizosferini veya endorhizosferini kolonize ederek çeşitli doğrudan ve dolaylı mekanizmalar aracılığıyla bitki büyümesini teşvik eder (Nia ve ark., 2012; Ramadoss ve ark., 2013). Önceki çalışmalar, PGPR kullanımının tuzluluğun neden olduğu bitki stresini hafifletmek için umut verici bir alternatif haline geldiğini (Yao ve ark., 2010) ve biyotik ile abiyotik streslerin yönetiminde mikroorganizmaların rolünün giderek daha fazla önem kazandığını göstermektedir. PGPR'nin abiyotik streslere karşı tolerans oluşturma potansiyeli son yıllarda kapsamlı bir şekilde incelenmiştir (Dodd ve Pérez-Alfocea, 2012).

Bu nedenle, bu incelemenin amacı toprakta biriken tuzun bitki büyümesi ve gelişimi üzerindeki olumsuz etkilerini ele almak, tarımsal üretimi sürdürülebilir kılmak için uygulanabilecek yönetim stratejilerini değerlendirmek ve tuzluluğun neden olduğu stresin bitkiler üzerindeki zararlı etkilerini azaltmaya yönelik yöntemleri, özellikle bitki büyümesini teşvik eden bakterilerin (PGPR) kullanımını da içerecek şekilde araştırmaktır.

Mikrobiyal Ajanların Tuzluluk Stresine Karşı Rolü

Mikroorganizmaların tuzlu topraklarda yaşamak için özel adaptasyon stratejileri vardır (Otlewska ve ark., 2020). Çeşitli çalışmalar, PGPR'lerin antioksidanlar, ACC deaminaz, indol-3-asetik asit (IAA), hücre dışı polimerik madde (EPS) ve uçucu organik bileşiklerin üretimi yoluyla bitkilerde tuz toleransını artırabileceğini ve besin maddelerinin emilimini iyileştirebileceğini ortaya koymuştur.

Tuzluluk Stresine Karşı Bitki Büyümesini Teşvik Edici Rizobakteriler (PGPR)

Toprak, bitki ve mikroorganizmalar arasındaki etkileşim, bitkilerin tüm yaşam döngüsü boyunca devam ederek bitki gelişimini teşvik eder, konukçu bitkide patojenlere karşı sistemik direnç oluşturur ve tuzluluk stresini hafifletir (Ha-Tran ve ark., 2021; Ruiu, 2020). Bitki büyümesini teşvik edici bakteriler (PGPR), bitki büyümesini teşvik etmek, besin maddesi alımında ve abiyotik stresleri tolere edebilmek için yıllardır kullanılmaktadır.

Bitki büyümesini teşvik eden rizobakteriler (PGPR), bitki kök bölgesi olan rizosferde bulunan bir bakteri grubudur. Bu terim bitki büyümesini arttıran bitki köklerini kolonize eden bakterileri ifade eder (Vejan ve ark., 2016). Rizosfer, bitki köklerinin bulunduğu toprak ortamıdır ve en yüksek mikrobiyal aktivitenin gerçekleştiği bölgedir. Bu bölgede, bitkiler için gerekli olan makro ve mikro besin elementleri, çeşitli rizobakteriler yoğun bir şekilde bitki tarafından alınabilir formlara dönüştürürler. Bunun yanı sıra, PGPR, yalnızca besin alımını iyileştirmekle kalmaz, aynı zamanda bitkilerin abiyotik streslere, özellikle de tuzluluk stresine karşı toleransını artırmada önemli roller üstlenir. PGPR'nin bitki tuzluluk toleransını artırmadaki rolü göz önüne alındığında, bu bakteriler yüksek tuz konsantrasyonlarının bitkiler üzerindeki morfolojik, fizyolojik ve biyokimyasal süreçlere olan toksik etkilerini azaltmada faydalı özellikler sergileyerek verim kaybının önemli ölçüde önlenmesini sağlar. Son on yılda *Rhizobium*, *Bacillus*, *Pseudomonas*, *Pantoea*, *Paenibacillus*, *Burkholderia*, *Achromobacter*, *Azospirillum*, *Microbacterium*, *Methylobacterium*, *Variovorax*, *Enterobacter* vb. dahil olmak üzere farklı cinslere ait bakterilerin farklı abiyotik stres ortamlarında konuk bitkilere tolerans sağladığı bildirilmiştir (Grover ve ark., 2011). Örneğin, *Azospirillum* aşılansız marul tohumları, NaCl'ye maruz kaldığında aşılansız tohumlara kıyasla daha iyi çimlenme oranları ve vejetatif büyüme göstermiştir (Barassi ve ark., 2006).

Heidari ve ark. (2011) kuraklık stresi koşulları altında *Pseudomonas* sp. ile aşılansız *Ocimum basilicum*'in bitki büyümesinin yanı sıra oksin ve protein içeriğinin de arttığını bildirmiştir. Golpayegani ve Tilebeni (2011) fesleğenin *Pseudomonas* sp. ve *Bacillus lentus* ile aşılansız tuzluluğun büyüme, fotosentez, mineral içeriği ve antioksidan enzimler üzerindeki etkilerini hafiflettiğini gözlemlemiştir. Dardanelli ve ark. (2008) *Azospirillum brasilense*'nin fasulye fidesi köklerinde kök dallanmasını desteklediğini ve flavonoidler ile lipokito oligosakkaritlerin salgılanmasını artırdığını gözlemlemiştir.

Tuzluluk Stresine Karşı Besin Alımını Artırma (Azot Fiksasyonu, Fosfat Çözücü Bakteriler)

Azot bitki büyümesi ve gelişmesi için oldukça önemli olan bir elementtir. Azot; DNA, RNA ve proteinler için gereklidir. Ayrıca fotosentez için gerekli bir pigment olan klorofilin yapısı için önemli bir bileşendir. Ayrıca ATP ve nükleik asitlerin yapı taşında da azot bulunmaktadır. Dünya atmosferinde bol miktarda azot (N₂) bulunmasına rağmen, bitkiler bu azotun sadece indirgenmiş formlarını kullanabilmektedirler (Smil, 1999; Wagner, 2011). Bitkiler azotu iki formda kullanmaktadırlar: a) ticari gübreler, gübreleme veya organik maddenin minerilizasyonu yoluyla topraktan, b) biyolojik N₂ fiksasyonu (BNF) yoluyla (Smil, 1999; Wagner, 2011).

Kimyasal gübreleme yoluyla verilen azotta bitki tarafından alınmayan azot toprak yüzeyinde çok hızlı kaybolabilir ve yer altı sularına karışabilir. Aşırı N gübrelemesi sonucu içme suyunda

biriken NO₃, bebeklerde ve küçük çocuklarda methemoglobin anemisine neden olmaktadır (Galloway ve ark., 1995; Vance, 2001). Yoğun tarım yapılan arazilerde azot birikmesi küresel bir sorun haline gelmekte ve ekolojik dengeyi olumsuz etkilemektedir. Sebze yetiştiricileri, maksimum verim elde etmek için uzun süredir zirai kimyasallar kullanmaktadır (Guertal 2009). Bununla birlikte, sebze üretiminde yaygın olarak kullanılan gübre kullanımı, çevresel sorunlara ve tüketici sağlığına olumsuz etkilere neden olmaktadır. Toprak mikroorganizmaları, özellikle PGPR, ucuz olmaları ve çevreye dost olmaları nedeniyle sebze yetiştiriciliği uygulamalarında önemli hale gelmektedir. Mekanik olarak, azot fikse edici PGPR, fitohormon olan indol asetik asit veya enzim ACC deaminaz gibi bileşikler üreterek sebzelerin büyümesini ve gelişmesini sağlamaktadır.

Biyolojik azot fikasayonunda atmosferdeki (N₂) azotu fikse eden mikroorganizmalar tarafından bitkilerin yararlanabileceği forma (amonyağa) dönüştürülmektedir (Ahemad ve Kibret, 2014; Kim ve Rees, 1994). Nispeten sınırlı sayıda bakteri türü N₂'yi indirgesede, filogenetik ve fizyolojik olarak çok çeşitli farklı niş türleri bulunmaktadır. Birçok PGPR ve endofitik bakteri diazotrofik olarak gelişebilir ve simbiyotik veya non-simbiyotik olabilir (Siddiqui, 2006).

Rhizobium ve Azotobacter gibi cinslerin, su ve besin alımını artırarak bitkilerin tuz stresini daha iyi tolere etmesini sağladığı kanıtlanmıştır (Shome ve ark., 2022).

Tuzluluk stresi, bitkilerin büyüme ve gelişmeleri üzerinde olumsuz etkilere yol açan önemli bir çevresel faktördür. Ancak bu stresi azaltmak ve bitkilerin besin alımını artırmak için fosfat çözücü bakterilerin (PSB) kullanımı umut verici bir yöntemdir. Fosfat çözücü bakteriler, bitkisel büyüme destekleyici bakteriler (PGPR) arasında önemli bir kategori oluşturmaktadır ve bitkilere besin maddesi sağlamada kritik bir rol oynar (Qingwei ve ark., 2023).

Fosfatın bitkiler tarafından emilebilir forma dönüştürülmesi, PSB'lerin en önemli fonksiyonlarından biridir. Bu süreç, bu bakterilerin ürettiği organik asitler aracılığıyla gerçekleşir. Örneğin, birçok PSB, inorganik fosfatları çözüdüren asetik, sitrik, gluconik, malik ve oksalik gibi farklı organik asitleri üretmektedir (Suleman ve ark., 2018). Bu asitlerin üretimi, bakterilerin toprağa olan etkisini artırarak fosfatın biyoyararlılığını yükseltir. Özellikle *Bacillus* ve *Pseudomonas* cinsinden bakteriler, tuz stresine karşı dayanıklılıklarını artırırken aynı zamanda fosfat çözünürlüğünü de artırabilmektedir (Tchakounté ve ark., 2020).

Tuzluluk Stresine Karşı ACC Deaminaz Üretimi

ACC (1-aminosiklopropan-1-karboksilik asit) deaminaz üreten bitki büyümesini teşvik eden kök bakterileri (PGPR), bitki büyümesini ve stres toleransını artırmada önemli bir rol oynar. Bu bakteriler, stres koşulları altında bitki büyümesini engelleyebilen bir hormon olan etilenin öncüsü olan ACC'yi parçalar. ACC deaminaz üreten PGPR, etilen seviyelerini azaltarak tuzluluk, kuraklık ve ısı stresi dahil olmak üzere çeşitli abiyotik stresleri hafifletebilir ve böylece genel bitki sağlığını ve üretkenliğini artırır. Araştırmalar, ACC deaminaz aktivitesine sahip PGPR'nin stres koşulları altında bitki büyüme parametrelerini önemli ölçüde iyileştirebileceğini göstermiştir. Örneğin, yapılan bir çalışmada ACC deaminaz içeren PGPR ile aşılamanın, askorbat peroksidaz (APX), katalaz (CAT) ve süperoksit dismutaz (SOD) gibi reaktif oksijen türleri (ROS) temizleyen enzimlerin aktivitelerini artırarak tuzlu koşullar altında bamyada bitkilerinin büyümesini artırdığını göstermiştir (Habib ve ark., 2016). Barnawal ve ark. (2014), ACC deaminaz aktivitesine sahip *Arthrobacter protophormiae*'nin bezelye bitkilerini tuzluluk stresine karşı koruduğunu bildirmiştir.

Tablo1. PGPR içeren ACC deaminazın bitkiler üzerindeki tuzluluk stresinin etkilerini hafifletmesi

Bitki türleri	Bakteriyel Suş	Kaynaklar
Bezelye (<i>Pisum sativum</i> L.)	<i>Arthrobacter protophormiae</i>	Barnawalve ark. (2014)
Domates (<i>Solanum lycopersicum</i> L.)	<i>Streptomyces</i> türü.	Palaniyandi ve ark. (2014)
Biber (<i>Capsicum annuum</i> L.)	<i>Bacillus</i> türü.	Wenfei Wang ve ark. (2018)
Yeşil fasulye (<i>Vigna radiata</i> L.)	<i>Pantoea</i> türü, <i>Enterococcus</i> türü. <i>Pseudomonas syringae</i> <i>Pseudomonas fluorescens</i> <i>Rhizobium phaseoli</i>	(Ahmad ve ark., 2011); Panwar ve ark. (2016)

Fitohormon Sentezi (IAA, Gibberellin)

PGPR tarafından indol asetik asit, gibberellinler ve tam olarak açıklanmamış bazı bileşenlerin üretimi, kök uzunluğunun, kök yüzey alanının ve kök ucu sayısının artmasına neden olur. Bu durum besin maddesi alımının daha iyi alınmasına yardımcı olur (Egamberdieva ve Kucharova, 2009; Grover ve ark., 2011). Fitohormonlar, bitkilerdeki biyokimyasal, fizyolojik ve morfolojik süreçler üzerinde etki gösteren son derece düşük miktarlarda bulunan organik maddelerdir; sentezleri düzgün bir şekilde düzenlenir. Bitkiler tarafından doğal olarak sentezlenmeyen ancak doğal ve sentetik yollarla ekzojen olarak sentezlenen fitohormonlar bitki büyüme düzenleyicileri olarak bilinir.

Oksinler, özellikle indol asetik asit (IAA), bitki sürgününde üretilir ve bazipetal olarak kök uçlarına taşınır (Martin ve Elliott, 1984), burada düşük konsantrasyonlarda hücre uzamasını artırarak gelişmiş kök büyümesiyle sonuçlanır. Ayrıca, oksin yanıl köklerin başlamasını destekler. Ancak kök uçlarındaki daha yüksek oksin konsantrasyonları kök büyümesi üzerinde engelleyici bir etkiye sahiptir. Bu engelleme, IAA ile etilen öncüsü 1-aminosiklopropan-1-karboksilik asit (ACC) (Glick ve ark., 1998) arasındaki ilişki göz önünde bulundurulduğunda, etilen sentezini teşvik ederek doğrudan veya dolaylı olabilir (Eliasson ve ark., 1989). İlginç bir şekilde, kök bölgesini kolonize eden bakteriler de kök büyümesindeki bu tür değişiklikleri etkileyebilir. *Azotobacter chroococcum* gibi EPS üreten bakterilerle muamele edilen bakla (*Vicia faba* cv. Giza3) tohumlarının bitki dokularında Na⁺ ve Cl⁻ konsantrasyonlarının azaldığı, buna karşılık N, P ve K konsantrasyonlarının arttığı belirlenmiştir (El-Ghany ve ark., 2020).

Tablo2. PGPR içeren bitkilerin tuzluluk stresinin etkilerini hafifletmesi

Bitki türleri	Bakteriyel Suş	Kaynaklar
Bakla (<i>Vicia faba</i> cv. Giza3)	<i>Azotobacter chroococcum</i>	(El-Ghany ve ark., 2020).
Domates (<i>Solanum lycopersicum</i> L.)	<i>Akromobakter piechaudii</i>	(Mayak ve ark., 2004)
Domates (<i>Solanum lycopersicum</i> L.)	<i>Enterobacter cloacae</i> , <i>Pseudomonas putida</i>	(Grichko ve Glick, 2001)
Biber (<i>Capsicum annuum</i>)	<i>Arthrobacter sp.</i> , <i>Bacillus sp.</i>	(Sziderics ve ark., 2007)

Sonuç ve Değerlendirme

Tarımsal üretim sistemleri, küresel ısınma ve değişen çevre koşullarının etkisi altındadır. Bu durum, bitkilerin biyotik ve abiyotik stres faktörlerine karşı dayanıklılığını artıran stratejilerin geliştirilmesini zorunlu kılmıştır. Yapılan araştırmalar, bitki büyümesini teşvik eden rizobakterilerin (PGPR), konukçu bitkilerde biyotik stres faktörlerine karşı direnç sağladığını ve aynı zamanda abiyotik stres koşullarına toleransı artırabildiğini ortaya koymaktadır.

Tuzluluk stresi, dünya genelinde tarımsal üretimde önemli verim kayıplarına yol açan başlıca abiyotik stres faktörlerinden biridir. Bitkilerde tuz toleransını artırmaya yönelik çeşitli ıslah teknikleri geliştirilmiş olsa da, bu yöntemler çoğu zaman yüksek maliyetli olup hedef bitkilere tuz toleransını başarılı bir şekilde aktarmada yetersiz kalabilmektedir. Bitkilerin tuzluluk stresine adaptasyonu, fizyolojik ve metabolik ağları içeren karmaşık mekanizmalarla gerçekleşmektedir.

PGPR, bitkilerde tuz stresinin olumsuz etkilerini azaltmak için çeşitli mekanizmalar kullanarak önemli bir rol oynamaktadır. Dolayısıyla, tuz toleransını ve ürün verimliliğini artırmada mikrobiyal ajanların biyogübre olarak kullanımı, sürdürülebilir tarıma yönelik stratejik bir yaklaşım olarak öne çıkmaktadır. Özellikle bakteriyel ajanların mikrokapsül teknolojisiyle ticarileştirilmesi ve çiftçilere ulaştırılması, tarımsal üretimde verim artışı sağlayarak organik tarım ürünlerinin miktarını ve kalitesini yükseltebilir.

Bununla birlikte, PGPR'nin topraklarda uzun vadeli tuz toleransını nasıl sağladığı ve bu bakterilerin toprak mikroflorası ile karmaşık çevresel faktörlerle nasıl etkileşime girdiği daha ayrıntılı araştırmalarla ortaya konmalıdır. Bu tür çalışmalar, PGPR'nin tarımsal üretimde daha etkin kullanımını sağlamak ve tuzlu topraklarda sürdürülebilir tarım uygulamalarını geliştirmek açısından kritik bir rol oynamaktadır.

Kaynaklar

Ahemad, M., & Kibret, M. (2014). Mechanisms and applications of plant growth promoting rhizobacteria: current perspective. *Journal of King Saud Univ Science*, 26(1), 1-20.

Ahmad, M., Zahir, Z. A., Asghar, H. N., & Asghar, M. (2011). Inducing salt tolerance in mung bean through coinoculation with rhizobia and plant-growth-promoting rhizobacteria containing 1-aminocyclopropane-1-carboxylate deaminase. *Canadian Journal of Microbiology*, 57(7), 578-589.

Barassi, C., Ayrault, G., Creus, C., Sueldo, R., & Sobrero, M. (2006). Seed inoculation with *Azospirillum mitigates* NaCl effects on lettuce. *Scientia Horticulturae*, 109(1), 8-14.

Barnawal, D., Bharti, N., Maji, D., Chanotiya, C. S., & Kalra, A. (2014). ACC deaminase-containing *Arthrobacter protophormiae* induces NaCl stress tolerance through reduced ACC oxidase activity and ethylene production resulting in improved nodulation and mycorrhization in *Pisum sativum*. *Journal of Plant Physiology*, 171(11), 884-894.

Çetin, M., Kaman, H., & Sesveren, S. (2021). Yoğun sulama uygulamalarının taban suyu kalitesi ve derinlikleri üzerine olan zamansal ve mekansal etkilerinin değerlendirilmesi: akarsu sulama birliği sahası örneği. *Tekirdağ Ziraat Fakültesi Dergisi*, 18, 649-659. doi:<https://doi.org/10.33462/jotaf.879491>

Dardanelli, M. S., de Cordoba, F. J. F., Espuny, M. R., Carvajal, M. A. R., Díaz, M. E. S., Serrano, A. M. G., . . . Megías, M. (2008). Effect of *Azospirillum brasilense* coinoculated with *Rhizobium* on *Phaseolus vulgaris* flavonoids and Nod factor production under salt stress. *Soil Biology and Biochemistry*, 40(11), 2713-2721.

- Dimkpa, C., Weinand, T., & Asch, F. (2009). Plant–rhizobacteria interactions alleviate abiotic stress conditions. *Plant, cell & environment*, 32(12), 1682-1694.
- Dodd, I. C., & Pérez-Alfocea, F. (2012). Microbial amelioration of crop salinity stress. *Journal of experimental botany*, 63(9), 3415-3428.
- Egamberdieva, D., & Kucharova, Z. (2009). Selection for root colonizing bacteria stimulating wheat growth in saline soils. *Biol Fert Soil* doi:10.1007/s00374-009-0366-y
- El-Ghany, A., F, M., & Attia, M. (2020). Effect of exopolysaccharide-producing bacteria and melatonin on faba bean production in saline and non-saline soil. *Agronomy*, 10(3), 316.
- Eliasson, L., Bertell, G., & Bolander, E. (1989). Inhibitory Action of Auxin on Root Elongation Not Mediated by Ethylene. *Plant Physiol.*, 91, 310-314.
- FAO. (2024). *Global status of salt-affected soils*: FAO.
- Galloway, J. N., Schlesinger, W. H., Levy, H., Michaels, A., & Schnoor, J. L. (1995). Nitrogen fixation: Anthropogenic enhancement-environmental response. *Global biogeochemical cycles*, 9(2), 235-252.
- Glick, B. R., Penrose, D. M., & Li, J. (1998). A model for the lowering of plant ethylene concentrations by plant growth-promoting bacteria. *Journal of theoretical biology*, 190(1), 63-68.
- Golpayegani, A., & Tilebeni, H. (2011). Effect of biological fertilizers on biochemical and physiological parameters of Basil (*Ocimum basilicum* L.) *Medicine Plant. Am–Eur J Agric. Environ Sci*, 11, 411-416.
- Grichko, V. P., & Glick, B. R. (2001). Amelioration of flooding stress by ACC deaminase-containing plant growth-promoting bacteria. *Plant Physiology and Biochemistry*, 39(1), 11-17.
- Grover, M., Ali, S. Z., Sandhya, V., Rasul, A., & Venkateswarlu, B. (2011). Role of microorganisms in adaptation of agriculture crops to abiotic stresses. *World Journal of Microbiology and Biotechnology*, 27, 1231-1240.
- Gül, V. (2022). *Bitkilerde tuz stresinin etkileri, savunma cevapları ve sinyal iletim yolu* (B. S. Dinler & V. Gül Eds.). Ankara/Türkiye: IKSAD.
- Ha-Tran, D. M., Nguyen, T. T. M., Hung, S.-H., & Huang, C.-C. (2021). Roles of Plant Growth-Promoting Rhizobacteria (PGPR) in Stimulating Salinity Stress Defense in Plants: A Review. *International Journal of Molecular Sciences*, 22, 2-39. doi:10.3390/ijms22063154
- Habib, S. H., Kausar, H., & Saud, H. M. (2016). Plant Growth-Promoting Rhizobacteria Enhance Salinity Stress Tolerance in Okra through ROS-Scavenging Enzymes. *BioMed Research International*. doi:10.1155/2016/6284547
- Heidari, M., Mousavinik, S., & Golpayegani, A. (2011). Plant growth promoting rhizobacteria (PGPR) effect on physiological parameters and mineral uptake in basil (*Ocimum basilicum* L.) under water stress. *J Agr Biol Sci*, 6, 6-11.
- Hussain, N., Sarwar, G., Schmeisky, H., Al-Rawahy, S., & Ahmad, M. (2010). Salinity and Drought Management in Legume Crops. In *Climate Change and Drought Management in Cool Season Grain Legume Crops*
- Kim, J., & Rees, D. C. (1994). Nitrogenase and Biological Nitrogen Fixation. *Perspectives in Biochemistry*, 33(2), 389-397.
- Läuchli, A., & Epstein, E. (1990). Plant responses to saline and sodic conditions. *Agricultural salinity assessment and management*, 71, 113-137.

- Manchanda, G., & Garg, N. (2008). Salinity and its effects on the functional biology of legumes. *Acta Physiologiae Plantarum*, 30, 595-618.
- Martin, H. V., & Elliott, M. C. (1984). Ontogenetic changes in the transport of indol-3-yl-acetic acid into maize roots from the shoot and caryopsis. *Plant physiology*, 74(4), 971-974.
- Mayak, S., Tirosh, T., & Glick, B. R. (2004). Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiology and Biochemistry*, 42(6), 565-572.
- Nia, S. H., Zarea, M. J., Rejali, F., & Varma, A. (2012). Yield and yield components of wheat as affected by salinity and inoculation with *Azospirillum* strains from saline or non-saline soil. *Journal of the Saudi Society of Agricultural Sciences*, 11(2), 113-121.
- Otlewska, A., Migliore, M., Dybka-Stepień, K., Manfredini, A., Struszczyk-Świta, K., Napoli, R., . . . Pinzari, F. (2020). When salt meddles between plant, soil, and microorganisms. *Frontiers in Plant Science*, 11, 553087.
- Palaniyandi, S., Damodharan, K., Yang, S., & Suh, J. (2014). *Streptomyces* sp. strain PGPA39 alleviates salt stress and promotes growth of 'Micro Tom' tomato plants. *Journal of applied microbiology*, 117(3), 766-773.
- Panwar, M., Tewari, R., & Nayyar, H. (2016). Native halo-tolerant plant growth promoting rhizobacteria *Enterococcus* and *Pantoea* sp. improve seed yield of Mungbean (*Vigna radiata* L.) under soil salinity by reducing sodium uptake and stress injury. *Physiology and Molecular Biology of Plants*, 22, 445-459.
- Qingwei, Z., Lushi, T., Yu, Z., Yu, S., Wanting, W., Jiangchuan, W., . . . Bilal, M. (2023). Isolation and characterization of phosphate-solubilizing bacteria from rhizosphere of poplar on road verge and their antagonistic potential against various phytopathogens. *BMC microbiology*, 23(1), 221.
- Ramadoss, D., Lakkineni, V. K., Bose, P., Ali, S., & Annapurna, K. (2013). Mitigation of salt stress in wheat seedlings by halotolerant bacteria isolated from saline habitats. *SpringerPlus*, 2, 1-7.
- Ruii, L. (2020). Plant-growth-promoting bacteria (PGPB) against insects and other agricultural pests. *Agronomy*, 10(6), 861.
- Shome, S., Barman, A., & Solaiman, Z. M. (2022). Rhizobium and phosphate solubilizing bacteria influence the soil nutrient availability, growth, yield, and quality of soybean. *Agriculture*, 12(8), 1136.
- Shrivastava, P., & Kumar, R. (2015). Soil salinity: A serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi Journal of Biological Sciences*, 22, 123-131.
- Siddiqui, Z. A. (2006). *PGPR: biocontrol and biofertilization* (Vol. 313): Springer.
- Singh, A. (2018). Alternative management options for irrigation-induced salinization and waterlogging under different climatic conditions. *Ecological Indicators*, 90, 184-192.
- Singh, A. (2022). Soil salinity: A global threat to sustainable development. *Soil Use and Management*, 38(1), 39-67. doi:<https://doi.org/10.1111/sum.12772>
- Smil, V. (1999). Nitrogen in crop production: An account of global flows. *GLOBAL BIOGEOCHEMICAL CYCLES*, 13(2), 647-662.
- Snapp, S., Shennan, C., & Van Bruggen, A. (1991). Effects of salinity on severity of infection by *Phytophthora parasitica* Dast., ion concentrations and growth of tomato, *Lycopersicon esculentum* Mill. *New Phytologist*, 119(2), 275-284.

- Suleman, M., Yasmin, S., Rasul, M., Yahya, M., Atta, B. M., & Mirza, M. S. (2018). Phosphate solubilizing bacteria with glucose dehydrogenase gene for phosphorus uptake and beneficial effects on wheat. *PloS one*, *13*(9), e0204408.
- Sziderics, A. H., Rasche, F., Trognitz, F., Sessitsch, A., & Wilhelm, E. (2007). Bacterial endophytes contribute to abiotic stress adaptation in pepper plants (*Capsicum annuum* L.). *Canadian Journal of Microbiology*, *53*(11), 1195-1202.
- Taşan, S., & Demir, Y. (2019). Toprakların tuzluluk ve sodikliliğinin alansal ve zamansal değişiminin jeostatistiksel yöntemlerle değerlendirilmesi: bafra ovası örneği. *Anadolu Journal of Agricultural Sciences*, *34*, 336-350. doi:<https://doi.org/10.7161/omuanajas.557601>
- Tchakounté, G. V. T., Berger, B., Patz, S., Becker, M., Fankem, H., Taffouo, V. D., & Ruppel, S. (2020). Selected rhizosphere bacteria help tomato plants cope with combined phosphorus and salt stresses. *Microorganisms*, *8*(11), 1844.
- Vance, C. P. (2001). Symbiotic Nitrogen Fixation and Phosphorus Acquisition. *Plant Nutrition in a World of Declining Renewable Resources. American Society of Plant Biologists*, *127*, 390-397.
- Vejan, P., Abdullah, R., Khadiran, T., Ismail, S., & Nasrulhaq Boyce, A. (2016). Role of plant growth promoting rhizobacteria in agricultural sustainability—a review. *Molecules*, *21*(5), 573.
- Wagner, S. C. (2011). Biological Nitrogen Fixation. *Nature Education*, *3*(10), 15.
- Wang, W., Vinocur, B., & Altman, A. (2003). Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta*, *218*, 1-14.
- Wang, W., Wu, Z., He, Y., Huang, Y., Li, X., & Ye, B.-C. (2018). Plant growth promotion and alleviation of salinity stress in *Capsicum annuum* L. by *Bacillus* isolated from saline soil in Xinjiang. *Ecotoxicology and Environmental Safety*, *164*, 520-529.
- Yadav, S., Irfan, M., Ahmad, A., & Hayat, S. (2011). Causes of salinity and plant manifestations to salt stress: a review. *Journal of environmental biology*, *32*(5), 667.
- Yao, L., Wu, Z., Zheng, Y., Kaleem, I., & Li, C. (2010). Growth promotion and protection against salt stress by *Pseudomonas putida* Rs-198 on cotton. *European Journal of Soil Biology*, *46*(1), 49-54.
- Yeo, A. (1983). Salinity resistance: physiologies and prices. *Physiologia plantarum*, *58*(2).
- Zörb, C., Geilfus, C.-M., & Dietz, K.-J. (2018). Salinity and crop yield. doi:10.1111/plb.12884

DETERMINATION OF EUNIS HABITAT TYPES OF SOME NATURAL PROTECTED AREAS IN ADANA (TÜRKİYE) PROVINCE

Ergün ÖZUSLU

Iğdır University, Postgraduate Education Institute, Agricultural Sciences Department, Iğdır, Türkiye

²Gaziantep Islam Science and Technology University, Faculty of Engineering and Natural Sciences, Gaziantep, Türkiye

Ahmet Zafer TEL

Iğdır University, Faculty of Agriculture, Department of Agricultural Biotechnology, Iğdır, Türkiye

ABSTRACT

Conservation of biodiversity should generally be done by considering the balance of conservation and utilization. One of the main ways of establishing this balance is to know the habitat characteristics. Today, areas where species are densely populated are under threat due to human activities. The European Union has developed the European Nature Information System (EUNIS) to identify and protect habitats. EUNIS Habitat classification aims to provide a common definition and hierarchical classification of all units. In this study, it was aimed to determine the EUNIS habitat types of seven natural protected areas in the Eastern Mediterranean region of Türkiye, namely Agyatan, Tuzla, Yumurtalık and Akyatan Lagoon, Yer Köprü Natural Formation, Şekerpınar and Tatarlı Village Ancient Remains. In this study, the habitat types of Agyatan, Tuzla, Yumurtalık and Akyatan Lagoon, Yer Köprü Natural Formation, Şekerpınar and Tatarlı Village Ancient Remains in the Eastern Mediterranean region of Türkiye were determined using observations and EUNIS Habitat Classification version 2021 and 2022. According to the findings, seven Level 1 and sixteen Level 2 habitat types with EUNIS habitat type codes N12, N14, N1B, MA651, S4, S5, S51, V12, MA5, MA55, C2, C2.44, C3.1, J1.2, R1, R1D3 were identified. These results contribute to the literature on the habitat types of the natural protected areas of Agyatan, Tuzla, Yumurtalık and Akyatan Lagoon, Yer Köprü Natural Formation, Şekerpınar and Tatarlı Village Ancient Remains in the Eastern Mediterranean region of Türkiye.

Keywords: Adana, EUNIS Habitat Type, Türkiye.

INTRODUCTION

Türkiye has a rich plant diversity. This diversity is due to Turkey's geographical location, characteristics of plant geography, climatic and edaphic differences and habitat diversity (Davis, 1965-1985; Davis and Hedge, 1975). This study was carried out to determine the EUNIS habitat types in natural and cultural sites in Adana province, which is part of the Eastern Mediterranean Region of Türkiye. The study covers the protected areas of Agyatan Lagoon, Tuzla Lagoon, Yumurtalık Lagoon, Yer Köprü Natural Formation, Akyatan Lagoon,

Şekerpınar and Tatarlı Village Ancient Ruins (Figure 1). General characteristics and coordinates of the study areas are given in Table 1.



Figure 1: Location of the study areas in Adana (Türkiye)

Adana province is located in the East Mediterranean region in terms of phytogeographically. The research area is in C5 grid-square system according to Davis's grid system. Research areas have a rich and unique place in terms of plant diversity. This is due to the fact that Turkey has three different climates, landforms and rich plant diversity (Davis, 1965-1985; Davis & Hedge, 1975).

The conservation of biological diversity is generally done by considering the balance of conservation and utilisation. One of the main ways of establishing this balance is to know the habitat characteristics. Today, areas where species are densely populated are under threat due to human activities. The European Union has developed the European Nature Information System (EUNIS) to identify and protect habitats. EUNIS Habitat classification aims to provide a common definition and hierarchical classification of all units.

In recent years, habitats are in danger of destruction and extinction due to human activities (Arslan et al., 2012). Therefore, habitat classification is important for the sustainability of natural resources (Moss & Roy, 1998). The European Union has established the European Nature Information System (EUNIS) (Davies et al., 2004; Anonymous, 2025) to utilize existing natural resources better, compare different habitat types, analyze habitats in more detail, create a common classification system, and build a database. Today, species and habitats are constantly threatened by extinction due to intensive use (Arslan et al., 2012). Therefore, the EUNIS classification system records habitat data comparably and provides a reference for the conservation of natural resources. Davies et al. (2004) and Anonymous, (2025) ranked EUNIS habitat types hierarchically. The system is currently organized into 9 main categories and their

subheadings. The EUNIS habitat classification review is on-going. Four groups are pending review: Inland waters, Wetlands, Constructed, industrial and other artificial habitats and Complexes. [M : Marine benthic habitats \(MA-MG\)](#), [MH : Pelagic water column](#), [MJ : Ice-associated marine habitats](#), [N : Coastal habitats](#), [R : Grasslands and lands dominated by forbs, mosses or lichens](#), [S : Heathland, scrub and tundra](#), [T : Forest and other wooded land](#), [U : Inland habitats with no or little soil and mostly with sparse vegetation](#), [V : Vegetated man-made habitats](#) (Anonymous, 2025).

EUNIS habitat types is a module that includes habitats listed in Annex I of the EU Habitats Directive and the Bern Convention. EUNIS habitat types is a program that covers all habitat types, from natural to built, terrestrial to marine, and is a more harmonized data collection of the Pan-European system. Habitat types include plant and animal communities and abiotic factors that characterize living environmental elements. All the factors in the definition form the outline of habitat classification and EUNIS classifications are hierarchically limited to 3 levels (Level 4 in marine habitats). Level 4 and below (Level 5 and below in marine habitats) are in combination with other classification systems and are considered in a common framework.).

Studies on habitat classification in Turkey are few and include Karaömerlioğlu (2007), Karaömerlioğlu & Düzenli (2008), Arslan & Arslantürk (2009), Arslan et al. (2014), Ulu et al. (2014), Mergen & Karacaoğlu (2015), Çiftçi (2015), Erdoğan (2016), Geven et al. (2016), Şahin & Karavelioğlu (2018a), Şahin & Karavelioğlu (2018b), Tug et al. (2018), Seyfe (2019), Özen & Ürker (2020), Çakmak & Aytaç (2020), Çakmak & Aytaç (2021) and Demir et al.

It is necessary to precisely determine the legally binding protected habitat types in EUNIS (Arslan et al. 2012). In Türkiye, although not at the habitat type level, some species or specific plant communities are protected on-site in areas with conservation statuses such as national parks, nature conservation areas, genetic conservation forests, research forests, etc. In this way, one or more habitat types are preserved in these areas (Arslan et al. 2012). Determination of habitat types will contribute to both biodiversity studies and the studies of different disciplines.

As a result of the work carried out on EUNIS in Türkiye between 2011-2020; a total of 140 EUNIS habitat types were identified in the 3rd Level. In addition, it has been determined that there are 26 new habitat types that are not defined in EUNIS, without any level restrictions (Çakmak & Aytaç, 2021). Tak and Tel (2024) identified 37 habitat types belonging to the European Nature Information System (EUNIS) in Akdağ (Malatya).

Within the scope of the National Biodiversity Inventory and Monitoring Project, 10 basic habitat types were identified across the provinces where the project was completed, and a total of 257 habitat records were given from 46 different habitat types across 25 provinces (Terzioğlu et al., 2015).

Table 1: The Study areas and general characteristics

No	Study Area	Field width (m ²)	Coordinate - Altitude
1	Ağyatan Lagoon	6.514.102,68	36° 39' 37. 67" North 35° 17' 06. 25" East, 1 m.
2	Tuzla Lagoon	17.460.549,73	36° 42' 32. 88" North 35° 03' 20. 56" East, 1 m.
3	Yumurталık Lagoon	90.208.440,02	36° 41' 38. 59" North 35° 33' 14. 37" East, 1 m.
4	Akyatan Lagoon	83.796.264,3	36° 38' 23. 81" North 35° 15' 57. 17" East, 1 m.
5	Yer Köprü Natural Formation	20.775,67	37° 40'56. 04" North 35° 29' 34. 83" East, 771 m.
6	Şekerpinar	31.671,49	37° 28'18. 48" North 34° 51' 42. 63" East, 834 m.
7	Tatarlı Village Antique Period Ruins	6.653,85	37° 07'24. 90" North 36° 03' 20. 12" East, 37 m.

Most countries have not yet fully developed the EUNIS habitat classification and generally use their own habitat classification types on a country-by-country basis. Turkey is ahead of other countries in this context and has reached the stage of determining EUNIS habitat types with the National Biodiversity Inventory and Monitoring Project (Terzioğlu et al., 2015). The existence of this situation, the knowledge of Turkey's habitat richness, its healthy use and obtaining EUNIS habitat types for the whole of Turkey will provide more comprehensive information about the habitat and areal data of our country in the future.

There are no previous studies on habitat classification in the research area. In this study, it was aimed to determine the EUNIS habitat types of seven natural protected areas in the Eastern Mediterranean region of Türkiye, namely Ağyatan, Tuzla, Yumurталık and Akyatan Lagoon, Yer Köprü Natural Formation, Şekerpinar and Tatarlı Village Ancient Remains.

CONCEPTUAL FRAMEWORK

General characteristics of the study areas

Ağyatan Lagoon is located in Karataş district of Adana province. Ağyatan Lagoon is a typical lagoon consisting of alluvium formed by delta sediments and dunes with the change of bed of Seyhan river. It is a I. Degree Natural Site. It is also protected as a wetland. Natural landscape elements are the sea, beach, dunes, Dalyan region, and Ağyatan lake (Figure 2). The area has some sand dunes between the lake and the sea, which has a flat land structure in terms of geomorphology (Figure 3; 4; 5). The size of the area is 6.514.102,68 m².




Figure 2: Satellite image of the Agyatan Lagoon protected area



Figure 3: Habitat type view (C3:1 Species-rich helophyte beds)



Figure 4: Habitat type view (N14: Mediterranean, Macaronesian and Black Sea shifting coastal dune)

This lagoon shrinks significantly during the summer, and vast mud flats appear. The lake is connected to the sea by a canal from the southwest. In periods when lake water level are high, water flows from the lake to the sea through the canal and from the sea to the lake in the low periods. With the effect of rainfall and water carried by drainage channels in winter and spring, the lake water becomes sweeter. In summer, salinity increases due to high evaporation and water ingress from the sea to the lake. The average annual rainfall of the research area is 769,9 mm. Average maximum and minimum temperatures respectively are 27,36 °C and 10 °C in August and May.



Figure 5: Habitat type view (C3:1 Species-rich helophyte beds)

Tuzla Lagoon is located in Karataş district of Adana province. This area is the I. Degree Natural Protected Area and Wildlife Development Area. It is a wetland, and the area is under protection. The size of Tuzla Lagoon is 17.460.549,73 m². The area is located in the southern part of the Çukurova Delta, where the Seyhan and Ceyhan rivers flow into the Mediterranean. There is a Salt Lake close to the sea and an inner lake (Figure 6). Tuzla lake is filled with precipitation in winter and dries up in summer. It has a saltwater habitat (Figure 7). The area has a coastline of approximately 15.5 km and generally shows a natural landscape feature. It has a geomorphologically flat land structure. There are dunes between the lake and the sea.




Figure 6: Satellite image of the Tuzla Lagoon protected area



Figure 7: Habitat type view (Tuzla lake)

Yumurtalık Lagoon is located in Karataş and Yumurtalık districts of Adana province. It is a natural site and Turkey's eleventh Ramsar area (Figure 8; 9; 10; 11). It is also a Nature Protection Area, and its size of 90.208.440,02 m².




Figure 8: Satellite image of the Yumurtalık Lagoon protected area



Figure 9: Habitat type view (C2.4: Tidally influenced rivers, upstream currents from river mouths.)



Figure 10: Habitat type view (N1B: Mediterranean and Black Sea coastal dune scrub)



Figure 11: Habitat type view (C3:1 Species-rich helophyte beds)

Akyatan Lagoon is located in Karataş district of Adana province. The area is I. and II. Degree Natural Site Area. The area is also a Wildlife Development Area. Akyatan Lake has a special topography due to its alluvial structures. Akyatan lake is the largest lagoon lake formed by the Seyhan and Ceyhan rivers in the Çukurova delta (Figure 12; 13; 14). The area has a coastline of approximately 20 km.




Figure 12: Satellite image of the Akyatan Lagoon protected area



Figure 13: Habitat type view (C2.4: Tidally influenced rivers, upstream currents from river mouths.)



Figure 14: Habitat type view (C2: Surface streams (macrophytic associations associated with permanent or seasonal streams are the representatives of this habitat type in the area)

Yer Köprü Natural Formation is located in Aladağ district of Adana province. The area is also a Degree I Natural Site Area. The area is also a wetland and is under protection. Yer Köprü Natural Formation covers an area of 20.775,67 m². The area and its surroundings are located in a valley (Figure 16; 17). Altitude of the area is 500 m at the bottom of the valley.




Figure 15: Satellite image of Yer Köprü Natural Formation



Figure 16: Habitat type view (S51: Mediterranean maquis and arborescent matorral)



Figure 17: Habitat type view (S51: Mediterranean maquis and arborescent matorral)

Şekerpınar is located in Pozantı district of Adana province. The area is II. Degree Natural Site Area. It has a size of 31.671,49 m². In addition, the area is a wetland (Figure 18; 19).




Figure 18: Satellite image of the Şekerpınar protected area



Figure 19: Habitat type view (J1.2: Dwellings in peri-urban and rural areas)

Tatarlı Village Antique Period Ruins is located in the Ceyhan district of Adana province. It is a I. Degree Natural and Archaeological Site. It has a size of 6.653,85 m². The area contains tree, shrub, and a grass vegetation layer (Figure 20). This wetland is also a residential area. The area consists of flat plains with an altitude of about 40 meters.



Figure 20: Habitat type view (J1.2: Dwellings in peri-urban and rural areas)

METHODOLOGY

The material of this study consists of seven natural and cultural protected areas in Adana province, located in the Eastern Mediterranean region of Turkey. Habitat types in the study areas were recorded observationally using the EUNIS habitat type hierarchical view (Marine version 2022 and terrestrial version 2021) (Anonymous, 2025).

CONCLUSION AND DISCUSSION

The study conducted by Çakmak & Aytaç (2021) in Türkiye, it was determined that out of the 326 third-level habitat types in the EUNIS habitat classification, 138 are found in Turkey. In this study, 22 habitat types at Level 2 were identified related to 9 habitat types at Level 1 (Çakmak & Aytaç, 2021). Turkey is a country with high biological and habitat diversity (Kanca et al., 2019). Therefore, in the future, data obtained from determining all habitat types in Turkey and establishing geographic information systems can be utilized by various disciplines. Many countries have completed the EUNIS habitat classification and EUNIS habitat types of nearly 30 provinces in Turkey have been determined (Terzioğlu et al., 2015; Kanca et al., 2019). All of these provinces have inland water habitats, grasslands and non-grass herbaceous habitats, forests, agricultural areas and man-made habitats (Terzioğlu et al., 2015). This situation will ensure that Turkey's habitat richness is known, utilized properly and EUNIS habitat types are obtained for the whole of Turkey. When the literature studies are examined, it is seen that marine habitats, marshes and peatland habitats are represented at very low levels, while heathland, scrubland and tundra are highly represented (Terzioğlu et al., 2015; Kanca et al., 2019; Çakmak & Aytaç, 2021). It is thought that this is due to the fact that most of the studies were conducted in terrestrial areas and the data on heathland, shrubland and tundra are high.

In addition, it was determined that the most common EUNIS habitat types at level 3 in Turkey are E1.2 (11 studies), G3.5 (9 studies) and G1.7 (8 studies) (Çakmak & Aytaç, 2021). In this study, only T3A5 Pinus brutia forest habitat was not encountered among the habitat types mentioned above because the study area has maquis and coastal vegetation. The reason for this is thought to be due to the abundance of studies conducted in areas with similar habitat types (Şahin & Karavelioğlu (2018a), Şahin & Karavelioğlu (2018b), Seyfe (2019), Özen & Ürker (2020), Çakmak & Aytaç (2020), Çakmak & Aytaç (2021) and Demir et al. (2022).

According to Terzioğlu et al. (2015), the habitats found in all study provinces in Turkey are C (Inland water habitats), E (Grasslands and non-grass herbaceous habitats), G (Forests), I (Agricultural areas) and J (Man-made habitats). Of these habitats, J (Man-made habitats) and G (Forests) are the most densely populated habitats. In line with the information received from the provinces, all habitat types were identified in Adana province in terms of level 1 habitats, while 6 provinces have the lowest number of level 1 habitat types (Ağrı, Isparta, Rize, Sivas, Tunceli and Karabük). In Ağrı province, the number of 2nd level EUNIS habitat types is quite high and this value is 16. Although the number of habitat classes is generally low, there is an increase in the number of level 2 habitats, probably due to the differentiation caused by habitat fragmentation. Adana and Edirne provinces also have the highest number of level 2 EUNIS habitat types, while this value is the lowest in Tokat and Tunceli provinces (Terzioğlu et al., 2015).

In our study, as a result of the classification of habitat types in the study areas according to EUNIS habitat types (Davies et al., 2004; Anonymous, 2025); it was determined that there were seven Level 1 and sixteen Level 2 habitat types at different levels (Table 2).

According to the findings, seven Level 1 and sixteen Level 2 habitat types with EUNIS habitat type codes N12, N14, N1B, MA651, S4, S5, S51, V12, MA5, MA55, C2, C2.44, C3.1, J1.2, R1, R1D3 were identified.

In this study, as a result of the classification of habitat types in the study areas according to EUNIS habitat types (Davies et al., 2004; Anonymous, 2025); it was determined that there were seven Level 1 and sixteen Level 2 habitat types at different levels (Table 2). When the study areas are evaluated separately; Three Level 1 and eight Level 2 habitat types in Agyatan Lagoon, four Level 1 and nine Level 2 habitat types in Tuzla Lagoon, four Level 1 and nine Level 2 habitat types in Yumurталık Lagoon, five Level 1 and twelve Level 2 habitat types in Akyatan Lagoon, There are 2 Level 1 and 5 Level 2 habitat types in Yer Köprü Natural Formation, two Level 1 and two Level 2 habitat types in Şekerpınar and two Level 1 and two Level 2 habitat types in Tatarlı Village Ancient Ruins (Table 2).

Table 2: EUNIS habitat types of the study areas

Area	EUNIS Classification Code	Habitat Name	EUNIS subclassification code and description
Agyatan Lagoon	M	Marine benthic habitats (MA-MG)	MA5 : Littoral sand, MA55: Mediterranean littoral sand
	N	Coastal habitats	N12: Mediterranean and Black Sea sand beach, N14: Mediterranean, Macaronesian and Black Sea shifting coastal dune, N1B: Mediterranean and Black Sea coastal dune scrub
	C	Inland waters	C2: Surface streams (macrophytic associations associated with permanent or seasonal streams are the representatives of this habitat type in the area). C2.4: Tidally influenced rivers, upstream currents from river mouths. C3.1: Species-rich helophyte beds
Tuzla Lagoon	M	Marine benthic habitats (MA-MG)	MA5 : Littoral sand, MA55: Mediterranean littoral sand
	N	Coastal habitats	N12: Mediterranean and Black Sea sand beach, N14: Mediterranean, Macaronesian and Black Sea shifting coastal dune, N1B: Mediterranean and Black Sea coastal dune scrub
	V	Vegetated man-made habitats	V12: Mixed crops of market gardens and horticulture
	C	Inland waters	C2: Surface streams (macrophytic associations associated with permanent or seasonal streams are the representatives of this habitat type in the area). C2.4: Tidally influenced rivers, upstream currents from river

			mouths. C3.1: Species-rich helophyte beds
Yumurtalık Lagoon	M	Marine benthic habitats (MA-MG)	MA5 : Littoral sand, MA55: Mediterranean littoral sand, MA651: Biocenosis of Mediterranean upper bathyal muds)
	N	Coastal habitats	N12: Mediterranean and Black Sea sand beach, N14: Mediterranean, Macaronesian and Black Sea shifting coastal dune, N1B: Mediterranean and Black Sea coastal dune scrub
	C	Inland waters	C2: Surface streams (macrophytic associations associated with permanent or seasonal streams are the representatives of this habitat type in the area). C2.4: Tidally influenced rivers, upstream currents from river mouths. C3.1: Species-rich helophyte beds
	V	Vegetated man-made habitats	V12: Mixed crops of market gardens and horticulture
Akyatan Lagoon	M	Marine benthic habitats (MA-MG)	MA5 : Littoral sand, MA55: Mediterranean littoral sand, MA651: Biocenosis of Mediterranean upper bathyal muds
	N	Coastal habitats	N12: Mediterranean and Black Sea sand beach, N14: Mediterranean, Macaronesian and Black Sea shifting coastal dune, N1B: Mediterranean and Black Sea coastal dune scrub
	S	Heathland, scrub and tundra	S4: Temperate shrub heathland S5: Maquis, arborescent matorral and thermo-Mediterranean scrub; S51: Mediterranean maquis and arborescent matorral
	C	Inland waters	C2: Surface streams (macrophytic associations associated with permanent or seasonal streams are the representatives of this habitat type in the area). C2.4: Tidally influenced rivers, upstream currents from river mouths. C3.1: Species-rich helophyte beds
	V	Vegetated man-made habitats	V12: Mixed crops of market gardens and horticulture
Yer Köprü Natural Formation	S	Heathland, scrub and tundra	S4: Temperate shrub heathland S5: Maquis, arborescent matorral and thermo-Mediterranean scrub; S51: Mediterranean maquis and arborescent matorral

	R	Grasslands and lands dominated by forbs, mosses or lichens	R1: Dry grasslands; R1D3: East Mediterranean xeric grassland
Şekerpınar	V	Vegetated man-made habitats	V12: Mixed crops of market gardens and horticulture
	J	Constructed, industrial and other artificial habitats	J1.2: Dwellings in peri-urban and rural areas
Tatarlı Village Antique Period Ruins	V	Vegetated man-made habitats	V12: Mixed crops of market gardens and horticulture
	J	Constructed, industrial and other artificial habitats	J1.2: Dwellings in peri-urban and rural areas

Within the scope of this information, it is planned to develop habitat and regional monitoring sections of the National Biodiversity Inventory and Monitoring Project. The project will contribute to the monitoring of species, habitats and landscape maps of provinces by determining the habitat types of Turkey in detail. In addition, especially for the monitoring section, EUNIS habitat types should be determined and maps should be prepared in digital environment. These maps will gain a different dimension by revealing habitat types. This study contributed to the determination of EUNIS habitat types in our country.

REFERENCES

- Anonymous, (2025). *Eunis habitat type hierarchical view (Marine version 2022 and terrestrial version 2021)*. Access address <https://eunis.eea.europa.eu> Last Access 01.03.2025.
- Arslan, M. & Arslantürk, N. (2009). Avrupa doğa bilgi sistemi (EUNIS) habitat sınıflandırması. *Orman Mühendisliği*. 46 (1-2-3). 48-51.
- Arslan, M., Bingöl, M. Ü., & Erdoğan, N. (2012). Avrupa doğa bilgi sistemi (EUNIS) habitat sınıflandırması ve Türkiye batı Öksin alanındaki Doğu Kayını (*Fagus orientalis* Lipsky) ormanları örneği. *Artvin Çoruh Üniversitesi Orman Fakültesi Dergisi*, 13 (2), 278-290.
- Çakmak, M. H. & Aytaç, Z. (2020). Determination and mapping of EUNIS habitat types of Mamak District (Ankara) Turkey. *Acta Biologica Turcica*. 33(4), 227-236.
- Çakmak, M. H. & Aytaç, Z. (2021). EUNIS Habitat sınıflandırmasının Türkiye durum değerlendirmesi. *Bilge International Journal of Science and Technology Research*. 5 (2), 157-163. <https://doi.org/10.30516/bilgesci.888297>
- Çiftçi, D. (2015). Sündiken Dağları Staphylinine (Coleoptera: Staphylinidae) grubunun tür çeşitliliği ve EUNIS habitatları ile ilişkisi (Tez no 395753). [Doktora Tezi, Gazi Üniversitesi Fen Bilimleri Enstitüsü Biyoloji Ana Bilim Dalı]. Yükseköğretim Kurulu Ulusal Tez Merkezi.
- Davies, C. E., Moss, D., & Hill, M.O. (2004). *EUNIS Habitat classification revised 2004*. http://Eunis.Eea.Eu.Int/Upload/Eunis_2004_Report.Pdf. Last Access 01.03.2025.
- Davis PH (1965-1985). *Flora of Turkey and the East Aegean Islands*, Vol. 1-9. Edinburgh: Edinburgh University Press.
- Davis PH, Hedge IC (1975). *Flora of Turkey Past, Present and Future*. Candollea, Edinburgh, UK.

- Demir, O., Kızılırmak, A., Bozdağ, C. M., & Cabi, E. (2022). Determination of terrestrial EUNIS habitat types of Mount Ganos (Işıklar), Tekirdağ, Türkiye. *Frontiers in Life Sciences and Related Technologies*. 3 (2), 69-74.
- Erdoğan, İ. (2016). Coğrafi bilgi sistemleri ve uzaktan algılama ile Aşağı Kelkit Havzası EUNIS habitat tiplerinin tanımlanması ve potansiyel ürün yetiştirme alanlarının tespiti, (Tez no 446385). [Yüksek Lisans Tezi, Gaziosmanpaşa Üniversitesi Fen Bilimleri Enstitüsü Biyoloji Ana Bilim Dalı]. Yükseköğretim Kurulu Ulusal Tez Merkezi.
- Geven, F., Özdeniz, E., Kurt, L., Bölükbaşı, A., Özbey, B.G., Özcan, A.U., & Turan, Ü. (2016). Habitat classification and evaluation of the Köyceğiz-Dalyan Special Protected Area (Muğla/Turkey). *Rendiconti Lincei. Scienze Fisiche e Naturali*. 27, 509-519. <https://doi.org/10.1007/s12210-016-0510-1>
- Kanca, H., Terzioğlu, E., Adıgüzel, U., Erbaş, S., & Erdoğan, E. (2019). *Türkiye'nin biyolojik çeşitliliği*. T.C. Tarım ve Orman Bakanlığı Doğa Koruma ve Milli Parklar Genel Müdürlüğü, Ankara.
- Karaömerlioğlu, D. (2007). Göksu deltasındaki (silifke) doğal ekosistemlerin bitki ekolojisi yönünden araştırılması, (Tez no 212532). [Doktora Tezi, Çukurova Üniversitesi Fen Bilimleri Enstitüsü Biyoloji Ana Bilim Dalı]. Yükseköğretim Kurulu Ulusal Tez Merkezi.
- Karaömerlioğlu, D. & Düzenli, A. (2008). Göksu deltası (silifke) doğal alanlarında ana habitat tiplerinin araştırılması. *Çukurova Üniversitesi Fen Bilimleri Enstitüsü Dergisi*. 17 (2), 2.
- Mergen, O., & Karacaoğlu, C. (2015). Tuz Lake Special Environment Protection Area, Central Anatolia, Turkey: The EUNIS habitat classification and habitat change detection between 1987 and 2007. *Ekoloji*. 24(95), 1-9. <https://doi.org/10.5053/ekoloji.2015.06>
- Moss, D. & Roy, D. (1998). *Towards a European habitat classification*. European Environment Agency, Copenhagen.
- Özen, A. & Ürker, O. (2020). Avrupa doğa bilgi sistemi (EUNIS) habitat sınıflandırmasını kullanarak Işıklı Gölü ve Gököl Sulak Alanlarında habitat değişimlerinin belirlenmesi. *Erzincan Üniversitesi Fen Bilimleri Enstitüsü Dergisi*. 13 (2), 518-531, <https://doi.org/10.18185/erzifbed.64607>.
- Seyfe, M. (2019). Kazan tepeleri (Kahramankazan/Ankara) sürüngen türlerinin EUNIS habitat tiplerine göre tercih ve dağılımları, (Tez no 598674). [Yüksek Lisans Tezi, Gazi Üniversitesi Fen Bilimleri Enstitüsü Biyoloji Ana Bilim Dalı]. Yükseköğretim Kurulu Ulusal Tez Merkezi.
- Şahin, B. & Karavelioğulları, F. A. (2018a). Erzincan ili EUNIS Habitat Tipleri ve Haritalandırılması (Poster). 1st International Congress on Plant Biology, Konya, Türkiye 10-12 Mayıs 2018, ss. 440.
- Şahin, B. & Karavelioğulları, F. A. (2018b). EUNIS Habitat Types and Mapping of Bayburt (Poster). International Ecology 2018 Symposium, Kastamonu, Türkiye, 19-23 Haziran 2023, ss. 1081.
- Tak, M. & Tel, A.Z. (2024). Determination of Akdağ (Adıyaman/Malatya) Habitat Diversity According to EUNIS Habitat Classification System (Editors: Mehmet Hakkı Alma & Sefa Altıkat). International Congress of High Value Added Agricultural Products, 01-03 December 2024, Iğdır, Türkiye. ISBN: 978-625-378-039-5.
- Terzioğlu, E., Güvendiren, A. D., Erdoğan, E., Erdoğan, N. M. & Ekmen Nural, İ. (2015). *Biyolojik Çeşitliliği İzleme ve Değerlendirme Raporu 2013-2014*. T.C. Orman ve Su İşleri Bakanlığı, Doğa Koruma ve Milli Parklar Genel Müdürlüğü, Ankara.
- Ulu Ağır, S., Kutbay, H.G., Karaer, F., & Surmen, B. (2014). The Classification of coastal dune vegetation in Central Black Sea Region of Turkey by numerical methods and EU habitat types. *Rendiconti Lincei. Scienze Fisiche e Naturali*. 25, 453-460. <https://doi.org/10.1007/s12210-014-0328-7>.

IS VERTICAL FARMING SUSTAINABLE?**DIKEY TARIM SÜRDÜRÜLEBİLİR Mİ?****Dr. Öğr. Üyesi Berna BAŞ**

Gaziantep Üniversitesi, Fen Edebiyat Fakültesi, Biyoloji Bölümü

ORCID ID: <https://orcid.org/0000-0003-2455-2849>**ÖZET**

Giriş ve Amaç: İklim değişimi, küresel ısınma, su kaynaklarının azalması, hızla artan nüfus, konvansiyonel tarımla yıpranan topraklar, tarım nüfusunun azalması gibi sorunlar gıda güvencesinin geleceğini tehlikeye sokmaktadır. Tarımsal verimi artırmak amacıyla bitkiler arasında gen-nakli, gen-düzenlenmeleri, nanoteknolojiye dayalı gübre-pestisid araştırmaları, topraksız dikey tarım (DT) gibi yeni çözüm arayışları da devam etmektedir. Dünyada ve ülkemizde on yılı aşkın süreden beri akademik ve ticari amaçlı DT faaliyetleri yürütülmektedir. Dikey veya vertikal tarımla, iklimden ve mevsimden bağımsız bütün zaman dilimlerinde yüksek teknoloji donanımlı çok katlı sera benzeri tesislerde belli bir alandan açık arazi üretimine nazaran çok fazla bitkisel üretim amaçlanmaktadır. DT, bir dönüm açık araziden elde edilecek verimden onlarca, yüzlerce kez fazla hasılat eldesiyle yüksek ticari kazanç hedefleyen ticari bir yatırımdır. Sera benzeri binalar tamamen kapalı ve ışıkla aydınlatılmaktadır veya çevresi gün boyu ışık alacak tarzda camla çevrili cemeke binalardır. Açık arazi uygulamalarına kıyasla DT’da daha az su tüketimi, LED aydınlatma sistemleriyle daha az enerji tüketimi, konvansiyonel tarımda kullanılan bütün kimyasal girdilerin uygunluğu, minimal ekosistem zararı, sel, kuraklık gibi risklerin olmaması DT’ı cazip hale getirmektedir. DT mekanlarının yıl boyu belli bir ısı seviyesinde kalması için ısıtma-soğutma ve havalandırma sistemlerinin enerji tüketimi bölgelere bağlı olarak yüksek maliyet getirebilir. Su kıtlığı olan orta doğu ülkelerinde ucuz enerji, kuzey ülkelerinde doğal rüzgar enerjisi maliyeti düşürebilir. Açık alan yetiştiriciliğinde makro-besin elementleri sentetik gübre olarak bitkilere verilirken mikro-besin elementleri grubundaki mineralleri bitkiler doğal olarak topraktan almaktadır. DT’da bitki gelişimi tamamen özel olarak hazırlanacak besin solüsyonlarına bağlıdır. Bazı çalışmalarda DT’la üretilen bitkilerin besin değerinin yüksek olduğu iddia edilmekte olup halk sağlığı açısından henüz risk bildirilmemiştir. Ele alınan derlemede DT’la ilgili son akademik ve ticari çalışmalar özetlenmektedir. **Materyal ve Metot:** Google Akademik uygulama üzerinden çeşitli yayınlar derlenmiştir. **Sonuç:** İlk kurulum maliyeti oldukça yüksek olan sistemin yıl boyu enerji tüketimi, bitkisel üretim maliyetini de yükseltecektir. Besin solüsyonları sentetik kimyasallarla hazırlandığı için sağlık riskleri açısından orta ve uzun vadede takibe alınmalıdır.

Anahtar Kelimeler: Dikey Tarım, Kapalı Alan Tarımı, Sürdürülebilirlik, Uygulanabilirlik**ABSTRACT**

Introduction and Purpose: Problems such as climate change, global warming, depletion of water resources, rapidly increasing population, soils worn out by conventional agricultural practices, and decreasing agricultural population endanger the future of food security. New solution searches such as gene transfer between plants, gene-editing, nanotechnology-based fertilizer-pesticide research, and soilless vertical farming (VF) are also ongoing to boost

agricultural yield. VF activities have been carried out for more than ten years in the world and in our country for academic and commercial purposes. In vertical farming, it is aimed to produce many more plant yields from a certain area in multi storey greenhouse-like facilities equipped with advanced technology in all periods independent of climate and season, compared to conventional farming methods. VF is a commercial investment that aims to obtain high commercial profit by obtaining tens or hundreds of times more yield than the yield obtained from one acre of open land. Greenhouse-like buildings are completely closed and illuminated or are buildings with glass-surround windows so that receive light throughout the day. In contrast to open field applications, VF has less water consumption, less energy consumption with LED lighting systems, the suitability of all chemical inputs used in conventional agriculture, minimal ecosystem damage, and the absence of risks such as floods and droughts, making VF attractive. The energy consumption of heating-cooling and ventilation systems for VF spaces to remain at a certain temperature level throughout the year may be expensive depending on the region. Cheap petrol in Middle Eastern countries with water scarcity and natural wind power in northern countries may reduce energy consumption prices. In open-field cultivation, macronutrients are donated to plants as synthetic fertilizers, while the minerals in the micronutrient category are absorbed naturally from the soil by plant roots. Plant development in VF depends entirely on specially formulated nutrient solutions. Although some studies stated that the nutritional value of plants generated with VF is higher, but no risk concerning public health has yet been reported. The present study outlines the latest academic and commercial studies on VF.


Materials and Methods: Various publications provided through the Google Scholar application have been reviewed.

Conclusion: The system incurs a higher startup installation cost, and it will consequently drive up the cost of crop production due to its constant energy consumption throughout the year. Since nutrient solutions contain based-synthetic chemicals, they should be traced in the medium- and long-term periods for a hazard to the people.

Key Words: Vertical Farming, Indoor Agriculture, Sustainability, Feasibility

GİRİŞ

Dikey tarımın (DT) kökleri ilk olarak Babil 'in asma bahçelerine dayandırılmakta olup, günümüzde modern anlamda ilk ticari dikey çiftlik 2012 yılında Singapur 'da kurulmuştur (Fei & ark., 2025). Ülkemizde de on yılı aşkın süredir akademik ve ticari çalışmalar yapılmaktadır. İsminden anlaşılacağı üzere dikey tarım veya vertikal tarım çiftçiliği, tarım arazisinin yanlara doğru değil de yukarı doğru genişletilmesiyle yapılmaktadır. Yani dikey çiftlikler, basit bir seranın çok katlı şekilde tesis edildiği ancak yüksek teknolojik donanımlı ve modern yapılardır (Şekil 1). Topraksız tarım olan dikey tarım yetiştiriciliğinde hidroponik, akuaponik, aeroponik gibi sistemler kullanılarak doğal ortama benzer optimal bitki yetiştirme koşullarında kontrollü bir yapay ortam yaratılmaktadır (Sabry, 2021). Dar bir alanda az miktarda girdi kullanarak yüksek verim amaçlanmaktadır. Bitkilerin yetiştirileceği raflar/katlar horizontal pozisyonda ve raflar/katlar üst üste vertikal pozisyonda inşa edilmekte olup bitkilerin ihtiyaç duyduğu ısı, ışık, nem, organik ve inorganik besin maddeleri yapay olarak temin edilmektedir. Özellikle geleneksel tarım sistemlerinde kullanılan büyük bir kısmı sentetik olan makro ve mikro besin elementleri içeren gübrelerin aynısı kullanılmaktadır ancak miktarları ve içerikleri bitkilere ve kullanılan sistemlere göre değişebilmektedir (Ferrarezi & ark., 2024; UKERC, 2024). Dolayısıyla bu sistemle geleneksel tarım veya sera tarımına göre birim alandan daha yüksek verimle daha fazla ticari kazanç hedeflendiği için halk sağlığı, çevresel yararlar, sürdürülebilir gıda güvenliği gibi konularla ilgili araştırma çalışmaları daha geri planda bırakılmıştır.



Şekil 1. Çeşitli dikey tarım tesisleri. **a)**UL Solutions, 2025 **b)**Plantagon's Vertical Farm, 2025 **c)**Vertical, 2025 **d)**Soilless Agriculture Illustrations & Vectors, 2025.

Ülkemizde Çevre, Şehircilik ve İklim Değişikliği Bakanlığı tarafından yayınlanan belgede DT uygulaması için gereken yapının özellikleri "ışık geçirmeyen duvarlarla kaplı, ısıl olarak iyi yalıtılmış ve neredeyse hava geçirmez özellikte depo benzeri bir yapı" şeklinde belirtilmektedir (Coğrafi Bilgi Sistemleri Genel Müdürlüğü, 2024). Oysa şekil 1b 'de görüldüğü gibi Plantagon firması tarafından İsveç'te yapılan çok katlı kübik camekan tesis gün-boyu doğal ışığı ve güneşi çeşitli düzeylerde alacak şekilde tasarlanmıştır. Ele alınan derlemede DT 'la ilgili son akademik ve ticari çalışmalar özetlenmektedir.

DİKEY TARIMIN POTANSİYELİ

DT modeliyle uygun bir tarım arazisinden elde edilecek ürün/verim potansiyelini aynı arazinin çok katlı izdüşümüyle seraya benzer tarzda çok katlı yüksek bina tesis ederek tarım alanını artırmak ve böylece yüksek hasılat amaçlanmaktadır. Dikey yetiştiriciliğin başarısında altı adet temel yapısal unsurun çok iyi tasarlanması gerekir (Kozai & Niu, 2016); 1)LED aydınlatma sistemine sahip çok raflı veya katlı düzenek, 2)sıvı-faz veya katı-faz sistem seçimi, 3)ısıtma-soğutma klima ve havalandırma ünitesi, 4)CO₂ gübreleme, 5)ısı yalıtımı iyi ve hava geçirmez duvar yapısı, 6)Sistemi kontrol eden otomasyon ünitesi (ısı, ışık, nem, CO₂ ve hava akımı gibi çevreyle ilgili ve besin solüsyonunun elektrik iletkenliği, pH, O₂, kök-zonu ısı kontrolü gibi faktörler sıkı kontrol altında tutulmalıdır). Ancak sürdürülebilirliği hâlâ tartışmalı olan dikey tarımsal yetiştiriciliğin avantajları ve dezavantajları ekonomik, çevresel, sosyal ve politik boyutlarıyla ele alınarak değerlendirilebilir (Benke & Tomkins, 2017).

Ekonomik Avantajları ve Dezavantajları

İlk başlangıçta kurulum maliyeti yüksek olan dikey tarım yetiştiriciliği vejetasyonu kısa olan toprak-üstü ve kısa boylu salata bitkileri, mikroyeşillikler, otsu sebzeler, bazı süs bitkileri vb. için daha ideal görünmekte olup hem teknik olarak hem de bedeli yüksek bir yetiştiricilik sistemi olduğundan dolayı her türlü bitki yetiştiriciliğine uygun değildir (Benke & Tomkins,

2017; Beacham, Vickers & Monaghan, 2019; Kozai & Niu, 2020). Günümüzdeki mevcut sistemle örneğin juvenil periyotları uzun olan ve büyük odunsu ağaçlarda yetişen meyveler, toprak altı yetişen bitki grupları gibi ürünlerin yetiştiriciliği için ideal bir model değildir. Boş bir alan üzerine çok katlı apartman modelinde inşa edilen sera yapı şeklindeki DT binaları veya tesisleri, birim alandan edinilecek verimi kat veya raf sayısına bağlı artırarak toplam hasılatı çok yükseklerle çıkaracaktır. Örneğin 1 m² vertikal alanda marul yetiştirilirse sera yetiştiriciliğine göre 12 kat, açık araziye göre 80 kat daha fazla hasılat elde edilebileceği hesaplanmıştır (Graamans & ark., 2018; Hallikainen, 2019; Wildeman, 2020). Bu şekilde lokal üretimle özellikle popülasyonun yoğun olduğu şehirlere yakın bölgelere tesis kurulmasıyla veya uygun bir binanın DT 'a uygun şekilde yeniden tasarlanmasıyla sağlanabilecek avantajlar; gıda transport giderleri ile CO₂ emisyonu azalacak, kolaylıkla bozulabilecek yiyeceklerde atık miktarı azalacak, kontrollü koşullar sağlanacağı için sel, kuraklık, hastalık-zararlı hasarları minimuma indirgenecek, traktör veya çiftlik makinelerinin kullandığı fosil türü yakıt kullanımı asgariye indirilecek, enerji tasarrufu da sağlanacaktır. Bu sistemle kutuplarda ve subtropikal bölgelerde mevsime bağlı yetiştiricilik sorun olmayacaktır. Ancak elektrik enerjisi üretim kaynağına bağlı olarak solar enerji, rüzgar enerjisi gibi doğal enerji kaynakları veya fosil kaynak enerjisi kullanımına göre enerji tüketim bedeli değişkenlik gösterebilir.

Dikey tarımda ısıtma, aydınlatma, havalandırma, klima düzeneklerinin enerji tüketimi en önemli gider kısmını oluşturmaktadır (Graamans & ark., 2018). Ülkemizde 2024 verilerine göre elektrik üretimimizin, %35,2'si kömürden, %18,9'u doğal gazdan, %21,5'i hidrolik enerjiden, %10,5'i rüzgardan, %7,5'i güneşten, %3,2'si jeotermal enerjiden ve %3,2'si diğer kaynaklardan elde edildiği rapor edilmiştir (TC Enerji ve Tabii Kaynaklar Bakanlığı, 2025). Kömürden enerji üretiminin yaratacağı karbon kirliliği, doğalgaza ödenen döviz, hızla azalan su kaynakları nedeniyle hidroelektrik enerjisinin de azalması gibi konular ele alınarak düşünülürse DK, ülkemiz açısından yüksek enerji maliyeti getirecektir, dolayısıyla ürün birim fiyatı da artacaktır. DT ile beraber karbon kaynağı nötr olan uygun maliyetli rüzgar enerjisi, dalga enerjisi, güneş enerjisi gibi sürdürülebilir temiz enerji kaynaklarına da ihtiyaç bulunmaktadır. Çevresi günboyu ısı-ışık alacak şekilde tamamen camla çevrili tasarlanan DT yapılarında kullanılacak malzemeye bağlı olarak ısı-ışık-enerji giderlerinden tasarruf sağlanabilir. Örneğin anti-reflective güneş panelleriyle klima kullanmadan soğutma sağlanabilir, antirefle filmle kaplı camlar kullanarak hem enerji tüketimi azaltılabilir hem de bu tarz camlardan enerji üretimi sağlanabilir (YUHUA, 2025), böylece enerji tüketim bedeli minimuma düşürülebilecektir. Dünya su kaynaklarının % 40 - % 70 gibi önemli miktarı tarımsal faaliyetler için kullanılmaktadır (World Bank Blogs, 2023). DT 'da su tüketimi açık arazi yetiştiriciliğine nazaran % 95 oranında tasarruf sağlamaktadır (World Economic Forum, 2023). Günümüzde açık alan yetiştiriciliğinde halen salma sulama sistemi de kullanılmakta ve çok fazla miktarlarda su harcanmaktadır, DT ile olağanüstü bir su tasarrufu sağlanacaktır.

Çevresel Avantajları ve Dezavantajları

DT uygulamalarıyla topraklar kimyasal girdilerle kirlenmeyecek ve organik yetiştiriciliğe benzer kalite de besin değeri yüksek ürün alınabilecek, lokal bir üretim olduğu için klasik tarım makineleri kullanılmayacak ve köyden kente gıda transferi de olmayacağı için harcanan yakıttan hem tasarruf sağlanacak hem de bu araçların kullandığı fosil yakıtların karbon emisyonu azalacaktır. DT ürünleri talep üzerine yetiştirildiğinde uzun süre raflarda/depolarda muhafazaya gerek kalmayacak ve dolaylı olarak enerji tasarrufu da sağlanacaktır. Böylece çevresel sürdürülebilirliği destekleyen DT 'ın ekosistem hasarını azaltacağı beklenmektedir.

Hidroponik sistemle DT 'ın başarıya ulaşmasında önemli faktörlerden biri su kalitesidir, çünkü gübre veya besin maddeleri su içinde çözülerek sisteme yüklenmektedir (London Grow, 2024). Bu amaçla suyun pH, ısı, besin solüsyon içeriği, elektrik kondaktivitesi, pest-mikrobiyel

bulaşma, alg-yosun gelişimi, tortu birikimi gibi su kalite standartlarını ilgilendiren fiziksel, kimyasal ve biyolojik özelliklerine atfedilen testler sürekli yapılarak su kalitesinin kontrol altında tutulması gerekmektedir (Auxgrow, 2024; London Grow, 2024). Hidroponik sulama sistemlerinde kullanılacak olan suda fark edilir düzeyde herhangi bir sorun olmasa bile temel kural olarak suyun 2 - 3 haftada bir değiştirilmesi önerilmektedir (London Grow, 2024). DT sulama suyu kimyevi gübre ve/veya pestisid içermekte olup kalite standartlarında değişimler olması durumunda drenajla kanalizasyona veya doğaya salınacaktır. Sonuçta sulak alanlara kadar ulaşabilecek olan kirli su ekosisteme de zarar verebilecektir. Ancak klasik tarıma nazaran DT atık suları çok daha az miktardadır ve teknolojik çözümlerle bu sorun ortadan kaldırılabılır niteliktedir. Örneğin filtrasyon sistemleriyle yapılacak drenajla çevresel hasar minimuma düşürülebilir.

Sosyal Boyutu

DT 'ın önemli eksiklerinden biri de kalifiye eleman yetersizliğidir. Konuyla ilgili uzmanların sayısı zamanla artacak, yapı mühendisliği gibi yeni iş olanakları ortaya çıkacak, biyoteknoloji 'nin kapsama alanı daha artacaktır. Halihazırda pahalı bir üretim olduğu için DT bitkileri yüksek gelir grubuna hitap etse de ilerleyen zamanda yeni gelişen teknolojilerle daha makul fiyatlı üretim imkan dahilinde görünmektedir. Dezavantajı ise kırsal kesimde yaşayan çiftçi nüfusu azaltabilir ve kentlere göçle çarpık şehirleşmeyi artırabilir. Bu nedenle bölgeye uygun devlet destekli yeni ürünlerin yetiştiriciliği, DT için uygun olmayan odunsu taksonlara ait hortikültür bitkileri, etnobotanik vb. içerikli bitki yetiştiriciliği gibi ulusal ekonomiye katkı sunabilecek projelerle çiftçiler ve yetenekleri güvenceye alınmalıdır. Bu konu aynı zamanda politik bir avantaj da sunabilir.

Politik Boyutu

Doğal ortamlarında bitkiler sürekli hastalık ve zararlı riski altındayken, kontrollü bir çevrede yapılacak üretim de bu risk faktörü nispeten azalacağı için DT biyogüvenlik yasalarıyla daha uyumlu olacaktır. 2016 yılında yapılan Birleşmiş Milletlerin İklim Değişim Konferansında kararlaştırılan Paris Antlaşmasının taahhütlerini yerine getirebilecek potansiyele sahip olması DT 'ın gelişmesine en önemli katkı sağlayacak bir politik sonuçtur. Diğer politik avantajı da ucuz enerji üretmek amacıyla küçük bölgesel birçok doğal enerji üretim santrallerinin kurulmasını destekleyecek ve enerji fiyatlarını düşürebilecek sonuçlara neden olma ihtimalidir. Az sayıdaki büyük enerji santrallerine bağımlılığı da azaltacaktır.

DIKEY TARIMIN SÜRDÜRÜLEBİLİRLİĞİ

Yapılan bazı çalışmalarda DT ile yetiştirilen buğdayın verimi klasik yöntem yetiştiriciliğiyle kıyaslanmıştır. LED aydınlatma sistemi ve %100 rüzgar enerjisinin kullanıldığı bir denemede, enerji verimliliği, kimyasal girdiler, alt-yapı sistemleri, üretim verileri gibi parametreler değerlendirildiğinde konvansiyonel üretim verimliliği DT sistemine göre önemli ölçüde üstünlük sağlamıştır (Sørensen, Olsen & Colley, 2021). Bunun nedenin buğdayın enerji yoğunluğunun yüksekliği ve fotosentezle enerji dönüşüm etkinliğinin düşük olmasından kaynaklandığı bildirilmiştir. Buğday örneğindeki gibi dikey üretimi uygun olmayan bu tarz bitkiler yeterli tarım alanı olmayan, ekosistemin önemli ölçüde dejenere olduğu, herhangi bir sebeple temel gıda güvencesinin zorunlu olduğu, maliyet sorunu olmayan veya düşük olduğu lokasyonlarda alternatif bir üretim modeli olarak uygulanabilir. Bunun yanısıra buğdayla ilgili yapılan başka bir araştırmada ise kontrollü bir çevrede 10 katlı dikey bir tesiste, bir hektar açık alan yetiştiriciliğine nazaran dünya ortalama yıllık buğday verimi olan 3,2 t/ha 'ın 220 ile 600

katı bir verim tahmini rapor edilmiştir (Asseng & ark., 2020). Ancak küçük bir alanda yapılan araştırmanın matematiksel hesaplamaları çok katlı ve daha geniş bir alana transfer edildiği zaman enerji verimliliği, sistem stabilitesinin devamlılığı, prosedürün geçerliliği, kontaminasyon riskinin yönetimi gibi risk olasılıkları ticari kazanç yönünden benzer sonuçlar vermeyebilir.

Bitkilerin gelişimi güneşten gelen fotonlar yardımıyla gerçekleşen fotosenteze bağlıdır. Dış çevreden tamamen izole edilen kapalı bir alanda DT sistemlerinde LED aydınlatma kullanılmakta olup kültüre alınacak bitkilerin fotosentez için gerek duyduğu ışık yoğunluk ve şiddeti bitki gelişimi ve verimini doğrudan etkileyecektir. Yetiştirilecek bitkilere uygun ışık yoğunluğu fotosentez verimliliğini en üst seviyeye çıkarmakta ve özellikle sebze grubunda yer alan bitkilerin nutrasötik kalitesini de artırmaktadır (Saengtharatip & ark., 2021). Sebze grubu için en ideal ışık yoğunluğu bitki çeşidine göre değişmekle beraber 150 ve 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ aralığındadır (Nájera & ark., 2022). Genellikle LED aydınlatma spektrumu 450 - 490 nm aralığındaki mavi ışık bitkilerin gelişimi ve büyüme verimliliğini desteklemekte olup, bitkilerde klorofil, nitrat, antioksidanlar ve toplam fenoliklerin miktarında artışlara neden olmaktadır (Saengtharatip & ark., 2021; Nájera & ark., 2022). Vertikal çilek üretiminde gün ışığının simüle edildiği yapay ışık yoğunluğunun $\geq 100 \mu\text{mol s}^{-1} \text{m}^{-2}$ ve maruziyet süresinin 6-8 saat arası ayarlanan denemede, kontrol grubu olarak pencereden gelen doğal ışık koşullarında yetişen çileklere göre, meyve tutumu, meyve ağırlığı ve verim üzerine belirgin bir şekilde olumlu yansıdığı rapor edilmiştir (Verma & ark., 2024). Ancak enerji tüketimi DT için önemli dezavantajlardan olup araştırmacılar yenilenebilir enerji kaynaklarına olan ihtiyacı belirtmişlerdir. Çok katlı dikey raflar şeklinde tesis edilen serada özellikle geniş yapraklı yeşilliklerin üretiminde raflar ve yapraklar güneş ışığının mekan içinde özellikle alt raflara homojen dağılımını engelleyebilir. Bununla ilgili yapılan bir denemede güneş ışınlarının yoğun geldiği bölgelere ışığı yönlendiren yansıtıcı paneller yerleştirilerek koyu gölge olan bölgelere doğal ışığın ulaşması sağlanmıştır (Lee & ark., 2024). Böylece çalışılan bitkilerin fotosentez aktivitesi %10 - %12 oranında artırılmıştır.

Portekiz 'de yapılan bir araştırmada katı-faz sistemle hidroponik yetiştirilen domates ve çilek tarımının katı atık substratları hindistan cevizi kabuğu ve hindistan cevizi kabuğu+çam kabuğu karışımından ibaret olup doğal olarak ağır metal elementleriyle bulaşık (Cd, Cr, Ni, Cu, Pb, Hg, Sb ve As gibi) bir maden ocağı çevresindeki topraklara uygulanmıştır (Gonçalves & ark., 2024). Elde edilen verilere göre, 30 günlük bir inkübasyondan sonra bu atık substratların topraktaki ağır metalleri ve metaloidleri %20 - %91 arasında değişen oranlarda immobil yaptığı, toprak toksisitesini toksik olmayan seviyelere düşürerek simbiyotik bir bakteri olan *Aliivrio fischeri* için daha yaşanır bir mikro-çevreye dönüşmesine katkı sunmuştur. Bu çalışma sonuçları yeni anlayışlara kapı açmaktadır. Kirlenmiş toprakların rehabilitasyonuna önemli destek sağlayan DT katı faz atıklarının valörizasyonu son derece ekonomik, kolay uygulanır ve çevre dostu sürdürülebilir bir çözüm sunulmaktadır. Örneğin patojenlerle bulaşık tarım topraklarında kimyasal pestisid kullanmaksızın DT atığı benzer substratlarla patojenler için elzem olan besin maddelerinin de alıkonmasıyla toprakların patojenlerden arındırılması da ihtimal dahilinde görülmektedir.

DT verimliliğinde diğer önemli faktörlerden birisi de gübreler ve toprak simbiyotik organizmalarıdır. Atık su arıtma tesislerinde bir yan ürün olarak oluşan mineral strüvit N, P ve Mg kaynağı olup yavaş salınan bir gübre olarak kullanılmaktadır (Ahmed & ark., 2018). Substrat olarak sadece perlit kullanılan topraksız bir ortamda yetişen fasulye tohumları nodülasyon bakterisi *Rhizobium* türleriyle muamele edilerek doğal toprağın simüle edildiği bir araştırmada fidelere damlama sulama sistemiyle strüvit gübreleme yapılmıştır (Arcas-Pilz & ark., 2021). Strüvit ve N₂-fikse eden bakteri kombinasyonu fasulyenin gelişimi sırasında bitkilerin N ihtiyaçları karşılanmıştır.

Kuzey Avrupa şartlarında, iklim deęişim etkisi ve çevresel sürdürülebilirlik perspektifinden marulun vertikal ve klasik sera yetiştiricilik performansları karşılaştırılmıştır. Finlandiya 'da yapılan çalışmada hem sera hem DT için iklim deęişim etkisi, toplam enerji ihtiyacı, fosil enerji kullanımı, mineral ve metal kaynaklarının kullanımı, alan kullanımı ve su kıtlığı parametreleri denemenin etki kategorilerini oluşturmuştur (Joensuu & ark., 2024). Çalışmada yenilenebilir enerji ve atık ısının geri kazanımı da dahil olmak üzere iklim deęişimi ve bütün etki kategorilerinin DT yetiştiriciliğinde en düşük seviyede kaldığı bildirilmiştir. Enerji kullanımı sera üretimine göre DT için oldukça düşük seviyede kalmakla beraber, ısının geri kazanımı olmazsa DT 'ın ortalama enerji tüketimi daha yüksek çıkmaktadır. Yani Finlandiya koşullarında yenilenebilir enerji dikkate alınmazsa, iklim deęişim etkisi ve kaynak kullanım verimliliği açısından DT ve sera üretim performansı birbirine benzemektedir.

Coğrafi yapıyı nedeniyle Japonya 'da ekilebilir tarım toprakları yaklaşık %20 - %30 dolaylarında olup tarım ve ormancılık devlet tarafından sübvansede edilmektedir (Trewartha, 1950). Bundan dolayı Japonya 'da DT yetiştiriciliği büyük ilgi odağı olmuş ve bugün 200 'den fazla DT çiftliklerinde dolma biber, brokkoli, salatalık domates, çilek gibi tarım ürünleri yetiştirilmektedir (Liu & ark., 2022). Japonya 'nın doğusundaki 2011 yılındaki büyük depremde sonra o bölgede bir vaka çalışması yapılmış olup bazı ithal sebzeler DT ile üretilmiş ve geleneksel tarım üretiminin çevresel etki sonuçları karşılaştırılmıştır (Liu & ark., 2022). Buna göre DT sebzelerinde çevreye salınan N (azot) miktarında yıllık %37, P (fosfor) miktarında ise %36 azalma olduğu rapor edilmiştir. Böylece Japonya gibi akuatik bir çevrede aşırı gübre kirliliği ve su kalitesindeki düşüş azaltılmıştır.

DT ile İlgili Çeşitli Ticari Şirket Haberleri

DT maliyeti işgal edilen alan, ürün çeşidi, ihtiyaç duyulan enerji, su, besin maddeleri vb. kaynaklara göre deęişebilir, ancak DT bitkileri sera ve geleneksel yetiştiricilikten 3 - 5 kat daha pahalıya mâl olmaktadır (Tasgal, 2019).

iFarm bir Finlandiya firmasıdır (iFarm, 2025). Şirket Suudi Arabistan, Birleşik Arap Emirlikleri, Rusya, Fransa, İsviçre, İtalya gibi çeşitli ülkelerde DT çiftlikleri kurmuştur ve fesleğen, marul, pazı, roka gibi yeşillikler üretmektedir. Firma tarafından DT 'la üretilen ürün fiyatının yaklaşık 17 dolar/kg olduğu ve çiftliğin toplam geliri 57.800 \$ olarak rapor edilmektedir. Finansal giderleriyle ilgili olarak da emek, kaynaklar, kamu hizmetleri, kira ve diğer işletme giderlerine yapılan aylık harcamaların yaklaşık 32.600 \$ 'a ulaştığı belirtilmektedir. Yani çiftliğin aylık getirisi yaklaşık 25.000 \$ civarındadır, yaklaşık 3,6 yıl sonra tüm yatırım maliyetinin telafi edileceği de tahmin edilmektedir.

Bir Kanada firması olan Eden Green çeşitli DT çiftlik tesisleri kurmaktadır (Eden Green Technology, 2025). Şirketin kurduğu tesislerde hidroponik sistemle marul, ıspanak, nane, kale, lavanta, fesleğen, biber, kişniş, dereotu, frenk soğanı gibi 50 'den fazla bitki yetiştiriciliği yapılmaktadır. Çilek, salatalık, kereviz, domates gibi bitkiler de ilerde kültüre alınmak üzere deneme aşamasındadır. Firmanın kurduğu seralar doğal güneş ışığını alacak tarzda tasarlanmakta ve diğer vertikal sistemlere göre % 90 daha az enerji kullanılmaktadırlar.

UNS Vertical Farms firması Dubai 'de kurulmuştur ve halen online satış yapmaktadır (UNS Vertical Farms, 2024). Diğer firma Plenty, ABD şirketi olup marul, roka, hardal yeşillikleri, ıspanak, kale gibi yeşillikleri üretmekte ve temiz, kontrollü kapalı bir mekanda hiçbir pestisid kullanmadıkları için bitkilerde hiç bir ilaç kalıntısının da bulunmadığı ifade edilmektedir (Plenty, 2025).

Bustanica, dünyanın en büyük kapalı hidroponik DT firmasıdır ve menşei Dubai olup çeşitli Avrupa, Asya ülkeleri ve ABD 'de yerleşkesi bulunmaktadır (Bustanica, 2024). Dubai yerleşkesinde bulunan üç katlı olan mekan yaklaşık 31.000 m² alana sahiptir ve yıllık 1.000.000

kg yapraklı yeşillikler üretilmektedir. Tamamen topraksız ortamda yapılan bitki yetiştiriciliğinde 250 milyon litre su tasarrufuyla konvansiyonel sistemden %95 daha az su kullanılmaktadır. 2022 yılında 3,5 milyon dolar yatırım maliyetiyle kurulan firma için 5 yıl içinde 138 milyon dolar gelir beklentisi hesaplanmaktadır.

SONUÇ

Klasik tarıma kıyasla inovatif teknolojilerin yoğun kullanıldığı DT sınırlı tarımsal kaynaklarla (toprak, kimyasal girdiler, su gibi) sağlıklı, güvenilir, verimli ürün temininde daha sürdürülebilir bir sistem olma yolunda ilerlemektedir. Kontrollü bir klima çemberinde az materyalle çok üretim sağlayan DT 'da, kimyasal girdilerin tamamının sentetik olması, uzman eleman yetersizliği, ilk yatırım maliyetinin çok yüksek olması, her bitki çeşidi için uygun olmaması, üretimin pahalı olması, bitkilerin beslendiği kimyasal besin solüsyon atıklarının doğaya salınması DT 'ın en önemli sorunları olarak görülmektedir. Özellikle bitkilerin doğal olarak topraktan aldığı doğal minerallerin yerine kullanılan sentetik kimyasalların insan sağlığına kronik zararlar ilgili etkisi bilinmemektedir. Sağlık üzerine etkilerini araştırmak için uzun süreli takip gereklidir. Diğer sorunlar ise çözülebilir nitelikte olup, zamanla deneyim kazanarak uzmanların yetişmesi, teknolojik gelişmelerle daha düşük maliyetli sistem kurulumu, klasik tarıma göre daha ucuz bitki üretim sistemleri veya prosedürleri, doğaya salınan gübre-pestisid içerikli atık solüsyonların filtrasyonla doğaya kontrollü bir şekilde bırakılarak ekosistemin korunması, bugün için yetiştirilmesi uygun olmayan bitkilerin de zamanla üretilebileceği yeni sistemlerin geliştirilmesiyle DT gıda güvencesinin sürdürülebilirliği için umut vermektedir. Ülkemizde enerji üretim maliyetinin yüksekliği nedeniyle dikey tarım için yenilenebilir doğal enerji kaynaklarıyla üretim yapılmadığı sürece, DT belli zamanlarda örneğin savaş, kıtlık, küresel ısınma gibi doğal afetler ve geçici felaket dönemlerinde veya tarıma elverişsiz çorak topraklarda, kurak bölgelerde vertikal tarımsal yetiştiricilik bütün tarım ürünleri için değil fakat vejetasyon süresi kısa olan bazı bitkiler için klasik tarıma alternatif değil fakat destekleyici olarak düşünülebilir.

KAYNAKLAR

Ahmed, N., Shim, S., Won, S. & Ra, C. (2018). Struvite recovered from various types of wastewaters: Characteristics, soil leaching behaviour, and plant growth. *Land Degradation & Development*, 29(9), 2864-2879.

Arcas-Pilz, V., Parada, F., Villalba, G., Rufi-Salis, M., Rosell-Melé, A. & Gabarrell Durany, X. (2021). Improving the fertigation of soilless urban vertical agriculture through the combination of struvite and rhizobia inoculation in *Phaseolus vulgaris*. *Frontiers in Plant science*, 12, 649304.

Asseng, S., Guarin, J. R., Raman, M., Monje, O., Kiss, G., Despommier, D. D. Meggers, F. M. & Gauthier, P. P. (2020). Wheat yield potential in controlled-environment vertical farms. *Proceedings of the National Academy of Sciences*, 117(32), 19131-19135.

Auxgrow, (2024). <https://auxgrow.com/tr/what-to-monitor-in-hydroponic-farming/> (Erişim tarihi: 11.02.2025)

Beacham, A. M., Vickers, L. H. & Monaghan, J. M. (2019). Vertical farming: a summary of approaches to growing skywards. *The Journal of Horticultural Science and Biotechnology*, 94(3), 277-283.

Benke, K. & Tomkins, B. (2017). Future food-production systems: vertical farming and controlled-environment agriculture. *Sustainability: Science, Practice and Policy*, 13(1), 13-26.

- Bustanica, (2024). <https://www.bustanica.ae/> (Erişim tarihi: 22.02.2025)
- Coğrafi Bilgi Sistemleri Genel Müdürlüğü, (2024). <https://www.akillisehirler.gov.tr/wp-content/uploads/fizibilite-rapor/10-Dikey%20Tar%C4%B1m.pdf>. (Erişim tarihi: 20.02.2025)
- Eden Green Technology, (2025). <https://www.edengreen.com/blog-collection/what-is-vertical-farming> (Erişim tarihi: 18.02.2025)
- Fei, S., Wu, R., Liu, H., Yang, F. & Wang, N. (2025). Technological Innovations in Urban and Peri-Urban Agriculture: Pathways to Sustainable Food Systems in Metropolises. *Horticulturae*, 11(2), 212.
- Ferrarezi, R. S., Qin, K., Hazard, C., Gatard, E., Gastaldo, T. B., Housley, M. J. Nieters, C. E. & Mesquita, M. (2024). Airflow, fertilizer solution recipes, and calcium concentrations influence lettuce and spinach growth in an indoor vertical farm. *Scientia Horticulturae*, 328, 112948.
- Gonçalves, J., Araújo, A., Pedron, T., Santos, R., Bouguerra, S., Ribeiro, J. A. Pereira, R., Pereira, C. M. & Azenha, M. (2024). Discarded substrates from soilless hydroponic horticulture as potential amendments for metal-contaminated soils. *Chemosphere*, 364, 143127.
- Graamans, L., Baeza, E., Van Den Dobbelsteen, A., Tsafaras, I. & Stanghellini, C. (2018). Plant factories versus greenhouses: Comparison of resource use efficiency. *Agricultural Systems*, 160, 31-43.
- Hallikainen, E. (2019). Life cycle assessment on vertical farming.
- iFarm, (2025). <https://ifarm.fi/technologies/ifarm-leafy-greens> (Erişim tarihi: 18.02.2025)
- Joensuu, K., Kotilainen, T., Räsänen, K., Rantanen, M., Usva, K. & Silvenius, F. (2024). Assessment of climate change impact and resource-use efficiency of lettuce production in vertical farming and greenhouse production in Finland: a case study. *The International Journal of Life Cycle Assessment*, 29(10), 1932-1944.
- Kozai, T. & Niu, G. (2016). Plant factory as a resource-efficient closed plant production system. In *Plant Factory* (pp. 69-90). Academic Press.
- Kozai, T. & Niu, G. (2020). Role of the plant factory with artificial lighting (PFAL) in urban areas. In *Plant Factory* (pp. 7-34). Academic Press.
- Lee, Y. Y., Yik, J., Koay, E. H. H. & Lau, H. H. (2024). Application of light redirecting panels to improve sunlight utilization in multi-tiered vertical farming systems. *Clean Technologies and Environmental Policy*, 26(2), 499-510.
- Liu, J., Oita, A., Hayashi, K. & Matsubae, K. (2022). Sustainability of vertical farming in comparison with conventional farming: A case study in Miyagi Prefecture, Japan, on nitrogen and phosphorus footprint. *Sustainability*, 14(2), 1042.
- London Grow, (2024). <https://www.londongrow.com/blogs/grow-tips/how-often-to-change-hydroponics-water-london-grow#:~:text=If%20your%20water%20quality%20checks,every%20two%20to%20three%20weeks>. (Erişim tarihi: 11.02.2025)
- Nájera, C., Gallegos-Cedillo, V. M., Ros, M. & Pascual, J. A. (2022). LED lighting in vertical farming systems enhances bioactive compounds and productivity of vegetables crops. In *Biology and Life Sciences Forum* (Vol. 16, No. 1, p. 24). MDPI.
- Plantagon's Vertical Farm, (2025). <https://www.archdaily.com/28903/plantagons-vertical-farm/plantagon-ed001> (Erişim tarihi: 22.02.2025)
- Plenty, (2025). <https://www.plenty.ag/about/> (Erişim tarihi: 22.02.2025)

- Sabry, F. (2021). *Vertical Farming: How shall we feed the three more billion people by 2050?* (Vol. 4). One Billion Knowledgeable.
- Saengtharapip, S., Joshi, J., Zhang, G., Takagaki, M., Kozai, T. & Yamori, W. (2021). Optimal light wavelength for a novel cultivation system with a supplemental upward lighting in plant factory with artificial lighting. *Environmental Control in Biology*, 59(1), 21-27.
- Soilless Agriculture Illustrations & Vectors, (2025). <https://www.dreamstime.com/illustration/soilless-agriculture.html> (Erişim tarihi: 22.02.2025)
- Sørensen, M. G., Olsen, S. I. & Colley, T. (2021). Comparing the Environmental Sustainability of Vertical and Conventional Wheat Farming Using Life Cycle Assessment.
- Tasgal, P. (2019). The Economics of Local Vertical and Greenhouse Farming Are Getting Competitive. Retrieved from The Economics of Local Vertical and Greenhouse Farming Are Getting Competitive: <https://agfundernews.com/the-economics-of-local-vertical-and-greenhouse-farming-are-getting-competitive.html>.
- TC Enerji ve Tabii Kaynaklar Bakanlığı, (2025). <https://enerji.gov.tr/bilgi-merkezi-enerji-elektrik#:~:text=2024%20y%C4%B1%C4%B1%20Aral%20Ay%C4%B1%20sonu,%C3%BC%20ise%20di%C4%9Fer%20kaynaklar%20%C5%9Feklindedir.> (Erişim tarihi: 19.02.2025).
- Trewartha, G. T. (1950). Land reform and land reclamation in Japan. *Geographical Review*, 40(3), 376-396.
- UKERC, (2024). UK Energy Research Centre. <https://ukerc.ac.uk/news/four-myths-about-vertical-farming-debunked-by-an-expert/> (Erişim tarihi: 23.02.2025).
- UL Solutions, (2025). <https://www.ul.com/news/trusted-partner-assessment-vertical-farming-systems> (Erişim tarihi: 23.02.2025)
- UNS Vertical Farms, (2024). <https://unsfarms.com/blog/post/everything-you-need-to-know-about-sustainable-farming-in-dubai> (Erişim tarihi: 22.02.2025)
- Verma, P., Singh, G., Singh, S. K., Mirza, A. A., Bakshi, M. & Kumar, L. (2024). Modulating Productivity of Strawberries (*Fragaria x ananassa* Duch.) Through Artificial Full-Spectrum Light in Indoor Vertical Farming. *Journal of Soil Science and Plant Nutrition*, 1-15.
- Vertical, (2025) <https://vertical.mt/what-is-vertical-farming/> (Erişim tarihi: 22.02.2025)
- Wildeman, R. (2020). Vertical farming: a future perspective or a mere conceptual idea. *University of Twente: Enschede, The Netherlands*.
- World Bank Blogs, (2023). <https://blogs.worldbank.org/en/opendata/strains-freshwater-resources-impact-food-production-water-consumption>. (Erişim tarihi: 20.02.2025)
- World Economic Forum, (2023). <https://www.weforum.org/stories/2023/06/how-vertical-farming-can-save-water-and-support-food-security/#:~:text=Vertical%20farms%20use%2095%25%20less,LED%20lights%20rather%20than%20sunshine.> (Erişim tarihi: 22.02.2025)
- YUHUA, (2025). <https://www.yuhua-glass.com/article/anti-reflective-glass-for-glass-greenhouses.html> Son Erişim 05 Şubat 2025.

**HARMFUL INSECTS OF THE HETEROPTERA (HEMIPTERA) SUBORDER SEEN
ON FRUIT TREES IN İĞDIR PROVINCE**

**İĞDIR İLİ MEYVE AĞAÇLARINDA GÖRÜLEN HETEROPTERA (HEMIPTERA)
ALTTAKIMINA AİT ZARARLI BÖCEKLER**

Arş. Gör. Dilek DOĞAN

Iğdır Üniversitesi, Ziraat Fakültesi

ORCID ID: <https://orcid.org/0000-0001-9692-7211>

Prof. Dr. Celalettin GÖZÜAÇIK

Iğdır Üniversitesi, Ziraat Fakültesi

ORCID ID: <https://orcid.org/0000-0002-6543-7663>

ÖZET

Türkiye bulunduğu coğrafi koşullar göz önüne alındığında meyvecilik anlamında subtropik ve ılıman iklim meyvelerinin üretimi konusunda oldukça verimli bir bölgede bulunmaktadır. Iğdır ili coğrafik konumu ve iklimi iklimi sahip olması sebebiyle Doğu Anadolu Bölgesi'nde meyve ve sebze yetiştiriciliği bakımından büyük öneme sahip illerinden birisidir. Bölge ekonomisi açısından büyük bir öneme sahip olan meyve üretimi, çeşitli etkenler nedeniyle hem miktar hem de kalite açısından kayıplara uğramaktadır. Özellikle meyve yetiştiriciliğinde birçok zararlı ile mücadele etmek zorunda kalınması durumu üzerinde durulması gereken önemli bir konudur. Iğdır ili meyve ağaçlarında tespit edilen Heteroptera (Hemiptera) alttakımına ait türlerden *Lygaeus equestris* (Linnaeus, 1758) *Apodiphus amygdali* (Germar, 1817), *Dolycoris baccarum* (Linnaeus, 1758), *Eurydema ornata* (Linnaeus, 1758), *Holcostethus vernalis* (Wolff, 1804), *Piezodorus lituratus* (Fabricius, 1794), *Rhaphigaster nebulosa* (Poda, 1761), *Monosteira uncostata* (Mulsant and Rey, 1852) ve *Stephanitis pyri* (Fabricius, 1775) zarar oluşturdukları belirlenmiştir. Sonuç olarak Iğdır ilinde Heteroptera alttakımına ait türler meyve ağaçlarında verdikleri zararlar nedeniyle dikkat çekmektedirler. Bu zararlılarla mücadelede kullanılan insektisitlerin hem çevreye hem de ekosistemde bulunan faydalı böceklere zarar verdiği bilinmektedir. Bu olumsuz etkiler göz önünde bulundurulduğunda, kimyasal mücadeleye alternatif olarak geliştirilen diğer mücadele yöntemleri yaygın şekilde uygulanmaya çalışılmaktadır. Ekonomik açıdan tehlike arz eden türlerin biyoloji ve ekolojilerinin iyi bilinmesinin yanında, dağılımlarının da belirlenmesi zararlı ile mücadelede fayda sağlayacaktır.

Anahtar Kelimeler: Meyve, Heteroptera, Iğdır, Zararlı

ABSTRACT

Turkey is located in a very fertile region for the production of subtropical and temperate climate fruits in terms of fruit growing considering its geographical conditions. Iğdır province is one of the provinces in the Eastern Anatolia Region, which has great importance in terms of fruit and

vegetable cultivation due to its geographical location and microclimate climate. Fruit production, which is of great importance for the economy of the region, suffers losses in terms of both quantity and quality due to various factors. Especially in fruit growing, the situation of having to fight with many pests is an important issue that needs to be emphasised. Among the species belonging to the suborder Heteroptera (Hemiptera) detected in fruit trees in Iğdır province, *Lygaeus equestris* (Linnaeus, 1758), *Apodiphus amygdali* (Germar, 1817), *Dolycoris baccarum* (Linnaeus, 1758), *Eurydema ornata* (Linnaeus, 1758), *Holcostethus vernalis* (Wolff, 1804), *Piezodorus lituratus* (Fabricius, 1794) *Rhaphigaster nebulosa* (Poda, 1761), *Monosteira unicastata* (Mulsant and Rey, 1852) and *Stephanitis pyri* (Fabricius, 1775) it has been determined that they cause damage. In conclusion in Iğdır province, species belonging to the suborder Heteroptera attract attention due to the damages they cause on fruit trees. It is known that insecticides used in the control of these pests harm both the environment and beneficial insects in the ecosystem. Considering these negative effects, other control methods developed as an alternative to chemical control are being widely applied. In addition to knowing the biology and ecology of economically hazardous species, determining their distribution will be beneficial in pest control.

Key Words: Fruit, Heteroptera, Iğdır, Harmful

GİRİŞ

Tarımsal üretim çeşitliliği bakımından oldukça zengin olan ülkemizde, meyve üretimi önemli bir yer teşkil etmektedir. Dünyada üzerinde tarımı gerçekleştirilen yaklaşık 140 meyve türünden 80'inden daha fazlası türünün ülkemizde ekonomik olarak yetiştirilebilmektedir. (Anonim, 2022). Toplam tarım ve orman alanlarının yaklaşık %5,5'inde meyve yetiştiriciliği yapılmaktadır. Yumuşak ve taş çekirdekli meyve ağaçları arasında, ağaç sayısı ve üretim açısından ilk sırada elma yer almakta, bunu sırasıyla armut, şeftali, kayısı ve erik izlemektedir. (Anonim, 1999). Ülkemizde bulunan toplam 26,6 milyon hektarlık tarım arazisinin yaklaşık %10'unu oluşturan 2,7 milyon hektarlık bölümünde meyve yetiştiriciliği gerçekleştirilmektedir (Anonim, 2004a). Doğu Anadolu Bölgesi'nde konumlanan Iğdır Ovası, sahip olduğu mikroklima özelliği sayesinde diğer bölgelere kıyasla daha ılıman bir iklim yapısına sahiptir. Iğdır Ovası ve çevresinde, Akdeniz iklimine benzeyen belirgin karasal iklim koşulları etkili olmaktadır. Iğdır Merkez, Karakoyunlu ve Aralık ilçelerinde kış ayları nispeten serin, yaz ayları ise sıcak ve kurak geçmekte olup, bölgenin yıllık yağış miktarı oldukça düşük seviyelerdedir. İlçelerden biri olan Tuzluca'da ise rakımın daha yüksek olması sebebiyle, Iğdır şehir merkezi ve diğer ilçelere kıyasla sıcaklıklar biraz daha düşük seyretmektedir. Iğdır ilinde yıllık ortalama sıcaklık 11,6°C olup, kış aylarında hava sıcaklığı -30°C'ye kadar düşerken, yaz aylarında 41°C'nin üzerine çıkabilmektedir. Bölgenin sahip olduğu yeryüzü şekilleri ve iklim özellikleri, pek çok farklı tarım ürününün yetişmesine imkân tanımaktadır. Bu geniş çeşitlilik, Iğdır'da tarımsal faaliyetlerin gelişmesine katkı sağlamış ve şehri, çevresindeki il ve ilçelere tarım ürünleri tedarik eden, onların ihtiyaçlarını karşılayan önemli meyve yetiştiriciliği merkezi konumuna getirmiştir (Anonim, 2013).



Şekil. 1. Iğdır İli ve İlçeleri

Iğdır'da 2004 yılı itibariyle yaklaşık 85.833 hektar olan tarım arazisinin, meyveciliğe ayrılan 2.104 hektar (% 2.45)'lık kısmında toplam 195.820 adet meyve ağacı bulunmaktadır. Bu ağaçlardan 112.700 adedini (% 57.55) kayısı, 34.446 adedini (% 17.59) elma, 27.300 adedini (% 13.94) şeftali, 3.674 adedini (% 1.87) armut, 11.700 adedini (% 5.97) asma, 2.100 adedini (% 1.07) vişne, 1.900 adedini (% 0.97) kiraz, 1.450 adedini (% 0.75) ise erik oluşturmaktadır (Anonim, 2004b). Meyve üretiminde ekonomik kayıplara yol açan zararlılar arasında, Heteroptera alttakımına ait türlerin de belirli bir payı bulunmaktadır. Iğdır ilinde meyve ağaçlarında bulunan Heteroptera alttakımına ait zararlı türler *Lygaeus equestris* (Linnaeus, 1758), *Apodiphus amygdali* (Germar, 1817), Büyük ağaç pentatomidi, *Dolycoris baccarum* (Linnaeus, 1758), Dut kımılı, *Eurydema ornata* (Linnaeus, 1758), *Holcostethus vernalis* (Wolff, 1804), *Piezodorus lituratus* (Fabricius, 1794), *Rhaphigaster nebulosa* (Poda, 1761), Ağaç küçük pentatomidi, *Monosteira unicostata* (Mulsant and Rey, 1852), Kavak kaplanı, *Stephanitis pyri* (Fabricius, 1775), Armut kaplanı bulunmaktadır. Meyve ağaçlarına zarar veren bu önemli zararlı türlerin kesin olarak tespit edilmesi, etkili mücadele yöntemlerinin geliştirilmesi açısından büyük bir gereklilik arz etmektedir. Iğdır ilinde meyveciliğin giderek daha fazla yaygınlaşması ve tarımsal üretimde önemli bir konuma yükselmesi ile birlikte, meyve ağaçlarında ciddi zararlar oluşturan ve verim kayıplarına neden olan Heteroptera alttakımına ait türlerin belirlenmesi büyük önem taşımaktadır. Bu araştırma kapsamında, söz konusu zararlı türlerin hangilerinin bölgede bulunduğu ve bunlardan hangilerinin daha yaygın olarak görüldüğü detaylı bir şekilde ortaya konulmuştur. Tarım ürünlerinde meydana gelen verim kayıplarını önlemek amacıyla yaygın şekilde kullanılan kimyasal pestisitlerin ise çeşitli olumsuz sonuçlara neden olduğu bilinmektedir. Bu olumsuzluklar arasında en önemlileri, insan sağlığına yönelik riskler ile ekosistem üzerindeki zararlı etkileridir. Kimyasal mücadele yöntemlerinin yol açtığı bu olumsuzlukları ortadan kaldırmak amacıyla, son yıllarda "entegre mücadele" adı verilen bir yöntem benimsenmiştir. Bu yaklaşımda, zararlı türlerin popülasyon dinamikleri ile doğal çevre ile olan etkileşimleri dikkate alınarak, farklı mücadele teknikleri ve yöntemleri birbirleriyle uyumlu şekilde uygulanmakta ve zararlıların popülasyon yoğunluğu ekonomik zarar eşiğinin altında tutulmaya çalışılmaktadır. Bu yöntemin başarılı olması için zararlı organizmaların doğru şekilde tanımlanması ve düzenli olarak izlenmesi büyük önem taşımaktadır. Özellikle zararlı türlerin biyolojik özellikleri, yaşam döngüsü, beslendiği bitki türleri ve doğal düşmanları detaylı bir şekilde bilinmelidir (Polat & Tozlu, 2010).

BULGULAR**İğdir İlinde Meyve Ağaçlarında Görülen Zararlı Heteroptera Türleri****Familiya: Lygaeidae*****Lygaeus equestris* (Linnaeus, 1758)**

Lygaeus equestris, Heteroptera alttakımına, Hemiptera takımına, Lygaeoidea üstfamilyasına ve Lygaeidae familyasına ait bir böcek türüdür. Bu türün bireyleri yaklaşık 11-12 milimetre uzunluğundadır. Kendine özgü siyah ve kırmızı desenleri, gelişmiş kanat yapıları ve güçlü bacakları ile dikkat çeker. Ön kanatlarındaki çapraz bantlar, zar üzerindeki beyaz noktaya ve kenar kısımlara kadar uzanır. Scutellum yapısı kılsız olup, küçük tüyler barındırır. Siyah bantlar, gözlere doğru yaklaştıkça genişler. Üzerindeki kırmızı-siyah desenler, böceğin korunmasına yardımcı olan caydırıcı bir etkiye sahiptir. Beslendiği bitkilerden toksik maddeler biriktirerek, potansiyel avcılar için hoş olmayan ve itici bir tat oluşturur (Demirkol, 2016). Türlerinin çoğu bitkilerle, özellikle de bitki tohumları ile beslenirler. Genellikle toprak yüzeyinde, taş ve döküntülerin altında ve alçak bitkiler üzerinde bulunurlar (Lodos & Önder, 1986).



Şekil. 2. *Lygaeus equestris* (Linnaeus, 1758) Ergini (Hillewaert, 2025)

Bu türün çok sayıda konukçu bitkisi bulunmakta olup, bazı tarım bitkilerinde yoğun popülasyonuna bağlı olarak ciddi zararlar oluşturduğu da literatürde belirtilmiştir. İğdir ilinde, “*L. equestris*, elma (*Malus domestica*), kayısı (*Prunus armeniaca* L.) ve kiraz eriği (*Prunus cerasifera*)” ağaçlarında tespit edilmiştir. Nisan ayından Ekim ayına kadar geçen süre boyunca elma ve kayısı üzerinde bulunduğu, ancak bu dönemde yoğun bir popülasyona ulaşmadığı tespit edilmiştir (Korkmaz, 2017).

Familiya: Pentatomidae***Apodiphus amygdali* (Germar, 1817), Büyük Ağaç Pentatomidi**

Baş ve vücut, kırmızımsı veya sarımsı kahverenginin farklı tonlarında olup, yoğun gözenekli bir dokuya sahiptir. Baş, önden bakıldığında dikdörtgen şeklinde görünmekte olup, gena bölgesi ön tarafta yukarı doğru kıvrımlıdır. Tylus'un ucu açık renkli, orta kısmı ise soluk sarı tonlarında belirginleşmektedir. Vücut yüzeyi, rastgele yerleşmiş çıkıntılar ve küçük çukurlarla kaplanmış bir yapı sergilemektedir. Pronotumun kenarları, ön tarafa yaklaştıkça küçük diken benzeri çıkıntılarla bezenmiştir. Hortum, kırmızımsı kahverengi olup, abdomenin ikinci segmentine kadar uzanmaktadır. Bacaklar siyah renkte olup, vücudun alt (ventral) kısmı açık

tonlarda ve desenli bir görünüme sahiptir. Stigmalar siyah olup, böceğin toplam uzunluğu 18 ila 19,5 mm arasında değişiklik göstermektedir (Külekçi, 2008).



Şekil 3. *Apodiphus amygdali* (Germar, 1817) Ergini

Polifag bir tür olan *Apodiphus amygdali*'nin ergin bireyleri ve nimfleri, stylet adı verilen özel ağız yapılarıyla ağaçların dal ve gövdelerine nüfuz ederek bitki özsuyunu emmek suretiyle beslenmektedir. Ancak, bu beslenme süreci sonucunda etkilenen ağaçlarda kayda değer bir zarar meydana gelmemektedir. Öte yandan, bazı ağaç türlerinde, zararlının sokup özsuyunu emdiği bölgelerde, bitki kendini koruma mekanizması olarak zambak salgılamaktadır (Lodos vd., 1978).

Anadolu ve Marmara bölgelerinde yayılım gösteren *Apodiphus amygdali*, dut, şeftali, kayısı, erik, badem, elma, armut, ceviz gibi birçok meyve ağacının yanı sıra çeşitli orman ağaçlarında, özellikle de çınar ağaçlarında yaşamaktadır. Erginleri ve nimfleri, bu meyve ağaçlarının dal, sürgün ve meyvelerinin özsuyunu emerek zarar oluşturmaktadır (Lodos, 1986; Bolu et al., 2006).

***Dolycoris baccarum* (Linnaeus, 1758), Dut Kıvılcığı**

Vücut, geniş ve yassı bir formda olup, genellikle koyu kahverengimsi veya soluk sarımsı bir renge sahiptir ve yüzeyi yoğun siyah beneklerle kaplanmıştır. Corium bölgesi çoğunlukla, bazen ise tüm vücut morumsu bir parlaklık sergileyebilir. Scutellum'un uç kısmında ve taban köşelerinde küçük kabartılı yapılar bulunmaktadır. Hem scutellum hem de pronotum üzerinde uzun ve dik tüyler yer almaktadır. Antenin üçüncü ve beşinci segmentleri siyah renkte olup, taban kısmı beyazdır. Hortum sarımsı bir tona sahip olup, son segmentinin arka yarısı siyah renkte ve coxae bölgesine kadar uzanmaktadır. Bacaklar sarı, stigmalar ise siyah renklidir. Kötü koku salgılayan bezin açıklığı kısa ve yarık şeklindedir. Connexivum, birbirini takip eden siyah bantlarla çevrilmiştir. Bireylerin vücut uzunluğu 10 ile 12 mm arasında değişmektedir (Aba, 2024).



Şekil. 4. *Dolycoris baccarum* (Linnaeus, 1758), Ergini (Awill, 2011)

Dolycoris baccarum, dönem dönem tarım alanlarında önemli ekonomik kayıplara neden olabilen, geniş bir konukçu yelpazesine sahip polifag bir zararlı türüdür. Bunun yanı sıra, bu türün kış mevsimini yüksek rakımlı bölgelerde geçirdiği ve ülkemizde mart ve nisan aylarından itibaren alçak kesimlerdeki tarım arazilerine göç ederek bitkiler üzerinde zarar oluşturduğu bildirilmektedir (Lodos, 1982). Bu zararlı, bitki yapraklarında buruşma, çukurumsu deformasyonlar ve büzüşme gibi çeşitli fiziksel bozulmalara yol açarak, meyvelerin hem kalitesini hem de verimini olumsuz şekilde etkilemektedir (Arslan, 2007).

Iğdır ilinde, *D. baccarum*, kayısı (*Prunus armeniaca*), şeftali (*Prunus persica*) ve kiraz (*Prunus avium*) ağaçlarında tespit edilmiştir (Korkmaz, 2017).

Eurydema ornata (Linnaeus, 1758)

Baş, kırmızı renkte olup üzerinde siyah lekeler barındırır. Pronotum, alt kısmında altı siyah leke taşıyabilir ya da siyah lekeler iki ayrı alanda yer alabilir. Scutellum, kırmızı renkte olup, alt kısmında siyah benekler bulunur. Abdomen siyah renkte olup, connexivum ise kırmızı ve siyah renklerin birleşiminden oluşur. Membran, açık kahverengimsi tonlarda olup, stigmaların etrafı siyah lekelerle çevrilidir. Bireylerin vücut uzunluğu 6,5 ile 8,5 mm arasında değişmektedir (Korkmaz, 2017).



Şekil. 5. *Eurydema ornata* (Linnaeus, 1758), Ergini

Lahana kokulu böceği olarak da bilinen *Eurydema ornatum*, lahana ve karnabahar fidelerine zarar veren bir türdür. Atalay ve Çağlayan (1990), tarafından yapılan araştırmaya göre, Ege Bölgesi'nde *Eurydema* cinsine ait yedi farklı tür bulunmakta ve bu türler genellikle Cruciferae familyasına ait bitkilerin yapraklarını ve tohumlarını emerek zarar vermektedir. Baloğlu vd.

(2005), ise *E. ornatum*'un kayısı ağaçlarında bulunan zararlı türler arasında yer aldığını, ancak ekonomik açıdan önemli bir zarar oluşturmadığını belirtmişlerdir. Iğdır ilinde, *E. ornata*, kayısı (*Prunus armeniaca*) ve şeftali (*Prunus persica*) ağaçlarında tespit edilmiştir (Korkmaz, 2017).

***Holcostethus vernalis* (Wolff, 1804)**

Baş ve vücut koyu kahverenginden olup, yüzeyi yaygın siyah çukurlarla kaplanmıştır. Başın yan kenarları aşağıya doğru düz bir hat şeklindedir. Antenin ilk üç segmenti sarımsı kahverengi tonlarında, diğer segmentleri ise sarımsı kırmızı renkte görülür. Hortum sarı renkte olup, ucu siyah renktedir. Pronotum, çukur bir yapıya sahip olup, rengi sarımsı kahverengi veya sarı tonlarındadır. Scutellum kahverengi renkte olup, yüzeyi siyah çukurlar ile kaplanmış ve uç kısmı sarıdır. Connexivum sarımsı kahverengi bir görünüm sergiler. Bacaklar sarımsı tonlarda olup, subcoxae bölgesinde siyah lekeler bulunmaktadır. Bireylerin vücut uzunluğu 8,5 ile 10 mm arasında değişmektedir (Külekçi, 2008).



Şekil. 6. *Holcostethus vernalis* (Wolff, 1804), Ergini

Ege ve Marmara bölgelerinde yaygın ve bol bulunduğu, baklagiller ve birçok ağaç ve otsu bitki türleri üzerinde beslendiği bildirilmektedir (Lodos et al., 1978). Yapılan başka bir çalışmada Baloğlu vd. (2005), Kayısı ağaçlarında bu türe rastlandığını ancak yüksek popülasyonlara ulaşmadığı ve bu yüzden herhangi bir ekonomik zarara sebep olmadığı belirtilmiştir.

***Piezodorus lituratus* (Fabricius, 1794)**

Kış mevsiminde kahverengimsi, yaz mevsiminde ise yeşil renkte görülür. Antenleri sarımsı veya kırmızımsı tonlardadır. Hortumu açık sarı renkte olup, orta coxae kadar uzanır. Vücudun alt bölgesi ve bacaklar yeşilimsi veya sarımsı bir renkteki tonları yansıtır. Bireylerin vücut uzunluğu 10 ile 12 mm arasında değişmektedir (Aba, 2024).



Şekil. 7. *Piezodorus lituratus* (Fabricius, 1794), Ergini

Arslan (2007), tarafından yapılan bir araştırmada, bu zararlı, mercimek tanelerinde buruşma, çukurumsu çökmeler ve büzüşme gibi fiziksel deformasyonlara neden olarak, kabuk altında soluk beyaz bir yapı oluşumuna yol açmıştır. Bu durum, tanelerin kalite ve verimini olumsuz yönde etkileyerek, mercimek ihracatında üreticiler için büyük bir engel oluşturmuştur.

Özgen ve ark. (2005), *Piezodorus lituratus* türünün Diyarbakır ve Gaziantep illerinde kayısı ve kiraz ağaçlarında tespit edildiğini ifade etmişlerdir. Ayrıca, bu türün kayısı ağaçlarında popülasyon yoğunluğunun zamanla artarak ilerleyen yıllarda ana zararlı konumuna gelebileceğini öngörmüşlerdir.

Rhaphigaster nebulosa (Poda, 1761), Ağaç Küçük Pentatomidi

Erginler, kahverengimsi esmerden gri kahverengiye kadar değişen renk tonlarına sahiptir. Vücutları, belirgin siyah beneklerle kaplanmıştır. Scutellum, üçgen şeklinde olup, uç kısımlarında her iki tarafta birer siyah leke yer alır. Vücut uzunluğu ise 13 ile 16 mm arasında değişir (Öncül, 2006).



Şekil. 8. *Rhaphigaster nebulosa* (Poda, 1761), Ergini

Rhaphigaster nebulosa'nın ağaçlarda tomurcuk, sürgün, meyve ve çiçekleri sokup emerek beslendiğini bildirmiştir (Lodos, 1986). Farklı bir araştırmacıda *R. nebulosa*'nın kiraz da zararlı olduğunu ayrıca birçok meyve ve orman ağaçları ile bazı süs bitkilerinde tomurcuk, sürgün, meyve ve çiçeklerde zarar yaptığını kaydetmişlerdir (Özbek ve ark., 1996). Başta dut, fındık, zeytin dahil diğer birçok meyve ağaçları ile çınar, söğüt, *Tamarix* ve diğer orman, süs ağaç ve ağaçcıklarında bulunur (Lodos, 1986).

Iğdır ilinde, *R. nebulosa*, kayısı (*Prunus armeniaca*) ve vişne (*Prunus cerasus*), ağaçlarında tespit edilmiştir (Korkmaz, 2017).

Familiya: Tingidae

Monosteira unicastata (Mulsant and Rey, 1852), Kavak Kaplanı

Antenler, baş, pronotum ve hemielytra kahverengi tonlarında olup, pronotum ve hemielytra yüzeyi dantel benzeri desenlerle süslenmiştir. Pronotum hafifçe kabarık bir yapıdadır ve üzerinde yanal karinalar bulunmamaktadır. Canlının vücut uzunluğu ise 3 ila 4 mm arasında değişmektedir (Korkmaz, 2017).




Şekil. 9. *Monosteira unicastata* (Mulsant and Rey, 1852), Ergini

Bouali (2024), tarafından yapılan bir araştırmada, beslenme alışkanlıkları açısından *M. unicastata*, öncelikle konukçuyu delme ve emme davranışları sergileyerek, badem ağaçları gibi bitkilerden özsu emer. Bu beslenme davranışı, özellikle nimf ve ergin evrelerinde görülür ve böcek, beslenmek amacıyla bitki dokularını hedef alır. Bu tür beslenme etkinliği, genellikle konak bitkilerde benekler, renk değişiklikleri ve yaprak bozulmaları gibi belirgin semptomlara yol açar. Ülkemizin hemen her tarafında bulunduğunu, genellikle kavak, söğüt, elma, armut ve bademlerde görüldüğünü, bunların içinde en çok kavak üzerinde bulunduğunu ve zarar yaptığını belirtmiştir (Lodos & Önder, 1983). Korkmaz, 2017), kayısı (*Prunus armeniaca*) ve vişne (*Prunus cerasus*) ağaçlarında tespit etmiştir.

Stephanitis pyri (Fabricius, 1775), Armut Kaplanı

Vücut yassı ve geniş, sarımsı esmer renkte ve üzeri siyah lekeli; pronotum öne doğru uzamış ve başı örtmüş, iki yandan genişleyerek yarım daire şeklinde dışarıya taşar. Pronotum ve hemielitra üstten dantela görünümünde; bacaklar koyu sarı renktedir. Vücut 3-4 mm boyundadır (Korkmaz, 2017).



Şekil. 10. *Stephanitis pyri* (Fabricius, 1775), Armut kaplanı Zararı ve Ergini

Stephanitis pyri taş ve yumuşak çekirdekli meyve ağaçları ile Rosaceae familyasına bağlı süs bitkilerinde zarar yapmaktadır. Zararı ergin ve nimflerinin yaprağın alt yüzeyinde beslenmesi sonucu ortaya çıkar (Aysal & Kıvan, 2008). Zarar sonucu yaprağın üst yüzeyinde klorotik lekeler oluşur. Yaprtağın alt yüzeyi ise ergin ve nimflerin beslenmesi sonucu ürettikleri atıklar nedeniyle siyah veya koyu kahverengi renk alır. Zarar gören yapraklarda fotosentez ve solunum faaliyetlerinde düşüş görülür. Bunun sonucu olarak yapraklarda kahverengileşme ve erken dökülme görülebilir (Göksu, 1964; Lodos, 1982). Armut dantel böceği, *Stephanitis pyri* (F., 1775) (Heteroptera: Tingidae), Türkiye'de özellikle elma ve kiraz olmak üzere meyve ağaçlarının önemli bir zararlısı olarak bilinmektedir (Göksu, 1964; Lodos, 1982; Gulpercin & Önder, 1999; Aysal & Kıvan, 2008). Bunun yanı sıra gül, ayva, armut, erik, kayısı, alle üzümü, kavak, çınar, karaağaç, muşmula, kestane, fındık ve cevizde rastlanmaktadır (Toros, 1988; Anonim, 2011; Ertop & Özpınar, 2011).

TARTIŞMA VE SONUÇ

Meyve yetiştiriciliği, kültürümüzde ve ülke ekonomisinde büyük bir öneme sahip olup, her geçen yıl bu alandaki üretim miktarında ve bu ürünlerin dış ticaretinde artış yaşanmaktadır (Sarı & Yıldırım, 2021).

Yapılan çalışmalar sonucunda Heteroptera alttakımına ait türlerden *Dolycoris baccarum* (Linnaeus, 1758), *Eurydema ornata* (Linnaeus, 1758), *Holcostethus vernalis* (Wolff, 1804), *Piezodorus lituratus* (Fabricius, 1794), *Rhaphigaster nebulosa* (Poda, 1761), *Lygaeus equestris* (Linnaeus, 1758), *Apodiphus amygdali* (Germar, 1817), *Monosteira uncostata* (Mulsant and Rey, 1852), *Stephanitis pyri* (Fabricius, 1775) meyve ağaçlarında önemli zararlı konumunda oldukları tespit edilmiştir.

Türkiye'de meyve ağaçlarında bulunan heteroptera türlerinin oluşturduğu zararlılarla ilgili yapılan araştırmalar, meyve üretiminde entegre zararlı yönetimi (IPM) yöntemlerinin daha etkili ve uzun vadede sürdürülebilir bir yaklaşım olduğunu göstermektedir. Zararlılarla mücadelede erken tespitin yanı sıra, kimyasal ve biyolojik kontrol yöntemlerinin doğru bir şekilde ve zamanında birleştirilerek uygulanması, böceklerle yapılan mücadelenin başarı oranını önemli ölçüde artırmıştır. Özellikle bölgesel ekosistem koşullarına uygun zararlı yönetim stratejileri, hem ürün verimliliğini artırmış hem de çevreye zarar vermeyen, doğa dostu bir üretim süreci oluşturulmasına yardımcı olmuştur. Sonuç olarak, mücadelede kimyasal ve biyolojik yöntemlerin dengeli bir biçimde kullanılması gerektiği vurgulanmış, bu yöntemlerin meyve üreticilerine ekonomik fayda sağladığı ve zararlı popülasyonlarının uzun vadede kontrol altına alınmasında etkili olduğu tespit edilmiştir. Tarımda sürdürülebilir yaklaşımların

benimsenmesi ve çevre dostu uygulamalara yönelmek, meyve üretiminin hem verimliliğini hem de sağlıklı gelişimini olumlu bir şekilde etkilemektedir.

KAYNAKLAR

- Aba, V. (2024). *Şanlıurfa İlindeki Badem Bahçelerinde Kışlayan Böcek (Arthropoda: Insecta) Faunasının Belirlenmesi* (Doctoral Dissertation).
- Anonim, (1999). *Tarımsal Yapı 1997*. Başbakanlık Devlet İstatistik Enstitüsü, No: 2234, Ankara. (Erişim tarihi: 01.02.1999)
- Anonim, (2004a). *Tarımsal Yapı ve Üretim*. Başbakanlık Devlet İstatistik Enstitüsü Yayınları, Ankara. (Erişim tarihi: 05.04.2004).
- Anonim, (2004b). *Iğdır İli Master Planı*. T.C. Tarım Bakanlığı, Ankara. (Erişim tarihi:06.02.2004)
- Anonim, (2022). TUİK verileri. www.tuik.gov.tr (Erişim tarihi: 10.02.2024).
- Anonim, (2011). *Gül-Tahıl Yaprakbiti*. http://www.rothamsted.ac.uk/insectsurvey/STMetopolophium_dirhodum.php (Erişim tarihi: (03.12.2011).
- Arslan, M. (2007). <http://bahcesel.net/anasayfa/bahceselozel-haber/1593-kirmizi-mercimekte-tebesirlesmeihracativurdu.html>. (Erişim tarihi: 21.12.2014).
- Atalay, R., & Çağlayan, L. (1990). Investigations on the population-injury relationships of *Eurydema ornatum* L. (Heteroptera, Pentatomidae) which harmful on cabbage and cauliflower seedlings. *Turkish Journal of Entomology*, 14(2).
- Aysal, T., & Kıvanç, M. (2008). Development and population growth of *Stephanitis pyri* (F.) (Heteroptera: Tingidae) at five temperatures. *Journal of Pest Science*, 81, 135-141.
- Baloğlu, S., Yılmaz, A., Uygun, N., Uygur, N., Erkıılıç, A., Ulusoy, R., Elekçioğlu, H., Erkıılıç, L., Aysan, Y., Sipahioğlu, M., Özgönen, H., Öztürk, N. ve Ölmez, S. (2005). Malatya ve civarında kayısılarda bitki koruma sorunlarının saptanması ve entegre mücadeleye yönelik çözüm yollarının belirlenmesi. Tübitak Projesi Nihai Raporu, Proje No: 2573/3, 106 s, Ankara.
- Bolu, H., İ. Özgen & M. Fent. (2006). Diyarbakır, Elazığ ve Mardin illeri badem ağaçlarında bulunan Pentatomidae (Heteroptera) türleri. Yüzüncü Yıl Üniversitesi, Ziraat Fakültesi, *Tarım Bilimleri Dergisi*, 16 (1): 25-28.
- Bouali, Z., Assouguem, A., Joutei, A. B., Ullah, R., Alqahtani, A. S., & Lahlali, R. (2024). The false tiger of almond, *Monosteira unicostata* (Hemiptera: Tingidae): Biology, ecology, and control methods. *Open Agriculture*, 9(1), 20220310.
- Bulak Korkmaz, Y. (2017). Iğdır ve Kars (Kağızman) illerinde yumuşak ve sert çekirdekli meyve türlerinde bulunan zararlı böcek türleri. Doktora Tezi.
- Ertop, S. & A. Özpınar. (2011). Çanakkale ili kiraz ağaçlarındaki fitofag ve yararlı türler ile bazı önemli zararlıların popülasyon değişimi. *Türkiye Entomoloji Bülteni*. 1(2): 109-118.
- Göksu ME. (1964). Research on biology and control of pear lace bug (*Stephanitis pyri* Fabr.) in Sakarya ve Kocaeli. *Göztepe Zir. Muc. Enst. Yay. No: 160*, Istanbul, Turkey.
- Gulpercin, N. & F. Önder. (1999). Bornova koşullarında *Stephanitis pyri* (F.) (Heteroptera: Tingidae)'nin biyolojisi ve doğal düşmanları üzerinde çalışmalar. *Türkiye Entomoloji Dergisi*, 23: 51-56.

- Kıvan M, Aysal T. (2009). Tekirdağ'da Armut Kaplanının Yeni Bir Doğal Düşmanı: *Stethoconus pyri* (Mella) (Hemiptera: Miridae). Türkiye III. *Bitki Koruma Kongresi*, 358, Van.
- Külekçi, G. (2008). Erzurum ili Pentatomidae (heteroptera) türleri; üzerinde faunistik ve sistematik çalışmalar. Yüksek Lisans Tezi.
- Lodos, N. (1986). *Türkiye Entomolojisi II. (Genel, Uygulamalı, Faunistik)*. Ege Üniversitesi Ziraat Fakültesi Yayınları, No: 429, Bornova, İzmir, 580 s.
- Lodos, N. (1982). *Türkiye entomolojisi (Genel, uygulamalı ve faunistik cilt II)*. E. Ü. Zir. Fak. Yay. No: 429, 591 s.
- Lodos, N., Önder, F., Pehlivan, E., Atalay, R. (1978). Ege ve Marmara Bölgesinin zararlı bölge faunasının tespiti üzerine çalışmalar. T.C. Gıda-Tarım ve Hayvancılık Bakanlığı Zirai Mücadele ve Zirai Karantina Genel Müdürlüğü, Ankara, 135- 136.
- Öncül, T. (2006). Edremit (Balıkesir) Yöresi Zeytin Bahçelerindeki Heteroptera Faunasının Kışlak Tuzaklarla Belirlenmesi Üzerinde Araştırmalar. Yüksek Lisans Tezi.
- Polat, A., & Tozlu, G. (2010). Erzurum'da *Archips rosana* (Linnaeus, 1758) (Lepidoptera: Tortricidae)'nın kısa biyolojisi, konukçuları ve parazitoitleri üzerinde araştırmalar. *Turkish Journal of Entomology*, 34(4), 529-542.
- Sarı, E. D., & Yıldırım, E. (2021). Gülnar (Mersin) İlçesi elma ağaçlarındaki zararlı ve yararlı Arthropoda türlerinin tespiti ve bazı biyoekolojik gözlemler. *Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi*, 24(6), 1247-1262.
- Schaefer, C. W., & Panizzi, A. R. (Eds.). (2000). *Heteroptera of economic importance*. CRC press.
- Toros, S. (1988). Park ve süs bitkileri zararlıları. *Peyzaj Mimarlığı Derneği Yayınları*, Ankara.

THE ANTHELMINTIC EFFECT OF PREPARATIVE FORMS OF *TANACERUM VULGARE* L. ON GASTROINTESTINAL NEMATODES IN SHEEP**KOYUNLARDA GASTROİNTESTİNAL NEMATODLAR ÜZERİNE TANACERUM VULGARE L.'nin İLAC FORMLARININ ANTELMİNTİK ETKİSİ****Prof. Dr. Etibar MEMMEDOV**

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of Veterinary Medicine, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0004-6397-5662>**ABSTRACT**

Introduction and Purpose: At present, a large number of drugs belonging to different chemical and pharmacological groups are used against parasitic diseases of animals, especially helminthic diseases. The search, study and application in veterinary practice of sufficiently effective, ecologically clean antihelminthic agents derived from plants that do not have toxic effects on the animal organism is relevant. The anthelmintic effect of Tansy (*Tanacerum vulgare* L.) on gastrointestinal nematodes in sheep was studied in this study. The pathogenicity of gastrointestinal nematodes varies according to species. Ostertagia species cause morphological and functional disorders of the gastric glands of the abomasal gland. Trichostrongylus and Nematodirus species cause villous atrophy of the intestine, while Oesophogastromum spp and Ccooperia spp cause haemorrhage in the large intestine. Tansy is one of the common medicinal plants used against intestinal helminths in animals since ancient times.

Material and methods: Experiments were carried out in the Sadarak, Sharur and Kangarli districts of Araz river basin of Nakhchivan Autonomous Republic to study the effect of medicinal forms of Tansy plant against gastrointestinal nematodes infection in 1-2 year-old sheep. Before the experiments, fecal samples were taken from the rectums of animals and coprological examination was carried out according to Fulleborn's method. It was found that sheep in the experimental groups were %100 infected with gastrointestinal nematodes. There were 4 groups of 10 animals in each experimental group. Group 1 animals were given 5 g of herbal plant meal per animal for 5 days; group 2 animals were given 5 ml/kg of herbal infusion; group 3 animals were given decoction at a dose of 5 ml/kg. All medicinal forms were given twice a day with an interval of 12 hours. Group 4 were controls and did not receive any substances. The animals were kept under the same nutritional conditions. In 5 days after drug of the plant, were taken fecal samples from the rectum of animals of experimental and control groups and coprological examination was performed.

Results: The animals were monitored visually during the experiment. None of them showed signs of poisoning, changes in feed and water consumption. According to the results of coprological examination, the effectiveness of treatment in the group receiving vegetable meal was %60; %40 effectiveness in the group receiving infusion and %50 in the group receiving decoction. The animals of the control group had %100 infection as at the beginning of the experiment.

Discussion and Conclusion: As a result of the conducted experiment, it was found that Tansy has an antihelminthic effect against gastrointestinal nematodes.. Thus, in animals infected with gastrointestinal nematodes, the antihelminthic efficacy is % 60 after application of Tansy herb flour for 5 days; this was % 50in decoction and %40 in infusion. Having determined the effect of Tansy on sheep gastrointestinal nematodes., it is very promising to use it as an herbal antihelminthic agent.

Key words: Tansy, Gastrointestinal nematodes Medicinal plant, Herbal flour, Infusion, Decoction.

Özet

Giriş ve Amaç: Bitkilerden elde edilen, hayvan vücudu üzerinde toksik etkisi olmayan, yeterince etkili, ekolojik açıdan temiz antelmintiklerin araştırılması ve veteriner pratiğinde uygulanması önemlidir. Bu çalışmada Nahçıvan Özerk Cumhuriyeti'nin Araz nehri havzasında bulunan Sadarak, Şarur ve Kangarlı bölgelerindeki özel çiftliklerde 1-2 yaşındaki koyunlarda Solucanotu (*Tanacerum vulgare* L.) bitkilerinin gastrointestinal nematodlar üzerine etkisini incelemek için deneyler yapıldı. Bazı şifalı bitkilerin koyunlarda nematod enfeksiyonları üzerindeki antelmintik etkisi incelenmiştir. Gastrointestinal nematodların patojenitesi türlere göre değişkendir. Ostertagiya türleri abomazumun gastrik bezlerinde morfolojik ve fomksiyonel bozukluklulara neden olur. Trichostrongylus ve Nematodirus türleri bağırsaqlarda villöz atrofiye, Oesophogastromum spp ve Ccooperia spp ise kalın bağırsaqlarda kanamaya neden olur. Solucanotu (*Tanacerum vulgare* L.) hayvanlarda bağırsak helmintlerine karşı eski çağlardan beri kullanılan yaygın şifalı bitkilerdir biridir.

Gereç ve yöntemler: Bu şifalı bitkinin gastrointestinal nematodları ile enfekte koyunlarda antelmintik etkisini incelemek için deneyler yapıldı. Deneme gruplarındaki koyunların %100 nematodlar ile enfekte olduğu belirlendi. Denemelerin yapıldığı ilçelerde her grupta 10 hayvan olacak şekilde 4 grup (toplam 40 hayvan) düzenlenmiştir. Grup 1'deki hayvanlara 5 gün boyunca hayvan başına 5 g bitkisel bitki unu; grup 2'deki hayvanlara 5 ml/kg bitkisel infüzyon; grup 3'teki hayvanlara 5 ml/kg dozunda dekoksasyon verildi. Tüm ilaç formlar 12 saat arayla günde iki kez verilmiştir. Grup 4 kontrol grubudur ve herhangi bir madde almamıştır. Hayvanlar aynı beslenme koşulları altında tutuldu. Bitkinin uygulanmasından 5 gün sonra, deney ve kontrol gruplarındaki hayvanların rektumlarından dışkı örnekleri alınmış ve koprolojik muayene yapılmıştır.

Bulgular: Deney sırasında hayvanlar görsel olarak izlendi. Hiçbirinde zehirlenme belirtisi, yem ve su tüketiminde değişiklik görülmedi. Koprolojik inceleme sonuçlarına göre Solucanotu'nun ot unu alan grupta tedavinin etkinliği %60; İnfüzyon uygulanan grupta %40, dekoksasyon uygulanan grupta ise %50 etkililik olmuştur. Kontrol grubundaki hayvanlarda deneyin başlangıcında olduğu gibi %100 enfeksiyon vardı.

Tartışma ve Sonuç: Yapılan deney sonucunda Solucanotu'nun gastrointestinal nematodlara karşı antihelminthik etkiye sahip olduğu tespit edilmiştir. Böylece gastrointestinal nematodlar ile enfekte hayvanlarda Solucanotu'nun otununun 5 gün süreyle uygulanmasından sonra antihelminthik etkinlik %60 olur; bu dekoksasyon %50 ve infüzyonda %40 idi. Solucanotu'nun koyunların gastrointestinal nematodlara karşı daha güçlü etkisinin belirlenmesi, gelecekte bitkisel bir antelmintik olarak kullanılması oldukça umut vericidir.

Anahtar kelimeler: Solucanotu, Gastrointestinal nematodlar, Şifalı bitki, Bitkisel un, İnfüzyon, Dekoksasyon

EFFECT OF HIGH TEMPERATURE ON WALNUT CULTIVATION

YÜKSEK SICAKLIĞIN CEVİZ YETİŞTİRİCİLİĞİNE ETKİSİ

Şeyma KAYA

Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Horticulture,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0009-0007-3708-0023>

Feride MERCAN

Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Horticulture,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0009-0001-7137-4582>

Akide ÖZCAN

Kahramanmaraş Sütçü İmam University, Göksun Vocational school,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0000-0001-7182-3811>

Mehmet SÜTYEMEZ

Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Horticulture,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0000-0003-2417-8009>

ABSTRACT

Introduction and Purpose: Kahramanmaraş has an important place in terms of walnut cultivation. In recent years, increasing temperatures and irregular rainfall have negatively affected fruit growing. It is a case of losses in yield and fruit quality in some fruit species as a result of climate change. Walnut is the most important source of agricultural income for Kahramanmaraş. The province is famous throughout Türkiye for its quality and delicious walnuts. However, in recent years, losses in productivity and quality have been observed due to changing climatic conditions. This situation increases the risk of limiting walnut cultivation in the region in the coming years. This study was conducted to determine the fruit quality performances of some walnut varieties grown in Kahramanmaraş.

Materials and Methods: This research was conducted on Chandler and Kaman 1 varieties in the application gardens of Kahramanmaraş Sütçü İmam University in 2023. Measurements and calculations of pomological properties of fruits of each variety were made. Measurements were made on 100 fruit samples collected separately from 3 plants of each variety, representing each tree. For each fruit, fruit dimensions (width, length, height) were measured with a caliper, and shell weight and internal weight were weighed with a precision scale. Fruit kernel ratio was calculated by dividing the weight of the fruit inside by the weight of the fruit with skin and multiplying by 100. **Results:** It was determined that in the Chandler walnut variety, the fruit weight varied between 8.69g and 13.16g, the internal ratio ranged between 36% and 39%, and the values of the same characteristics in the Kaman 1 variety varied between 9.94g and 16.42g, and the internal ratio ranged between 38% and 55%, respectively. It is known that the inner color of the fruit of the Chandler variety is very light. It has been observed that Chandler is

significantly affected by sunburn under Kahramanmaraş conditions. In the Kaman 1 variety, it was observed that the loss was less in terms of both internal color and internal ratio.

Discussion and Conclusion: According to the data obtained from this study, it was determined that the varieties had pomological results well below the standard quality and yield values. This is a very striking result. It is clear that long-lasting, extremely hot days in the Kahramanmaraş region in 2023 will cause quality and yield loss due to sunburn. In addition, it is seen that cultural works cannot be carried out on time or at all due to the deficiencies in labor services due to the earthquake that occurred in the province, which triggered the decrease in tree yield and quality values. Therefore, it has been confirmed that cultural maintenance works, especially irrigation, should be done on time and at an adequate level during long summer temperatures.

Key Words: Walnut; high temperature; Sustainability

ÖZET

Giriş ve Amaç: Kahramanmaraş ceviz yetiştiriciliği bakımından önemli bir yere sahiptir. Son yıllarda artan sıcaklıklar ve düzensiz yağışlar meyve yetiştiriciliğini olumsuz etkilemektedir. Bazı meyve türlerinde iklim değişikliğinin sonucu verim ve meyve kalitesinde kayıpların olduğu bir vakadır. Ceviz, Kahramanmaraş için en önemli tarımsal gelir kaynağıdır. İl, Türkiye genelinde kaliteli ve lezzetli cevizleriyle meşhurdur. Ancak son yıllarda değişen iklim şartlarından dolayı verim ve kalitede kayıplar görülmektedir. Bu durum bölgede gelecek yıllarda ceviz yetiştiriciliğini sınırlandırma riskini artırmaktadır. Bu çalışma, Kahramanmaraş'ta yetiştirilen bazı ceviz çeşitlerinin meyve kalite performanslarının belirlenmesi amacıyla yapılmıştır.

Gereç ve Yöntem: Bu araştırma, 2023 yılında Kahramanmaraş Sütçü İmam üniversitesi uygulama bahçelerinde bulunan Chandler ve Kaman 1 çeşitleri üzerinde yürütülmüştür. Her çeşide ait meyveler de pomolojik özelliklerin ölçümleri ve hesaplamaları yapılmıştır. Her çeşide ait 3 bitkiden, her ağacı temsil edecek şekilde ayrı ayrı toplanan 100 er meyve örneği üzerinde ölçümler yapılmıştır. Her meyve için kumpas ile meyve boyutları (en, boy, yükseklik), ölçülmüş ve hassas teraziyle kabuklu ağırlık ve iç ağırlık tartımı yapılmıştır. Meyve iç oranı iç meyve ağırlığının kabuklu meyve ağırlığına bölünerek 100 ile çarpılmasıyla hesaplanmıştır.

Bulgular: Chandler ceviz çeşidinde meyve ağırlığı 8.69g ile 13.16g iç oranı %36 ile %39 arasında Kaman 1 çeşidinde aynı özelliklere ait değerlerin sırasıyla 9.94g ile 16.42g iç oranı %38 ile %55 arasında değiştiği belirlenmiştir. Chandler çeşidinin meyve iç rengi çok açık olduğu bilinmektedir. Kahramanmaraş şartlarında Chandler'ın güneş yanıklığından önemli seviyede etkilendiği gözlemlenmiştir. Kaman 1 çeşidinde ise hem iç rengi hemde iç oranı bakımından kaybın daha az olduğu görülmüştür.

Tartışma ve Sonuç: Bu çalışmadan elde edilen verilere göre, çeşitlerin standart kalite ve verim değerlerinin çok altında pomolojik sonuçlara sahip oldukları belirlenmiştir. Bu çok dikkat çekici bir sonuçtur. Kahramanmaraş bölgesinde 2023 yılında uzun süren aşırı sıcak günlerin güneş yanıklığı sebebiyle kalite ve verim kaybına sebep olduğu açıktır. Ayrıca ilde meydana gelen deprem nedeniyle işçilik hizmetlerinde görülen eksilikler nedeniyle kültürel işlerin zamanında ya da hiç yapılamaması da ağaç verim ve kalite değerlerinin düşmesini tetiklediği görülmektedir. Bundan dolayı, uzun süren yaz sıcaklıklarında başta sulama olmak üzere kültürel bakım işlerinin zamanında ve yeterli düzeyde yapılması gerektiği teyit edilmiştir.

Anahtar Kelimeler: Ceviz; Yüksek sıcaklık, Sürdürülebilirlik

SUSTAINABLE AGRICULTURE AND SUGAR BEET: AN ANALYSIS OF CARBON FOOTPRINT, WATER USAGE, AND EFFICIENCY

SÜRDÜRÜLEBİLİR TARIM VE ŞEKER PANCARI: KARBON AYAK İZİ, SU KULLANIMI VE VERİMLİLİK ÜZERİNE İNCELEME

Halit Burak YÜKSEKDAĞ

Kayseri Sugar, Sustainable Agriculture Engineer, Kayseri, Turkey

ORCID ID: <https://orcid.org/0009-0004-1251-6087>

Hayrullah ÖZBEK

Kayseri Sugar, Sustainable Agriculture Engineer, Kayseri, Turkey

ORCID ID: <https://orcid.org/0009-0005-2699-6234>

Mustafa Can SARIOĞLU

Kayseri Sugar, Plant Protection and Plant Nutrition Engineer, Kayseri Turkey

ORCID ID: <https://orcid.org/0009-0005-2417-3832>

Mustafa ARSLAN

Kayseri Sugar, Sustainable Agriculture Chef, Kayseri, Turkey

ORCID ID: <https://orcid.org/0009-0006-8162-6241>

Mustafa BÖLÜK

Kayseri Sugar, Chef of Plant Protection and Plant Nutrition , Kayseri Turkey

ORCID ID: <https://orcid.org/0009-0001-6543-2483>

ABSTRACT

Introduction and Purpose: Globally, 22% of sugar production comes from sugar beets. Turkey ranks fifth with a 9.1% share in production. Sugar beet is considered a strategic crop. Sustainable agriculture is an agricultural system that promotes environmentally friendly farming practices while preserving natural resources, aiming to maintain economic, social, and ecological balance. This approach seeks to sustain agricultural productivity while minimizing environmental damage and ensuring the long-term sustainability of the sugar sector.

Materials and Methods: This study was initiated in 2022 to evaluate the long-term benefits and economic returns of sustainable agriculture and was monitored regularly until 2024 to analyze the results. Conducted with the same 65 farmers each year, the research focused on assessing the impact of sustainable farming practices on agricultural production, environmental sustainability, and economic gains. The farmers involved in the study operate within the Kayseri Şeker cultivation area. The amount of sustainably produced sugar, sugar beet

cultivation areas, farmer records, net beet amounts delivered by region, the share of sustainable agriculture in total sugar production over the years, and data on fertilizers, pesticides, irrigation, and emissions were systematically recorded. This comprehensive monitoring process provides crucial data on reducing environmental impacts, enhancing agricultural productivity, and determining the long-term economic contributions of sustainable farming.

Results: Compared to 2022, in 2023, fertilizer use decreased by 0.51%, pesticide use by 6.8%, irrigation by 1%, and carbon emissions by 53%. By 2024, these reductions reached 1.3% for fertilizers, 8.6% for pesticides, 20% for irrigation, and 67% for carbon emissions.

Discussion and Conclusion: The results demonstrate that sustainable agricultural practices offer both environmental and economic advantages. Savings in water and fertilizer usage, maintaining productivity, and reducing carbon emissions clearly highlight the necessity of sustainable farming. Expanding sustainable agriculture is critical for preserving natural resources and ensuring future food security.

Keywords: Sustainable Agriculture, Environmental Sustainability, Agricultural Production, Economic Development, Carbon Emissions

Giriş ve Amaç: Dünya genelinde şeker üretiminin %22'si şeker pancarından yapılmaktadır. Türkiye %9,1'lik üretim payıyla beşinci sırada yer almaktadır. Şeker pancarı stratejik bir bitki konumundadır. Sürdürülebilir tarım, doğal kaynakları koruyarak çevre dostu tarım uygulamalarını teşvik eden ve ekonomik, sosyal ve ekolojik dengenin sağlanmasını hedefleyen bir tarımsal sistemdir. Bu yaklaşım, tarımsal üretimde verimliliği korurken, çevresel zararları en aza indirmeyi ve uzun vadede şeker sektörünün sürdürülebilirliğini sağlamayı amaçlamaktadır.

Gereç ve Yöntem: Bu çalışma, sürdürülebilir tarımın uzun vadeli faydalarını ve ekonomik getirilerini değerlendirmek amacıyla 2022 yılında başlatılmıştır ve 2024 yılına kadar düzenli olarak takip edilerek sonuçlar analiz edilmiştir. Her yıl aynı 65 çiftçi ile yürütülen bu araştırma, sürdürülebilir tarım uygulamalarının tarımsal üretim, çevresel sürdürülebilirlik ve ekonomik kazançlar üzerindeki etkilerini belirlemeye odaklanmıştır. Çalışmada yer alan çiftçiler, Kayseri Şeker ekim sahasında faaliyet göstermektedir. Sürdürülebilir tarım kapsamında üretilen şeker miktarı, şeker pancarı üretim alanları ve çiftçi kayıtları, bölgeler bazında yatırılan net pancar miktarları, yıllara göre sürdürülebilir tarımın toplam şeker üretimindeki payı, gübre, ilaçlama, sulama ve emisyon verileri sistematik olarak kayıt altına alınmıştır. Bu kapsamlı takip süreci, sürdürülebilir tarımın çevresel etkilerini azaltma, tarımsal verimliliği artırma ve uzun vadeli ekonomik katkılarını belirleme açısından önemli veriler sağlamaktadır.

Bulgular: 2022 yılına göre 2023 yılında %0,51 gübreleme, %6.8 ilaçlama, %1 sulama ve %53 karbon emisyonunun, 2024 yılında ise %1.3 gübreleme, %8.6 ilaçlama, %20 sulama ve %67 karbon emisyonunun düştüğü gözlemlenmiştir. **Tartışma ve Sonuç:** Elde edilen sonuçlar, sürdürülebilir tarım uygulamalarının hem çevresel hem de ekonomik açıdan avantajlar sağladığını göstermektedir. Su ve gübre kullanımında sağlanan tasarruflar, verimliliğin korunması ve karbon salınımının azaltılması, sürdürülebilir tarımın gerekliliğini açıkça ortaya koymaktadır. Sürdürülebilir tarım uygulamalarının genişletilmesi, doğal kaynakların korunması ve gelecekteki gıda güvenliğinin sağlanması için kritik bir öneme sahiptir.

Anahtar Kelimeler: Sürdürülebilir Tarım, Çevresel Sürdürülebilirlik, Tarımsal Üretim, Ekonomik Kalkınma, Karbon emisyonu

**DETERMINATION OF ORANGE FRUIT RIPENESS LEVELS USING DEEP
LEARNING-BASED INSTANCE SEGMENTATION METHOD**

Assist. Prof. Dr. Tahsin UYGUN

Burdur Mehmet Akif Ersoy University, Faculty of Engineering and Architecture, Department
of Electrical and Electronics Engineering, Burdur, Türkiye

ORCID ID: <https://orcid.org/0000-0002-9625-9513>

Assist. Prof. Dr. Mesut ÇOŞLU

Burdur Mehmet Akif Ersoy University, Faculty of Engineering and Architecture, Department
of Landscape Architecture, Burdur, Türkiye

ORCID ID: <https://orcid.org/0000-0003-3952-6563>

Assist. Prof. Dr. Mete YİĞİT

Burdur Mehmet Akif Ersoy University, Faculty of Engineering and Architecture, Department
of Mechanical Engineering, Burdur, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6500-2002>

Assoc. Prof. Dr. İlker ÜNAL

Burdur Mehmet Akif Ersoy University, Faculty of Engineering and Architecture, Department
of Mechanical Engineering, Burdur, Türkiye

ORCID ID: <https://orcid.org/0000-0002-5188-4438>

ABSTRACT

Accurately determining fruit ripeness levels is critical in agricultural productivity, quality control, and harvest scheduling. The precise detection of orange fruit ripeness helps reduce yield losses, providing economic benefits to producers while ensuring higher-quality products for consumers. Traditional methods primarily rely on manual inspections, which are time-consuming and prone to human error. Therefore, deep learning-based approaches offer more automated, fast, and precise solutions for fruit ripeness detection. This study aims to evaluate the effectiveness of instance segmentation-based deep learning models for detecting the ripeness levels of orange fruit. In this study, an instance segmentation approach was employed to determine the ripeness levels of orange fruit. Three state-of-the-art single-stage deep learning models YOLOv8n-seg, YOLOv9c-seg, and YOLO11n-seg were trained and their performances were compared. The dataset consisted of 1000 images, comprising 500 unripe and 500 full ripened orange fruit images collected under real-world conditions. The images were divided into training, validation, and test sets, ensuring a balanced distribution of ripeness levels. Standard deep learning techniques such as data augmentation and hyperparameter tuning were applied to enhance model performance. The models were evaluated using mean Average Precision (mAP), Precision, and Recall metrics. The results demonstrated that the highest performance was achieved using the YOLO11n-seg model. This model obtained 0.984 mAP@0.5, 0.96 mAP@0.5-0.95, 0.991 Precision, and 0.938 Recall, outperforming the other

models. These findings indicate that YOLO11n-seg is the most effective deep learning model for detecting orange fruit ripeness levels. The findings of this study are expected to contribute to the development of automated quality control systems in agricultural production and assist in optimizing harvest timing based on ripeness levels. Furthermore, integrating this approach with autonomous agricultural technologies, such as drones and robotic systems, is anticipated to facilitate real-time applications in the field. Future studies will focus on improving the generalization capability of the model by utilizing larger datasets and developing low-cost, portable, and scalable systems for practical implementation in precision agriculture.

Keywords: Artificial Intelligence, Citrus, Deep Learning, Instance Segmentation, State-of-the-Art (SOTA) Model.

INTRODUCTION

The world population is expected to exceed 9 billion in the next 25 years. This population growth will also require an increase of around 70% in agricultural production (Ben Ayed and Hanana, 2021). Agriculture, which makes significant contributions to human and animal nutrition, is a critical component of the ecosystem. The most important challenges of agricultural production are the increasing world population, climate change and food security. While struggling with these challenges, producers need to pay attention to issues such as environmental pollution and water conservation, as well as achieving minimum cost and maximum income. In this context, precision agriculture has come to the fore in recent years, especially with the use of information and communication technologies in agriculture (Viana et al., 2022; Aslan et al., 2022).

Citrus fruits are an important evergreen plant group belonging to the Aurantioideae subfamily of the Rutaceae family, including high economic value varieties such as orange, tangerine, lemon, grapefruit, bitter orange, kumquat, citron, shaddock and bergamot. Citrus fruits, which are native to Southeast Asia, are cultivated in temperate and subtropical climates. Approximately 70% of the production is carried out in the northern hemisphere, mostly in the Mediterranean Region and the United States. Citrus fruits, which have great economic importance, contain high levels of vitamin C. In addition to being consumed fresh, the fruits of these plants are also used for juice, jam and medicinal purposes. In addition, citrus fruit wastes are also used as biofuel today (TEPGE, 2024; Hamid et al., 2024). Citrus fruits are an important agricultural product on a global scale due to their wide variety of species, storage properties and use in different industries. The most commonly used citrus fruits commercially are orange, tangerine, lemon and grapefruit. These fruits are important agricultural products and also contribute significantly to the economy (Turgut and Ersöz, 2024).

The increase in citrus cultivation, which is an agricultural product that brings high economic profit, has brought with it some difficulties in recent years. In this context, growers may face significant difficulties such as the expansion of citrus production areas, the spread of diseases and pests due to climate change, and the excessive use of chemicals in control (Yu et al., 2024). Citrus pests pose a significant threat to the plant in terms of both yield and fruit quality. Early prevention of damage to citrus is very important in terms of sustainable cultivation, reducing costs, and especially contributing to the reduction of environmental pollution (Wu et al., 2024).

Deep learning techniques have been widely used in agricultural fields in recent years with the development of image sensors and computer vision technology. Especially in citrus cultivation, studies on the detection of diseases and pests and the damage they cause to the plant and the determination of the maturity levels of fruits come to the fore (Xing et al., 2019; Wu et al., 2024; Yu et al., 2024). The use of this technology has made significant contributions to agricultural applications by resulting in cost savings, increased detection efficiency and

profitability. Popular deep learning architectures used in the detection of pests are deep autoencoders (DAE) and convolutional neural networks (CNN) (Wu et al., 2024).

This study aims to evaluate the effectiveness of instance segmentation-based deep learning models for detecting the ripeness levels of orange fruit. In this study, an instance segmentation approach was employed to determine the ripeness levels of orange fruit. Three state-of-the-art single-stage deep learning models YOLOv8n-seg, YOLOv9c-seg, and YOLO11n-seg were trained and their performances were compared.

Related Works

The identification and categorization of fruit ripeness have historically been important fields of study in the agricultural and food sectors. Conventional approaches have frequently been time-consuming, expensive, and labor-intensive, constraining their industrial applicability (Taniwaki et al, 2009). As a result, contemporary methods like non-destructive testing and computer vision technologies have been progressively utilized for evaluating fruit maturity (Nishani et al., 2017).

The detection of fruit ripeness and quality assessment are extensively studied domains utilizing computer vision and deep learning methodologies. Computer vision is a field that examines picture and video data to execute tasks such object detection, categorization, and location. These operations are generally executed utilizing bounding boxes. Deep learning models, especially convolutional neural networks (CNNs), have shown elevated accuracy levels in object detection tasks.

Liu et al. (2019) employed HOG (Histogram of Oriented Gradients) descriptors to train SVM (Support Vector Machine) classifiers, mitigating the impact of fluctuating lighting conditions on tomato recognition. Li et al. (2017) integrated Fast Normalized Cross Correlation (FNCC) with Hough Transform detection to facilitate the identification of green tomatoes when the stem background color was analogous. Kurtulmuş et al. (2011) utilized color characteristics and Gabor filtering techniques for the identification of immature citrus fruits. Ashtiani et al. (2021) employed transfer learning to optimize a CNN model for detecting mulberry ripeness, yet the model exhibited prolonged and inefficient detection times. Appe et al. (2023) utilized a VGG16-based DCNN model for tomato ripeness detection but identified challenges with image detection in intricate backgrounds and emphasized the necessity to enhance the model's generalization ability.

YOLO (You Only Look Once) is a notable algorithm in object detection. YOLO analyzes the complete image simultaneously to forecast bounding boxes and class probabilities (Majumder and Wilmot, 2023). This methodology enhances YOLO's speed and efficiency relative to two-stage detectors such as Faster R-CNN (Diwan et al., 2023). Various iterations of YOLO have been created over time, integrating architectural enhancements and novel functionalities. For example, YOLOv3 attained enhanced object detection efficacy by multi-scale detection and a more profound architecture (Redmon et al., 2016). YOLOv4, on the CSPDarknet53 backbone, provided enhanced accuracy and velocity (Bochkovskiy et al., 2020). These developments have facilitated the extensive application of YOLO in agriculture, including plant disease diagnosis, harvest optimization, and fruit freshness assessment.

Research on many fruit varieties has shown the object identification and classification abilities of YOLO-based deep learning models. Zhou et al. (2024) introduced YOLOv5-NMM (YOLOv5 with Navel Orange Measure Model), an object detection model based on YOLOv5, designed for the rapid and precise assessment of ripeness levels in Gannan navel oranges within natural settings. The model, augmented with enhancements to the original YOLOv5, seeks to attain superior accuracy and performance in identifying orange ripeness stages. The research

tackled issues including occlusion, dense distribution, tiny target dimensions, inclement weather, and fluctuations in lighting within natural settings. Experimental findings indicated that the YOLOv5-NMM model attained an accuracy of 93.2%, a recall of 89.6%, and a mean average precision (mAP) of 94.2%. The model surpassed other prominent models, including Faster R-CNN, YOLOv3, original YOLOv5, and YOLOv7-tiny, regarding accuracy and parameter efficiency.

The YOLOv7 model attained a mean average precision (mAP) of 78.1% in detecting banana ripeness (Antony and Fajar, 2024). In a separate investigation employing the YOLOv9 model for detecting avocado ripeness, the model attained a mean Average Precision (mAP) of 84.3% (Widiati and Haryanto, 2024). Likewise, the YOLOv5 and DeepSORT algorithms were employed to assess the ripeness levels and quantify sweet peppers in greenhouses, attaining a mean Average Precision (mAP) of 80.3% (Escamilla et al., 2024). A separate investigation on sugar apple fruit with the DeepLabv3+ model attained an accuracy rate of 96.6% (Xie et al., 2024). Furthermore, the integration of YOLOv9 with the Swin Transformer model for the identification of strawberry maturity achieved a mean Average Precision (mAP) of 87.3% (Wang et al., 2024).

Equitable data dissemination and the application of data preparation techniques are prevalent elements in all investigations. Methods for image data augmentation and annotation were implemented, and model training processes were refined (Antony and Fajar, 2024; Xie et al., 2024). Nonetheless, obstacles persist in the extensive implementation of computer vision and deep learning models in agriculture. These factors encompass uneven datasets, fluctuating ambient conditions (e.g., variations in illumination and background), and the identification of diminutive items. Subsequent study needs to concentrate on employing larger and more equitable datasets to enhance model efficacy, investigating adaptability to various fruit varieties, and creating lightweight models for real-time applications (Xie et al., 2024; Widiati and Haryanto, 2024).

MATERIALS AND METHODS

Data Acquisition

In this study, the dataset was collected between December 14-15 and December 21 from orange orchards located in Manavgat, Antalya. A total of 1000 images were acquired using Xiaomi POCO X3 Pro and Xiaomi Redmi Note 8 Pro smartphones. The images were stored in .jpeg format with resolutions of 3000×3000 and 3472×3472 pixels, maintaining a 1:1 aspect ratio. To enhance the success and generalization ability of the deep learning algorithm, images were captured under various natural lighting conditions. The image acquisition process was conducted in carefully selected orchard areas where no visible damage or external factors could affect the dataset. Furthermore, no pesticides or chemical treatments had been applied in these orchards before the image collection process.

Data Preprocessing and Annotation

In this study, Roboflow was used for image preprocessing, annotation, and data augmentation. The dataset consists of unripe and full ripened orange images, as shown in Figure 1. Two annotation classes were defined: “Unripe” and “Full Ripened”, each containing 500 images. Before training, all images with original resolutions of 3000×3000 and 3472×3472 pixels were resized to 640×640 pixels using Roboflow. The dataset was then split into training, validation, and test sets with a ratio of 7:2:1. After annotation and partitioning, the dataset was exported in compressed (zip) YOLO format for model training.




Fig. 1. The Dataset Structure Generated in The Study

The image labeling process was performed manually in the initial stage using a bounding box annotation tool. To evaluate the accuracy and distribution of the labeled dataset, statistical analyses were conducted, and the results are presented in Figure 2. Figure 2a shows the number of instances per class, ensuring a balanced representation of the "Unripe" and "Full Ripened" categories. Figure 2b illustrates the bounding box sizes, offering insight into the variations in object dimensions. Figure 2c presents the normalized position distribution of labeled objects, providing a visualization of how annotations are spread across the images. Figure 2d depicts the normalized label size distribution, revealing a linear correlation between width and height, with a higher density of smaller bounding boxes. These analyses help assess dataset consistency and ensure the annotations align with model training requirements.




Fig. 2. Statistical analysis of the bounding box labels in the dataset includes a) the total number of labels per class, b) the distribution of label sizes, c) the normalized location map of labeled targets, and d) the normalized size distribution of the labeled objects

The accuracy and distribution of the bounding box labels were further analyzed using a correlogram, as shown in Figure 3. This visualization presents the relationships between label position (x , y) and label size (width, height) through scatter plots and histograms. The diagonal histograms illustrate the individual distributions of each variable, while the scatter plots reveal the correlations between them. The strong positive correlation between width and height suggests that bounding boxes tend to maintain proportional scaling. The label positions (x , y) exhibit a dense and centered distribution, indicating that most annotations are concentrated in specific regions of the images. These results confirm that the dataset is well-structured, consistently labeled, and does not contain irregular or misaligned annotations, ensuring its quality for deep learning model training.




Fig. 3. Correlation distribution graphs of bounding box labels in the dataset

Single-Stage Instance Segmentation Models

The proposed model, YOLO11 (Jocher and Qiu, 2024), developed by Ultralytics, builds upon the capabilities of YOLOv8, supporting functions such as instance segmentation, pose estimation, and oriented object detection, while offering five scalable versions tailored for different applications. Instead of the C2f block used in YOLOv8, YOLO11 integrates the more efficient C3K2 block, enhancing overall performance without compromising computational speed. Additionally, it incorporates the C2PSA (Cross Stage Partial with Spatial Attention) module, which strengthens spatial feature representation, particularly improving accuracy in detecting small and overlapping objects. (Jegham et al., 2024). The general architectural structure of the YOLO11-seg model is presented in Figure 4, providing an overview of its components and functionality.




Fig. 4. General Structure of the YOLO11-seg Model Architecture (Jocher and Qiu, 2024; Tsoi, 2025)

Experimental Environment and Parameters

To assess the performance of the study, a MacBook Pro 2012 (Apple Inc., Cupertino, California, USA) and a Google Colab (Mountain View, California, USA) notebook were utilized as the deep learning environment. The experimental setup was implemented within Colab, leveraging a virtual Nvidia Tesla T4 GPU (Santa Clara, California, USA) for accelerated computations. The detailed configuration of the experimental environment is presented in Table 1, while the hyperparameters used in the training phase are outlined in Table 2.

Table 1. Configuration and training environment

Device & Software	Environmental Parameter	Value
Apple MacBook Pro 2012	Operating system	Windows 10
	CPU	Intel Core i5
	Ram	16 G- LPDDR3L
Google Colab	Deep learning framework	Pytorch
	Programming language	Python3.10
	Virtual ram	90 G
	Virtual storage	250 G
Virtual GPU (Nvidia Tesla T4)	Memory	16G
	Bandwidth	320 G/s
	Cuda Core	2560

Table 2. Hyperparametric configuration

Parameters	Value
Image-size	640
Epochs	25
Batch-size	8
Optimizer	SGD
Momentum	0.937
lr	0.01
Weight_decay	0.0005
Warmup_epochs	3
Warmup_momentum	0.8
Warmup_bias_lr	0.1

Model Evaluation Indicators

In the experimental phase of the study, the model's box and mask accuracy was evaluated using $mAP@0.5$ and $mAP@0.5-0.95$ metrics. The calculations were performed based on Equation (1) for $mAP@0.5$ and Equation (2) for $mAP@0.5-0.95$, ensuring a comprehensive assessment of detection and segmentation performance (Wu et al., 2023; Zhang et al., 2023)

$$mAP_{0.5} = \frac{1}{n_c} \int_0^1 P(R) dR \quad (1)$$

$$mAP_{0.5-0.95} = avg(mAP_i), i = 0.5:0.05:0.95 \quad (2)$$

In the equation, "nc" denotes the number of classes, "P" represents precision, and "R" stands for recall. The precision (P) is computed using Equation (3), while the recall (R) is determined based on Equation (4) (Zhang et al., 2023).

$$P = \frac{TP}{TP + FP} \quad (3)$$

$$R = \frac{TP}{TP + FN} \quad (4)$$

TP (True Positive) refers to correctly detected objects, representing the number of predicted bounding boxes with $\text{IoU} > 0.5$. FP (False Positive) indicates incorrectly detected objects, including prediction boxes with an $\text{IoU} \leq 0.5$. FN (False Negative) represents undetected labels, indicating missed objects in the images. These metrics are essential for evaluating the model's effectiveness in accurately detecting and segmenting objects (Mohana Sri et al., 2023). Figure 5 presents a detailed workflow diagram of the study. As shown in Figure 5, the processes of data acquisition, data uploading, annotation, and preprocessing have been carried out. In the next section, model training and detection performance analysis results are provided in detail.




Fig. 5. Workflow of the Study

RESULTS AND ANALYSIS

As shown in Figure 6, the class-based evaluation results indicate that the YOLO11n-seg model demonstrated high precision (P), recall (R), and mean Average Precision (mAP) values, achieving strong performance in both Unripe and Full Ripened categories. The Unripe class achieved a precision of 1.000, indicating that all detected instances were correctly classified. The recall value of 0.962 showed that only a few instances were missed. Additionally, $\text{mAP}@0.5$ and $\text{mAP}@0.5-0.95$ values were calculated as 0.985 and 0.971, respectively, confirming the model's high accuracy in distinguishing unripe oranges. The evaluation of the Full Ripened class showed a precision of 0.982, while the recall was 0.915, indicating that some ripened oranges were not detected. However, $\text{mAP}@0.5$ and $\text{mAP}@0.5-0.95$ values were calculated as 0.983 and 0.950, respectively, demonstrating that the model still performed well in segmenting ripened fruits. When all classes were evaluated together, $\text{mAP}@0.5$ and

mAP@0.5-0.95 values were calculated as 0.984 and 0.960, respectively. These values indicate that the model provides high reliability in terms of overall accuracy and successfully detects all orange ripeness levels.




Fig. 6. Class-Based Evaluation Results of the YOLO11n-seg Model

As shown in Figure 7, the training performance of the YOLOv8n-seg, YOLOv9c-seg, and YOLO11n-seg models was evaluated based on mAP@0.5 and segmentation loss across epochs. The mAP@0.5 curve, as shown in Figure 7a, demonstrates that all three models have reached high accuracy levels, with YOLO11n-seg achieving the most stable and consistent performance throughout training. While YOLOv9c-seg started with a lower initial accuracy, it gradually improved over the epochs, catching up with the other models. As presented in Figure 7b, the segmentation loss curve on the right shows that all models experienced a rapid decrease in loss within the initial epochs, indicating effective learning. YOLO11n-seg maintained the lowest and most stable loss values, confirming its strong convergence and robustness. Although YOLOv9c-seg exhibited higher loss in the early stages, it progressively reduced over time, but with more fluctuations compared to the other models. According to the obtained results, YOLO11n-seg outperforms the other models in both accuracy and stability, making it the most reliable model for orange fruit ripeness segmentation. The stable mAP@0.5 and low segmentation loss values indicate that YOLO11n-seg effectively balances precision and learning efficiency, demonstrating superior generalization in the training process.




Fig. 7. (a) mAP@0.5 (seg) values across epochs, (b) segmentation loss across epochs for different models

As shown in Figure 8, the overall performance of the YOLOv8n-seg, YOLOv9c-seg, and YOLO11n-seg models was evaluated using Precision (P), Recall (R), mAP@0.5, and mAP@0.5-0.95 metrics. YOLO11n-seg achieved the highest performance, with precision, recall, mAP@0.5, and mAP@0.5-0.95 values calculated as 0.991, 0.938, 0.984, and 0.960, respectively, demonstrating its strong detection and segmentation capabilities. YOLOv9c-seg followed closely with 0.998, 0.952, 0.979, and 0.965, while YOLOv8n-seg obtained 0.988, 0.937, 0.977, and 0.964, respectively. These results indicate that, while all three models performed well, YOLO11n-seg provided the most balanced and stable accuracy across all metrics, confirming its effectiveness in detecting orange fruit ripeness levels.




Fig. 8. Performance Evaluation Results of Different Models

As shown in Figure 9, the GPU usage and inference time of the YOLOv8n-seg, YOLOv9c-seg, and YOLO11n-seg models were analyzed to evaluate computational efficiency. YOLO11n-Seg demonstrated the lowest GPU usage and inference time, making it the most efficient model in terms of resource consumption. GPU usage for YOLO11n-seg, YOLOv8n-seg, and YOLOv9c-seg was recorded as 2.7G, 3.0G, and 11.9G, respectively, indicating that YOLOv9c-seg required significantly more memory, while YOLO11n-seg had the most optimized memory consumption. Similarly, inference time was measured as 6.6 ms, 6.9 ms, and 7.7 ms, respectively, showing that YOLO11n-seg provided slightly faster inference compared to the other models. These results suggest that YOLO11n-seg achieves an optimal balance between

computational efficiency and accuracy, making it the most suitable model for real-time applications with limited hardware resources.




Fig. 9. GPU Usage and Inference Time Comparison of Different Models

As shown in Figure 10, the YOLO11n-seg model has successfully detected oranges in the test dataset, even in challenging conditions where the fruit has been partially visible or obscured behind leaves. The top row has presented the original images, while the bottom row has illustrated the model's instance segmentation results. The model has accurately identified both unripe and full ripened oranges, demonstrating their robustness in detecting fruits with varying lighting conditions, occlusions, and different orientations. Some detections have highlighted the model's ability to recognize oranges even when parts of the fruit have been hidden behind leaves or only partially visible in the frame.



Fig. 10. Examples of Detections from the Test Dataset Using the YOLO11n-seg Model

DISCUSSION

This study demonstrates the effectiveness of YOLO11n-seg in detecting orange fruit ripeness levels using instance segmentation, achieving high accuracy while maintaining computational efficiency. Compared to the YOLOv5-NMM model proposed by Zhou et al. (2024), which was trained on 3000 images and achieved precision, recall, and mAP@0.5 values of 0.932, 0.896, and 0.942, respectively, our method was trained on a smaller dataset of 1000 images but

outperformed it with 0.991, 0.938, and 0.984, respectively. The YOLO11n-seg model's higher accuracy highlights the advantage of using instance segmentation over traditional object detection, allowing for better localization of partially occluded and overlapping fruits. When compared to the YOLO-CIT model combined with the R-LBP algorithm (Wang et al., 2024), which achieved precision, recall, and mAP@0.5 values of 0.881, 0.931, and 0.858, respectively, with an inference speed of 6.1ms, our proposed method, YOLO11n-seg, demonstrated superior performance with precision, recall, and mAP@0.5 values of 0.991, 0.938, and 0.984, respectively, while maintaining a faster inference time of 6.6ms. This indicates that YOLO11n-seg not only improves detection accuracy but also maintains computational efficiency, making it suitable for real-time applications in agricultural automation.

CONCLUSION

This study presents a deep learning-based instance segmentation approach for detecting orange fruit ripeness levels, achieving high accuracy and computational efficiency. The YOLO11n-seg model demonstrated superior performance compared to other state-of-the-art models, achieving precision, recall, mAP@0.5, and mAP@0.5-0.95 values of 0.991, 0.938, 0.984, and 0.960, respectively, while maintaining an efficient inference time of 6.6ms. Unlike traditional object detection methods that rely on bounding boxes, this study utilized instance segmentation, enabling more precise localization and detection of partially occluded and overlapping fruits. The dataset used in this study, consisting of 1000 images of unripe and fully ripened oranges captured under natural conditions, ensured a balanced and diverse representation for training, improving the model's generalization ability. The inclusion of small, overlapping, and partially visible fruits further enhanced the model's robustness. The experimental results confirm that YOLO11n-seg effectively detects fruit ripeness with high precision, making it suitable for real-time agricultural applications. Future research should focus on expanding the dataset with more fruit varieties and environmental conditions to further enhance model adaptability. Additionally, optimizing the model for mobile and edge computing devices will facilitate real-time deployment in precision agriculture and automated harvesting systems. The findings of this study reinforce the importance of deep learning-based segmentation models in improving fruit quality assessment and harvest optimization, contributing to the advancement of smart agricultural technologies.

REFERENCES

- Antony, D., & Fajar, A. N. (2024). Object detection using convolutional neural network YOLOv7 to detect banana ripeness. *Journal of Theoretical and Applied Information Technology*, 31(6).
- Appa, S. R. N., Arulselvi, G., & Balaji, G. (2023). Tomato ripeness detection and classification using VGG based CNN models. *International Journal of Intelligent Systems and Applications in Engineering*, 11, 296–302.
- Aslan, M. F., Durdu, A., Sabanci, K., Ropelewska, E., & Gültekin, S. S. (2022). A Comprehensive Survey of Recent Studies with UAV for Precision Agriculture in Open Fields and Greenhouses. *Applied Sciences*, 12(3), 1047. <https://doi.org/10.3390/app12031047>
- Ashtiani, S. H. M., Javanmardi, S., Jahanbanifard, M., Martynenko, A., & Verbeek, F. J. (2021). Detection of mulberry ripeness stages using deep learning models. *IEEE Access*, 9, 100380–100394.

- Aytaş, M., Yumruktepe, R., & Mart, C. (2001). Using Pheromone Traps to Control California Red Scale *Aonidiella aurantii* (Maskell)(Hom.: Diaspididae) in the Eastern Mediterranean Region. *Turkish Journal of Agriculture and Forestry*, 25(2), 97-110.
- Ben Ayed, R., & Hanana, M. (2021). Artificial intelligence to improve the food and agriculture sector. *Journal of Food Quality*, 2021(1), 5584754. <https://doi.org/10.1155/2021/5584754>
- Bochkovskiy, A., Wang, C. Y., & Liao, H. Y. M. (2020). YOLOv4: Optimal speed and accuracy of object detection. *arXiv preprint arXiv:2004.10934*.
- Diwan, T., Anirudh, G., & Tembhurne, J. V. (2023). Object detection using YOLO: Challenges, architectural successors, applications. *Multimedia Tools and Applications*, 82(6), 9243–9275.
- Fonte, A., Garcerá, C., Tena, A., & Chueca, P. (2021). Volume Rate Adjustment for Pesticide Applications against *Aonidiella aurantii* in Citrus: Validation of CitrusVol in the Growers' Practice. *Agronomy*, 11(7), 1350. <https://doi.org/10.3390/agronomy11071350>
- García Morales, M., Denno, B. D., Miller, D. R., Miller, G. L., Ben-Dov, Y., & Hardy, N. B. (2016). ScaleNet: a literature-based model of scale insect biology and systematics. *Database*, 2016, bav118.
- Golan, K., Kot, I., Kmiec, K., & Górska-Drabik, E. (2023). Approaches to Integrated Pest Management in Orchards: *Comstockaspis pernicios*a (Comstock) Case Study. *Agriculture*, 13(1), 131. <https://doi.org/10.3390/agriculture13010131>
- Hamid, S., Sharma, K., Kumar, K., & Thakur, A. (2024). Types and Cultivation of Citrus Fruits. In *Citrus Fruits and Juice: Processing and Quality Profiling* (pp. 17-43). Singapore: Springer Nature Singapore. https://doi.org/10.1007/978-981-99-8699-6_2
- Jacas, J. A., Karamaouna, F., Vercher, R., & Zappalà, L. (2010). Citrus pest management in the northern Mediterranean basin (Spain, Italy and Greece). *Integrated management of arthropod pests and insect borne diseases*, 3-27.
- Jegham, N., Koh, C.Y., Abdelatti, M., Hendawi, A., (2024). Evaluating the evolution of YOLO (You Only Look Once) models: A comprehensive benchmark study of YOLO11 and its predecessors. *arXiv (Cornell University)*. doi:10.48550/arxiv.2411.00201.
- Jocher, G., & Qiu, J. (2024). Ultralytics YOLO11 (Version 11.0.0) [Software]. Retrieved from <https://github.com/ultralytics/ultralytics>.
- Kurtulmuş, F., Lee, W. S., & Vardar, A. (2011). Green citrus detection using 'eigenfruit', color and circular Gabor texture features under natural outdoor conditions. *Computers and Electronics in Agriculture*, 78, 140–149.
- Li, H., Zhang, M., Gao, Y., Li, M., & Ji, Y. (2017). Green ripe tomato detection method based on machine vision in greenhouse. *Transactions of the Chinese Society of Agricultural Engineering*, 33, 328–334.
- Liu, L., Li, Z., Lan, Y., Shi, Y., & Cui, Y. (2019). Design of a tomato classifier based on machine vision. *PLoS ONE*, 14(e0219803).
- Majumder, M., & Wilmot, C. (2023). Automated vehicle counting from pre-recorded video using You Only Look Once (YOLO) object detection model. *Journal of Imaging*, 9(7), 131.
- Mohana Sri, S., Swethaa, S., Aouthithiye Barathwaj, S. R. Y., & Sai Ganesh, C. S. (2023). Intelligent Debris Mass Estimation Model for Autonomous Underwater Vehicle. *Computer Vision and Pattern Recognition*. <https://doi.org/10.48550/arXiv.2309.10617>

- Nishani, S., Deshpande, S., & Gundewadi, G. (2017). Use of acoustics as non-destructive techniques: A review. *International Journal of Current Microbiology and Applied Sciences*, 6, 2468–2476. <https://doi.org/10.20546/ijcmas.2017.606.293>
- Normark, B. B., Morse, G. E., Krewinski, A., & Okusu, A. (2014). Armored scale insects (Hemiptera: Diaspididae) of San Lorenzo National Park, Panama, with descriptions of two new species. *Annals of the Entomological Society of America*, 107(1), 37-49.
- Tsoi, P. P. (2024). YOLO11: The cutting-edge evolution in object detection—A brief review of the latest in the YOLO series. <https://medium.com>. (Accessed date: 2025-03-3).
- Pekas, A., Aguilar, A., Tena, A., & Garcia-Marí, F. (2010). Influence of host size on parasitism by *Aphytis chrysomphali* and *A. melinus* (Hymenoptera: Aphelinidae) in Mediterranean populations of California red scale *Aonidiella aurantii* (Hemiptera: Diaspididae). *Biological Control*, 55(2), 132-140. <https://doi.org/10.1016/j.biocontrol.2010.07.010>
- Pekas, A., Navarro-Llopis, V., Garcia-Marí, F., Primo, J., & Vacas, S. (2015). Effect of the California red scale *Aonidiella aurantii* sex pheromone on the natural parasitism by *Aphytis* spp. in Mediterranean citrus. *Biological Control*, 90, 61-66. <https://doi.org/10.1016/j.biocontrol.2015.05.016>
- Pina, T. (2007). Control biológico del piojo rojo de California, *Aonidiella aurantia* (Maskell)(Hemiptera: Diaspididae) y estrategias reproductivas de su principal enemigo natural *Aphytis chrysomphali* (Mercet)(Hymenoptera: Aphelinidae) (Doctoral dissertation, Universitat de València).
- Redmon, J., Divvala, S., Girshick, R., & Farhadi, A. (2016). You only look once: Unified, real-time object detection. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition* (pp. 779–788). Las Vegas, NV, USA.
- Roelofs, W. L., Gieselmann, M. J., Cardé, A. M., Tashiro, H., Moreno, D. S., Henrick, C. A., & Anderson, R. J. (1977). Sex pheromone of the California red scale, *Aonidiella aurantii*. *Nature*, 267(5613), 698-699.
- Rodrigo, E., Troncho, P., & García-Marí, F. (1996). Parasitoids (Hym.: Aphelinidae) of three scale insects (Hom.: Diaspididae) in a citrus grove in Valencia, Spain. <https://doi.org/10.1007/BF02893295>
- Taniwaki, M., Takahashi, M., & Sakurai, N. (2009). Determination of optimum ripeness for edibility of postharvest melons using nondestructive vibration. *Food Research International*, 42(1), 137–141. <https://doi.org/10.1016/j.foodres.2008.09.007>
- TEPGE (2024). Turunçgiller Ürün Raporu. <https://arastirma.tarimorman.gov.tr/tepge/Belgeler/PDF%20C3%9Cr%C3%BCn%20Raporlar%C4%B1/2024%20C3%9Cr%C3%BCn%20Raporlar%C4%B1/Turun%C3%A7giller%20C3%9Cr%C3%BCn%20Raporu%202024-408%20TEPGE.pdf> (Access date: 2025-02-6)
- Turgut, M., & Ersöz, C. (2024). Narenciye Sektörünün İhracat Performansı: Hedef Pazar Analizi. *Tarsus Üniversitesi Uygulamalı Bilimler Fakültesi Dergisi*, 4(1), 62-79.
- Vacas, S., Alfaro, C., Primo, J., & Navarro-Llopis, V. (2015). Deployment of mating disruption dispensers before and after first seasonal male flights for the control of *Aonidiella aurantii* in citrus. *Journal of Pest Science*, 88, 321-329.
- Vanaclocha, P., Urbaneja, A., & Verdú, M. J. (2009). Mortalidad natural del piojo rojo de California, *Aonidiella aurantii*, en cítricos de la Comunidad Valenciana y sus parasitoides asociados. *Boletín de sanidad vegetal. Plagas*, 35(1), 59-71.

- Viana, C. M., Freire, D., Abrantes, P., Rocha, J., & Pereira, P. (2022). Agricultural land systems importance for supporting food security and sustainable development goals: A systematic review. *Science of the total environment*, 806, 150718. <https://doi.org/10.1016/j.scitotenv.2021.150718>
- Viveros Escamilla, L. D., Gómez-Espinosa, A., Escobedo Cabello, J. A., & Cantoral-Ceballos, J. A. (2024). Maturity recognition and fruit counting for sweet peppers in greenhouses using deep learning neural networks. *Agriculture*, 14, 331. <https://doi.org/10.3390/agriculture14030331>
- Widiati, W., & Haryanto, T. (2024). Deep learning for automatic classification of avocado fruit maturity. *Jurnal Pilar Nusa Mandiri*, 20(1), 75–80. <https://doi.org/10.33480/pilar.v20i1.5043>
- Wu, Y., Han, Q., Jin, Q., Li, J., & Zhang, Y. (2023). LCA-YOLOv8-Seg: An Improved Lightweight YOLOv8-Seg for Real-Time Pixel-Level Crack Detection of Dams and Bridges. *Appl. Sci.*, 13, 10583. <https://doi.org/10.3390/app131910583>
- Wu, X., Liang, J., Yang, Y., Li, Z., Jia, X., Pu, H., & Zhu, P. (2024). SAW-YOLO: A Multi-Scale YOLO for Small Target Citrus Pests Detection. *Agronomy*, 14(7), 1571. <https://doi.org/10.3390/agronomy14071571>
- Xie, Z., Ke, Z., Chen, K., Wang, Y., Tang, Y., & Wang, W. (2024). A lightweight deep learning semantic segmentation model for optical image-based post-harvest fruit ripeness analysis of sugar apples (*Annona squamosa*). *Agriculture*, 14, 591. <https://doi.org/10.3390/agriculture14040591>
- Xing, S., Lee, M., & Lee, K.-k. (2019). Citrus Pests and Diseases Recognition Model Using Weakly Dense Connected Convolution Network. *Sensors*, 19(14), 3195. <https://doi.org/10.3390/s19143195>
- Yu, Y.-N., Xiong, C.-L., Yan, J.-C., Mo, Y.-B., Dou, S.-Q., Wu, Z.-H., & Yang, R.-F. (2024). Citrus Pest Identification Model Based on Improved ShuffleNet. *Applied Sciences*, 14(11), 4437. <https://doi.org/10.3390/app14114437>
- Zhang, L., Ding, G., Li, C., & Li, D. (2023) DCF-Yolov8: An Improved Algorithm for Aggregating Low-Level Features to Detect Agricultural Pests and Diseases. *Agronomy*, 13, 2012. <https://doi.org/10.3390/agronomy13082012>.
- Zhou, B., Wu, K., & Chen, M. (2024). Detection of Gannan navel orange ripeness in natural environment based on YOLOv5-NMM. *Agronomy*, 14, 910. <https://doi.org/10.3390/agronomy14050910>

**THE ROLE OF LOCAL GOVERNMENTS IN SUSTAINABLE AGRICULTURE:
THE ISTANBUL CASE**

**SÜRDÜRÜLEBİLİR TARIMDA YEREL YÖNETİMLERİN ROLÜ: İSTANBUL
ÖRNEĞİ**

İlknur İŞÇİ

Ondokuz Mayıs University, Graduate Education Institute, Department of Field Crops,
Samsun, Turkey

ORCID ID: <https://orcid.org/0009-0005-7481-7202>

Gülcan KAYMAK BAYRAM

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: <https://orcid.org/0000-0002-0915-0529>

Zeki ACAR

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: : <https://orcid.org/0000-0002-0484-1961>

İlknur AYAN

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: : <https://orcid.org/0000-0002-5097-9013>

ÖZET

Tarımsal üretim insanlığın geleceği açısından ikamesi olmayan bir faaliyettir. Ancak yapılan gözlemlere ve eldeki verilere göre çok uzak olmayan bir gelecekte üretimde ciddi sorunların yaşanacağı görülmektedir. Bunun gerçekleşmesini önlemek için de tutarlı üretimin sürdürülebilir hale getirilmesi gerekmektedir. Dünya nüfusunun hızlıca artması ve teknolojik gelişmelerin baş döndürücü bir hızla ilerlemesi tarımsal üretimde insanoğlunu, birim alandan yüksek verim almaya yöneltmiştir. Buna bağlı olarak kullanılan sistemler ve mücadele yöntemlerinde yüksek oranda kimyasal kullanılmaya başlanmıştır. Özellikle bitki besleme ve tarımsal mücadelede önemli oranda yapay kimyasal kullanımı yaygınlaşmıştır. Bu durum verimi önemli oranda artırmış fakat çevre ve sağlık açısından riskler ortaya çıkmaya başlamıştır. Aşırı suni gübre ve kimyasal ilaçların kullanımı çevre problemlerini de beraberinde getirmiştir. Birçok yerde toprak verimliliğini yitirmiş, çoraklaşmış, kimyasal gübreler ve ilaçlar sebebiyle elde edilen ürünler insan sağlığını tehdit eder duruma gelmiştir. Özellikle gelişmekte olan ülkeler başta olmak üzere, tarımsal girdi fiyatlarının aşırı yükselmesi, buna karşın ürün fiyatlarının yeterince artmaması ve sağlanan desteklerin yetersiz kalması nedeniyle çok sayıda

küçük üretici tarımdan kopmuş ve büyük kentlerin kenar mahallelerinde tüketici konumuna gelmiştir. Bu gelişmeler tarımda sürdürülebilirlik kavramını ön plana çıkarmıştır.

Yerel yönetimlerin tarımsal kalkınmadaki ve tarımın sürdürülebilirliğindeki rolü büyüktür. Yerel yönetimler, ildeki tarım sektörünü destekleyerek hem şehrin hem de ülkenin ekonomisine katkı sağlar. Yerel yönetimler ayrıca çevreyi koruyarak sürdürülebilir bir tarım modeli oluşturur. Yerel yönetimlerin vizyon ve stratejilerini belirlerken tarımsal faaliyetleri göz önünde bulundurmaları gerekir. Yerel yönetimlerin tarımsal kalkınma faaliyetleri için belediyeler, valilik ve il özel idareleri arasında iş birliği yapmaları gerekir. Yerel yönetimlerin tarımsal kalkınma faaliyetleri hem üreticiyi hem de tüketiciyi korur. Kaynakların daha etkin kullanılabilmesi ve başarılı olabilmek adına, yerel yönetimlerin üreticilerin kuracakları kooperatifler ve birlikler aracılığıyla faaliyetlerini yürütmesi daha başarılı sonuçlar alınmasını sağlayacaktır. Kooperatifler, çiftçilerin birlikte hareket etmesini, ortak sorunlarını çözmesini, üretim ve pazarlama gücünü artırmasını sağlar. Yerel yönetimler de kooperatiflerin ürünlerini piyasa fiyatından satın alarak halka uygun fiyata dağıtabilir. Böylece hem üreticiyi hem de tüketiciyi korumuş olur.

Anahtar Kelimeler: Sürdürülebilir Tarım, Yerel Yönetim, Destekler, Ekoloji

ABSTRACT

Agricultural production is an activity that has no substitute for the future of humanity. However, according to observations and available data, it is seen that serious problems will be experienced in production in the not too distant future. In order to prevent this from happening, consistent production must be made sustainable. The increases in the world population and the dizzying pace of technological developments have led humanity to obtain high yields from unit area in agricultural production. Accordingly, high amounts of chemicals have begun to be used in the systems and control methods used. The use of artificial chemicals has become widespread, especially in plant nutrition and agricultural control. This situation has significantly increased yields, but risks have begun to emerge in terms of environment and health. Excessive use of artificial fertilizers and chemical drugs has also brought environmental problems. In many places, the soil has lost its fertility, has become barren, and the products obtained due to chemical fertilizers and drugs have become a threat to human health. Especially in developing countries, due to the excessive increase in agricultural input prices, the insufficient increase in product prices and the inadequacy of the provided supports, many small producers have broken away from agriculture and have become consumers in the slums of big cities. These developments have brought the concept of sustainability in agriculture to the forefront.

The role of local governments in agricultural development and sustainability of agriculture is great. Local governments contribute to the economy of both the city and the country by supporting the agricultural sector in the province. Local governments also create a sustainable agricultural model by protecting the environment. Local governments need to take agricultural activities into consideration when determining their vision and strategies. Local governments need to cooperate with municipalities, governorships and provincial administrations for agricultural development activities. Agricultural development activities of local governments protect both the producer and the consumer. In order to use resources more effectively and be successful, local governments carrying out their activities through cooperatives and unions to be established by producers will provide more successful results. Cooperatives enable farmers to act together, solve their common problems and increase their production and marketing power. Local governments can also purchase the products of cooperatives at market prices and distribute them to the public at affordable prices. In this way, both the producer and the consumer are protected.

Key Words: Sustainable Agriculture, Local Government, Subsidies, Ecology

GİRİŞ

Doğadaki bütün canlı varlıklar başta olmak üzere bitkisel ve hayvansal üretimin devamı için sürdürülebilirlik kavramı en çok kullandığımız kelimedir. Tarımda, gıda üretiminde, ekonomide, eğitimde ve daha birçok alanda sürdürülebilirlik kavramı son yüzyılda en çok konuşulan, zaman zaman korkulan ve gelecekte neler yapılabilirliğini tartıştığımız bir kavram haline gelmiştir. İnsanların, hayvanların ve hatta doğada yaşayan tüm canlıların yaşamsal döngülerini sağlayan en önemli temel dayanak tarımsal üretimdir. Tarımsal üretim açısından vazgeçilmez unsur ise sürdürülebilir tarımdır. Peki sürdürülebilir tarım nedir? Birçok literatüre göre; Gelecek nesillerin ihtiyaçlarını karşılama yeteneğinden ödün vermeden toplumun mevcut gıda ve tekstil ihtiyaçlarını karşılayacak sürdürülebilir yollarla yapılan tarım anlamına gelir. Tarımın sürdürülebilirliğini artırmanın birçok yöntemi olmakla beraber temel hedef; tarım sistemlerinin değişen çevre koşullarında büyüyen bir nüfusu beslemesini sağlamak için potansiyel bir çözüm sunmaktır.

Dünyanın gelişmiş ülkelerinde olduğu gibi Türkiye'de de bir yandan sentetik üretim girdileri denetimsizce kullanılırken, bir yandan da işleme teknik ve teknolojilerinin oluşturdukları olumsuz sonuçlar düşünülmeden kullanılan klasik/konvansiyonel tarımsal üretim yapılmaktadır. Günümüzde bu uygulamalar doğal dengenin bozulmasına olan etkileri ve besin zinciri yoluyla insanın yanı sıra tüm canlılara ulaşabilen yaşam tehlikesi yaratma özelliği ile artık sanayi ya da kentsel kirlilikler kadar dikkat çekmeye başlamıştır. Sürdürülebilir tarım, uzun dönemde doğal kaynakların korunmasının yanı sıra çevreye zarar vermeyen tarımsal teknolojilerin kullanıldığı bir tarımsal yapının oluşturulmasıdır (Turhan, 2005). Sürdürülebilir tarımın amaçlarına bakacak olursak sırasıyla, insanların yiyecek ve giyecek ihtiyaçlarını karşılamak, doğal kaynakları korumak ve toprak ve su kalitesinin azalmasını engellemek, yenilenemeyen kaynakları etkili bir şekilde kullanmak, doğal biyolojik döngüleri kullanmak ve kontrol altına almak, tarımın ekonomik canlılığını, üreticilerin ve ailelerin yaşam kalitesini sağlamaktır. Sürdürülebilir tarım sistemleri; mono kültür yerine sistem üzerine yoğunlaşmasıyla, ürün ve hayvan işletmeciliği arasındaki bağı artırmasıyla, biyolojik olayların öneminin anlaşılmasıyla ve sistem içinde besin madde döngüsünün artırılmasıyla diğer tarım sistemleri ve konvansiyonel tarımdan ayırt edilmektedir.

Sürdürülebilir kalkınma, üç temel konu eşzamanlı dikkate alınarak gerçekleştirilebilmektedir. Bu temel kavramlar ekonomik, sosyal ve ekolojik sürdürülebilirliktir. Ancak ekonomik kalkınmanın giderek daha fazla önem kazandığı düşük gelirli bölge ve ülkelerde ekolojik sürdürülebilirliğin zaman zaman göz ardı edildiği görülmektedir. Bu durum, ekolojik sürdürülebilirlikle ilgili genel ve kabul gören değerlendirme parametrelerinin ekolojik sürdürülebilirliği takip edebilecek yeterlikte olmamasından dolayı gerçek fayda/masraf analizinin yapılamamasından kaynaklanmaktadır. Doğal kaynakların piyasa fiyatına sahip olmaması da fayda/masraf analizinin yapılmasını zorlaştırmakta ve çoğu zaman yalnızca oluşan çevre kirliliğinin ortadan kaldırılması için yapılan masraflar çevre maliyeti olarak nitelendirilmektedir. Bu eksik değerlendirme sorununu ortadan kaldırmak amacıyla yapılan çalışmalarda doğal kaynakların fiyatlandırılması ise, sürdürülebilirliğin temel felsefesi ile bütünüyle çelişmektedir. Tarımsal üretim sırasında ortaya çıkan çevre kirliliği ve toprak bozunumu gibi nedenlerle pek çok ülkede tarım kırsal alanla birlikte değerlendirilerek, tarımın ekonomik açıdan ve kırsalda istihdamın sürdürülebilirliği amacıyla farklı uygulama yöntemleri geliştirilmiştir (Pezikoğlu, 2012).

Dünya nüfusuna paralel olarak artan tarım ürünleri ihtiyacı, yoğun kimyasal girdi uygulamalarıyla verimde artış sağlanarak karşılanmaya çalışılmıştır. Ancak tarımsal faaliyetlerde gereğinden fazla kullanılan girdilerin, insan sağlığı ve çevre üzerindeki olumsuz

etkileri zamanla artan oranda hissedilmeye başlanmıştır. Bunun sonucu olarak, sadece verim artışını değil, aynı zamanda insan sağlığını ve çevreyi korumayı esas alan sürdürülebilir tarım sistemleri önem kazanmıştır (Aydın Eryılmaz vd., 2019).

Kentleşme ve Yerel Yönetimler

Dünya coğrafyasında %3 yer kaplamasına rağmen, kentli nüfus dünya genelindeki CO₂ salınımının %70'inden sorumludur; buna ek olarak, toplam enerjinin üçte ikisini de kentli nüfus harcamaktadır. Benzer şekilde, üretilen gıdanın %70'ini tüketmektedir. Buna rağmen gıdaya erişimde yaşanan adaletsizlikler yetersiz beslenme ve beslenme bozuklukları, aşırı kilo, obezite ve bunlara bağlı sorunlar önemli toplum sağlığı sorunları haline gelmiştir. Giderek derinleşen ve bir krize dönüşen bu sorunlar tarihsel bir birikimin sonucudur. Kentler büyürken konut, endüstri, altyapı ve arazi talebi konusunda yaşanan rekabet ve gıda dışındaki sektörlerde yaşanan hızlı ekonomik büyüme kentlerde tarıma ayrılan alanın giderek küçülmesine neden olmuş ve kentleri gıda temininde uzun tedarik zincirlerine ve küresel pazarlara mahkûm olan tüketim merkezleri haline getirmiştir. Uzun yıllar boyunca göz ardı edilen kent gıda politikaları 2000'li yılların başından itibaren önemli gündem konularından biri haline gelmiştir. 2007-2008 yıllarındaki artan gıda fiyatları krizi ve onun akabinde ortaya çıkan finansal/ekonomik kriz ile gıda ve tarım politikaları, özellikle gelişmiş ülkelerdeki kentlerde merkezi hükümetlerden bağımsız olarak kent gündeminde yer almaya başlamıştır (Anonim, 2021).

Son yıllarda iklim değişikliğinin tetiklediği doğal afetlerin artmasıyla kentlerin kırılganlığı artmış ve artmaya da devam edecektir. Dünya, iklim krizinin tetiklediği doğal afetlerin yanı sıra, 2020 yılında SARS-CoV-2 virüsü ile küresel ölçekte etkili yeni bir kriz ile karşı karşıya gelinmiştir. Pandeminin açığa çıkardığı sağlık krizi bir gıda krizini de gözler önüne sermiştir. Gıda krizinden özellikle tarım ve gıda üretiminden uzak büyük kentlerde yaşayan halklar etkilenmiş ve kent halkı uzun bir dönem sonra belki ilk defa gıdanın sofraya nereden geldiği, kimlerin ürettiği ve kimlerin taşıyıp pazara-marketlere getirdiği sorusu ile yüz yüze kalmıştır. Uzun bir zamandır dile getiriliyor olsa da gıda üretim-tüketim zincirinin beklenmedik olaylar ya da afetler karşısında ne kadar kırılgan olabileceği belki de ilk kez bu kadar derinden fark edilmiştir (Anonim, 2021).

Büyükşehir Yasası ile köyler mahalle statüsüne alınmış ve büyükşehir belediyelerinin yetki alanına girmiştir. Bu da büyükşehir belediyelerinin sorumluluklarını ve faaliyet çeşitliliğini artırmıştır. Mahalle statüsüne alınan köylerde yaşayan insanların ihtiyaçları kentlilerden farklıdır. Bu ihtiyaçları karşılamak için büyükşehir belediyeleri kırsal kalkınma müdürlükleri kurmuştur. Ülkemizdeki her şehrin, büyük veya küçük, bir tarımsal potansiyeli vardır. Bu potansiyeli değerlendirmek hem şehrin ekonomisine hem de ülkenin gıda güvenliğine katkı sağlar. Büyükşehir Yasası ile ildeki tarımsal faaliyetleri desteklemek, çiftçileri üretime teşvik etmek, tarımsal faaliyetlerin sürdürülebilirliğini sağlamak ve geliştirmek Büyükşehir Belediyelerinin asli görevlerinden biridir. Belediyeler ildeki halka ucuz tarım ürünleri sunmak istiyorsa, kendi şehrinde üretim yapmalı, yapılmasını desteklemelidir. Aksi takdirde, taşıma, lojistik ve aracı maliyetleri nedeniyle, bu temel gıda ürünlerine tarladan sofraya gelene kadar 3-4 kat daha pahalıya sahip olacaktır. Dolayısıyla, her şehir en azından temel gıda konusunda kendi kendine yetebilir olmalıdır. Bunun için belediyeler önlemler almalı, gıda arzı güvenliği konularında tarımda sürdürülebilir projeler üretmelidir (Anonim, 2024).

İstanbul Örneği

Bir başka açıdan bakıldığında ise yerel yönetimlerin son yıllarda kenti kırsalla bütünleştiren, açlık ve yoksullukla mücadeleyi, kentli nüfusun sağlıklı ve besleyici gıdalara erişimini kolaylaştırmayı hedefleyen, insanlara aş, çalışanlara iş üreten, doğayı ve kültürel değerleri

koruyan gıda stratejileri ile sürdürülebilir tarımsal üretim arayışları içinde oldukları da söylenebilmektedir. Türkiye’de de bazı belediyeler tarımın sürdürülebilmesi ve küçük üreticilerin tarıma devam edebilmesi noktasında, üreticilerin ürünlerinin satışı ve pazarlaması noktasında, çiftçilere girdi sağlama noktasında destek sağlamaktadır. Bu bağlamda İstanbul Büyükşehir Belediyesi, İstanbul kenti için coğrafyası, tarihi, ekonomisi, sosyal yapısı ve kültürüne uygun, geniş bir katılımcı kitlenin görüşleri doğrultusunda, uygulanabilir, gerçekçi ve öncelikli bazı alanların tespiti ile kısa, orta ve uzun vadede eylem planlarını içeren bir gıda strateji belgesi hazırlamıştır.

İstanbul Gıda Strateji Belgesi (İGSB) ekonomik, sosyal ve ekolojik öncelikleri göz önüne alan, tarımda sürdürülebilirlikle sağlıklı ve iyi gıdaya herkesin erişebileceği bir gıda sistemine geçişi amaçlamaktadır. Bu hedeflere ulaşmak için sağlık, eğitim, çevre, adalet, insan hakları, sosyal ve ekonomik kalkınma politikaları birlikte değerlendirilmektedir. İGSB; Gıda tedarik zincirinde doğa ve insan sağlığına öncelik veren, üreticileri ve tüketicilerin haklarını adil bir şekilde koruyan, gelir ve toplumsal cinsiyet eşitliğini gözetten, ekolojik ilkeleri ve hayvan refahını gözetten, şeffaflık prensiplerine uyan, konunun tüm tarafları ile diyalog ve iş birliği içinde çalışan bir strateji belgesidir (Anonim, 2021).

İstanbul Gıda Stratejileri ve Sürdürülebilir Tarım ile Öne Çıkan Hedefler;

- İstanbul’un kırsal alanlarının korunması, mümkünse yenilerinin ilave edilmesi,
- İstanbul’un maviyle yeşili buluşturan özellikleri öne çıkarılarak, deniz ürünlerinin korunması ve geliştirilmesi,
- Giderek azalmakta olan çiftçiler, özellikle küçük ölçekli üreticiler ile kadın çiftçi ve girişimcilerin sosyal ve ekonomik olarak desteklenmesi,
- Üreticilerin, katma değerli ürünlere yönlendirilmesi,
- Mevcut kentsel gıda politikalarında söz sahibi olamayan kırılgan grupların belirlenmesi,
- Kırılgan gruplara öncelik vererek gıda kooperatifleri ve topluluk destekli tarım gibi oluşumların desteklenmesi,
- Küçük aile işletmelerinin ürünlerinin pazarlara erişiminin kolaylaştırılması,
- Küçük çiftçilere girdi desteğinin sağlanması,
- Tarım işçilerinin ve mevsimlik işçilerin haklarının gözetilmesi,
- Gıda ve üretici pazarları için altyapı desteğinin sağlanması,
- Mahalle lokantaları ve üreticiden tüketiciye doğrudan satış marketlerinin açılması,
- Açlık ve yetersiz beslenme sorunları yaşayan kesimlere destek sağlanması,
- Halk Ekmeğ üretim kapasitesinin artırılması ve kaliteli ekmeğin ucuz fiyata yoksul-kırılgan kesimlere ulaştırılması,
- Halk Süt programı ile dağıtılan süt miktarının artırılması,
- Çocukların şeker içeriği yüksek, besin ögesi içeriği zayıf, aşırı işlenmiş gıda ürünlerine erişimini azaltacak adımların atılması,
- Park ve bahçelerde sıcaklara daha dayanıklı ve su tüketimi az bitkilerin kullanılması,
- “Azalt, Yeniden Değerlendir ve Geri Dönüştür” esaslarının benimsenmesi,
- Gıda toptancı hallerinin gıdaların bozulmasına, zarar görmesine ya da zayıf olmasına yol açacak teknik ve hijyenik koşullarının kontrol edilmesi ve iyileştirilmesi,
- Kentte doğa dostu küçükbaş hayvancılık, arıcılık ve balıkçılık uygulamalarını desteklemek,
- Hayvancılık sektörünün doğaya olan ağır etkilerini azaltmak için (metan gazı salımı, aşırı su kullanımı gibi) alternatif beslenme modelleri hakkında eğitim vermek,
- Kooperatifleşmeyi yaygınlaştırmak,
- Gıda üretimine temel teşkil edecek tohum ve fidan üretmek.

Uluslararası alanda her 10 yılda bir yapılan Dünya Gıda Zirveleri, Birleşmiş Milletler Binyıl Kalkınma Hedefleri (2000-2015) ve onu takip eden Sürdürülebilir Kalkınma Hedefleri (2015-2030) açlığın ve beslenme yetersizliğinin ortadan kaldırılması ve gıda güvencesinin sağlanmasının önemini devamlı gündeme getirmiştir. 2008’de yaşanan ve 2011’de tekrar eden gıda krizi ve 2020’den beri yaşamakta olduğumuz pandemi hem ulusal hem de küresel ölçekte konunun önemini gündemde tutmaktadır. Bugün pek çok uluslararası kurumun (FAO, Dünya Gıda Güvencesi Komitesi CFS, IFAD, Dünya Gıda Programı WFP, Avrupa Topluluğu Ekonomik ve Sosyal Komitesi, Birleşmiş Milletler Çevre Programı UNEP gibi) sürdürülebilir, etkin, besleyici ve sağlıklı gıda sistemlerini yaşama geçirecek politikaların gereğinden söz etmesi bu konudaki küresel endişeleri yansıtmaktadır (Anonim, 2021).

Ancak bu kurumlarda sürdürülebilirlik ve gıda güvencesinin önemi konusunda fikir birliği olsa da bu kavramların tanımlanmasında ve önceliklerin belirlenmesi noktasında oldukça farklı yaklaşımlar gözlenmektedir. Dünya pazarlarına egemen olan çok uluslu şirketler ve uluslararası kurumlar çözümü üretim kapasitesinin ve verimliliğin artırılmasında, sulama sistemlerinin modernizasyonu, melez veya GDO’lu tohumlar, tarım ilaçları, yapay gübre, akıllı tarım, dikey tarım, yapay proteinler, dijital tarım türü teknolojik çözümlerde ve küresel ticarete görmektedir. Bu yaklaşım, türlü nedenlerle gıdaya erişemeyenlere de gıda bankaları, hayır kurumları ve uluslararası yardım kurumları aracılığı ile yardım edilerek çözüme ulaşılabileceğini savunmaktadır. Bütün bu yaklaşımlar iklim krizini derinleştirme, kimyasal kirliliği ve biyolojik çeşitlilik kaybını artırma gibi olumsuz sonuçlara yol açmaktadır. Dolayısıyla, doğal hayatın istikrarını gözeten, doğal hayattaki canlılar arasındaki ilişkilerin devamlılığını esas alan, kamusal refahı sağlamayı ve toplumsal barışı korumayı amaçlayan yaklaşımlara ihtiyaç vardır.

Dünyada birçok kentte uygulanan kent tarımı İstanbul için önemli bir değişim ve olumlu bir gelecek için başlangıç olacaktır. Küçük projelerden başlayarak mahallelerdeki sahipsiz ve kullanılmayan alanlarla, balkon, arka bahçe, teras gibi apartman ve kent yaşamına uygun ama şimdiye kadar kullanılmamış alanların tarımsal üretime açılması, mahalle bostanları oluşturulması sürdürülebilir tarımda küçük ancak etkisi büyük ilk adımlar olarak görülebilir.

Kent tarımı yerel gıda sistemlerinin bir parçası olup, gıdanın üretildiği, pazarlandığı ve tüketildiği yerin kent olduğu bir sistemdir. Bu sistem işlenmemiş, atıl kalmış kent topraklarını kent sakinleri için sosyal ve ekonomik getirisi olan alanlara dönüştürür. Kentlerde olanaklar elverdiği ölçüde hayvancılık, arıcılık, balıkçılık, balık çiftlikleri, bitkisel ürün yetiştirme, çiçekçilik, fide ve tohum üretimi yapılabilmektedir. Sağlıklı gıdaya erişim için atılacak ilk adım yerel tarıma öncelik vermektir.

Kentlerde uygun mahallerde oluşturulacak kent bostanları çok amaçlı politikalara hizmet etmektedir. Özellikle dar gelirli kent sakinlerinin sağlıklı taze sebze ve meyveye erişimini arttıran, yerel üretimi desteklediği için istihdam sorununa çözüm öneren ve tedarik zincirini kısaltarak ekolojik etkileri en aza indiren, gıda üretimine yönelik kişisel becerileri artıran bir üretim modelidir. Kentliden kentliye, komşudan komşuya yardım ve eğitim projeleri ile vatandaşların dayanışma bilincinin ve sosyal barışın artırılmasını da desteklemektedir. Bir yandan küçük üreticilerin mahsullerini kent pazarına aktarıp devamlı ve güvenilir gelir ile ekonomik kalkınmaları sağlanacak, diğer yandan kentli nüfus sağlıklı gıdaya, kısa tedarik zinciriyle doğrudan ulaşacaktır. Bunun yanında her mahallede kompost yerleri kurulması ile tarımın en önemli meselelerinden biri olan gıda atıklarının değerlendirilmesi sağlanacaktır. Atıkların dönüşüm yoluyla tekrar kullanımının sağlanması çevre korumadan, kimyasal gübreye son vermeye, atık yönetiminden ucuz ve toprağı çok yormayan onarıcı tarıma geçişe kadar birçok amaca da hizmet edecektir. Ancak kent tarımında su kullanımı önemli bir boyuttur. Kent Tarımı adını verdiğimiz tarım şeklinde kullanılacak tatlı su içme suyu hattından kullanılacağından bu suyun iyi yönetilmesi gerekmektedir. Su ihtiyacı düşük olan bitkilere öncelik verilmeli ve sulama suyu ihtiyaçları binaların çatılarına düşen yağmur sularının

depolanması ile sağlanmalıdır. Şehrin içme ve kullanma suyu kaynakları bu amaçla minimum düzeyde kullanılmalıdır.

İstanbul Büyükşehir Belediyesi'nin sürdürülebilir tarımı destekleyen örnek çalışmaları;

Fide Desteği; Çiftçilere toplamda 54,4 milyon adet sebze fidesi desteğinde bulunulmuştur.

Fide Üretimi; İstanbul çiftçisinin girdi maliyetlerini azaltmak ve fide taleplerini kendi tesislerinden karşılamak amacıyla fide üretimine başlamışlardır.

Biyoteknik Mücadele; Seralarda pestisit kullanımını azaltmak amacıyla çiftçilere biyoteknik mücadele araçları sağlamaktadırlar.


Gübre Desteği; İstanbul'da sebze üretimi yapan ve Belediye'nin yazlık fide desteğinden yararlanan çiftçilere gübre desteği vermeye başlamışlardır. Destek kapsamında 222,65 ton gübre dağıtımı gerçekleştirmişlerdir.

Malç Naylonu Desteği; İstanbullu çiftçilerin girdi maliyetlerini azaltmak, tarıma teşvik etmek amacıyla malç naylonu desteğine başlamışlardır. Destek kapsamında 2.507 top malç naylonu dağıtılmıştır.

Damlama Sulama Hortumu Desteği; İstanbul Büyükşehir Belediyesi'nden yazlık fide desteği alan ve seralarda üretim yapan çiftçiler bu destekten faydalanabilmektedir. Destek kapsamında, 361,6 kilometre damla sulama hortumu verilmiştir.

Çilek Üretim Denemeleri; Çiftçilerin, üretimlerini çeşitlendirmeleri için 72.000 adet çilek fidesi dağıtımı yapılmıştır.

Deneme Ekimleri; İstanbul'da yaygın olarak ekimi yapılan bitkilere ait yerli çeşitlerin deneme ekimlerini yapılmaktadır. Yüksek verim elde edilen çeşitlerin tarımsal destekleme kapsamında çiftçiler tarafından kullanılması sağlanmaktadır.



Ayçiçeği Tohumu Desteği; Destek kapsamında 2 binden fazla çiftçiye 8.934 torba ayçiçeği tohumu dağıtılmıştır.

Buğday Tohumu Desteği; Üretim maliyetini düşürmek ve yüksek verimli yerli tohum ekimini yaygınlaştırmak için, 2022 yılında toplam 1.548.375 kg tohum desteğinde bulunuldu. 2024 yılında ise toplam 1.530.500 kg tohum desteği verildi.

Arpa ve Yulaf Tohumu Desteği; İstanbul Büyükşehir Belediyesi, tarımsal üretimden çekilen çiftçileri geri kazanmak amacıyla 2023 yılında ilk kez, 1.197.830 kg arpa ve yulaf tohumu desteği sağladı. 2024 yılında ise 323.300 kg arpa ve 136.920 kg yulaf tohumu desteği verildi.

Halk Ekmek Buğday Alım Projesi; Çiftçilerin üretimlerini teşvik etmek amacıyla İstanbullu çiftçilerden Halk Ekmek yapımı için, ekmeklik buğday alımı gerçekleştirildi.

Mazot Desteği; 2022 yılında 157.347 litre, 2023 yılında 308.603 litre, 2024 yılında ise 242.655 litre mazot desteği verilmiştir.

Makine ve Ekipman Desteđi; Üretim alanlarını analiz ederek, uygun makine ve ekipman temini yoluyla üretimlerini desteklemektedirler.

Sera Desteđi; İstanbul'un pek çok yöresinde üreticilere destek olmak amacıyla, seralarda sebze üretimi yapılmaktadır.

Yaş Meyve ve Sebze Hali Artıklarının Kullanımı; Fermantasyon işlemi sonrasında elde edilen artıklar kompost üretiminde kullanılmaktadır. Yine fermantasyon işlemi sırasında elde edilen metan gazı ise yakılarak elektrik üretilmektedir. Elektrik üretim santralinin soğutulmasında kullanılan su, fide seralarımızın ısıtılmasında kullanılmakta ve bu seralarda üretilen sebze fideleri destek kapsamında çiftçilere dağıtılmaktadır.

Toprak ve Su Analizi Desteđi; Çiftçilerin tarlasına ekip diktiđi bitkileri doğru beslemeleri, eksik olan bitki besin elementlerini takviye etmeleri amacıyla çiftçilerin toprak ve su analizlerini ücretsiz olarak yapmaktadırlar. Toplamda, 589 adet toprak ve 181 adet su analizi yapılmıştır.

Sığır ve Manda Süt Yemi Desteđi; 2022 yılında başlayan proje kapsamında çiftçilere 1.249 ton, 2023 yılında 1.640, 2024 yılında ise 1.590 ton sığır süt yemi dağıtılmıştır.

Kuzu Besi Yemi Desteđi; 2021 yılında başlayan proje kapsamında çiftçilere 1.181 ton, 2022 yılında 1.115 ton, 2023 yılında 1.104 ton, 2024 yılında yaklaşık 1.598 ton yem desteđi sağlanmıştır.

Silajlık Mısır Tohumu Desteđi; Hayvancılıkla geçimini sağlayan çiftçilere silajlık mısır tohumu desteđi sağlamaktadırlar. 2021 yılında başlatılan bu destek 2022 yılında da devam ettirilmiştir. Bu süreçte toplam 2.217 çuval silajlık mısır tohumu dağıtılmıştır. 2023 yılında 1.788 torba, 2024 yılında ise 1.713 torba silajlık mısır tohumu dağıtılmıştır.

Kurbanlık Hayvan Gübresinin Kullanımı; Kurban hizmeti verilen alanlardan toplanan gübreler fermente edilerek talep eden çiftçilere ücretsiz verilmektedir.

Üretici/Kooperatif Pazarları; Üreten İstanbul için çiftçiler destekleniyor. 2020 yılında Kadıköy ve 2021 yılında da Beşiktaş'ta çiftçi/kooperatif pazarları kuruldu. İstanbullu çiftçiler ve ülkenin her bir köşesinde tarımsal üretim yapan kooperatifler ürettikleri tarımsal ürünleri bu pazarlarımızda aracısız satmaktadır

Çiftçi Eğitimleri; İldeki çiftçilere, hayvan yetiştiricilerine ve balıkçılara konusunda uzman ziraat mühendisleri, veteriner hekimler, su ürünleri mühendisleri ve su bilimleri mühendisleri tarafından aşağıdaki konularda eğitimler verilmektedir:

- Toprak hazırlığı ve gübreleme,
- Fide dikimi ve bakımı,
- Bitki besleme,
- Hastalık ve zararlılar ile mücadele,
- Buzađı bakımı ve beslenmesi,
- Süt hijyeni ve sanitasyonu,
- Satış yerlerinde su ürünleri hijyeni,
- Sürdürülebilir balıkçılık.
- Kooperatifçilik ve kooperatifçiliđin önemi,
- Topraktan Fileye İzleme Modeli Temel Yeterlilik Eğitimi.

Öğrenci Eğitimleri; Devlet okullarında ve özel okullarda tarım ve su ürünleri konularında farkındalık eğitimleri verilmektedir. Amaç, çocuklara tarımı ve balıkçılığı sevdirmek, tarımsal ve sucul alanları korumalarını sağlamak, balık tüketimini özendirmek, su ve tarım ürünlerinin faydalarını öğretmektir.

İSTKA Projesi; İstanbul Kalkınma Ajansı 2021 yılı, Yenilikçi İstanbul Mali Destek Programı kapsamında hibe almaya hak kazanan ve proje yürütme süreci devam eden Toprakten Fileye Gıda İzleme Modeli'nde;

- Üretici ve tüketicinin aracısız buluşabildiği Üretici Pazarı sisteminin teknolojik olanaklar kullanılarak modernize edilmesi,
- Toprakten pazara kadar izlenilebilir bir gıda sistemi kurulması,
- Çiftçi teknoloji okur-yazarlığının artırılması,
- Tüketicilerin pazara gitmeden, ürün ve üreticilerle ilgili bilgilere ulaşabilmesi,
- Stoklar ve fiyatların takip edilerek, çeşitli iletişim araçlarında yayınlanması ile birlikte şeffaflık sağlanması,
- Üretici-Tüketici bağının güçlendirilmesi amaçlanmaktadır.

Dijital Pazarlama; İstanbul Büyükşehir Belediyesi, Halk Market e-ticaret projesi vasıtasıyla çiftçi kooperatiflerinin ürünlerinin pazarlanmasına yardımcı olmaktadır.

Çiftçi Kooperatiflerinin Güçlendirilmesi; İstanbul Büyükşehir Belediyesi, ihtiyaç sahibi ailelere yardım paketleri dağıtmaktadır. Bu paketlerin içinde bulunan temel gıda maddeleri Türkiye'nin çeşitli yörelerinde faaliyet gösteren çiftçi kooperatiflerinden temin edilmektedir.

SONUÇ

Yeni bir yüzyıla giren dünyamızda, gelecekle ilgili kaygılar daha yoğun bir biçimde yaşanmaya başlamıştır. Dünya nüfusunun hızla artması ve gelecekle ilgili birçok araştırmanın yapılması, "ileride bu kadar insanı beslemeye yetebilecek kadar çok besin maddesini üretebilecek miyiz?" sorusunu aklımıza getirmektedir. Yapılan araştırmaların pek çoğunun sonucu gösteriyor ki uygulanan yoğun tarımsal üretim programlarıyla sürdürülemez bir gelişmenin eşiğine yaklaşılmıştır. Sonuçta, artık tarımsal üretimin doğaya zarar vermeden artması gereği karşımıza çıkmaktadır. Bunu sağlayabilmek için erozyonu, toprağın tuzlulaşmasını, su kaynaklarının kirlenmesini ve diğer zararları en aza indirgeyen sürdürülebilir tarım tekniklerinin geliştirilmesi gereklidir. Hem gelişmiş hem de gelişmekte olan ülkeler besin üretimini artırmanın yollarını ararken, tarımda kullanılan doğal kaynakları da güvence altına alacak yeni yöntemler geliştirme zorunluluğuyla karşı karşıyadırlar. Özellikle, uygulanan yanlış ve yoğun girdili yetiştirme teknikleri tarımsal alanlarımızda geri getiremeyeceğimiz mikroorganizma kayıplarına neden olmaktadır. Kuşaklardır süren tarımsal mücadele ilaçlarının da kontrol altına alınmadığı gözlenmektedir. Bu uygulama, insan sağlığında yarattığı tehlike, diğer türleri tehdit etmesi ve tarımsal mücadele ilacına direncin artması nedeniyle kontrol altına alınması gereklidir. Dünyada ve ülkemizde sürdürülebilir yaşamın bir gereği olarak önlemler alınmadığı takdirde ileride aç insanların sayısı artacak, hatta felaket boyutuna ulaşacaktır. Bu sorunlarla karşı karşıya olan dünyamızda üretim için doğal kaynaklar açısından tehlike oluşturmayan tarım sistemi insan, çevre ve ekonomik olarak sürdürülebilir tarımsal üretimi gerçekleştiren bir yaklaşımdır.

Sonuç olarak günümüzde, gelecek nesilleri tehlikeye atmadan gereksinimlerini karşılayan bir toplum yaratmak açısından sürdürülebilir tarım son derece önemli bir kavramdır. Toprak ve su kaynaklarının korunması, doğal kaynakların korunması, erozyon ve orman yangınları ile mücadele, biyolojik çeşitliliğin sağlanması, entegre ilaç yönetimi tarımda uygun yetiştirme

tekniklerinin kullanılarak tarımsal arazilerimizin verimliliğinin artırılması son yıllarda önemini gittikçe arttıran sürdürülebilir yaşam ve sürdürülebilir bir tarım için önemli bir gerek olarak karşımıza çıkmaktadır. Sürdürülebilir tarımda güvenilir gıda üretimi tek amaç değildir. Sürdürülebilir tarım, gıda üretimi ile birlikte kırsal kalkınmayı sağlayan, çiftçilerin ekonomik düzeylerini ve yaşam kalitesini artıran bir sistemdir. Sürdürülebilir tarım için birçok gösterge belirlenmiştir. Farklı çalışmalarda gösterge kıstaslarında değişkenlik olsa da bu çalışmada temelde sürdürülebilir tarıma ait göstergeler; doğal ortama ait göstergeler, sosyal ve ekonomik göstergeler olarak sınıflandırılmıştır. Doğal ortama ait göstergelerin bilinmesi ve bu potansiyel doğrultusunda hedef ve politikaların belirlenmesi esas olmalıdır. Çünkü doğal ortam özelliklerine aykırı uygulamaların yapılması, uzun vadede yapısal sorunların oluşmasına yol açmaktadır. Günümüzde görülen tarımsal sorunların kaynağında da doğal ortam özellikleriyle uyuşmayan uygulamaların yapılması bulunmaktadır. Bununla beraber tek başına tarımsal üretimin doğru yöntem ve uygulamalarla eksiksiz bir şekilde yapılması sürdürülebilirliğin gerçekleştiği anlamına gelmemektedir. Üreticinin ürettiği ürünlerin bir değer ifade etmesi adına karşılığını alması gerekmektedir. Tarlada kalmış satılmayan ürünlerde sürdürülebilirliği riske eden sosyal ve ekonomik göstergelerdir. Bu yönüyle sürdürülebilir tarım ve sürdürülebilir tarımsal uygulamalarda bütüncül bir yaklaşım bulunmaktadır. Tarımsal faaliyetlerde ilk aşama olan ürünlerin üretilmesinden tüketiciye ulaştırıldığı son aşamaya kadar tüm aşamaların eksiksiz işlemesi, sürdürülebilirliğin tam anlamıyla gerçekleştiğinin göstergesidir. Sürdürülebilir tarım ve sürdürülebilir tarımsal uygulamaların hedefinde; toprak, su gibi doğal kaynakları devamlı kılmak için yapılan iyileştirmelerle çevre kalitesini sağlamak, güvenli gıda üretimini gerçekleştirmek, tarımsal ekonominin dinamizmini devamlı kılmak, tarımsal kesimin hayat standartlarını geliştirmek bulunmaktadır. Tarımın sürdürülebilirliğinin sağlanmasında merkezi idare yanında, yerel yönetimlerin de aktif rol üstlenmesi yaşamsal önem taşımaktadır. İstanbul örneği bu anlamda çok önemli bir rehber oluşturmaktadır.

Kaynaklar

Anonim, (2021): https://tarim.ibt.istanbul/img/1551435102021_7928277850i.pdf, Erişim Tarihi: 01.02.2025

Anonim, (2024): <https://kurumsaltarim.com/125-tarimsal-kalkinmada-yerel-yonetimlerin-rolu.html>, Erişim Tarihi: 03.02.2025.

Aydın Eryılmaz, G., Kılıç, O. ve Boz, İ. (2019). *Türkiye'de Organik Tarım ve İyi Tarım Uygulamalarının Ekonomik, Sosyal ve Çevresel Sürdürülebilirlik Açısından Değerlendirilmesi*. Yüzüncü Yıl Üniversitesi Tarım Bilimleri Dergisi, 29(2), 352-361. <https://doi.org/10.29133/yyutbd.446002>

Pezikoğlu, F. (2012). *Sürdürülebilir Tarım ve Kırsal Kalkınma Kavramı İçinde Tarım-Turizm-Kırsal Alan İlişkisi ve Sonuçları*. Karamanoğlu Mehmetbey Üniversitesi Sosyal Ve Ekonomik Araştırmalar Dergisi, 2012(1), 83-92.

Turhan, Ş. (2005). *Tarımda Sürdürülebilirlik ve Organik Tarım*. Tarım Ekonomisi Dergisi, 11(1 ve 2), 13-24.

**DETERMINATION OF YIELD AND QUALITY PARAMETERS OF SUGAR BEET
PARENTAL LINES UNDER LIMITED IRRIGATION**

**ŞEKER PANCARI EBEVEYN HATLARIN KISITLI SULAMA İLE VERİM VE
KALİTE PARAMETRELERİNİN BELİRLENMESİ**

İlyas KILINÇER

Kayseri Şeker R&D Center, Researcher, Kayseri, Turkey

ORCID ID: <https://orcid.org/0000-0002-7109-3772>

Dr. Özgür ÖZMEN

Kayseri Şeker R&D Center, R&D Agricultural Breeding Chief, Kayseri, Turkey

ORCID ID: <https://orcid.org/0000-0003-4188-3252>

Dr. Anıl Mehmet BALTACI

Kayseri Şeker R&D Center, Sustainable Agriculture Manager, Kayseri, Turkey

ORCID ID: <https://orcid.org/0000-0003-1890-131X>

Prof. Dr. Mehmet ARSLAN

Erciyes University, Faculty of Agriculture, Agricultural Biotechnology, Kayseri, Turkey

ORCID ID: <https://orcid.org/0000-0002-0530-157X>

ABSTRACT

Introduction and Objective: Sugar, a fundamental nutrient in human nutrition, holds a significant position among industrial crops and is primarily produced from sugar beet in our Türkiye. Sugar beet belongs to the Amaranthaceae family. Achieving high yields in sugar beet cultivation requires adequate rainfall and regular irrigation. The annual water consumption of sugar beet is approximately 700 mm. Inadequate irrigation leads to various stress factors, ultimately resulting in yield reduction. Therefore, institutions and private companies engaged in sugar beet breeding place great emphasis on developing varieties and parental lines that are not only high-yielding and disease-resistant but also tolerant to drought stress. One of the key strategies for achieving high yields with minimal irrigation is the development of drought-tolerant parental lines.

Materials and Methods: In this study, conducted in 2022 and 2023, 22 genotypes and four commercial controls within a sugar beet breeding population were evaluated for yield and quality parameters under different levels of water deficit. Water deficit levels of 30%, 60%, and 100% were applied. The responses of the genotypes to drought stress were assessed based on parameters such as yield, sugar content, sodium content, α -amino nitrogen content, and potassium content. **Results:** According to variance analysis performed using the SAS program, the effects of genotype and irrigation on yield, polar, and potassium values were found to be

highly significant. However, for sodium and nitrogen, only the effect of genotype was determined to be highly significant. Additionally, the irrigation*genotype interaction was found to be statistically significant. LSD tests were applied for yield, polar, sodium, potassium, and α -amino nitrogen parameters.

Discussion and Conclusion: Based on the results related to irrigation constraints, yield, polar, and potassium were categorized into three groups. According to the LSD test, the highest-yielding genotypes were Picasso, 19, 9, 23, and Dozer, while the lowest-yielding genotypes were 33, 55, and 41. In the LSD table, the genotypes with the highest polar values, following the four commercial control groups, were 36 and 17, whereas the lowest polar values were observed in genotypes 55, 102, and 244. This study, which aims to identify high-yielding and high-quality lines under limited irrigation conditions, represents a significant step toward sustainable agriculture and water resource conservation. Additionally, it aims to contribute to breeding efforts focused on enhancing drought tolerance in sugar beet.

Key Words: Drought Stress; Sugar Beet; Abiotic Stress; Limited Irrigation

ÖZET

Giriş ve Amaç: Endüstri bitkileri arasında önemli bir yere sahip olan ve insan beslenmesinin temel besin maddesi olan şeker, Türkiye’de ağırlıklı olarak şeker pancarından üretilmektedir. Şeker pancarı, *Amaranthaceae* familyasına ait bir bitkidir. Şeker pancarında yüksek verim elde edebilmek için yeterli yağış ve düzenli sulama gereklidir. Şeker pancarının yıllık su tüketimi yaklaşık 700 mm’dir, yeterli sulama yapılmadığında çeşitli stres faktörleri ortaya çıkmakta ve verim düşmektedir. Bu nedenle, şeker pancarı ıslahı yapan firmalar verim, kalite ve hastalıklara dayanıklılığın yanı sıra kuraklık stresine dayanıklı çeşitler ve ebeveyn hatlar geliştirmeye önem vermektedir. Minimum sulama ile yüksek verim elde etmenin yollarından biri kuraklık stresine dayanıklı ebeveyn hatlarının geliştirilmesidir.

Gereç ve Yöntem: 2022 ve 2023 yıllarında yapılan çalışmada, şeker pancarı ıslah popülasyonu içerisinde 22 genotip ve 4 ticari kontrol kullanılarak farklı su kısıtı seviyelerindeki verim ve kalite parametreleri belirlenmiştir. Çalışmada %30, %60 ve %100 su kısıtı seviyesi uygulanmıştır. Ekilen genotiplerin kuraklık stresine karşı gösterdiği tepkileri belirlemek amacıyla verim, polar, sodyum, α -amino azot ve potasyum oranları değerlendirilmiştir.

Bulgular: SAS programında yapılan varyans analiz sonuçlarına göre, genotip ve sulamanın verim, polar, potasyum değerleri üzerindeki etkisi çok önemli bulunmuştur. Sodyum, azot açısından ise genotip etkisinin çok önemli olduğu belirlenmiştir. Ayrıca sulama*genotip interaksiyonun istatistiksel olarak çok önemli olduğu bulunmuştur. Verim, polar, sodyum, potasyum ve α -amino azot parametreleri için LSD testleri uygulanmıştır.

Tartışma ve Sonuç: Sulama kısıtı yönünden sonuçları yorumladığımızda verim, polar ve potasyum 3 gruba ayrılmıştır. LSD testine göre en yüksek verim Picasso, 19, 9, 23 ve Dozer genotiplerinde iken en düşük ise 33, 55, 41 genotipleri olmuştur. En yüksek polar 4 adet ticari kontrol grubundan sonra 36 ve 17 genotipleri çıkarken en düşük ise 55, 102, 244 genotipleri olduğu belirlenmiştir. Kısıtlı sulama imkanlarıyla yüksek verim ve kaliteye sahip hatların belirlenmesi üzerine yapılan bu çalışma sürdürülebilir tarım ve su kaynakların korunması için atılan önemli bir adımdır. Ayrıca şeker pancarının kuraklık stresine dayanıklılığını artırmaya yönelik ıslah çalışmalarına katkı sağlamayı hedeflemektedir.

Anahtar Kelimeler: Kuraklık Stresi; Şeker Pancarı; Abiyotik Stres; Kısıtlı Sulama

**IN THE REPUBLIC OF AZERBAIJAN LEGAL BASIS OF SWITCHING TO
ORGANIC BEEKEEPING**

**AZERBAIJAN CUMHURİYETİ'NDE ORGANİK ARICILIĞA GEÇİŞİN HUKUKİ
TEMELİ**

Yunus RUSTAMLI

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Veterinary
Department, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0000-0786-0597>

Ali TAHIROV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Veterinary
Department, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0006-1280-7292>

ABSTRACT

Introduction and Purpose: Ancient of Azerbaijan Republic The culture and rich nature of beekeeping forms the basis for the production and processing of high-quality and safe beekeeping products that enable the treatment and prevention of "civilized" diseases and other pathologies, as well as the application of these products in dietetics. Therefore, amateur and professional beekeepers should switch from traditional beekeeping to organic beekeeping while protecting the agricultural ecosystem. **Materials and Methods:** Based on legal literature, observation, analysis and personal research, we found that the legal basis of the transition to organic beekeeping in the Republic of Azerbaijan is regulated by the Laws of the Republic of Azerbaijan "On Ecological Clean Agriculture", "On Beekeeping" and "On Food Safety". "Rules of ecological supervision in ecologically clean agriculture and duties of accredited organizations" and "Production, processing, labeling and sale of ecologically clean agriculture and food products" approved by the decision of the Council of Ministers of the Republic of Azerbaijan dated January 8, 2009 and numbered 5. The State Standard of the Republic of Azerbaijan (AZS 921:2022) "Guide to the transition to organic beekeeping" and other regulatory legal regulations adopted in accordance with them also determine the legal basis for the transition to organic beekeeping. **Results:** According to international standards, the transition period for bee colonies to organic beekeeping is 1 year. During the transition period, the honeycombs and waxes of the colonies used in all stages of beekeeping product production must be replaced with the honeycombs and waxes from organically produced colonies. If it is not possible to replace the combs and waxes within one year, the transition period is extended until they are completely replaced by the certification body or the authorized supervision body. **Discussion and Conclusion:** In order to speed up the transition to organic beekeeping, awareness raising activities should be done with beekeepers.

Key words: Republic of Azerbaijan, Law, legal, organic beekeeping, transition period

ÖZET

Giriş ve Amaç: Azerbaycan Cumhuriyeti'nin kadim arıcılık kültürü ve zengin doğası, "medeniyet" hastalıkları ve diğer patolojilerin tedavisi ve önlenmesini mümkün kılan, ayrıca bu ürünlerin diyetetikte uygulanmasını sağlayan yüksek kaliteli ve güvenli arıcılık ürünlerinin üretimi ve işlenmesi için temel oluşturmaktadır. Bu nedenle amatör ve profesyonel arıcıların tarımsal ekosistemi koruyarak geleneksel arıcılıktan organik arıcılığa geçiş yapmaları gerekmektedir. **Gereç ve Yöntemler:** Yasal literatür, gözlem, analiz ve kişisel araştırmalara dayanarak Azerbaycan Cumhuriyeti'nde organik arıcılığa geçişin yasal dayanağının Azerbaycan Cumhuriyeti'nin "Ekolojik Temiz Tarım Hakkında", "Arıcılık Hakkında" ve "Gıda Güvenliği Hakkında" Kanunları ile düzenlendiğini tespit ettik. Azerbaycan Cumhuriyeti Bakanlar Kurulunun 8 Ocak 2009 tarihli ve 5 sayılı kararıyla onaylanan "Ekolojik olarak temiz tarımda ekolojik gözetim kuralları ve akredite kuruluşların görevleri" ve "Ekolojik olarak temiz tarım ve gıda ürünlerinin üretimi, işlenmesi, etiketlenmesi ve satışı". Azerbaycan Cumhuriyeti Devlet Standardı (AZS 921:2022) "Organik arıcılığa geçiş kılavuzu" ve bunlara uygun olarak kabul edilen diğer düzenleyici yasal düzenlemeler de organik arıcılığa geçişin hukuki dayanağını belirlemektedir. **Bulgular:** Uluslararası standartlara göre arı kolonileri için organik arıcılığa geçiş süresi -1 yıldır. Geçiş döneminde arıcılık ürünü üretiminin tüm aşamalarında kullanılan kolonilerin petek ve mumlarının mutlaka organik olarak üretilen kolonilerden gelen petek ve mumlarla değiştirilmesi gerekmektedir. Bir yıl içerisinde petek ve mumların değiştirilmesi mümkün olmazsa, geçiş süresi belgelendirme kuruluşu veya yetkili gözetim kuruluşu tarafından tamamen değiştirilinceye kadar uzatılır. **Tartışma ve Sonuç:** Organik arıcılığa geçişi hızlandırmak için arıcılarla bilinçlendirme çalışmaları yapılmalıdır.

Anahtar kelimeler: Azerbaycan Cumhuriyeti, Hukuk, yasal, organik arıcılık, geçiş dönemi

**EVALUATION OF THE FORAGE QUALITY AND NUTRITIONAL VALUE OF
THE PASTURE AREA OF ULUTAŞ NEIGHBORHOOD OF İSPİR DISTRICT IN
ERZURUM PROVINCE**

**ERZURUM İLİ İSPİR İLÇESİ ULUTAŞ MAHALLESİ MERA ALANININ OT
KALİTESİ VE BESİN DEĞERİNİN DEĞERLENDİRİLMESİ**

Arş. Gör. Dr. Muhammed İkbal ÇATAL

Recep Tayyip Erdoğan Üniversitesi, Ziraat Fakültesi, Tarla Bitkileri Bölümü, Rize, Türkiye.

ORCID ID: <https://orcid.org/0000-0002-4888-770X>

Prof. Dr. Adil BAKOĞLU

Recep Tayyip Erdoğan Üniversitesi, Pazar Meslek Yüksek Okulu, Bitkisel ve Hayvansal Üretim Bölümü, Rize, Türkiye.

ORCID ID: <https://orcid.org/0000-0003-1189-8461>

ÖZET

Bu çalışma, hayvancılıkta önemli bir yere sahip olan meraların ot kalitesinin ve besin değerinin belirlenmesinin öneminden yola çıkarak, Erzurum ili İspir ilçesi Ulutaş Mahallesi meralarının ot kalitesini ve besin değerini değerlendirmeyi amaçlamıştır. Meraların sürdürülebilir yönetimi ve hayvan beslemesi açısından bitki kompozisyonu ve besin değerinin doğru bir şekilde belirlenmesi kritik öneme sahiptir. Çalışma, 2020 yılı Temmuz ayında Ulutaş Mahallesi mera alanından seçilen 12 farklı noktadan bitki örnekleri toplanarak yürütülmüştür. Örnekleme, 50x50 cm boyutlarındaki kuadratlar kullanılarak gerçekleştirilmiştir. Toplanan örnekler kurutulup öğütüldükten sonra, kuru madde oranı (KMO), ham protein (HP), asit deterjan lifi (ADF), nötr deterjan lifi (NDF), asit deterjan proteini (ADP), fosfor (P), potasyum (K), kalsiyum (Ca) ve magnezyum (Mg) içerikleri NIRS cihazı ile belirlenmiştir. Ayrıca, ADF ve NDF değerleri kullanılarak kuru madde tüketimi (KMT), sindirilebilir kuru madde (SKM), nispi yem değeri (NYD), sindirilebilir enerji (SE) ve metabolik enerji (ME) değerleri hesaplanmıştır. Analizler sonucunda, ortalama kuru madde oranı %86,64, ham protein içeriği %11,23, ADF %44,09, NDF %64,30, SKM 54,55 ve NYD 78,92 olarak tespit edilmiştir. Ortalama SE değeri 2,60 Mcal/kg ve ME değeri 2,14 Mcal/kg olarak bulunmuştur. Mineral madde içerikleri ise ortalama olarak P %0,30, K %1,28, Ca %1,49 ve Mg %0,49 olarak belirlenmiştir. Ca/P oranı 4,97 ve K/(Ca+Mg) oranı 0,65 olarak hesaplanmıştır. Bulgular, Ulutaş merasının otlarının kuru madde içeriğinin yüksek olduğunu, ancak ham protein ve enerji içeriklerinin orta düzeyde, lif içeriklerinin ise nispeten yüksek olduğunu göstermektedir. Bu durum, mera otlarının besin değerini sınırlayan faktörler olarak değerlendirilebilir. Mineral madde içeriklerinin dengeli olduğu görülmektedir. Mera verimliliğini ve ot kalitesini artırmak için sürdürülebilir mera yönetim stratejilerinin uygulanması ve mera ıslah çalışmalarının yapılması önerilmektedir.

Anahtar Kelimeler: Ulutaş, Erzurum, Ot Kalitesi, Besin Değeri.

ABSTRACT

This study aimed to evaluate the forage quality and nutritional value of pasture area in Ulutaş Neighborhood, İspir District, Erzurum Province, based on the importance of determining forage quality and nutritional value of pastures, which have a significant place in animal husbandry. Accurate determination of plant composition and nutritional value is critical for sustainable pasture management and animal feeding. The study was conducted in July 2020 by collecting plant samples from 12 different points selected from the pasture area of Ulutaş Neighborhood. Sampling was carried out using 50x50 cm quadrats at each point. After drying and grinding the collected samples, dry matter content (DM), crude protein (CP), acid detergent fiber (ADF), neutral detergent fiber (NDF), acid detergent protein (ADP), phosphorus (P), potassium (K), calcium (Ca), and magnesium (Mg) contents were determined using a NIRS device. In addition, dry matter intake (DMI), digestible dry matter (DDM), relative feed value (RFV), digestible energy (DE), and metabolizable energy (ME) values were calculated using ADF and NDF values. As a result of the analyses, the average dry matter content was determined as 86.64%, crude protein content as 11.23%, ADF as 44.09%, NDF as 64.30%, DDM as 54.55, and RFV as 78.92. The average DE value was found to be 2.60 Mcal/kg and the ME value was found to be 2.14 Mcal/kg. The mineral matter contents were determined as P 0.30%, K 1.28%, Ca 1.49%, and Mg 0.49% on average. The Ca/P ratio was calculated as 4.97 and the K/(Ca+Mg) ratio as 0.65. The findings show that the dry matter content of the pasture grasses in Ulutaş is high, but the crude protein and energy contents are moderate, and the fiber contents are relatively high. This situation can be evaluated as limiting factors for the nutritional value of pasture grasses. Mineral matter contents appear to be balanced. It is recommended to implement sustainable pasture management strategies and carry out pasture improvement studies to increase pasture productivity and forage quality.

Keywords: Ulutaş, Erzurum, Forage Quality, Nutritional Value.

GİRİŞ

Hayvancılık sektöründe yem maliyetleri, üretim giderlerinin büyük bir kısmını oluşturarak işletmelerin kârlılığını doğrudan etkilemektedir. Bu bağlamda, sürdürülebilir hayvancılık için ekonomik kaba yem kaynakları büyük önem taşımakta olup, çayır ve meralar bu kaynakların başında gelmektedir. Dünya genelinde hayvancılık faaliyetlerinde kullanılan kaba yemin yaklaşık %70'inin meralardan sağlandığı bilinmektedir (Lund, 2007). Bu durum, meraların hayvancılık sektörü için vazgeçilmez bir kaynak olduğunu göstermektedir. Ancak, günümüzde artan nüfus, iklim değişikliğinin yol açtığı ekolojik sorunlar ve ekonomik dalgalanmalar, meralardan temin edilen kaliteli kaba yeme olan talebi sürekli artırmaktadır. Bu nedenle, meraların korunması ve sürdürülebilir bir şekilde kullanılması büyük önem taşımaktadır. Meraların tarım arazilerine dönüştürülmesi, yerleşim alanlarının genişletilmesi veya aşırı otlatma gibi amaç dışı kullanımlar, bu doğal kaynakların sunduğu faydaları ciddi şekilde azaltmaktadır. Dolayısıyla, meraların etkin bir şekilde korunması ve sürdürülebilir yönetimi, hem hayvancılık sektörünün geleceği hem de ekolojik denge açısından hayati bir gerekliliktir.

Aşırı otlatma, özellikle kurak ve yarı kurak bölgelerdeki mera alanlarında ciddi ve geri dönüşü zor hasarlara neden olan önemli bir sorundur (Snyman, 2005; Holeček vd., 2011). Bu durum, mera verimliliğini önemli ölçüde düşürerek ekolojik dengeyi bozmaktadır. Meraların sürdürülebilir bir şekilde kullanılabilmesi için, mera yönetim ilkelerine uygun stratejilerin benimsenmesi ve mera popülasyonlarının korunması büyük önem taşır. Ayrıca, risk altındaki bölgelerde rehabilitasyon çalışmalarının yapılması gerekmektedir. Bu hedeflere ulaşmak için, meraların mevcut durumunun doğru bir şekilde analiz edilmesi ve bozulmalarına yol açan faktörlerin belirlenmesi şarttır. Herhangi bir rehabilitasyon yönteminin geliştirilmesi veya uygulanması, vejetasyon özelliklerinin detaylı bir şekilde incelenmesini gerektirir. Aksi

takdirde, beklenen sonuçlar elde edilemeyebilir. Bu nedenle, mera ıslah çalışmalarına başlamadan önce, toprak yapısı, topografya ve bitki örtüsü açısından farklılık gösteren mera bölümlerinin botanik kompozisyonunun, verimlilik ve kalite durumlarının tam olarak ortaya konulması ve bu verilere dayalı olarak uygun ıslah uygulamalarının yapılması kritik bir öneme sahiptir (Çınar vd., 2014; Alay vd., 2016).

Ayrıca, çayır, mera ve yem bitkileri ekim alanlarından elde edilen yemlerin besin değeri, otun kalitesiyle doğrudan bağlantılıdır. Yemin lezzeti, hayvanlar tarafından tüketim oranı, sindirilebilirliği, zararlı madde içeriği, kimyasal ve morfolojik yapısı, enerji ve protein değeri gibi unsurlar yem kalitesini belirleyen temel faktörlerdir. Bununla birlikte, iklim koşulları (sıcaklık, yağış), mevsimsel değişiklikler, bitki örtüsündeki buğdaygil ve baklagil oranı, yükselti ve bakı gibi çevresel faktörler de yem kalitesini etkileyen önemli unsurlar arasında yer alır (Kirilov, 2001; Kaya, 2008).

Bu araştırma, Erzurum'un İspir ilçesinde yer alan Ulutaş Mahallesi meralarında yetişen bitki örtüsünün besin değerleri ve mineral bileşenlerini incelemeyi hedeflemektedir. Meraların sürdürülebilir kullanımı ve hayvan besleme verimliliği açısından, bitki örtüsünün yapısal özellikleri ve besin içeriğinin doğru bir şekilde analiz edilmesi kritik bir öneme sahiptir. Bu amaçla, Ulutaş Mahallesi'nden toplanan bitki örnekleri üzerinde kapsamlı laboratuvar analizleri gerçekleştirilmiştir. Yapılan incelemeler sonucunda, bitkilerin temel besin maddeleri ve mineral kompozisyonları belirlenmiştir. Elde edilen bulgular, bölgedeki hayvancılık faaliyetlerinin meralardan nasıl daha etkin bir şekilde faydalanabileceğini ve bu alanların taşıdığı potansiyeli ortaya koymaktadır. Ayrıca, bu çalışmanın, bölgeye özgü mera yönetimi stratejilerinin geliştirilmesine bilimsel bir temel oluşturması amaçlanmıştır.

MATERYAL VE METOD

Bu çalışma, 2020 yılı içerisinde Erzurum ilinin İspir ilçesine bağlı Ulutaş Mahallesi sınırları içerisinde yürütülmüştür. Araştırma alanı, coğrafi olarak 40° 34' 34" Kuzey enlemi ve 40° 54' 23" Doğu boylamı koordinatlarında konumlanmakta olup, deniz seviyesinden yaklaşık 2100 metre yüksekliktedir ve İspir ilçe merkezine yaklaşık 24 kilometre mesafededir. Çalışma alanının coğrafi konumu Şekil 1'de harita üzerinde gösterilmekte, bölgeden elde edilen bazı fotoğraflar ise Şekil 2'de sunulmaktadır. Erzurum iline ait 2020 yılı meteorolojik verilerine göre, toplam yağış miktarı 295,6 mm, ortalama sıcaklık 5,6 °C ve ortalama nisbi nem oranı %67 olarak kaydedilmiştir (Anonim, 2021).



Şekil 1. Çalışma alanının konumu (Google Earth)





Şekil 2. Çalışma alanının çekilen fotoğraflar

Bu çalışmada bitki örnekleme, 2020 yılının Temmuz ayında Ulutaş mahallesi mera alanından seçilen 12 farklı noktada gerçekleştirilmiştir. Örnekleme süreci, her bir noktada 50x50 cm boyutlarındaki kareler kullanılarak, bitkilerin toprak yüzeyinden kesilerek toplanması şeklinde yürütülmüştür. Toplanan ot örnekleri, bitkisel materyalin kimyasal analizlere hazırlanması amacıyla öncelikle 60 °C sıcaklıkta etüvde kurutulmuş ve ardından 1 mm elek çaplı bir değirmende öğütülerek homojen bir yapıya getirilmiştir. Bu şekilde hazırlanan bitki örneklerinde kuru madde oranı (KMO), ham protein (HP), asit deterjan lifi (ADF), nötr deterjan lifi (NDF), asit deterjan proteini (ADP), fosfor (P), potasyum (K), kalsiyum (Ca) ve magnezyum (Mg) içerikleri, Foss NIR Systems Model 6500 Win ISI II v1.5 model NIRS cihazı kullanılarak belirlenmiştir.

Buna ek olarak, her bir bitki örneği için ADF ve NDF değerleri temel alınarak kuru madde tüketimi (KMT), sindirilebilir kuru madde (SKM), nispi yem değeri (NYD), sindirilebilir enerji (SE) ve metabolik enerji (ME) değerleri, literatürde yer alan aşağıdaki formüller aracılığıyla hesaplanmıştır:

$$\text{Sindirilebilir Kuru Madde (SKM)} = 88,9 - (0,779 \times \% \text{ADF}) \text{ (Oddy vd., 1983)}$$

$$\text{Kuru Madde Tüketimi (KMT)} = 120 / (\% \text{NDF}) \text{ (Sheaffer vd., 1995)}$$

$$\text{Nispi Yem Değeri (NYD)} = (\text{SKM} \times \text{KMT}) / 1,29 \text{ (Sheaffer vd., 1995)}$$

$$\text{Sindirilebilir Enerji (SE)} = 0,27 + 0,0428 \times (\% \text{SKM}) \text{ (Fonnesbeck vd., 1984)}$$

$$\text{Metabolik Enerji (ME)} = 0,821 \times \text{SE (Mcal/kg)} \text{ (Khalil vd., 1986)}$$

Analiz sonuçlarının değerlendirilmesinde, makro element içerikleri arasındaki ilişkileri incelemek amacıyla Ca/P ve K/(Ca+Mg) oranları hesaplanmıştır. Elde edilen verilerin istatistiksel analizleri, JMP 13 istatistik yazılımı kullanılarak tanımlayıcı istatistik yöntemleri ile gerçekleştirilmiştir.

BULGULAR VE TARTIŞMA

Ulutaş Mahallesi mera alanından elde edilen ot örneklerinin kalite özelliklerine ilişkin analiz sonuçları Tablo 1'de detaylı olarak sunulmuştur. Tablo 1'in incelenmesiyle, analizlere tabi tutulan ot örneklerinin kuru madde oranları, ham protein içerikleri, lif oranları (ADF ve NDF), enerji değerleri (sindirilebilir enerji (SE) ve metabolik enerji (ME)) ve mineral madde kompozisyonları (fosfor (P), potasyum (K), kalsiyum (Ca) ve magnezyum (Mg)) ile bu mineraller arasındaki oranlar (Ca/P ve K/(Ca+Mg)) belirlenmiştir.

Tablo 1. Ulutaş Mahallesi merasının besin değerleri

Kalite Özellikleri	Ortalama±Std sapma (Minimum – Maksimum)
Kuru Madde Oranı (KM) (%)	86.64±3.28 (84.40 – 90.38)
Ham Protein Oranı (HP) (%)	11.23±1.23 (10.23 – 12.61)
Asit Deterjan Fiber (ADF) (%)	44.09±4.13 (41.32 – 48.84)
Nötr Deterjan Fiber (NDF) (%)	64.30±4.36 (61.29 - 69.30)
Asit Deterjan Protein (ADP) (%)	1.54±0.13 (1.46 – 1.69)
Sindirilebilir Kuru Madde (SKM)	54.55±3.22 (50.85 – 56.71)
Kuru Madde Tüketimi (KMT)	1.87±0.13 (1.71 – 1.96)
Nisbi Yem Değeri (NYD)	78.92±10.44 (66.95 – 86.07)
Sindirilebilir Enerji (SE) (Mcal/kg)	2.60±0.14 (2.45 – 2.70)
Metabolik Enerji (ME) (Mcal/kg)	2.14±0.11 (2.01 – 2.21)
Fosfor (P) (%)	0.30±0.08 (0.25 – 0.39)
Potasyum (K) (%)	1.28±0.09 (1.22 – 1.38)
Kalsiyum (Ca) (%)	1.49±0.08 (1.44 – 1.58)
Magnezyum (Mg) (%)	0.49±0.06 (0.45 – 0.56)
Ca/P	4.97±1.22 (3.56 – 5.76)
K/(Ca+Mg)	0.65±0.01 (0.64 – 0.65)

Tablo 1'e bakıldığında, **Kuru Madde Oranı (KM)**: Analiz edilen ot örneklerinin kuru madde oranı ortalama %86,64 olarak bulunmuştur. Standart sapma $\pm 3,28$ olup, minimum ve maksimum değerler sırasıyla %84,40 ve %90,38'dir. Bu sonuçlar, mera otlarının yüksek oranda kuru madde içerdiğini göstermektedir. Yüksek kuru madde oranı, otun depolanması ve muhafazası açısından avantaj sağlamaktadır.

Ham Protein Oranı (HP): Ot örneklerinin ortalama ham protein içeriği %11,23 olarak tespit edilmiştir. Standart sapma $\pm 1,23$ olup, en düşük ve en yüksek değerler sırasıyla %10,23 ve %12,61'dir. Bu değerler, mera otlarının hayvanların protein ihtiyacını karşılamada orta düzeyde bir potansiyele sahip olduğunu göstermektedir. Ancak, hayvan türü ve fizyolojik dönemine göre bu değerin yeterliliği değişebilir.

Asit Deterjan Fiber (ADF) ve Nötr Deterjan Fiber (NDF): Ot örneklerinin ortalama ADF değeri %44,09 ($\pm 4,13$) ve ortalama NDF değeri %64,30 ($\pm 4,36$) olarak belirlenmiştir. ADF ve NDF değerleri, bitkisel materyalin sindirilebilirliğini etkileyen önemli faktörlerdir. Yüksek

ADF ve NDF değerleri, otun sindirilebilirliğinin düşük olduğunu gösterir. Bu sonuçlar, mera otlarının orta düzeyde bir sindirilebilirliğe sahip olduğunu düşündürmektedir.

Asit Deterjan Protein (ADP): Örneklerin ortalama ADP içeriği %1,54 ($\pm 0,13$) olarak bulunmuştur. ADP, bitkideki bağlanmış proteini ifade eder ve sindirilebilirliği düşüktür. Düşük ADP değeri, otun protein kalitesinin nispeten iyi olduğunu göstermektedir.

Sindirilebilir Kuru Madde (SKM), Kuru Madde Tüketimi (KMT) ve Nispi Yem Değeri (NYD): Hesaplamalar sonucunda ortalama SKM %54,55 ($\pm 3,22$), ortalama KMT 1,87 ($\pm 0,13$) ve ortalama NYD 78,92 ($\pm 10,44$) olarak bulunmuştur. SKM, otun sindirilebilir kısmını, KMT hayvanın otu ne kadar tüketeceğini ve NYD ise otun genel yem kalitesini gösterir. Bu değerler, mera otlarının orta düzeyde bir yem kalitesine sahip olduğunu ve hayvanlar tarafından orta düzeyde tüketilebileceğini göstermektedir.

Sindirilebilir Enerji (SE) ve Metabolik Enerji (ME): Ot örneklerinin ortalama SE değeri 2,60 Mcal/kg ($\pm 0,14$) ve ortalama ME değeri 2,14 Mcal/kg ($\pm 0,11$) olarak tespit edilmiştir. Bu değerler, mera otlarının hayvanlara sağlayabileceği enerji miktarını göstermektedir. Elde edilen sonuçlar, mera otlarının orta düzeyde bir enerji içeriğine sahip olduğunu göstermektedir.

Mineral Madde İçerikleri (P, K, Ca ve Mg): Ot örneklerinin ortalama fosfor (P) içeriği %0,30 ($\pm 0,08$), potasyum (K) içeriği %1,28 ($\pm 0,09$), kalsiyum (Ca) içeriği %1,49 ($\pm 0,08$) ve magnezyum (Mg) içeriği %0,49 ($\pm 0,06$) olarak bulunmuştur. Bu mineraller, hayvanların sağlıklı gelişimi için gereklidir. Elde edilen değerler, mera otlarının bu mineraller açısından orta düzeyde bir zenginliğe sahip olduğunu göstermektedir.

Ca/P ve K/(Ca+Mg) Oranları: Mineral oranları incelendiğinde, ortalama Ca/P oranı 4,97 ($\pm 1,22$) ve ortalama K/(Ca+Mg) oranı 0,65 ($\pm 0,01$) olarak hesaplanmıştır. Bu oranlar, mineraller arasındaki dengenin hayvan sağlığı için uygun olup olmadığını değerlendirmede kullanılır. Elde edilen Ca/P oranının yüksek olması, rasyonda yeterli fosfor bulunup bulunmadığının kontrol edilmesini gerektirebilir. K/(Ca+Mg) oranının ise nispeten dar bir aralıkta seyrettiği görülmektedir.

Baykal vd., (2024) tarafından 2020 yılında Lup metodu kullanılarak aynı bölgede (Ulutaş Mahallesi'nde (İspir/Erzurum)) gerçekleştirilen çalışmada, mera alanının toprak örtüsü, botanik kompozisyonu, kalite derecesi ve durum sınıfı özellikleri incelenmiştir. Bu çalışmada, araştırma alanında 7 *Poaceae*, 12 *Fabaceae* ve diğer familyalara ait 49 olmak üzere toplam 68 takson tespit edilmiştir. Mera alanının toplam toprak örtüsü %84,75 olarak belirlenirken, botanik kompozisyon oranları *Poaceae* için %14,70, *Fabaceae* için %33,14 ve diğer familyalar için %52,16 olarak saptanmıştır. Elde edilen 3,59 mera kalite derecesi, meranın durumunun "Zayıf" olduğunu göstermektedir. Bu bulgular, Ulutaş Mahallesi merasının bitki örtüsü çeşitliliğinin nispeten yüksek olduğunu ancak baskın türlerin genellikle istenmeyen familyalardan oluştuğunu ve genel mera kalitesinin düşük olduğunu ortaya koymaktadır. Bu durum, mera otlarının besin değerlerini de etkileyebilecek bir faktör olarak değerlendirilmelidir.

Yapılan bu çalışmada, aynı mera alanından toplanan ot örneklerinin besin değerleri ve mineral madde içerikleri analiz edilerek, mera kalitesine ilişkin daha detaylı bilgiler elde edilmiştir (Tablo 1). Analiz sonuçlarına göre, ot örneklerinin ortalama kuru madde oranı %86,64 ($\pm 3,28$) olarak bulunmuştur. Bu yüksek kuru madde oranı, otun depolanması ve muhafazası için olumlu bir özellik olarak değerlendirilebilir. Ortalama ham protein içeriği %11,23 ($\pm 1,23$) olarak tespit edilmiştir. Bu değer, mera otlarının hayvanların protein ihtiyacını karşılama potansiyelinin orta düzeyde olduğunu göstermektedir. Lif oranları (ADF: %44,09 $\pm 4,13$; NDF: %64,30 $\pm 4,36$) ise otların sindirilebilirliğinin orta düzeyde olduğunu işaret etmektedir. Ayrıca, ortalama sindirilebilir enerji (SE) değeri 2,60 Mcal/kg ($\pm 0,14$) ve metabolik enerji (ME) değeri 2,14 Mcal/kg ($\pm 0,11$) olarak bulunmuştur. Mineral madde içerikleri incelendiğinde ise, ortalama fosfor (P) %0,30 ($\pm 0,08$), potasyum (K) %1,28 ($\pm 0,09$), kalsiyum (Ca) %1,49 ($\pm 0,08$) ve

magnezyum (Mg) %0,49 ($\pm 0,06$) olarak tespit edilmiştir. Ca/P oranı 4,97 ($\pm 1,22$) ve K/(Ca+Mg) oranı 0,65 ($\pm 0,01$) olarak hesaplanmıştır. Bu bulgular, Baykal vd. (2024)'ün mera durumunun "Zayıf" olduğu tespitiyle örtüşmektedir. Düşük mera kalite derecesi ve baskın istenmeyen türler, otların besin değerlerini ve dolayısıyla mera verimliliğini olumsuz etkilemektedir. Bu durum, mera ıslah çalışmalarının gerekliliğini ortaya koymaktadır.

Bu çalışmada Ulutaş Mahallesi merasından elde edilen ot örneklerinin besin değerleri, literatürdeki benzer çalışmalarla karşılaştırıldığında bazı farklılıklar ve benzerlikler göstermektedir. Tablo 1'de sunulan bulgulara göre, Ulutaş merasının ortalama ham protein (HP) içeriği %11,23 olarak bulunmuştur. Bu değer, Kılıç (2018)'in Trabzon'daki merada saptadığı %16,6 HP oranından düşüktür. Benzer şekilde, Şahinoğlu (2010)'nun Samsun'daki merada bulunduğu %16,33-18,64 aralığındaki HP değerlerinden ve Nadir (2010)'in Tokat'taki merada tespit ettiği %16,48-18,81 aralığındaki değerlerden de daha düşüktür. Ancak, Ulutaş merasının HP içeriği, Güllap (2010)'in Erzurum'daki farklı mera kesimlerinde bulunduğu %8,26-13,12 aralığına yakın ve Parlak vd. (2015)'nin Çanakkale'deki meralarda belirlediği %9,10-13,18 aralığı içinde yer almaktadır. Ayrıca, Tutar ve Kökten (2019)'in Bingöl'deki merada bulunduğu %9,9-12,9 aralığına da benzerlik göstermektedir. Aydın ve Başbağ (2017)'in Karacadağ meralarında saptadığı %19,19'luk HP değeri ise Ulutaş merasının oldukça üzerindedir. Bu farklılıklar, meraların bulunduğu bölgenin iklim koşulları, bitki türleri, toprak özellikleri ve otlatma yönetimi gibi faktörlerden kaynaklanabilir.

Lif oranları (ADF ve NDF) açısından bakıldığında, Ulutaş merasının ADF (%44,09) ve NDF (%64,30) değerleri, Şahinoğlu (2010), Nadir (2010) ve Aydın ve Başbağ (2017)'in çalışmalarında saptanan değerlerden daha yüksektir. Bu durum, Ulutaş merasının otlarının sindirilebilirliğinin diğer meralara göre daha düşük olabileceğini göstermektedir. Ancak, Güllap (2010)'in Erzurum'daki meralarda bulunduğu geniş ADF (%25,83-51,35) ve NDF (%43,57-50,28) aralıkları içinde bazı değerler Ulutaş merasının değerlerine yakın seyretmektedir. Benzer şekilde, Parlak vd. (2015)'nin Çanakkale'deki meralarda belirlediği NDF (%43,18-51,57) ve ADF (%29,40-31,73) oranları da Ulutaş merasına göre daha düşüktür. Tutar ve Kökten (2019) çalışmasında ise NDF oranları (% 52.5 ile % 62.7 arasında) Ulutaş merasına yakın değerler gösterirken, ADF oranları (%34.8 ile %37.4 arasında) daha düşüktür. Nispi yem değeri (NYD) açısından bakıldığında, Ulutaş merasının NYD değeri (78,92), Kılıç (2018)'in bulunduğu 112,9 ve Nadir (2010)'in bulunduğu 174,96-189,77 aralığından belirgin şekilde düşüktür. Aydın ve Başbağ (2017) çalışmasındaki 137.7 NYD değeri de Ulutaş merasının üzerindedir. Tutar ve Kökten (2019) çalışmasında ise NYD oranları (91.8 ile 109.4 arasında) Ulutaş merasına göre daha yüksek bulunmuştur. Bu farklılıklar, meraların bitki kompozisyonu, olgunluk dönemi ve çevresel faktörler gibi etkenlerden kaynaklanmaktadır. Mineral madde içerikleri açısından da benzer farklılıklar gözlenmektedir. Örneğin, Ulutaş merasının K içeriği Şahinoğlu (2010) ve Aydın ve Başbağ (2017)'in bulunduğu değerlerden daha düşüktür. Bu karşılaştırmalar, Ulutaş merasının besin değerlerinin ve mineral madde içeriğinin, diğer bölgelerdeki meralara göre bazı farklılıklar gösterdiğini ve bu farklılıkların bölgesel koşullar ve mera yönetimi uygulamalarından kaynaklandığını ortaya koymaktadır.

SONUÇ

Bu çalışmada, Erzurum ili İspir ilçesine bağlı Ulutaş Mahallesi meralarının ot kalitesi ve besin değerleri 2020 yılı Temmuz ayında toplanan bitki örnekleri üzerinden incelenmiştir. Bu mera alanından 12 farklı noktadan 50x50 cm'lik kareler kullanılarak ot örnekleri toplanmıştır. Örneklerin laboratuvar analizleri sonucunda, ortalama kuru madde oranını %86,64, ham protein içeriğini %11,23, ADF %44,09, NDF %64,30, SKM 54,55 ve NYD 78,92 olduğu tespit edilmiştir. Ayrıca, ortalama SE değeri 2,60 Mcal/kg ve ME değeri 2,14 Mcal/kg olarak belirlenmiştir. Mineral madde içerikleri ise ortalama olarak fosfor (P) %0,30, potasyum (K)

%1,28, kalsiyum (Ca) %1,49 ve magnezyum (Mg) %0,49 olarak bulunmuştur. Bu verilerden hesaplanan Ca/P oranı 4,97 ve K/(Ca+Mg) oranı 0,65 olarak tespit edilmiştir. Bu bulgular, Ulutaş merasının otlarının kuru madde içeriğinin yüksek olduğunu, ancak ham protein ve enerji içeriklerinin orta düzeyde, lif içeriklerinin ise nispeten yüksek olduğunu göstermektedir.

Elde edilen sonuçlara göre, Ulutaş Mahallesi meralarının hayvan beslemesi açısından orta düzeyde bir potansiyele sahip olduğunu ortaya koymaktadır. Yüksek kuru madde oranı, otun depolanması ve muhafazası açısından avantaj sağlarken, orta düzeydeki ham protein ve enerji içerikleri ile nispeten yüksek lif oranları, mera otlarının besin değerini sınırlayan faktörler olarak değerlendirilebilir. Mineral madde içeriklerinin ve oranlarının da dengeli olduğu görülmektedir. Ancak, mera yönetim uygulamaları ve bitki kompozisyonu gibi faktörlerin ot kalitesini etkilediği göz önünde bulundurulmalıdır. Bu nedenle, mera verimliliğini ve ot kalitesini artırmaya yönelik sürdürülebilir mera yönetim stratejilerinin uygulanması ve mera ıslah çalışmalarının yapılması önerilmektedir.

KAYNAKLAR

- Alay, F., İspirli, K., Uzun, F., Çınar, S., Aydın, İ. & Çankaya, N. (2016). Uzun süreli serbest otlatmanın doğal meralar üzerine etkileri. *Gaziosmanpaşa Üniversitesi Ziraat Fakültesi Dergisi*, 33(1): 116-124.
- Anonim. (2021). T.C. Başbakanlık Devlet Meteoroloji İşleri Genel Müdürlüğü, Erzurum İl Müdürlüğü Kayıtları.
- Aydın, A. & Başbağ, M. (2017). Karacadağ'ın farklı yükseltilerindeki meraların durumu ve ot kalitesinin belirlenmesi. *Anadolu Tarım Bilimleri Dergisi*, 32(1), 74-84.
- Baykal, H., Çatal, M.İ., & Bakoğlu, A. (2024). Ulutaş Mahallesi'nin (İspir/ERZURUM) Botanik Kompozisyonu ve Mera Durumunun Belirlenmesi. *Journal of Anatolian Environmental and Animal Sciences*, 9(4), 669-675.
- Çınar, S., Hatipoğlu, R., Avcı, M., İnal, İ., Yücel, C. & Avağ, A. (2014). Hatay ili Kırıkhan ilçesi taban meralarının vejetasyon yapısı üzerine bir araştırma. *Gaziosmanpaşa Üniversitesi Ziraat Fakültesi Dergisi*, 31(2): 52-60.
- Fonnesbeck, P.V., Clark, D.H., Garret, W. N. & Speth, C.F. (1984). Predicting energy utilization from alfalfa hay from the Western Region. *Proceeding of American Society of Animal Sciences (Western Section)*, 35: 305-308.
- Güllap, M.K. (2010). *Kargapazarı Dağında (Erzurum) farklı otlatma sistemi uygulamalarının mera bitki örtüsüne etkisi*. Doktora Tezi, Atatürk Üniversitesi, Erzurum.
- Holechek, J.L., Pieper, R.D. & Herbel, C.H. (2011). *Range management principles and practices*. Sixth edition. Prentice Hall, Upper Saddle River, New Jersey, USA, 444 sy.
- Kaya, Ş. (2008). Kaba yemlerin değerlendirilmesinde göreceli yem değeri ve göreceli kaba yem kalite indeksi. *Türk Bilimsel Derlemeler Dergisi*, 1(1): 59-64.
- Khalil, J.K., Sawaya, W.N. & Hyder, S.Z. (1986). Nutrient composition of Atriplex leaves grown in Saudi Arabia. *Journal of Range Management*, 39: 104-107.
- Kılıç, S. (2018). *Trabzon ili Düzköy ilçesi Beypınarı merasında farklı gübre uygulamalarının meranın verim, kalite ve botanik kompozisyonuna etkileri üzerine bir araştırma*. Yüksek Lisans Tezi, Tokat Gaziosmanpaşa Üniversitesi, Tokat.
- Kirilov, A. (2001). *Lucerne quality and possibilities for its estimation*, In : I. Delgado and J. Lloveras (Eds.), *Quality in lucerne and medics for animal production*. Zaragoza, CIHEAM, p. 231-234, <http://om.ciheam.org/om/pdf/a45/01600089.pdf> (Erişim tarihi: 23.10.2016).

- Lund, H.G. (2007). Accounting for the world's rangelands. *Rangelands*, 29(1), 3-10.
- Nadir, M. (2010). *Tokat ili Yeşilyurt köyü doğal merasının botanik kompozisyon, kuru madde verimi ve kalitesinin belirlenmesi*. Yüksek Lisans, Gaziosmanpaşa Üniversitesi, Tokat.
- Oddy, V.H., Robards, G.E. & Low, S.G. (1983). *Prediction of in vivo dry matter digestibility from the fiber nitrogen content of a feed*. In: Feed Information and Animal Production. (ed) Robards, G.E., Packham, R.G., Commonwealth Agricultural Bureau, Farnham Royal, UK, 395-398.
- Parlak, A.Ö., Parlak, M., Gökkuş, A. & Demiray, H.C. (2015). Akdeniz (Çanakkale) meralarının ot verimi ve kalitesi ile botanik kompozisyonu ve bazı toprak özellikleri. *Çanakkale On Sekiz Mart Üniversitesi Ziraat Fakültesi Dergisi*, 3(1), 99-108.
- Sheaffer, C.C., Peterson, M.A., Mccalin, M., Volene, J.J., Cherney, J.H., Johnson, K.D., Woodward, W.T. & Viands, D.R. (1995). Acid detergent fiber, neutral detergent fiber concentration and relative feed value. North American Alfalfa Improvement Conference, Minneapolis.
- Snyman, H.A. (2005). Rangeland degradation in a semi-arid South Africa. I: influence on seasonal root distribution, root/shoot ratios, and water-use efficiency. *Journal of Arid Environments*, 60, 457-481. <https://doi.org/10.1016/j.jaridenv.2004.06.006>
- Şahinoğlu, O. (2010). *Bafra ilçesi Koşu köyü merasında uygulanan farklı ıslah yöntemlerinin meranın ot verimi, yem kalitesi ve botanik kompozisyonu üzerine etkileri*. Doktora Tezi, Ondokuz Mayıs Üniversitesi, Samsun.
- Tutar, H. & Kökten, K. (2019). Mera vejetasyon özelliklerinin farklı yönelere göre değişimi. *Türkiye Tarımsal Araştırmalar Dergisi*, 6(3), 312-318.

**INNOVATIVE BIOCONTROL STRATEGIES FOR SUSTAINABLE WEED
MANAGEMENT**

**SÜRDÜRÜLEBİLİR YABANCI OT YÖNETİMİ İÇİN YENİLİKÇİ BİYOLOJİK
KONTROL STRATEJİLERİ**

Ayfer GÜNEY SARITAŞ

Iğdır University, Postgraduate Education Institute, Department of Agricultural Sciences, Iğdır,
Türkiye

ORCID ID: <https://orcid.org/0000-0001-7641-9711>

Ramazan GÜRBÜZ

Iğdır University, Faculty of Agriculture, Department of Plant Protection Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0003-3558-9823>

ABSTRACT

Weed management plays a key role in ensuring the sustainability of agriculture and has a direct impact on crop productivity, biodiversity and ecosystem health. Conventional chemical-based weed control methods, although widely used and effective, have attracted considerable criticism for their negative impacts, including environmental degradation, harm to non-target species and the alarming rise in herbicide-resistant weed populations. In response to these concerns, innovative biological control (biocontrol) strategies are emerging as viable, environmentally friendly alternatives. These strategies harness the potential of natural predators, pathogens and competing plant species to suppress weed growth in a targeted and sustainable manner. By minimising chemical inputs, reducing pollution and improving soil health, the implementation of biocontrol measures can promote ecological balance. Recent advances in biocontrol research have led to the identification of specific fungal pathogens, insects and allelopathic plants that show promise in controlling invasive and persistent weed species. Case studies such as the use of fungal pathogens to control *Parthenium hysterophorus* and the use of bioherbivores to control *Striga* spp. highlight the practical applications and potential scalability of biocontrol techniques. However, challenges such as the unpredictability of field performance, the risk of non-target effects and the need for a regulatory framework remain significant barriers to widespread adoption. This paper provides an overview of emerging trends, practical applications and the socio-economic implications of bio-control strategies in the sustainable management of weeds. By addressing current challenges and fostering interdisciplinary collaboration, biocontrol can become a cornerstone in achieving global agricultural sustainability.

Keywords: Biological control, Sustainable agriculture, Weed management, Herbicide resistance, Ecological balance

ÖZET

Yabancı ot yönetimi, tarımın sürdürülebilirliğini sağlamada önemli bir rol oynar ve ürün verimliliği, biyolojik çeşitlilik ve ekosistem sağlığı üzerinde doğrudan bir etkiye sahiptir. Yaygın olarak kullanılan ve etkili olan geleneksel kimyasal bazlı ot kontrol yöntemleri, çevresel

bozulma, hedef dışı türlere zarar verme ve herbisite dirençli ot popülasyonlarındaki endişe verici artış gibi olumsuz etkileri nedeniyle önemli eleştirilere maruz kalmıştır. Bu endişelere yanıt olarak, yenilikçi biyolojik kontrol (biyokontrol) stratejileri uygulanabilir, çevre dostu alternatifler olarak ortaya çıkmaktadır. Bu stratejiler, ot büyümesini hedefli ve sürdürülebilir bir şekilde bastırmak için doğal yırtıcıların, patojenlerin ve rekabet eden bitki türlerinin potansiyelinden yararlanır. Kimyasal girdileri en aza indirerek, kirliliği azaltarak ve toprak sağlığını iyileştirerek, biyokontrol önlemlerinin uygulanması ekolojik dengeyi teşvik edebilir. Biyokontrol araştırmalarındaki son gelişmeler, istilacı ve kalıcı ot türlerini kontrol etmede umut vadeden belirli mantar patojenlerinin, böceklerin ve allelopatik bitkilerin tanımlanmasına yol açmıştır. *Parthenium hysterophorus*'u kontrol etmek için mantar patojenlerinin kullanımı ve *Striga spp.*'yi kontrol etmek için biyoherbivorların kullanımı gibi vaka çalışmaları. biyolojik kontrol tekniklerinin pratik uygulamalarını ve potansiyel ölçeklenebilirliğini vurgulamak. Ancak, saha performansının öngörülemezliği, hedef dışı etki riski ve düzenleyici bir çerçeveye ihtiyaç duyulması gibi zorluklar yaygın bir şekilde benimsenmesinin önünde önemli engeller olmaya devam ediyor. Bu makale, yabancı otların sürdürülebilir yönetiminde biyolojik kontrol stratejilerinin ortaya çıkan eğilimlerine, pratik uygulamalarına ve sosyo-ekonomik etkilerine genel bir bakış sunmaktadır. Mevcut zorlukları ele alarak ve disiplinler arası iş birliğini teşvik ederek, biyolojik kontrol küresel tarımsal sürdürülebilirliğe ulaşmada bir temel taşı haline gelebilir.

Anahtar kelimeler: Biyolojik kontrol, Sürdürülebilir tarım, Yabancı ot yönetimi, Herbisit direnci, Ekolojik denge

GİRİŞ

Yabancı otlar, ürün verimini azaltmalarına rağmen tarımsal ekosistemlerin dengesi için önemli bir rol oynar. Bu sebeple, bu ot türlerinin yayılımlarını ve yoğunluklarını anlamak amacıyla, tarım sistemlerini incelemek ekolojik yöntemlerin kullanılması gereklidir (Boström, 2002; Tomita ve ark., 2003). Sürdürülebilir tarımda etkili bir yabancı ot yönetimi oluşturabilmek için, öncelikle yabancı ot biyolojisi üzerine temel araştırmaların yapılması gereklidir. Bu çalışmalar, popülasyon dinamikleri, büyüme ve gelişim süreçleri, biyotik ve abiyotik agroekosistem faktörleriyle etkileşimler, fenoloji, ürünlere müdahale, allelopati ve rekabet (tohum bankası) gibi konuları daha iyi anlamak gereklidir (Altieri and Nicholls 2000). Bu tür araştırmalar, yabancı ot yönetim stratejilerinin doğru bir şekilde uygulanmasını sağlayacaktır. Özellikle, yabancı otları azaltmanın belirli yöntemlerini kullanmak ve ürün kalitesi ile verim üzerinde olumsuz etkilerden kaçınmak önem arz etmektedir (Canizales, 2010). Yabancı otlar, toprağın kalitesini artırma ve biyolojik çeşitliliği destekleme gibi birçok ekosistem işlevi sağlar ve bu, uzun vadede tarımsal ekosistemlerin verimliliğini korumaya katkıda bulunmaktadır (MacLaren, 2020). Bu nedenle, sürdürülebilir tarımın doğal kaynakları koruma ve tarımı daha düşük maliyetle geliştirmesi sayesinde çevresel, sosyal ve ekonomik sebeplerden dolayı sürdürülebilir yabancı ot kontrolüne geçiş yapılması gerekmektedir (Gnanavel, 2015).

Yabancı otlar, tarımsal zararlılar içinde en maliyetli grubu oluşturur ve kültür bitkilerinde %45'ten fazla verim kaybına neden olmaktadır. Bu oran, ürünlerde hastalıklar (%25) ve böcek zararlıları (%20) ile kıyaslandığında oldukça yüksektir. Yabancı otların neden olduğu verim kayıpları; yabancı otların çıkış zamanı, yoğunluğu, türü gibi çeşitli faktörlere bağlıdır. Eğer yabancı otlar kontrol edilmezse, %100 verim kaybına yol açabilir (Peerzada, 2019). Genel anlamda, biyolojik çeşitliliği ve ekosistem işleyişini destekleyen tarımsal uygulamalar, tarımsal gelişmiş ekosistem işleyişini sürdürebilmek için önemli faydalar sağlar (MacLaren, 2020). Yabancı otlar, tür, topluluk ve ekosistem düzeylerinde önemli olumsuz ekolojik etkilere sebep olabilmektedirler (Vilà ve ark., 2011). Yabancı otlar, geniş tarım alanlarından meralara, su kütlelerine, doğal sistemlere ve rekreasyon alanlarına kadar birçok ekosistemde ciddi sorunlara

neden olmaktadır. Yabancı otların büyümesini engelleyebilen tüm canlı organizmalar genel olarak biyolojik kontrol ajanları olarak kabul edilir. Ancak, eklembecaklılar (böcekler, akarlar) ve bitki patojenleri gibi özel ajanlar, hedefli yabancı ot kontrolü için daha uygun olup yaygın olarak kullanılmaktadır (Charudattan, 2001; McFadyen, 1998; Schwarzländer ve ark., 2018; Winston ve ark., 2014).

Yabancı Otların Tarım ve Ekosistem Açısından Faydaları ve Rolü

Yabancı ot yönetimi, sürdürülebilir tarım uygulamalarının merkezinde yer alır. Yabancı otlar, ekinlerle rekabet ederek verimliliği düşürebilir, ancak uygun yönetimle bu sorunların önüne geçmek mümkündür. Uygun yabancı ot yönetimi, tarımda biyolojik çeşitliliği korur ve toprak sağlığını destekler (Altieri, 1999; Sheley ve Petroff, 1999). Yabancı otların, doğru yönetildiğinde ekosistem ve tarım üzerinde birçok olumlu etkisi vardır. Yabancı otlar, toprak erozyonunu önleyerek toprağı stabilize eder. Bazı yabancı ot türleri, toprağın organik madde içeriğini artırarak toprak sağlığını iyileştirir (Marshall ve Moonen, 2002). Yabancı otlar, çeşitli böcek ve hayvan türlerine yaşam alanı sağlar. Bu, ekosistemin genel sağlığını ve dirençliliğini artırır (Altieri, 1999). Birçok yabancı ot, arılar ve diğer tozlayıcılar için önemli bir nektar kaynağıdır. Bu, ekinlerin tozlaşma oranını ve dolayısıyla verimliliğini artırır (Marshall ve Moonen, 2002). Yabancı otlar, mikrohabitat oluşturarak faydalı organizmalara yaşam alanı sunar. Bu organizmalar zararlı böceklerle doğal yollarla mücadele eder (Sheley ve Petroff, 1999). Yabancı otlar, besin döngüsü ve toprak oluşumu gibi ekosistem hizmetlerine katkıda bulunur. Bu hizmetler, tarımsal sürdürülebilirliği ve uzun vadeli verimliliği artırır (Altieri, 1999).

Herbisitlerin Çevre Üzerindeki Olumsuz Etkileri

Tarımın temel hedefi, ekolojik dengenin korunarak, birim alandan en yüksek ve kaliteli ürünlerin elde edilmesidir. Bu hedefi kısıtlayan başlıca faktörler arasında hastalıklar, zararlılar ve yabancı otlar yer almaktadır (Topal, 2011a). Kültür bitkilerindeki üretim üzerinde en çok etkisi olan faktörlerden biri kesinlikle yabancı otlardır. Herbisitler, yabancı otlarla mücadelede en etkili ve hızlı çözüm olarak görülse de, bu her zaman beklenen sonuçları sağlamayabilir. Herbisitlerin çevre kirliliğine yol açması, insan sağlığına olumsuz etkilerinin bulunması, yabancı otlarda direnç gelişimi ve ülkemiz için önemli miktarda döviz kaybına neden olması gibi olumsuz sonuçlar göz önünde bulundurulmalıdır (DeBach, 1964; Zengin, 2013). Sürekli olarak yüksek miktarlarda sentetik herbisit kullanımı, çevreye zarar vermekte ve herbisitlere karşı dirençli yabancı otların artışına yol açmaktadır (Bhadoria, 2011). Bu durum ekonomik olarak zararlı olmayan türlerin zamanla baskın hale gelmesine, biyolojik çeşitliliğin azalmasına ve kültür bitkilerinde fitotoksiste gibi olumsuz etkilerin ortaya çıkmasına neden olabilmektedir. Bu gibi durumlar, birçok çevresel ve tarımsal soruna yol açmaktadır (Kitiş, 2009). Bu durum, tarımda herbisitlerin kullanımını ve yabancı otların yönetimini karmaşık hale getirir. Bu nedenle, daha sürdürülebilir ve etkili yöntemlerin geliştirilmesini zorunlu kılmaktadır.

Tüm bu nedenlerle, biyolojik kontrol üzerine yapılan araştırmaların önemi açıkça ortaya çıkmaktadır. Biyolojik yabancı ot kontrolü, bitkiyle beslenen veya patojen olan doğal düşmanların kullanımını içerir. Biyokontrolün amacı, yabancı otları tamamen yok etmek değil, popülasyon ekolojisi prensiplerine dayanarak yabancı ot yoğunluğunu tolere edilebilir seviyelerde tutmaktır. Pratikte biyolojik kontrol, faydalı bitki türlerine zarar vermeden, üzerinde bulunduğu konukçu bitkilerine doğrudan veya dolaylı olarak zarar veren ya da zayıflatan doğal düşmanların desteklenmesi ve korunmasını sağlamaktadır (DeBach, 1964; Zengin, 2013). Bu makale biyolojik yabancı ot kontrolünün yöntemlerini ve uygulamalarını detaylandırarak, tarımda daha sürdürülebilir ve çevre dostu yöntemlerin benimsenmesine katkıda bulunmayı amaçlamaktadır.

Yabancı Ot Mücadelesinde Biyokontrol Yöntemleri

Wapshere ve ark. (1989), göre biyolojik kontrol ajanlarının uygulanmasında dört yöntem bulunmaktadır. Uygulanan yöntemler klasik veya aşılama, artırma veya takviye, koruma ve geniş spektrumlu yöntemler olarak elle alınmaktadır.

Klasik veya Aşılama Yöntemi: Bu yöntemde, dış kaynaklı bir doğal düşman, hem dışarıdan gelen (DeBach, 1964) hem de yerel zararlılara (Carl, 1982) karşı kullanılır. Bu yöntem, kültür bitkileri ile yakın akraba olmayan yabancı otlarla sınırlıdır ve insan müdahalesinin az olduğu habitatlarda daha etkilidir (Wapshere ve ark., 1989). Bir bitki, doğal alanı dışında yabancı ot haline geldiğinde, bu yabancı otun anavatanında doğal düşmanları aranır, konukçu seçiciliği test edilerek ithal edilir ve zararlı bitkinin bulunduğu alanlara salınır. Bu yöntem, en yaygın ve başarılı metotlardan biridir (Batra, 1981; Julien ve ark., 1984). Klasik biyolojik kontrolde, dağılım ve dışta kalma, ekonomik ve sosyal açıdan oldukça önemlidir (Tisdell ve ark., 1984). Kontrol ajanı, sınırlı sayıda alana ithal edildikten sonra, konukçusunu arar, bulur ve diğer alanlara yayılır. Ekonomik olarak, kimyasal ve mekanik kontrol yöntemlerine göre avantaj sağlar, çünkü insan çabası ve ilave masraf gerektirmez. Bunun yanı sıra, biyolojik kontrol ajanı, ulaşılması zor alanlara da yayılabilir (Zimdahl, 1993; Zengin, 2013).

Artırma veya Takviye Yöntemi: Bu yöntem, yabancı otların istila ettiği alanlarda, biyolojik ajanların popülasyonlarını artırmadan uzun süre varlıklarını sürdürmediği durumlarda kullanılır. Laboratuvarlarda yetiştirilen biyolojik kontrol ajanları, mevcut biyolojik kontrolü başlatmak veya tamamlamak amacıyla salınır (DeBach, 1964; Ridgway ve Vinson, 1976). Bu yöntemde, kontrol ajanı ve hedef yabancı ot genellikle yerlidir. Etkili bir ajanın, hızlı popülasyon artışına izin veren yetiştirme metoduna sahip olması gerekir. Bu yöntemde, genellikle patojenler ve nematodlar kullanılır. Evrimin bir sonucu ve doğal dengenin yansıması olan takviye tekniğinin bilinçli kullanımı, nispeten yenidir (Zimdahl, 1993; Zengin, 2013).

Koruma Yöntemi: Bu yöntemde, yerli yabancı otların kontrol edilmesi, yerel parazit, predatör ve hastalıkların sayısının korunması veya artırılması ile sağlanır. Bu yaklaşıma göre, biyolojik kontrol potansiyeline sahip bir organizma üzerindeki baskıyı azaltarak, bu ajanın daha etkili çalışması sağlanabilir (Zimdahl, 1993; Zengin, 2013).

Geniş Spektrum Yöntemi: Bu yöntem, belirli yabancı otları kontrol etmek için doğal düşman popülasyonlarının suni olarak kullanılmasıdır. Örneğin, belirli otları seçici olarak otlayan hayvanların kullanılması bu kategoriye girer (Zimdahl, 1993; Zengin, 2013).

Yabancı Otların Biyokontrolünde Patojenlerin Rolü

Patojenler, bitki hastalıklarına neden olan zararlı mikroorganizmalar olarak tanımlanır. Patojenler, yabancı ot kontrolünde kullanılan biyolojik etmenler arasında önemli bir yere sahiptir. Yabancı ot patojenleri, yabancı otların büyümesini ve gelişmesini engelleyerek onların baskılanmasına yardımcı olur. Bu patojenler, mantarlar, bakteriler, virüsler ve nematodlar gibi çeşitli mikroorganizmalar olabilir. Patojenler, yabancı otlarda hastalık oluşturarak bitkilerin zayıflamasına ve ölmelerine neden olmaktadır. Örneğin, bazı mantar türleri yabancı otların köklerinde veya yapraklarında enfeksiyon yaparak onların büyümesini engellemektedirler. Yabancı otların büyüme hormonlarını etkileyerek onların gelişimini durdurabilirler. Bu, otların rekabet gücünü azaltmakta ve tarım bitkilerine daha fazla alan ve kaynak sağlamaktadır. Ayrıca yabancı ot popülasyonunu doğal yollarla kontrol ederek onların yayılmasını engellemektedir. Bu da, kimyasal herbisit kullanımını azaltarak çevre dostu bir kontrol yöntemi sunmaktadır.

Patojenler, yabancı otların biyolojik kontrolünde etkili biyokontrol ajanlarıdır. Bu patojenler, yabancı otların büyümesini ve çoğalmasını engelleyen zararlı küfler, bakteriler ve virüsler içerir. Örneğin, *Trichoderma* gibi bitki patojeni funguslar, yabancı otların büyümesini engelleyerek tarım alanlarında kullanılmaktadır (Duraçe ve ark., 2024). Bunun yanı sıra, *Puccinia chondrillina* gibi pas mantarları da yabancı otların yapraklarını enfekte ederek

fotosentezi engeller ve bitkiyi zayıflatır (Hasan ve Ayres, 1990). Yabancı otların biyokontrolü, bitkiyle beslenen veya patojen olan doğal düşmanların kullanımını içerir. Amaç, yabancı otları tamamen yok etmek değil, popülasyon ekolojisi prensiplerine dayanarak yabancı ot yoğunluğunu tolere edilebilir seviyelerde tutmaktır. Pratikte biyokontrol, faydalı bitki türlerine zarar vermeden, üzerinde bulunduğu konukçu bitkilerine doğrudan veya dolaylı olarak zarar veren ya da zayıflatan doğal düşmanların desteklenmesi ve korunmasını sağlar (DeBach, 1964; Zengin, 2013). Yabancı otların biyokontrolü, yabancı ot popülasyonlarını azaltmak veya gelişimlerini sınırlamak için doğal düşmanların kullanılmasıdır (Watson, 1993; Eken ve Demirci 2002). Bu amaçla kullanılan bitki patojenlerine Bioherbisit (biyolojik herbisit) denir (Charudattan ve Dinooor, 2000 Eken ve Demirci 2002).

Yabancı otların gelişimini engelleyen veya bunları öldüren fungal bitki patojenli formülasyonlara mikoherbisit denir (Templeton ve ark., 1979). İlk defa 1971 yılında Avustralya'da *Puccinia chondrillina* adlı bir fungusun, akhindiba (*Chondrilla juncea*) bitkisinin kontrolü için denenmesiyle bu alanda adım atılmıştır (Julien ve Griffiths, 1998; Barton, 2004). Mikoherbisitlerle yabancı otların kontrolü, bitki patojeni bir fungusun bir yabancı ot popülasyonuna uygulanarak, o yabancı ot popülasyonunu ekonomik zarar seviyesi altına düşürmesi veya bu seviyeye yakın tutması şeklinde düşünülmektedir (Greaves, 1991). Yerli ve yabancı orijinli funguslar, mikoherbisit olarak kullanılma potansiyeline sahiptirler ve bunlar, hem yerli, hem de yabancı orijinli yabancı otların kontrolünde kullanılabilirler. Mikoherbisitler, yıllık ürünlerdeki yabancı otları kontrol etmede, çok büyük potansiyele sahiptirler. Mikoherbisitler, meyve bahçelerindeki çok yıllık yabancı otlara (Ridings ve ark., 1976) ve aynı zamanda, herbisitlerin kullanılmadığı su ve meralardaki yabancı otlara karşı da kullanılma potansiyeline de sahiptirler (Williams, 1979).

Doğada hem yararlı hem de zararlı olabilen çeşitli bakteriler bulunur. Yabancı otlarla biyolojik mücadelede fitopatogen bakteriler kullanılır. Bu bakteriler, bitkinin toprakla temas eden kısımlarında ürettikleri fitotoksik maddelerle bitki büyümesini engeller (Suslow ve Schroth, 1982; Tranel ve ark., 1993; Çolak ve ark., 2019). Böcekler ise, yabancı otların çeşitli fizyolojik ve biyolojik süreçlerini hedef alarak onların büyümesini ve yayılmasını engeller. Ayrıca istilacı yabancı türlerin kontrolü için biyoherbivorlar, doğal düşmanları olarak tanımlanmıştır. Bu bitkiler, istilacı türlerin büyümesini ve yayılmasını engelleyerek ekosistem dengesini korumaya yardımcı olur. Türkiye'de yapılan çalışmalar, denizlerimizdeki istilacı türlerin tehdidini azaltmak için biyoherbivor bitkilerin kullanımını önerilmektedir (Blossey, 1999). Sonuç olarak, patojenler yabancı ot baskılamasında önemli bir rol oynar ve sürdürülebilir tarım uygulamalarının bir parçası olarak değerlendirilebilir. Bu yöntem, çevresel koruma ve tarım verimliliği açısından önemli avantajlar sunmaktadır.

Son yıllarda, mantar patojenlerinin kullanımı hakkında önemli gelişmeler yaşanmıştır. Bilim insanları, mantar enfeksiyonlarının antibiyotik direnci gibi ciddi sağlık sorunlarına yol açabileceğini belirtmişlerdir. Dünya Sağlık Örgütü (WHO), mantar patojenlerini sistematik olarak önceliklendirmeye yönelik ilk küresel çabayı başlatarak, insan sağlığı için en tehlikeli patojenler arasında *Aspergillus fumigatus*, *Candida*, *Nakaseomyces glabratus* ve *Trichophyton indotineae* gibi mantar türlerini listelemişlerdir (WHO, 2022).

Yabancı Otların Biyokontrolünde Doğal Yırtıcıların Rolü

Doğal yırtıcılar, yabancı otların biyolojik kontrolünde önemli bir rol oynar. Bu yırtıcılar, yabancı otları doğrudan hasar vererek veya patojenlerin yayılmasını engelleyerek bu otların baskısını azaltır. Örneğin, belirli böcekler ve arılar, yabancı otların çiçeklerini hasar vererek onların çoğalmasını engeller (Şentürk ve Günayar, 2019). Ayrıca, bazı predatör böcekler ve nematodlar, yabancı ot tohumlarını ve fidelerini hedef alarak popülasyonlarını kontrol eder (Blossey, 1999). Biyokontrol araştırmalarından biride büyükbaş ve küçükbaş hayvanların yabancı ot kontrolünde kullanılmasıdır. Ülkemizde mera alanlarında büyük sorun teşkil eden

yabancı ot türlerinden biri de "köygöçüren" olarak bilinen *Cirsium* spp. bitkisidir (Webb ve ark., 1988; Ender ve ark., 2020). Ülkemizde yaygın olarak yetiştirilen keçiler, bu yabancı otların tüm fenolojik dönemlerinde beslenmektedirler (Clark ve ark., 1982; Crouchley, 1983; Batten, 1984; Lamming, 2001; Ender ve ark., 2020).

Araştırmalarda çiftlik hayvanlarının kullanılması hem hayvanların beslenmesine katkı sağlar hem de istenmeyen bitkilerin vejetasyonunu azaltmaktadır. Bir yıllık buğdaygil ve otsu dikotiledonlarla mücadelede koyunlar etkili bir araçtır. Sığırlar, bazı odunsu bitkilerin yeni sürgünlerini otlayarak bu bitkilerin kontrolünü sağlamaktadır. Keçiler ise çalimsı ve odunsu türlerle mücadelede en başarılı biyolojik ajanlardır (Koç ve Gökkuş, 1993). Yeni Zelanda'da, meralardaki istenmeyen çalılar *Discaria toumatou* ve *Hieracium pilosella*, keçi ve koyun otlatması ile etkili bir şekilde kontrol edilmiştir (Cossens ve ark., 1989). Koyunlar ayrıca *Geranium*, *Taraxacum* ve *Senecio* gibi bitki türleriyle mücadelede başarılı olup yüksek rakımlardaki *Delphinium* türlerinin kontrolünde koyun otlatılarak sağlanmaktadır (Vallentine, 1989). Kaz, ördek ve tavuklar çilek, ahududu ve bazı sebzelerde dar yapraklı ve küçük geniş yapraklı yabancı otları kontrol edebilmektedirler. Tavuk ve kazlar, çeşitli ürünlerdeki *Cyperus* spp.'yi seçici olarak kontrol ederken, ördekler küçük havuzlardaki *Lemna* spp.'yi etkili bir şekilde temizler. Kazlar ise geniş yapraklı ürünlerdeki saz ve graminelerin seçici kontrolünde kullanılmaktadır (Conley ve Peterson, 1957).

Yabancı Otların Biyokontrolünde Allelopatinin Rolü

Allelopati, bir bitkinin kimyasal yolla çeşitli bileşikler salarak diğer bitkileri olumlu veya olumsuz etkilemesi olarak tanımlanabilir (Rice, 1984). Bu zararlı etkiden yararlanarak biyolojik yabancı ot kontrolü sağlanabilir (Narwal, 1994; Kohli ve ark., 1998). Çeşitli araştırmalar, allelopatik bitkilerin patojenleri azaltma ve yabancı ot oluşumunu etkileme potansiyeline dikkat çekmiştir. Xuan ve ark. (2005) tarafından Güneydoğu Asya ve Japonya ekosistemlerinde yapılan bir çalışmada, yüzlerce allelopatik bitki incelenmiş ve 30'dan fazla bitki türünün hektara 1-2 ton uygulanmasının yabancı ot biyokütlesini yaklaşık %70 oranında azalttığı, pirinç verimini ise %20 artırdığı görülmüştür (Ender ve ark., 2020). Yabancı otların biyokontrolü alanında çalışan bilim insanlarının temel amacı, çevreye daha az zarar veren ve doğrudan hedef odaklı uygulamaları hayata geçirmektir (Strong ve Pemberton, 2001; Louda ve ark., 2003; Balciunas, 2004). Allelopatik bitkiler, bitki kimyası ile çevresel bitkiler üzerindeki etkileriyle bilinir. Bu bitkiler, yabancı ot yönetimi için oldukça etkili araçlardır. Allelopatik bitkiler, yabancı otların büyümesini engelleyerek doğal bitki topluluklarını korumaya yardımcı olur (Teasdale, 1996).

İstilacı yabancı türlerin kontrolü için biyoherbivorlar, doğal düşmanları olarak tanımlanmıştır. Bu bitkiler, istilacı türlerin büyümesini ve yayılmasını engelleyerek ekosistem dengesini korumaya yardımcı olur. Türkiye'de yapılan çalışmalar, denizlerimizdeki istilacı türlerin tehdidini azaltmak için biyoherbivor bitkilerin kullanımını önerebilir (Blossey, 1999).

Rekabet eden bitki türlerinin kullanımı, yabancı otların baskılanmasında etkili bir yöntemdir. Bu bitki türleri, yabancı otların yerleşimini engelleyerek onların yayılmasını önler. Örneğin, bazı bitki türleri, yabancı otların büyümesi için gerekli olan besin maddelerini emer veya yabancı otlara karşı direnç gösterir (Bulut ve Dinler, 2021). Örtücü bitkileri olarak bilinen bu bitkiler, toprak yüzeyini kaplayarak yabancı otların ışık almasını engeller ve onların çimlenmesini azaltır (Teasdale, 1996).

Biyokontrol Yöntemlerinin Benimsenmesinin Sosyo-Ekonomik Sonuçları

Biyokontrol, tarımda yabancı ot, böcek ve diğer zararlıları yönetmek için doğal düşmanların kullanılmasıdır. Bu yöntem, sentetik kimyasallara dayalı geleneksel kontrol yöntemlerine alternatif olarak sunulur ve sürdürülebilir tarım için önemli olmaktadır. Biyokontrolün

ekonomik ve sosyal etkilerini anlamak, tarım politikalarının şekillendirilmesi ve çiftçilerin benimsemesi açısından kritik öneme sahiptir.

Ekonomik Etkiler: Biyokontrol, çiftçiler için maliyet etkinliği sağlar. Sentetik herbisitlerin ve pestisitlerin sürekli kullanımı yüksek maliyetlere yol açarken, biyokontrol yöntemleri daha düşük maliyetli ve uzun vadede daha sürdürülebilirdir. Örneğin, DeBach (1964) biyokontrol yöntemlerinin, kimyasal kontrol yöntemlerine göre daha düşük maliyetli olduğunu ve ekonomik getiri sağladığını belirtmiştir. Ayrıca, biyokontrol yöntemlerinin kullanımı, sentetik kimyasallara olan bağımlılığı azaltarak dışa bağımlılığı da azaltmaktadır.

İstihdam ve Gelir: Biyokontrol uygulamaları, yeni istihdam olanakları yaratmaktadır. Bu yöntemlerin geliştirilmesi, uygulanması ve izlenmesi için teknik bilgi ve uzmanlık gerekmektedir. Bhadoria (2011) çalışmasında, biyokontrol projelerinin tarım işçileri ve araştırmacılar için yeni istihdam fırsatları sunduğunu ve yerel ekonomilere katkı sağladığını vurgulamıştır. Ayrıca, biyokontrol uygulamalarının benimsenmesi, çiftçilerin gelir düzeyini artırmakta ve kırsal kalkınmayı desteklemektedir.

Sosyal Kabul ve Eğitim: Çiftçilerin ve toplulukların biyokontrol yöntemlerine adaptasyonu, eğitim ve farkındalık yaratma faaliyetleri ile desteklenmelidir. Topal (2011b), biyokontrol yöntemlerinin benimsenmesinin, çiftçilerin geleneksel bilgilere olan güvenini artırdığını ve çevresel farkındalığı güçlendirdiğini belirtmiştir. Eğitim programları ve saha çalışmaları, biyokontrolün etkinliğini artırmak için önemlidir.

Çevresel Etkiler: Biyokontrol, ekosistem hizmetlerine önemli katkılar sağlar. Doğal düşmanların kullanımı, biyolojik çeşitliliği artırır ve toprak verimliliğini iyileştirir. Ayrıca, biyokontrol yöntemleri, kimyasal kirliliği azaltarak çevresel sürdürülebilirliği desteklemektedir. Bu sayede, tarımsal faaliyetlerin çevresel etkileri minimize edilmekte ve ekosistem dengesi korunmaktadır.

Yapılan Çalışmalar: Farklı bölgelerde biyokontrol benimsenmesinin sosyo-ekonomik sonuçlarını inceleyen araştırmalar ve başarı hikayeleri, bu yöntemlerin etkinliğini ve uygulanabilirliğini göstermektedir. Örneğin, Bhadoria (2011) çalışmasında, biyokontrol projelerinin başarılı olduğu bölgelerde ekonomik ve sosyal faydaların önemli ölçüde arttığını rapor etmiştir.

Biyokontrol yöntemlerinin tarımda daha geniş çapta benimsenmesi, hem ekonomik hem de sosyal açıdan büyük faydalar sağlamaktadır. Bu yöntemlerin kullanımı, tarımsal sürdürülebilirlik ve kırsal kalkınma açısından umut verici bir strateji olarak öne çıkmaktadır. Çiftçiler, biyokontrol yöntemlerinin avantajlarını dikkate alarak bu uygulamaları teşvik etmeli ve desteklenmelidir.

SONUÇ

Biyokontrol, sürdürülebilir tarımda hayati bir role sahiptir. Bu yöntem, zararlı organizmaların biyolojik etmenler aracılığıyla kontrol edilmesi sürecini kapsamaktadır. Zararlılarla mücadelede kimyasal pestisitlerin kullanılmasını en aza indirerek, çevreye ve insan sağlığına olan olumsuz etkileri azaltmaktadır. Biyokontrolün sürdürülebilir tarıma olan katkıları; hedef zararlıları etkili bir şekilde kontrol ederken diğer organizmalara zarar vermez. Böylece ekosistemin dengesini korunmasını sağlar. Toprak yapısını koruyarak mikroorganizma çeşitliliğini korumaktadır. Daha sağlıklı ve güvenli ürünler elde edilir. Ayrıca uzun vadede daha etkili olma potansiyeli vardır. Bu nedenlerle, biyokontrol yöntemleri tarımda sürdürülebilirliğin sağlanması için önemli bir araçtır. Biyokontrol yöntemlerinin gelecekte daha etkili ve yaygın bir şekilde kullanılmasını sağlamak için önemli bir adım olarak değerlendirilmektedir. Biyokontrol yöntemlerinin sosyo-ekonomik etkilerini ele alarak, tarımsal sürdürülebilirlik ve

kırsal kalkınma açısından sunduğu fırsatlar açısından değerli olmaktadır. Biyokontrol araştırmalarının devam etmesi, sürdürülebilir tarımın geleceği için kritik öneme sahiptir. Çiftçilerin, araştırmacıların ve bitki koruma uzmanları bu yöntemleri desteklemesi, tarımın geleceği için önemlidir. Şu anda biyolojik kontrol uygulamalarını kolaylaştırmaya yönelik çalışmalar sınırlı sayıda ve bu teknolojik ilerlemeleri kullanarak artırılması gerekmektedir.

KAYNAKLAR

- Altieri, M. A. (1999). "The ecological role of weeds in agroecosystems." In *Weed Management in Agroecosystems*. CRC Press.
- Altieri, M. Nicholls, C. (2000). *Agroecología: teoría práctica para una agricultura sustentable* [en línea]. (ser. Textos básicos para la formación ambiental), vol. 4, Ed. Programa de las Naciones Unidas para el Medio Ambiente - Red de Formación Ambiental para América Latina el Caribe, 250 p., ISBN 978-968- 7913-07-0
- Balciunas, J. K. (2004). Are mono-specific agents necessarily safe? The need for pre-release assessment of probable impact of candidate biocontrol agents, with some examples. In XI International Symposium on Biological Control of Weeds, pp. 252-257**
- Barton, J. (2004). How good are we at predicting the field host-range of fungal pathogens used for classical biological control of weeds?. *Biological Control* 31(1): 99-122**
- Batra, S. W. T. (1981). Biological control of weeds: principles and prospects. Beltsville Symposium on Agricultural Research No. 5, Biological control in crop production. (ed. Papavizas, G.C.), Granada London, 45-59.
- Bhadoria, P. B. S. (2011). Allelopathy: A natural way towards weed management. *American Journal of Experimental Agriculture*, 1(1), 7-20.
- Blossey, B. (1999). Before, During and After: The Need for Long-term Monitoring in Invasive Plant Species Management. *Biological Invasions*, 1(4), 301-311.
- Boström, U. y Fogelfors, H. (2002). Long-term effects of herbicide application strategies on weeds and yield in spring-sown cereals. *Weed Science*, vol. 50, no. 2, marzo de pp. 196-203, ISSN 0043-1745, 1550-2759, DOI 10.1614/0043-1745
- Bulut, D., & Dinler, H. (2021). Biyolojik Mücadelede Trichoderma'lar ve Biyolojik Kontrol Mekanizmaları. *Uşak Üniversitesi Fen ve Doğa Bilimleri Dergisi*.
- Canizales, S. A.; Celemín, J. S. Mora-Delgado, J. (2010). Diversidad uso de arvenses en pasturas defincas ganaderas del Alto Magdalena (Tolima, Colombia). *Zootecnia Tropical*, vol. 28, no. 3, pp. 427-437, ISSN 0798-7269.
- Carl, K. P. (1982). Biological control of native pests by introduced natural enemies. *Biological Control News and Information* 3: 190-200.
- Charudattan R. (2001). Biological control of weeds by means of plant pathogens: significance for integrated weed management in modern agro-ecology. *BioControl* 46:229–60
- Charudattan, R., Dinooor, A. (2000). Biological control of weeds using plant pathogens: accomplishments and limitations. *Crop Protection*, 19: 691-695.
- Çolak, E. Ş., Yüksel, E., & Canhilal, R. (2019). Yabancı otların kontrolünde biyolojik mücadele. *Erciyes Tarım ve Hayvan Bilimleri Dergisi*, 2(3), 23-29.
- DeBach, P. (1964). Successes, trends and future possibilities. In *Biological Control of Insect Pests and Weeds* (Eds. P. DeBach and E.I. Schlinger), 673-713.
- Duraçe, E., Dikmetas, D. N., & Karbancıoğlu-Güler, F. (2024). Biyokontrol Yaklaşımı ile Küflerin Kontrolü. *ITU Journal of Food Science and Technology*.
- Eken, C., & Demirci, E. (2002). Colletotrichum türleri ile yabancı otların biyolojik kontrolü/biological control of weeds with Colletotrichum species. *Atatürk Üniversitesi Ziraat Fakültesi Dergisi*, 34(2).
- Gnanavel, I. (2015). Sürdürülebilir Tarım İçin Çevre Dostu Yabancı Ot Kontrol Seçenekleri. *Sci. Int.* 3, 37–47.

- Greaves, M. P. (1991). Integration of Biological Control Agents With Chemical Pesticides. In: Te Beest, D.O., Microbial Control of Weeds, Chapman and Hall, London, 189-208.
- Hasan, S., & Ayres, P. G. (1990). The Control of Plant Diseases by Fungal Antagonists. *Biological Reviews*, 65(1), 1-25.
- Julien, M. H., and Griffiths, M. W. (1998). Biological control of weeds: a world catalogue of agents and their target weeds. ed. 4. Cab International**
- Julien, M. H., J. Kerr, R. R. Chan, (1984). Biological control of weeds: an evaluation. *Protection Ecology*, 7: 3-25.
- Kitiş, Y.E. (2009). Çukurova Bölgesi Turunçgil Bahçelerinde Canlı ve Cansız Malç Uygulamalarının Entegre Yabancı Ot Kontrolü Açısından Değerlendirilmesi Çukurova Üniversitesi Fen Bilimleri Enstitüsü Bitki Koruma Anabilim Dalı.
- Kohli, R. K., Batish, D., and Singh, H. P. (1997). Allelopathy and its implications in agroecosystems. *Journal of Crop Production*, 1(1): 169-202
- Louda, S. M., Arnett, A. E., Rand, T. A., & Russell, F. L. (2003). Invasiveness of some biological control insects and adequacy of their ecological risk assessment and regulation. Conservation Biology 17(1): 73-82**
- MacLaren C, Storkey J, Menegat A, Metcalfe H, Dehnen-Schmutz K. (2020). An ecological future for weed science to sustain crop production and the environment. A review. *Agronomy for Sustainable Development*. 2020;40:1-29. Available: <https://doi.org/10.1007/s13593-020-00631-6>
- MacLaren, C.; Storkey, J.; Menegat, A. (2020). Bitkisel üretimi ve çevreyi sürdürmek için yabancı ot biliminin ekolojik geleceği. inceleme. *Agron. Sustain. Dev.* 40, 24.
- Marshall, E. J. P., & Moonen, A. C. (2002). "Field margins in northern Europe: their functions and interactions with agriculture." *Agriculture, Ecosystems & Environment*.
- McFadyen REC. (1998). Biological control of weeds. *Annu. Rev. Entomol.* 43:369–93112.
- Paynter Q, Overton JM, Hill RL, Bellgard SE, Dawson MI. 2012. Plant traits predict the success of weed biocontrol. *J. Appl. Ecol.* 49:1140–48
- Narwal, S.S. (1994). Allelopathy in Crop Production. Scientific Publisher, Jodhpur, India pp. 288**
- Peerzada AM, Bukhari SAH, Dawood M, Nawaz A, Ahmad S, Adkins S. (2019). Weed management for healthy crop production. *Agronomic Crops: Volume 2: Management Practices*. 225-256. Available: https://doi.org/10.1007/978-981-32-9783-8_13
- Rice, E.L., (1984). Allelopathy, Second ed. Academic Press Inc., Orlando, FL, pp. 422**
- Ridgway, R. L., S. B. Vinson, (1976). Biological Control by Augmentation of Natural Enemies: Insect and Mite Control with Parasites and Predators. Plenum Press, New York
- Ridings, W. H., D. J. Mitchell, C. L. Shoulties, N. E. El-Ghell, (1976). Biological control of milkweed vine in Florida citrus groves with a pathotype of *Phytophthora citrophthora*. In Proc. IV Int. Symp. Biol. Control of Weeds, ed. T.E. Freeman, 224-240.
- Schwarzländer M, Hinz HL, Winston RL, Day MD. (2018). Biological control of weeds: an analysis of introductions, rates of establishment and estimates of success, worldwide. *Biocontrol* 63:319–31
- Şentürk, Ş., & Günayar, Ö. A. (2019). Fungal Biyokontrol Ajanları ve Metabolitleri. *Mantar Dergisi*
- Sheley, R. L., & Petroff, J. K. (1999). "Biology and Management of Noxious Rangeland Weeds." Oregon State University Press.
- Strong, D.R., and Pemberton, R.W. (2001). Food webs, risks of alien enemies and reform of biological control. In: Wajnberg, E., Scott, J.K., Quimby, P.C. (Eds.), Evaluating Indirect Ecological Effects Of Biological Control. CABI Publishing, Wallingford, UK, pp. 57-79**
- Suslow, T. V., and Schroth, M. N. (1982). Role of deleterious rhizobacteria as minor pathogens in reducing crop growth. *Phytopathology* 72(1): 111-115

- Teasdale, J. R. (1996). Contribution of Cover Crops to Weed Management in Sustainable Agricultural Systems. *Journal of Production Agriculture*, 9(4), 475-479.
- Templeton, G. E., D. O. Te Beest, R. J. Jr. Smith, (1979). Biological weed control with mycoherbicides. *Ann. Rev. Phytopathol.* 17, 301-310.
- Tisdell, C. A., B. A. Auld, K. M. Menz, (1984). On assessing the value of biological control of weeds. *Prot. Ecol.*, 6, 169-179
- Tomita, S.; Nawata, E.; Kono, Y.; Nagata, Y.; Noichana, C.; Sributta, A. Inamura, T. (2003). "Differences in weed vegetation in response to cultivating methods and water conditions in rainfed paddy fields in north-east Thailand". *Weed Biology and Management*, vol. 3, no. 2, 1 de junio de 2003, pp. 117-127, ISSN 1445-6664, DOI 10.1046/j.1445-6664.2003.00093.x.
- Topal, S. (2011a). Allelokimyasalların herbisit etkileri, Dumlupınar Üniversitesi, Fen Bilimleri Enstitüsü Dergisi Sayı: 25 ıss1302-3055.
- Topal, Ş. (2011b). Biyolojik mücadele ve tarımsal ekosistemler. *Tarım ve Mühendislik*, 111(1), 29-32.
- Tranel, P.J., Gealy, D.R. and Kennedy, A.C. (1993). Inhibition of downy brome (*Bromus tectorum*) root growth by a phytotoxin from *Pseudomonas fluorescens* strain D7. *Weed Technol.*, 7: 134-139.
- Vilà M, Espinar JL, Hejda M, Hulme PE, Jarosik V. (2011). Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* 14:702–8
- Wapshere, A. J., E. S. Delfosse, J. M. Cullen, (1989). Recent developments in biological control of weeds. *Crop Prot.* 8: 227-250.
- Watson, A.K. (1993). Current Status of Bioherbicide Development and Prospects for Rice in Asia. Food and Fert. Tech. Centre, Taipei, Taiwan, Ext. Bull No: 365, 7 pp.
- Williams, P. H. (1979). Vegetable Crop Protection in the People's Republic of China. *Ann. Rev. Phytopathol.*, 17: 311-324.
- Winston RL, Schwarzländer M, Hinz HL, Day MD, Cock MJW, Julien MH, eds. (2014). *Biological Control of Weeds: A World Catalogue of Agents and their Target Weeds*. Morgantown, WV: USDA For. Serv. For. Health Technol. Enterp. Team. 5th ed.
- World Health Organization (WHO). (2022). Global Priority List of Fungal Pathogens.
- Xuan, T. D., Shinkichi, T., Khanh, T. D., and Chung, I. M. (2005). Biological control of weeds and plant pathogens in paddy rice by exploiting plant allelopathy: an overview. Crop protection, 24(3), 197-206**
- Zengin, H. (2013). Yabancı Otlarla Biyolojik Mücadele Yöntemleri. *Atatürk Üniversitesi Ziraat Fakültesi Dergisi*, 28(3).
- Zimdahl, R. L. (1993). *Fundamentals of Weed Science*. Academic Pres, Inc., New York, 191-205.

INTEGRATION OF COVER CROPS IN WEED SUPPRESSION AND SOIL FERTILITY

ÖRTÜCÜ BİTKİLERİNİN YABANCI OT BASKILAMA VE TOPRAK VERİMLİLİĞİNE ETKİLERİ

Ayfer GÜNEY SARITAŞ

Iğdır University, Postgraduate Education Institute, Department of Agricultural Sciences, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0001-7641-9711>

Ramazan GÜRBÜZ

Iğdır University, Faculty of Agriculture, Department of Plant Protection Iğdır, Türkiye²

ORCID ID: <https://orcid.org/0000-0003-3558-9823>

ABSTRACT

To control weeds and improve the sustainability of agricultural ecosystems, the integration of cover crops into agricultural systems can improve soil fertility and suppress weed growth, while making a significant contribution to soil productivity. Therefore, the use of cover crops is in great demand. Cover crops such as legumes, cereals and crucifers are increasingly being integrated into agricultural production areas to maintain soil structure and reduce dependency on chemical inputs. This review addresses the mechanisms by which cover crops contribute to weed suppression. However, allelopathy, light and nutrient competition and mulch systems that inhibit weed seed germination are also included. In addition, the role of cover crops in improving soil fertility is discussed, focusing on nitrogen fixation, organic matter enrichment and microbial activity. Findings from recent studies are synthesized to highlight best practices for cover crop selection, planting and mowing or soil incorporation or tilling times, and integration into various cropping systems. In particular, the ecological benefits associated with the use of cover crops, including the potential for competition with cultivated crops and management difficulties, were emphasized. In addition, the need for guiding studies to investigate varieties suitable for the climate and soil characteristics of a region was emphasized in this study. We believe that the topics discussed here will shed light on farmers, all stakeholders engaged in plant production and administrators making legal regulations.

Keywords: Cover crops, weed suppression, soil fertility, sustainable agriculture, nitrogen fixation

ÖZET

Yabancı otları kontrol etmek ve tarımsal ekosistemlerin sürdürülebilirliğini iyileştirmek için, örtücü bitkilerinin tarımsal sistemlere entegrasyonu, toprak verimliliğini iyileştirme ve yabancı otların gelişmelerini baskıladığı gibi aynı zamanda toprak verimliliğine de ciddi katkılar sunabilmektedir. Bundan dolayı örtücü bitkilerin kullanımına önemli rağbetler bulunmaktadır. Baklagiller, tahıllar ve turpgiller gibi örtücü bitkilerin, toprak yapısının korunması ve kimyasal

girdilere olan bağımlılığın azaltılması için giderek tarımsal üretim alanlarına entegre edilmektedir. Bu derlemede örtücü bitkilerinin yabancı ot baskılamasına katkıda bulunduğu mekanizmaları ele alınmıştır. Bununla birlikte allelopati, ışık ve besin rekabeti ve yabancı ot tohumu çimlenmesini engelleyen malç sistemleri de dahil edilmiştir. Ayrıca, örtücü bitkilerinin toprak verimliliğini iyileştirmedeki rolü, azot fiksasyonu, organik madde zenginleştirme ve mikrobiyal aktiviteye odaklanarak konular ele alınmıştır. Örtücü bitkilerin seçimi, ekimi ve biçme veya toprağa karıştırma yada yere yatırma zamanları ve çeşitli ekim sistemlerine entegrasyon için en iyi uygulamaları vurgulamak üzere son çalışmalardan elde edilen bulguları sentezlenmiştir. Özellikle, yetiştirilen kültür bitkileri ile rekabet potansiyeli ve yönetim zorlukları da dahil olmak üzere, örtücü bitkilerin kullanımıyla ilişkili ekolojik faydaların üzerinde durulmuştur. Ayrıca, bu çalışmada bir bölgenin ilkim ve toprak özelliklerine uygun çeşitlerin araştırılmasına yönelik yol gösterici çalışmaların yapılmasına olan ihtiyaçlara vurgu yapılmıştır. Burada ele alınan konular çiftçilere, bitkisel üretim yapan bütün paydaşlara ve yasal düzenlemeler yapan yöneticilere ışık tutacağı kanaatindeyiz.

Anahtar Kelimeler: örtücü bitkiler, yabancı ot baskılama, toprak verimliliği, sürdürülebilir tarım, azot fiksasyonu

GİRİŞ

Sürdürülebilir tarım, gıda üretimini, çevre yönetimini ve ekonomik sürdürülebilirliği dengelemeyi hedeflemektedir. Bu bağlamda, örtücü bitkileri, tarımsal ekosistemlerin dayanıklılığını ve verimliliğini artırmada önemli bir araç olarak kabul edilmektedir. Genellikle ürünlerin yetişmediği dönemlerde ekilen örtücü bitkileri, satış amacıyla değil, toprak sağlığını iyileştirmek, erozyonu azaltmak ve biyolojik çeşitliliği artırmak gibi temel ekolojik işlevleri yerine getirmek için kullanılmaktadır. Örtücü bitkilerini kullanarak, çiftçiler kimyasal girdilere olan bağımlılıklarını azaltmakta ve tarımsal faaliyetlerinin uzun vadeli sürdürülebilirliğini artırmaktadır (Altieri, 1995).

Örtücü bitkilerinin kullanımı, toprak bozulması, iklim değişikliği ve kaynakların tükenmesi gibi küresel sorunlara yanıt olarak artmaktadır. Bu bitkiler, her biri kendine özgü faydalar sunan baklagiller, çimenler ve turpgiller gibi çeşitli türleri içermektedir. Örneğin, baklagiller atmosferik nitrojeni bağlayarak toprak verimliliğini artırırken, çimenler erozyonu azaltmada ve toprak yapısını iyileştirmede etkilidir. Turpgiller ise bazı toprak kökenli patojenleri ve zararlıları baskılamaktadır (Drinkwater ve ark., 1998). Bu yararlar, dış girdileri en aza indirerek ve ekosistem hizmetlerini geliştirerek sürdürülebilir tarım ilkelerine katkıda bulunmaktadır.

Yabancı Otların Kontrolü ve Toprak Verimliliğinin Önemi

Sürdürülebilir tarımda, örtücü bitkilerinin iki önemli rolü yabancı ot kontrolü ve toprak verimliliğinin artırılmasıdır. Özellikle sentetik herbisit kullanımını azaltmayı hedefleyen sistemlerde, yabancı ot yönetimi çiftçiler için büyük bir zorluk olmaya devam etmektedir. Örtücü bitkileri, allelopati, ışık ve besin rekabeti ile yabancı ot tohumu çimlenmesini engelleyen fiziksel malç bariyerlerinin oluşturulması gibi mekanizmalarla yabancı ot kontrolüne katkı sağlamaktadır. Örneğin, çavdar (*Secale cereale*), yabancı ot çıkışını engelleyen ve yabancı ot büyümesini kısıtlayan allelopatik bileşikler salgılayarak yoğun bir malç tabakası oluşturmaktadır (Teasdale ve Mohler, 2000). Buna ek olarak, örtücü bitkileri, toprağın fiziksel, kimyasal ve biyolojik özelliklerini geliştirerek toprak verimliliğini artırmaktadır. Yonca (*Trifolium spp.*) ve fiğ (*Vicia spp.*) gibi baklagil örtücü bitkileri, atmosferdeki azotu sabitleyerek sonraki ana mahsuller için kullanılabilir hale getirmektedir. Bu doğal süreç, sentetik gübre ihtiyacını azaltmakta ve çevreye azot kaybını önlemektedir (Snapp ve ark., 2005). Ayrıca, örtücü bitkilerinin kalıntılarının ayrışması toprağa organik madde katarak su tutma kapasitesini, yapısını ve mikrobiyal aktivitesini iyileştirmektedir (Schipanski ve ark., 2014).

Örtücü bitkilerinin ekim sistemlerine entegre edilmesi, sadece verimliliği artırmakla kalmaz, aynı zamanda çevresel hedeflere de hizmet etmektedir. Toprak erozyonunu azaltarak, besin döngüsünü iyileştirerek ve sera gazı emisyonlarını düşürerek, örtücü bitkileri tarımsal alanların sürdürülebilirliği ve dayanıklılığına katkıda bulunmaktadır (Tilman ve ark., 2002). Ancak, bu sistemlerin benimsenmesi dikkatli planlama gerektirir. Bu, uygun türlerin seçimi, ekim ve sonlandırma zamanlaması ile ana mahsullerle rekabetin yönetilmesini içermektedir.

Bu makale, örtücü bitkilerinin yabancı ot kontrolü ve toprak verimliliğini artırmadaki rolünü incelemekte ve çiftçiler ile politika yapımcılar için pratik öneriler sunmak amacıyla mevcut araştırmaları sentezlemektedir. Bu önemli yönleri ele alarak, örtücü bitkileri sürdürülebilir tarımı ilerletmek için umut verici bir strateji olarak öne çıkmaktadır.

Örtücü Bitkilerin Yabancı Otları Baskılama Mekanizmaları

Örtücü bitkiler, tarımsal ekosistemlerde çeşitli ekolojik ve ekonomik faydalar sağlamaktadırlar. Özellikle yabancı ot baskılama konusunda önemli bir rol oynar. Örtücü bitkilerin yabancı ot baskılama mekanizmaları:

Alelopati: Örtücü bitkiler, allelopati adı verilen doğal kimyasal savunma mekanizmalarıyla yabancı otları baskılayabilir. Allelopatik bitkiler, toprakta ve bitki örtücüsünde bulunan kimyasal bileşikler salgılayarak yabancı ot tohumlarının çimlenmesini ve büyümesini engeller. Örneğin, çavdar (*Secale cereale*) örtücü bitkisi, allelopatik bileşikler salarak yabancı ot çıkışı azaltabilir (Teasdale ve Mohler, 2000). Bu kimyasal savunmalar, tarımsal alanlarda yabancı ot popülasyonlarının kontrol altında tutulmasına yardımcı olur.

Işık ve Besinler İçin Rekabet: Örtücü bitkiler, yabancı otlarla ışık ve besinler için rekabet ederek onların büyümesini engeller. Yoğun bir örtücü bitkisi tabakası, güneş ışığını bloke eder ve yabancı otların fotosentez yapmasını zorlaştırır. Aynı zamanda, örtücü bitkiler topraktaki besin maddelerini tüketerek yabancı otların ihtiyaç duyduğu kaynakları azaltır. Bu rekabet mekanizması, tarımsal alanlarda yabancı otların yayılmasını engellemeye yardımcı olur (Drinkwater ve ark., 1998).

Malç Uygulamaları: Örtücü bitkiler, toprağın yüzeyine malç katmanları oluşturarak yabancı ot tohumlarının çimlenmesini fiziksel olarak engeller. Yoğun bir malç tabakası, toprak yüzeyinde yabancı ot tohumlarının ulaşmasını zorlaştırır ve çimlenmelerini engeller. Bu mekanizma, tarımsal alanlarda yabancı ot popülasyonlarını azaltmaya yardımcı olur. Örtücü bitkilerin kalıntılarının toprakta ayrışması, toprak sağlığını iyileştirir ve toprağın su tutma kapasitesini artırır (Schipanski ve ark., 2014).

Örtücü bitkiler, tarımsal ekosistemlerde yabancı ot baskılama mekanizmaları olarak önemli bir rol oynar. Bu bitkilerin kullanımı, toprağın verimliliğini artırarak çiftçilere ekonomik fayda sağlar ve çevresel sürdürülebilirliği destekler. Örtücü bitkilerin seçimi ve uygulaması, tarımsal uygulamalarda sürdürülebilirlik ve ekonomik verimlilik sağlamak için kritik öneme sahiptir.

Örtücü Bitkilerinin Toprak Sağlığı ve Erozyon Kontrolündeki Rolü

Örtücü bitkileri, toprağı yerinde tutarak kabuklanmayı engeller ve rüzgar ile su erozyonuna karşı koruma sağlar. Bu bitkilerin toprak üstünde kalan kısımları, yağmur damlalarının toprağa doğrudan çarpmasını önleyerek erozyon etkilerini azaltır. Erozyon süreçleri üzerindeki etkileri, toprağın kopması ve taşınmasının ne kadar azaltıldığına bağlıdır. Örtücü bitkileri, canlı bitkiler veya bitki kalıntıları ile toprak örtücüsünün miktarını ve süresini artırarak erozyonu azaltır. Aynı zamanda akan suyun kayma kuvvetini azaltarak veya toprak parçacıklarının kopmaya karşı direncini artırarak dere içi ve dere dışı erozyonu önler. Bu bitkiler, yüzey sızdırmazlığını önleyip depolama kapasitesini artırarak ve toprak yapısını iyileştirerek oluşan sızmanın miktarını artırır. Bu süreç, akış suyunun kayma kuvvetini azaltır (Dabney, 1998). Ayrıca, örtücü bitkileri veya yüzeyde kalan kalıntılar akış hızını yavaşlatarak erozyonu daha da azaltabilir

(Brown ve Norton, 1994). Son olarak, örtücü bitkilerinin kökleri, yüzeydeki kalıntıları yerinde tutarak akış halindeki suyun toprağı kolayca hareket ettirmesini ve suyun kesme kuvvetlerinin toprağı zarar vermesini engeller (Parker, 1915). Örtücü bitkiler, erozyonu etkili bir şekilde kontrol eder ve çevrenin korunmasına katkı sağlar.

Organik madde kaynağı olan örtücü bitkiler, toprak biyolojik aktivitesi için substrat sağlayarak toprağın fiziksel özelliklerini etkiler. Çavdar (*Secale cereale* L.), yulaf (*Avena sativa* L.), arpa (*Hordeum vulgare*) gibi bazı kışlık buğdaygil türleri bu amaçla yaygın olarak kullanılır. Örtücü bitkilerinin kökleri, yüzeydeki kalıntıları yerinde tutarak, suyun toprak üzerinde kolayca hareket etmesini ve suyun kesme kuvvetlerine maruz bırakmasını engeller. Ayrıca, bazı bitkiler, toprak profilini fiziksel olarak değiştirebilir ve toprak yapısını iyileştirebilir (McVay ve ark., 1989). Örtücü bitkileri, makro gözenekler (örneğin solucan yuvaları) oluşmasını artırır. Böylece, su infiltrasyonunun artmasına ve toprağın erozyona karşı duyarlılığının azalmasına yol açar. Geniş ve derin kök sistemlerine sahip örtücü bitkileri, toprağın içinde kanallar oluşturarak toprak yapısını ve gözenekliliğini iyileştirir (Williams ve Weil, 2004). Çavdar, yulaf ve yonca gibi örtücü bitkilerinin kökleri, sıkıştırılmış toprak katmanlarına nüfuz edebilir, bu da toprak havalanmasını iyileştirir ve sonraki bitkilerin kök büyümesini kolaylaştırır (Chen ve Weil, 2010). Örtücü bitkisi kökleri ayrıştıkça, toprak parçacıklarını bir arada tutan ve toprak yapısını iyileştiren organik maddeyi biriktirirler (Abdollahi ve ark., 2014).

Örtücü bitkileri, toprak mikroorganizmaları için organik atıklar ve kök salgıları sağlayarak, topraktaki mikrobiyal topluluğı besler ve etkiler (Lehman ve ark., 2015). Bu artan mikrobiyal aktivite, organik maddelerin ayrışmasına ve kalıcı bir organik madde formu olan humusun oluşumuna katkıda bulunur. Ayrıca, örtücü bitkileri, solucanlar gibi toprak faunası için elverişli bir ortam yaratarak, organik maddelerin toprakta parçalanmasına ve dahil edilmesine yardımcı olur (Blouin ve ark., 2013). Örtücü bitkileri, N, P, K, Ca, Mg, S, vb. gibi besinlerinin geri dönüşümünde yardımcı olur. Yüksek konsantrasyonlarda mineralleri biriktirir. Örtücü bitkileri, özellikle azot gibi besin maddelerinin sonraki mahsuller için gereksinimini azaltma potansiyeline sahiptir. Bu bitkilerin kökleri, azotu topraktan çekerek nitratin yeraltı sularına sızmasını ve topraktan aşağı doğru hareket etmesini önler (Gabriel ve ark., 2013). Azot fiksasyonu, bitkilerin büyümesi için temel bir besin elementi olan azotun atmosferik gaz halinden toprağı dönüşmesi sürecidir. Bu süreç, bitkilerin köklerinde yaşayan azot bağlayan bakteriler tarafından gerçekleştirilir. Azot fiksasyonu, bitkilerin ihtiyaç duyduğu amino asit ve proteinlerin üretiminde önemli bir rol oynar (Haktanır ve Arcak 1997; Galloway ve ark., 2004)

Örtücü Bitkilerin Seçimi ve Kontrolü

Örtücü bitkilerinin seçimi, ekosistem sağlığı ve verimliliği için kritik bir rol oynar. Farklı örtücü bitkileri farklı yararlar sağlar ve bu nedenle baklagiller, otlar ve brassicalar gibi çeşitli türler arasından seçim yapmak önemlidir.

Baklagiller (Fabaceae) genellikle yüksek azot fiksasyonu yetenekleri nedeniyle tercih edilirler. Köklerindeki nodüller, atmosferdeki azotu bitkilerin kullanabileceği formda bağlar. Bu, toprak verimliliğini artırır ve sonraki ekinler için gerekli besin maddelerini sağlar. Örneğin, yonca (*Medicago sativa*) ve üçgül (*Trifolium* spp.) yaygın olarak kullanılan baklagil örtücü bitkileridir (Abdollahi ve Munkholm, 2014). Otlar (Poaceae) güçlü kök sistemlerine sahiptir ve bu özellikleriyle toprağı yerinde tutarak erozyonu önlerler. Çavdar (*Secale cereale*) ve yulaf (*Avena sativa*) gibi otlar, ayrıca organik madde sağlama ve toprak yapısını iyileştirme yetenekleriyle bilinirler (Blouin ve ark., 2013). Brassicalar (*Brassicaceae*) grubundan bazı bitkiler, nematod ve patojenlerle mücadelede yardımcı olabilir. Hardal (*Sinapis* spp.) ve kanola (*Brassica napus*) gibi bitkiler, köklerinden salgıladıkları glukozinolatlar sayesinde doğal biyofumigantlar olarak işlev görürler (Brown ve Norton, 1994).

Örtücü Bitkilerinin Ekim ve Sonlandırma Zamanlaması

Örtücü bitkilerinin ekim ve sonlandırma zamanlaması, örtücü bitkisinin amacına ve iklim koşullarına bağlı olarak değişir. Doğru zamanlama, örtücü bitkilerinin maksimum faydayı sağlaması için önemlidir.

Ekim Zamanlaması: Örtücü bitkileri genellikle ana ürünün hasadından hemen sonra ekilir. Bu, toprağın boş kalmaması ve erozyon riskinin azaltılması için önemlidir. Örneğin, kış Örtücü bitkileri yaz sonu veya sonbahar başında ekilir (Chen ve Weil, 2010).

Sonlandırma Zamanlaması: Örtücü bitkileri, ana ürünün ekiminden önce veya belli bir gelişim aşamasına ulaştıktan sonra sonlandırılır. Sonlandırma yöntemleri arasında biçme, otlatma, ve kimyasal veya mekanik yollarla öldürme yer alır. Çeşitli örtücü bitkileri, farklı bitkisel döngülerin ve iklim koşullarının gerektirdiği esnekliklere yanıt verebilirler (Dabney, 1998).

Örtücü Bitkilerinin Çeşitli Ekim Sistemlerine Entegrasyonu

Örtücü bitkilerinin tarımsal sistemlere entegrasyonu, sürdürülebilir tarım uygulamaları açısından hayati öneme sahiptir. Farklı ekim sistemleri, örtücü bitkilerinin sağladığı faydaları maksimum düzeye çıkarabilir.

Monokültür Sistemlerde: Tek bir ana ürünün yetiştirildiği sistemlerde, örtücü bitkileri genellikle ana ürünün büyüme sezonu dışındaki dönemlerde ekilir. Örneğin, mısır veya buğday tarlalarında, kış örtücü bitkileri olarak çavdar veya yulaf kullanılır (Gabriel ve ark., 2013).

Polikültür Sistemlerde: Çeşitli bitkilerin birlikte yetiştirildiği sistemlerde, örtücü bitkileri bir çeşit bitki çeşitliliği sağlar ve biyolojik çeşitliliği artırır. Bu, zararlı organizmaların kontrolünü kolaylaştırır ve toprak sağlığını iyileştirir (Lehman ve ark., 2015).

No-Till Sistemlerde: Toprak işlemez tarım sistemlerinde, örtücü bitkileri, toprak işleme ihtiyacını azaltarak toprağın yapısını korur ve erozyonu önler. Örneğin, kışlık baklagil ve ot karışımları, bahar ekiminden önce doğal olarak geri döner ve toprağın korunmasına katkı sağlar (McVay ve ark., 1989).

Mevcut Araştırma Boşlukları ve Gelecekteki Yönlendirmeler

Örtücü bitkileri ile ilgili araştırmalar, genel olarak geniş coğrafi alanlara yayılmış olsa da, bölgeye özgü kılavuzlara olan ihtiyaç hala önemini korumaktadır. Farklı iklim koşulları, toprak tipleri ve tarımsal uygulamalar, örtücü bitkilerinin etkisini doğrudan etkileyebilir. Bu nedenle, bölgeye özgü çalışmalara ve kılavuzlara ihtiyaç vardır (Lehman ve ark., 2015). Örtücü bitkilerinin optimal kullanımı ve yönetimi için yerel koşullara uygun çözümler geliştirilmelidir (Williams ve Weil, 2004).

Örtücü bitkilerinin toplam faydaları üzerine uzun vadeli çalışmalar sınırlıdır. Kısa vadeli araştırmalar, örtücü bitkilerinin toprağı koruma ve besin geri dönüşümünde sağladığı faydaları göstermiş olsa da, uzun vadeli etkiler hakkında daha fazla bilgiye ihtiyaç vardır (Gabriel ve ark., 2013). Uzun vadeli çalışmalar, örtücü bitkilerinin toprağın biyolojik aktivitesi ve yapısı üzerindeki kalıcı etkilerini daha iyi anlamamıza yardımcı olabilir (Blouin ve ark., 2013).

Kaplama mahsulü uygulamalarında yenilikler, örtücü bitkilerinin daha etkin ve verimli bir şekilde kullanılmasını sağlayabilir. Bu yenilikler, biyoteknolojik gelişmeler, toprak sağlığına yönelik yeni yaklaşımlar ve dijital tarım teknolojileri gibi alanlarda olabilir (Chen ve Weil, 2010). Örneğin, örtücü bitkilerinin genetik modifikasyonu, belirli zararlılara karşı direnç geliştirilmesi veya kök sistemlerinin güçlendirilmesi gibi yenilikçi yaklaşımlar, kaplama mahsulü sistemlerinde büyük faydalar sağlayabilir (McVay ve ark., 1989).

Sonuç

Kaplama mahsulleri, küresel tarımsal zorlukların ele alınmasında önemli bir rol oynamaktadır. Bu bitkiler, toprak sağlığını koruma, erozyonu önleme ve kimyasal girdilerin azaltılmasını sağlamada etkili araçlardır. Örtücü bitkileri, toprakta organik madde miktarını artırarak biyolojik aktiviteyi destekler ve toprak yapısını iyileştirir. Ayrıca, su tutma kapasitesini artırarak ve besin maddelerinin kaybını azaltarak sürdürülebilir tarım uygulamalarının yaygınlaştırılmasına yardımcı olurlar. Bu, dünya genelinde tarımsal üretim sistemlerinin dayanıklılığını artırır ve çevresel baskıları azaltır. Örtücü bitkilerinin sağladığı faydaların en üst düzeye çıkarılması için çiftçiler ve araştırmacılar arasında iş birliği büyük önem taşımaktadır. Örtücü bitkilerinin ekim ve sonlandırma zamanlaması konusunda dikkatli planlama yapılmalıdır. Bölgeye özgü örtücü bitkisi seçimleri yapılmalı ve yerel koşullara uygun uygulamalar geliştirilmelidir. Örtücü bitkilerinin uzun vadeli etkileri üzerine daha fazla çalışma yapılmalıdır. Kaplama mahsulü uygulamalarında yenilikçi yöntemler ve teknolojiler araştırılmalıdır. Sürdürülebilir tarım uygulamalarını teşvik eden politikalar geliştirilmelidir. Çiftçilerin örtücü bitkilerini kullanmalarını teşvik eden finansal ve teknik destek programları oluşturulmalıdır. Sonuç olarak, kaplama mahsulleri, küresel tarımsal zorlukların üstesinden gelinmesinde ve sürdürülebilir tarımın yaygınlaştırılmasında önemli bir araçtır.

Kaynaklar

Abdollahi L, Munkholm LJ, Garbout A, Heckrath G. (2014). The effects of organic matter application and intensive tillage and cover cropping on soil quality. *Soil Science Society of America Journal*. 78(3):897-905.

Abdollahi, L., & Munkholm, L. J. (2014). Tillage system and cover crop effects on soil quality: I. Chemical, mechanical, and biological properties. *Soil Science Society of America Journal*, 78(1), 262-270.

Altieri, M. A. (1995). *Agroecology: The science of sustainable agriculture*. CRC Press.

Blouin, M., Hodson, M. E., Delgado, E. A., Baker, G., Brussaard, L., Butt, K. R., ... & Peres, G. (2013). A review of earthworm impact on soil function and ecosystem services. *European Journal of Soil Science*, 64(2), 161-182.

Brown LC, Norton LD. (1994). Surface residue effects on soil erosion from ridges of different soils and formation *Transactions of the ASAE*. 37(5):1515-1524.

Brown, T. T., & Norton, U. (1994). Effects of cover crops on soil properties and weed dynamics in conservation tillage systems. *Agronomy Journal*, 86(1), 107-113.

Chen, G., & Weil, R. R. (2010). Penetration of cover crop roots through compacted soils. *Plant and Soil*, 331(1-2), 31-43.

Dabney, S. M. (1998). Cover crop impacts on watershed hydrology. *Journal of Soil and Water Conservation*, 53(3), 207-213.

Drinkwater, L. E., Wagoner, P., & Sarrantonio, M. (1998). Legume-based cropping systems have reduced carbon and nitrogen losses. *Nature*, 396(6708), 262-265.

Gabriel JL, Garrido A, Quemada M. (2013). Cover crops effect on farm benefits and nitrate leaching: Linking economic and environmental analysis. *Agricultural Systems*. Oct 1;121:23-32

Gabriel, J. L., Muñoz-Carpena, R., Quemada, M. (2013). The role of cover crops in irrigated systems: Water balance, nitrate leaching, and soil water retention. *Agricultural Water Management*, 121, 32-44.

Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., ... & Vöösmary, C. J. (2004). Nitrogen cycles: past, present, and future. *Biogeochemistry*, 70, 153-226.

Haktanır, K., & Arcaç, S. (1997). Toprak biyolojisi (Toprak ekosistemine giriş).

Lehman RM, Cambardella CA, Stott DE, Acosta-Martínez V, Manter DK, Buyer JS, (2015). Understanding and improving soil biological health: The solution for reversing soil degradation. *Sustainability*. 2015;7(1):988-1027

Lehman, R. M., Osborne, S. L., Duke, S. E., Cambardella, C. A., & Van Tassel, D. L. (2015). Cover crop effects on soil microbial communities and biochemical functions. *Applied Soil Ecology*, 91, 34-41.

McVay, K. A., Radcliffe, D. E., & Hargrove, W. L. (1989). Winter legume effects on soil properties and nitrogen fertilizer requirements. *Soil Science Society of America Journal*, 53(6), 1856-1862.

Parker, E. C. (1915). *Field management and crop rotation: planning and organizing farms; Crop rotation systems; soil amendment with fertilizers; relation of animal husbandry to soil productivity; and other important features of farm management*. Webb Publishing Company.

Schipanski, M. E., Barbercheck, M., Douglas, M. R., Finney, D. M., Haider, K., Kaye, J. P., & White, C. (2014). A framework for evaluating ecosystem services provided by cover crops in agroecosystems. *Agricultural Systems*, 125, 12-22.

Snapp, S. S., Swinton, S. M., Labarta, R., Mutch, D., Black, J. R., Leep, R., ... O'Neil, K. (2005). Evaluating cover crops for benefits, costs, and performance within cropping system niches. *Agronomy Journal*, 97(1), 322–332.

Teasdale, J. R., & Mohler, C. L. (2000). The quantitative relationship between weed emergence and the physical properties of mulches. *Weed Science*, 48(3), 385-392.

Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural sustainability and intensive production practices. *Nature*, 418(6898), 671–677.

Williams, S. M., & Weil, R. R. (2004). Cover crop root channels may alleviate soil compaction effects on soybean crop. *Soil Science Society of America Journal*, 68(4), 1403-1409.

AGRICULTURE IN KAHRAMANMARAŞ

KAHRAMANMARAŞ'TA TARIM

İsmail GÜVENÇ

Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Horticulture,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0000-0002-4686-9487>

Arzu AKBAŞ

Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Horticulture,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0009-0001-8242-7950>

Soner AKCAN

Kahramanmaraş Sütçü İmam University, Faculty of Agriculture, Department of Horticulture,
Kahramanmaraş, Turkey

ORCID ID: <https://orcid.org/0009-0001-9522-6875>

ABSTRACT

Introduction and Purpose: Kahramanmaraş is a province located in the Mediterranean Region of Turkey, known for its high agricultural potential. Its fertile lands and diverse climate conditions have made agriculture a cornerstone of the region's economy. This study aims to examine the agricultural activities of the province in terms of crop production, livestock, and aquaculture.

Materials and Methods: The study analyzed agricultural data from Kahramanmaraş. Information related to crop production, livestock, and aquaculture was compiled from official sources such as the Ministry of Agriculture and Forestry and the Turkish Statistical Institute (TÜİK). The province's agricultural areas, cultivated crops, livestock population, and aquaculture production were analyzed.

Result: Kahramanmaraş has 375,309 hectares of agricultural land. Field crops (74%), including wheat, sunflower, and corn, are prominent. Garden crops such as red pepper, grapes, and walnuts are economically significant. The province has 231,500 cattle and 990,593 small ruminants. Trout production reached 8,405 tons in 2023.

Discussion and Conclusion: Despite its high agricultural potential, the province faces challenges in productivity, modernization, and sustainability. These challenges can be overcome through technological investments and efficient resource management. Kahramanmaraş holds significant potential in the agricultural sector. Steps taken in crop production, livestock, and aquaculture will contribute substantially to the province's economic development.

Keywords: Kahramanmaraş, agriculture, crop production, livestock, aquaculture, sustainability, red pepper production

ÖZET

Giriş ve Amaç: Kahramanmaraş, Türkiye'nin Akdeniz Bölgesi'nde yer alan ve tarımsal potansiyeli yüksek bir ildir. Verimli toprakları ve çeşitli iklim koşulları, tarımı bölge ekonomisinin temel taşı haline getirmiştir. Bu çalışma, ilin tarımsal faaliyetlerini bitkisel üretim, hayvancılık ve su ürünleri açısından incelemeyi amaçlamaktadır.

Gereç ve Yöntem: Çalışmada, Kahramanmaraş'ın tarımsal verileri incelenmiştir. Bitkisel üretim, hayvancılık ve su ürünleriyle ilgili bilgiler, Tarım ve Orman Bakanlığı ve TÜİK gibi resmi kaynaklardan derlenmiştir. İlin tarım alanları, yetiştirilen ürünler, hayvan varlığı ve su ürünleri üretimi analiz edilmiştir.

Bulgular: Kahramanmaraş'ta 375.309 hektar tarım alanı bulunmaktadır. Tarla bitkileri (%74) arasında buğday, ayçiçeği ve mısır öne çıkmaktadır. Kırmızıbiber, üzüm ve ceviz gibi bahçe bitkileri ekonomik açıdan önemlidir. İlde 231.500 büyükbaş ve 990.593 küçükbaş hayvan bulunmaktadır. Alabalık üretimi ise 2023'te 8.405 tona ulaşmıştır.

Tartışma ve Sonuç: İl, tarımsal potansiyeli yüksek olmasına rağmen verimlilik, modernizasyon ve sürdürülebilirlik konularında zorluklarla karşı karşıyadır. Bu zorluklar, teknolojik yatırımlar ve etkin kaynak yönetimiyle aşılabılır. Kahramanmaraş, tarım sektöründe büyük bir potansiyele sahiptir. Bitkisel üretim, hayvancılık ve su ürünleri alanında atılacak adımlar, ilin ekonomik kalkınmasına önemli katkılar sağlayacaktır.

Anahtar Kelimeler: Kahramanmaraş, tarım, bitkisel üretim, hayvancılık, su ürünleri, sürdürülebilirlik, kırmızıbiber üretimi

USE OF THE ASTAXANTHIN IN TILAPIA (*OREOCHROMIS NILOTICUS*)

TİLAPİALARDA (*OREOCHROMIS NILOTICUS*) ASTAKSANTİN KULLANIMI

Ilayda Gizem AKIN

The Graduate School of Health Sciences of Ankara University, Ankara University Faculty of Veterinary Medicine, Department of Animal Nutrition and Nutritional Diseases, Ankara, Turkey

ORCID ID: <https://orcid.org/0009-0003-3260-4984>

Prof. Dr. Ozge SIZMAZ

Ankara University Faculty of Veterinary Medicine, Department of Animal Nutrition and Nutritional Diseases, Ankara, Turkey

ORCID ID: <https://orcid.org/0000-0002-2027-5074>

ABSTRACT

Astaxanthin, a natural carotenoid pigment, has significant effects on coloration, growth performance, and antioxidant capacity in fish. Its use, particularly in important aquaculture species such as tilapia, has gained increasing importance due to consumer demand. The meat quality of tilapia is directly related to improvements in skin color. In this context, the potential of astaxanthin to enhance the external appearance of fish presents a significant advantage for producers. The use of astaxanthin in tilapia farming is also noteworthy for improving growth performance. Fish supplemented with astaxanthin have shown improved feed conversion rates (FCR) and increased body weight gains. Moreover, astaxanthin supplementation enhances the antioxidant capacity of fish, reducing oxidative stress. This is especially beneficial in minimizing the effects of environmental factors such as disease and stress. The effects of astaxanthin are not limited to growth and health; they also result in noticeable improvements in tilapia pigmentation. This pigment strengthens the reddish hues in the skin and muscle tissues of tilapia, leading to more attractive and higher market-value products. Studies conducted with various dosages have demonstrated that astaxanthin is effective in enhancing pigmentation levels in tilapia. Depending on the dosage, more pronounced and desirable color changes can be observed in the fish. These findings suggest that astaxanthin can enhance both efficiency and quality in tilapia farming. However, it is emphasized that the optimal dosage and application duration need to be determined. Future research will help further explore these parameters and contribute to the development of more efficient production strategies in the industry. In conclusion, it is understood that the use of astaxanthin positively impacts growth, coloration, and antioxidant capacity in tilapia production. However, more comprehensive research is necessary to establish the appropriate application conditions and further enhance the effectiveness of astaxanthin.

Key Words: Tilapia; Astaxanthin; Pigmentation; Antioxidant effect

ÖZET

Astaksantin, doğal bir karotenoid pigmenti olup, balıklarda renklenme, büyüme performansı ve antioksidan kapasite üzerinde önemli etkiler göstermektedir. Özellikle tilapia gibi önemli su

ürünleri türlerinde, astaksantin kullanımını tüketici talebine bağlı olarak daha fazla önem kazanmaktadır. Tilapyaların et kalitesi, özellikle deri rengindeki iyileşmelerle doğrudan ilişkilidir. Bu bağlamda astaksantin, balıkların dış görünüşünü iyileştirme potansiyeli, üreticiler için büyük bir avantaj sunmaktadır. Tilapyaların yetiştirilmesinde astaksantin kullanımını, büyüme performansını artırma yönünden de dikkat çekicidir. Astaksantin takviyesi yapılan balıklarda, yemden yararlanma oranları (FCR) iyileşmiş ve canlı ağırlık kazançları artmıştır. Bunun yanında, astaksantin desteği, balıkların antioksidan kapasitesini artırarak oksidatif stresi azaltmaktadır. Bu, özellikle hastalık ve stres gibi çevresel faktörlerin etkilerini minimize etme açısından faydalıdır. Astaksantin etkisi sadece büyüme ve sağlıkla sınırlı değildir; aynı zamanda tilapyaların renklenmesi üzerinde de gözle görülür bir iyileşme sağlanmaktadır. Bu pigment, tilapyaların deri ve et dokularındaki kırmızımsı renk tonlarını güçlendirerek, daha cazip ve pazar değeri yüksek ürünler elde edilmesine olanak tanır. Farklı dozajlarla yapılan çalışmalar, astaksantin tilapyalarda pigmentasyon düzeyini artırmada etkili olduğunu göstermektedir. Doza bağlı olarak, balıklarda daha belirgin ve istenilen renk değişimlerine yol açmaktadır. Bu bulgular, astaksantin tilapya yetiştiriciliğinde verimlilik ve kaliteyi artırabileceğini ortaya koymaktadır. Ancak optimal dozajın ve uygulama süresinin belirlenmesi gerektiği vurgulanmaktadır. Gelecekte yapılacak araştırmalar, bu parametrelerin daha ayrıntılı bir şekilde incelenmesini sağlayarak, sektörde daha verimli üretim stratejilerinin geliştirilmesine katkı sağlayacaktır. Sonuç olarak, astaksantin kullanımının tilapya üretiminde büyüme, renklenme ve antioksidan kazanımı üzerinde olumlu etkileri olduğu anlaşılmaktadır. Bununla birlikte, doğru uygulama koşullarının belirlenmesi ve astaksantin etkinliğini artırmak adına daha kapsamlı araştırmalar yapılması gerekmektedir.

Anahtar Kelimeler: Tilapya; astaksantin; pigmentasyon; antioksidan etki

GİRİŞ

Son on yılda su ürünleri yetiştiriciliği alanında önemli ilerlemeler kaydedilmiştir (Yücel & Tolon, 2024). Dünya su ürünleri sektörü hem küresel gıda güvenliği hem de ekonomik kalkınma açısından giderek artan bir öneme sahiptir. Birleşmiş Milletler Gıda ve Tarım Örgütü (FAO) tarafından yayımlanan 2024 raporuna göre, 2022 yılında toplam su ürünleri üretimi 187 milyon tona ulaşmış olup, bunun %51'i yetiştiricilik yoluyla sağlanmıştır. Asya, %70'lik payıyla sektörde lider konumda bulunurken, Çin %35'lik payıyla en büyük üretici olarak öne çıkmaktadır (T.C. Tarım ve Orman Bakanlığı, 2024). Toplam su ürünleri üretiminin %89'u doğrudan insan tüketimi için kullanılmakta olup, bu durum, balıkçılık ve yetiştiricilik faaliyetlerinin küresel gıda güvenliği için kritik bir rol oynadığını göstermektedir (Fisheries, FAO, 2022).

Son yıllarda, pazardaki rekabet gücünü artırmak ve deniz balıklarına alternatif bir ürün sunmak amacıyla tilapyanın deniz ortamında yetiştirilmesi ve ticarileştirilmesi önemli bir potansiyel olarak gündeme gelmiştir. Tilapia yetiştiriciliğinde, diğer birçok balık türünde olduğu gibi, ürün kalitesi ve verimi; stok yoğunluğu, kullanılan yavru balıkların (fingerling) ortalama boyutları, tercih edilen tür, üretim sistemi ve çevresel koşullar gibi çeşitli faktörlerden etkilenmektedir (Dey, 2001). Tilapia, yetiştirilmesi kolay ve geniş çevresel koşullara uyum sağlayabilen bir balık türüdür (Sukri ve ark., 2022).

Afrika kökenli olan tilapia, istilacı tür olarak kabul edilmekte olup, tatlı sularda yaygın olarak bulunmaktadır. Uzun sırt yüzgeci ve dikenli yüzgeçleri ile karakterize edilen bu tür, yüksek et kalitesi, temel protein kaynağı olması ve çekici renk özellikleri nedeniyle pazarda büyük bir talep görmektedir (Arous ve ark., 2014). Ancak, kafes yetiştiriciliği ile üretilen tilapyalarda karşılaşılan başlıca sorunlardan biri, deri ve et renginin soluk olmasıdır. Bu durum, özellikle astaksantin eksikliğine bağlı olarak ortaya çıkmaktadır. Hayvanlar, vücutlarında karotenoid pigmentlerini sentezleyemedikleri için, bu bileşeni tamamen diyet yoluyla almak zorundadır (Galasso ve ark., 2017).

Astaksantin, karotenoid grubuna ait, dokuda kırmızı-turuncu renklenmeyi sağlayan bir pigmenttir. Kimyasal yapısı C₄₀H₅₂O₄ olup, merkezindeki konjuge çift bağlar nedeniyle kırmızı renge sahiptir (Higuera ve ark.,2006). Bu bağlar, serbest radikallerle reaksiyona girerek antioksidan özellik gösterir ve hücreleri oksidatif hasara karşı korur (Guerin ve ark., 2003; Yuan ve ark., 2011). Astaksantin, diğer antioksidanlarla kıyaslandığında, özellikle lipit ve DNA hasarını önlemede daha etkin bulunmuştur (Naguib, 2000). Aynı zamanda, su ürünleri yetiştiriciliğinde büyük önem taşıyan pigmentasyon, büyüme ve bağışıklık sistemi üzerindeki olumlu etkileri nedeniyle yaygın olarak kullanılmaktadır (Rahman ve ark., 2016; Jagruthi ve ark., 2014; Han ve ark., 2018).

Astaksantin, doğal olarak algler, mantarlar, mayalar ve bakteriler tarafından sentezlenmektedir (Elbahnaswy & Elshopakey, 2024). Ticari olarak sentetik astaksantin ise kimyasal sentez veya mikroorganizma fermentasyonu yoluyla üretilmektedir. Özellikle *Haematococcus pluvialis* mikroalgleri, astaksantin için en zengin doğal kaynaklardan biri olarak gösterilmektedir (Ambati ve ark., 2014; Zhao ve ark., 2019).

Karotenoidlerin büyüme performansını artırma mekanizmaları üzerine çeşitli hipotezler öne sürülmüştür. Bunlardan ilki, karotenoidlerin sindirim enzimlerinin aktivitesini artırarak besinlerin sindirimini ve emilimini desteklemesiyle yem tüketimini artırmasıdır (Baron ve ark., 2008; Zhang ve ark., 2013). İkinci olarak, karotenoidlerin kabuklu sucul organizmaların kabuk değiştirme döngülerini kısaltabileceği ve NADPH metabolizmasını yöneterek enerji tüketimini optimize edebileceği belirtilmiştir (Hertrampf & Piedad-Pascual, 2012; Mao ve ark., 2017). Son olarak, astaksantin bağırsak florası üzerinde olumlu etkileri olduğu ve sindirilemeyen bileşenlerin biyoyararlanımını artırarak ek besin desteği sağlayabileceği düşünülmektedir (Vasudevan ve ark., 2006). Astaksantin ile yapılan çalışmalar tablo 1 de verilmiştir.

Tablo 1: Balıklarda Astaksantin Kullanımı

BALIK TÜRÜ	DOZ	SONUÇ	REFERANS
Asian tiger shrimp (<i>Penaeus monodon</i>)	100 mg kg ⁻¹ AX*	Hipoksi koşullarına karşı ağırlık artışında azalma olmamış ve balıkların hayatta kalma oranında iyileşme görülmüştür.	Niu ve ark., 2014
Avrupa Levreği (<i>Dicentrarchus labrax</i>)	60,80 ve 100 mg kg ⁻¹ AX* dozu 60 gün boyunca	Büyüme performansında artış, osmotik strese karşı direnç gelişmiştir. Ayrıca hepatik Süperoksit dismutaz ve glutatyon peroksidazı artırdığı, bağırsak mukozal fagositik ve lizozim aktivitelerini artırdığı gözlenmiştir.	Kurnia ve ark., 2007
Chinese mitten crab (<i>Eriocheir sinensis</i>)	0,30,60,90 ve 120 mg kg ⁻¹ AX dozu, 40 gün	AX* takviyeli diyetler, kontrol grubuna göre daha iyi büyüme ve hayatta kalma oranı sağladığı 60-90 mg/kg AX* takviyesi olan Chinese mitten crab'de (<i>Eriocheir sinensis</i>) antioksidan kapasite, bağışıklık ve amonyak direncini artırdığı Hepatopankreas ve hemolimf dokularında en yüksek toplam	Jiang ve ark., 2020

		antioksidan kapasitesi ve en düşük oksidatif stres göstergeleri (MDA) Diyet 3 ve 4 ile elde edildiği görülmüştür.	
Diskus balıkları	10,50 ve 100 mg/kg dozlarında AX, 4 yumurtlama dönemi boyunca	Astaksantin'in yumurta kırmızılık derecesi üzerindeki etkisi, 100 mg/kg AX* içeren grupta kontrol grubuna göre %77'lik bir artış gösterdiği gözlenmiştir.	Eralp, 2011
Gökkuşuğu alabalığı	50 ppm dozunda AX*, 225 gün boyunca	AX*, kas ve karaciğer dokusunda Total oksidan kapasite ve Oksidatif stres indeksi düzeylerini önemli ölçüde düşürdüğü ve doymamış yağ asidi ile TAK düzeylerini değiştirmedeği Balıkların serumunda AST, ALT ve PON aktivitelerini ile LDL düzeyini düşürürken; HDL düzeyini arttırdığı görülmüştür.	Gökçeoğlu, 2022
Sarı prenses (<i>Labidochromis caeruleus</i>) ve Mavi prenses (<i>Pseudotropheus socolofi</i>)	Kadife çiçeği (<i>Tagetes erecta</i> , KÇÖ) özütü, Astaksantin (AX) ve Mannan-oligosakkarit (MOS) katkılı yemler, 30 gün boyunca	Her iki türde de en iyi büyüme ve renklenme değerlerinin %4 KÇÖ, 50 mg AX/kg yem ve %1 MOS katkılı gruplarda elde edildiği görülmüştür.	Çavdar,2020
Pasifik beyaz karidesi	50 ppm	Karideslerin ağırlık artışını ve büyüme hızını artırdığı gözlenmiştir.	Fang ve ark.,2021
Pasifik beyaz karidesi	25-150 mg/kg	Düşük çözünmüş oksijen stresine karşı hayatta kalma oranını yükselttiği ve oksidatif stresi azalttığı görülmüştür. Antioksidan enzim aktivitelerini artırmış ve inflamasyon ile apoptozla ilişkili gen ekspresyonlarında iyileşme sağladığı görülmüştür	Zhang ve ark.,2013
Pasifik beyaz karidesi	120-150 mg/kg	Oksidatif stresin azalttığı görülmüştür.	Ju ve ark.,2011

Kırmızı tilapialarda	0, %4, %8 ve %12 konsantrasyonlarında AX dozu 6 hafta boyunca	Astaksantin takviyesi ile balıkların renginin iyileştiği görülmüştür. Büyüme performansında belirgin bir fark gözlemlenmemiştir. En yüksek kızarıklık %12 astaksantin eklenen grupta görülmüştür. Astaksantin seviyesinin artması, daha fazla iyileşmeye yol açmamıştır	Harith ve ark., 2024
Tilapia (<i>Oreochromis spp</i>)	0, 0.5, 1.0, 1.5 g/kg 90 gün boyunca	1 ve 1.5 g/kg dozlarında büyüme performansı, immün yanıt ve antioksidan kapasitesinde artış FCR da iyileşme görülmüştür.	Eldessouki ve ark., 2024
Tilapia (<i>Oreochromis niloticus</i>)	0 g/kg (T1; kontrol), 5 g/kg (T2), 10 g/kg (T3) ve 20 g/kg (T4) 8 hafta Soğuk Stresi	T4 diyetiyle beslenen hem erkek hem de dişi tilapialarda kolesterol içeriğinde önemli bir artış gözlenmiştir. Kaslardaki toplam karotenoid içeriği hem erkek hem de dişi tilapia balıklarında T4 takviyesi ile beslenenlerde önemli ölçüde daha yüksek değerler görülmüştür.	Vongkampang ve ark., 2023
Tilapia (<i>Oreochromis niloticus</i>)	0 g/kg (T1), 5 g/kg (T2), 10 g/kg (T3), 20 g/kg (T4) 4 hafta	T4 grubunda kas karotenoid içeriğinde ve antioksidan aktivitede artış, Erkeklerde spermatozoa konsantrasyonu Dişilerde yumurta üretiminde iyileşme görülmüştür.	Panase ve ark., 2024

AX*: Astaksantin

SONUÇ

Astaksantin su ürünleri yetiştiriciliğine entegrasyonu, yalnızca balık sağlığını desteklemekle kalmayıp, aynı zamanda sektörün sürdürülebilirliğine yönelik yenilikçi çözümler sunmaktadır. Bu doğal pigment, biyolojik faydaları ve ekonomik potansiyeli ile su ürünleri sektörünün büyümesine katkıda bulunabilecek güçlü bir araçtır. Astaksantin doğru kullanımı hem balık sağlığını desteklemekte hem de çevresel etkileri azaltarak daha bilinçli ve bütüncül yetiştiricilik uygulamalarını teşvik etmektedir. Ayrıca, oksidatif stres ve balık sağlığı arasındaki dengeyi anlamak, hızla değişen çevresel koşullara uyum sağlamak ve türlerin korunmasını desteklemek açısından kritik bir öneme sahiptir. İnsan faaliyetlerinin sucul yaşam üzerindeki etkisinin giderek arttığı günümüzde, astaksantin gibi yenilikçi çözümler, su ürünleri yetiştiriciliğinde sürdürülebilir üretimi ve ekolojik dengeyi sağlamada kilit bir rol oynamaktadır. Bu bağlamda, astaksantin araştırmaları, akuakültürün geleceğine dair umut vadeden bir yol haritası sunmaktadır.

KAYNAKÇA

- Ambati, R. R., Phang, S. M., Ravi, S., & Aswathanarayana, R. G. (2014). Astaxanthin: Sources, extraction, stability, biological activities and its commercial applications—A review. *Marine drugs*, 12(1), 128-152.
- Arous, W. H., El-Bermawi, N. M., Shaltout, O. E., & Essa, M. A. (2014). Effect of adding different carotenoid sources on growth performance, pigmentation, stress response and quality in red tilapia (*Oreochromis spp.*). *Middle East J. Appl. Sci*, 4(4), 988-999.
- Baron, M., Davies, S., Alexander, L., Snellgrove, D., & Sloman, K. A. (2008). The effect of dietary pigments on the coloration and behaviour of flame-red dwarf gourami, *Colisa lalia*. *Animal Behaviour*, 75(3), 1041-1051.
- Çavdar, N. (2020). Kadife çiçeği (*Tagetes erecta*), astaksantin ve mannan-oligosakkarit'in *labidochromis caeruleus* ve *pseudotropheus socolofi* (Cichlidae Bonaparte, 1835) yavrularının büyüme, renklenme, karaciğer ve bağırsak histolojisi üzerine etkileri.
- Dey, M. M. 2001. *Tilapia Production In South Asia And Far East. Tilapia Production, Marketing And Technological Developments. S. Subasinghe And Tarlochan Singh (Editors).*
- Elbahnaswy, S., & Elshopakey, G. E. (2024). Recent progress in practical applications of a potential carotenoid astaxanthin in aquaculture industry: a review. *Fish Physiology and Biochemistry*, 50(1), 97-126.
- Eldessouki, E. A., Elshopakey, G. E., Elbahnaswy, S., Shakweer, M. S., Abdelwarith, A. A., Younis, E. M., ... & Eissa, E. S. H. (2024). Influence of astaxanthin-enriched *Haematococcus pluvialis* microalgae on the growth efficacy, immune response, antioxidant capacity, proinflammatory cytokines, and tissue histomorphology of hybrid red tilapia. *Aquaculture International*, 1-22.
- Eralp, H. (2011). *Diskus balıklarında (symphysodon spp.) anaç yemlerine eklenen farklı oranlarda astaksantin ilavesinin yumurta verimi, kalitesi, açılım oranı ve larvalar üzerine etkisinin belirlenmesi (Doctoral dissertation, SDÜ Fen Bilimleri Enstitüsü).*
- Fang, H., Xie, J., Zhao, W., Liu, Z., Liu, Y., Tian, L., & Niu, J. (2021). Study supplementation of astaxanthin in high-fat diet on growth performance, antioxidant ability, anti-inflammation, non-specific immunity and intestinal structure of juvenile *Trachinotus ovatus*. *Aquaculture Nutrition*, 27(6), 2575-2586.
- Fisheries, F. A. O. (2022). *The state of world fisheries and aquaculture towards blue transformation.*
- Galasso, C., Corinaldesi, C., & Sansone, C. Carotenoids from marine organisms: biological functions and industrial applications. *Antioxidants (Basel)*. 2017; 6 (4): 96.
- Gökçeoğlu, A. (2022). *Astaksantin Uygulamasının Gökkuşluğu Alabalığı (Oncorhynchus Mykiss) Oksidan, Antioksidan ve Yağ Asitleri Düzeyleri Üzerine Etkisi*
- Guerin, M., Huntley, M. E., & Olaizola, M. (2003). *Haematococcus astaxanthin: applications for human health and nutrition. TRENDS in Biotechnology*, 21(5), 210-216.
- Han, T., Li, X., Wang, J., Wang, C., Yang, M., & Zheng, P. (2018). Effects of dietary astaxanthin (AX) supplementation on pigmentation, antioxidant capacity and nutritional value of swimming crab, *Portunus trituberculatus*. *Aquaculture*, 490, 169-177.
- Harith, Z. T., Sukri, S. M., Remlee, N. F. S., Sabir, F. N. M., & Zakaria, N. N. A. (2024). Effects of dietary astaxanthin enrichment on enhancing the colour and growth of red tilapia, *Oreochromis sp.* *Aquaculture and Fisheries*, 9(1), 52-56.

- Hertrampf, J. W., & Piedad-Pascual, F. (2012). Handbook on ingredients for aquaculture feeds. Springer Science & Business Media.
- Higuera-Ciapara, I., Felix-Valenzuela, L., & Goycoolea, F. M. (2006). Astaxanthin: a review of its chemistry and applications. *Critical reviews in food science and nutrition*, 46(2), 185-196.
- Jagruthi, C., Yogeshwari, G., Anbazahan, S. M., Mari, L. S. S., Arockiaraj, J., Mariappan, P., ... & Harikrishnan, R. (2014). Effect of dietary astaxanthin against *Aeromonas hydrophila* infection in common carp, *Cyprinus carpio*. *Fish & shellfish immunology*, 41(2), 674-680.
- James, R., Sampath, K., Thangarathinam, R., & Vasudevan, I. (2006). Effect of dietary spirulina level on growth, fertility, coloration and leucocyte count in red swordtail, *Xiphophorus helleri*. *Israeli Journal of Aquaculture-Bamidgeh*, 58.
- Ju, Z. Y., Deng, D. F., Dominy, W. G., & Forster, I. P. (2011). Pigmentation of Pacific white shrimp, *Litopenaeus vannamei*, by dietary astaxanthin extracted from *Haematococcus pluvialis*. *Journal of the World Aquaculture Society*, 42(5), 633-644.
- Kurnia, A., Satoh, S., Kuramoto, D., & Hanzawa, S. (2007). Effect of different astaxanthin sources on skin pigmentation of red sea bream (*Pagrus major*). *Aquaculture Science*, 55(3), 441-447.
- Mao, X., Guo, N., Sun, J., & Xue, C. (2017). Comprehensive utilization of shrimp waste based on biotechnological methods: A review. *Journal of Cleaner Production*, 143, 814-823.
- Naguib, Y. M. (2000). Antioxidant activities of astaxanthin and related carotenoids. *Journal of agricultural and food chemistry*, 48(4), 1150-1154.
- Niu, J., Wen, H., Li, C. H., Liu, Y. J., Tian, L. X., Chen, X. U., ... & Lin, H. Z. (2014). Comparison effect of dietary astaxanthin and β -carotene in the presence and absence of cholesterol supplementation on growth performance, antioxidant capacity and gene expression of *Penaeus monodon* under normoxia and hypoxia condition. *Aquaculture*, 422, 8-17.
- Panase, P., Vongkampang, T., Wangkahart, E., & Sutthi, N. (2023). Impact of astaxanthin-enriched bacterium (*Paracoccus carotinifaciens*) on growth, immune response, and reproduction performance of broodstock Nile tilapia during winter season.
- Panase, P., Vongkampang, T., Wangkahart, E., & Sutthi, N. (2024). Impacts of astaxanthin-enriched *Paracoccus carotinifaciens* on growth, immune responses, and reproduction performance of broodstock Nile tilapia during winter season. *Fish Physiology and Biochemistry*, 1-20.
- Rahman, M. M., Khosravi, S., Chang, K. H., & Lee, S. M. (2016). Effects of dietary inclusion of astaxanthin on growth, muscle pigmentation and antioxidant capacity of juvenile rainbow trout (*Oncorhynchus mykiss*). *Preventive nutrition and food science*, 21(3), 281.
- Sukri, S. A. M., Andu, Y., Harith, Z. T., Sarijan, S., Pauzi, M. N. F., Wei, L. S., ... & Kari, Z. A. (2022). Effect of feeding pineapple waste on growth performance, texture quality and flesh colour of Nile tilapia (*Oreochromis niloticus*) fingerlings. *Saudi Journal of Biological Sciences*, 29(4), 2514-2519.
- T.C. Tarım ve Orman Bakanlığı, Tarım Reformu Genel Müdürlüğü, Tarımsal Ekonomi ve Politika Geliştirme Enstitüsü. (2024). Su ürünleri raporu (TEPGE Yayın No: 403). Araştırma Yayınları.
- Yuan, J. P., Peng, J., Yin, K., & Wang, J. H. (2011). Potential health-promoting effects of astaxanthin: a high-value carotenoid mostly from microalgae. *Molecular nutrition & food research*, 55(1), 150-165.
- YÜCEL, B., & TOLON, M. T. Su Ürünleri ve Ziraatte Yenilikçi Yaklaşımlar: Sürdürülebilirlik, Teknoloji ve Ekolojik Stratejiler, 2024.

Zhang, J., Liu, Y. J., Tian, L. X., Yang, H. J., Liang, G. Y., Yue, Y. R., & Xu, D. H. (2013). Effects of dietary astaxanthin on growth, antioxidant capacity and gene expression in Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture Nutrition*, 19(6), 917-927.

Zhang, J., Liu, Y. J., Tian, L. X., Yang, H. J., Liang, G. Y., Yue, Y. R., & Xu, D. H. (2013). Effects of dietary astaxanthin on growth, antioxidant capacity and gene expression in Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture Nutrition*, 19(6), 917-927.

Zhao, Y., Yue, C., Geng, S., Ning, D., Ma, T., & Yu, X. (2019). Role of media composition in biomass and astaxanthin production of *Haematococcus pluvialis* under two-stage cultivation. *Bioprocess and biosystems engineering*, 42, 593-602.

EFFECT OF BIOLOGICAL AND ORGANIC FERTILIZERS ON YIELD AND QUALITY IN ORGANIC SUGAR BEET FARMING

ORGANİK ŞEKER PANCARI TARIMINDA BİYOLOJİK VE ORGANİK GÜBRELERİN VERİM VE KALİTE ÜZERİNE ETKİSİ

Hayrullah OZBEK

Kayseri Sugar Sustainable Agriculture Engineer, Kayseri, Turkey

ORCID ID <https://orcid.org/0009-0005-2699-6234>

Halit Burak YUKSEKDAG

Kayseri Sugar Sustainable Agriculture Engineer, Kayseri, Turkey

ORCID ID: <https://orcid.org/0009-0004-1251-6087>

Oguzhan ULU

Kayseri Sugar Deputy General Manager (R&D), Kayseri, Turkey

ORCID ID: <https://orcid.org/0009-0002-6026-959X>

Anıl Mehmet BALTACI

Kayseri Sugar Sustainable Agriculture Manager, Kayseri, Turkey

ORCID ID: <https://orcid.org/0000-0003-1890-131X>

ABSTRACT

Introduction and Purpose: It is estimated that over 100 million tons of nitrogen, potassium, and phosphorus-based chemical fertilizers are used annually worldwide to increase plant production. Organic farming is an agricultural method that rejects synthetic fertilizers and pesticides, focusing instead on enhancing soil fertility through practices like crop rotation, green manure, composting, and biological pest control. This experiment was designed to prevent excessive and unconscious fertilization by sugar beet producers and to statistically compare the effects of organic fertilizers on yield and quality with standard production methods. **Materials and Methods:** The study aimed to identify the most suitable fertilizer for organic sugar beet cultivation. Experiments included fertilizers containing PGPR group bacteria, farm manure, solid worm fertilizer, vermicast tea, and leonardite-based applications. For each plot, 15 beets were randomly selected and sent to a quality control laboratory. Technological quality data were measured using a Betalyser device. Variance analysis was applied to the numerical data based on the experimental design. The F-test determined the significance of differences, and significant differences were grouped using the Least Significant Difference (LSD) method. **Results:** Statistical analysis revealed that the application containing PGPR group bacteria yielded the best results at 7.01 t/da, while the combination of PGPR bacteria with leonardite achieved the highest sugar content at 17.16%. **Discussion and Conclusion:** In this experiment,

PGPRs (Plant Growth-Promoting Rhizobacteria) improved nutrient uptake, activated hormonal activity, and suppressed harmful root microflora, neutralizing pollutants and regulating plant diseases. This resulted in better growth and higher yields. The efficiency of specific PGPRs can be further enhanced by optimizing them for local soil conditions. They are expected to replace, pesticides and chemical manures, which have negative impacts on sustainable agriculture.

Keywords: PGPR; Sugar Beet; Organic Agriculture

ÖZET

Giriş ve Amaç: Dünyada bitkisel üretimi arttırmak için yılda 100 milyon tondan fazla azot, potasyum ve fosforlu kimyasal gübrelere kullanıldığı tahmin edilmektedir. Organik tarım, sentetik gübre ve pestisit gibi kimyasal maddelerin kullanımını reddeden, toprak üretkenliğini arttırmaya odaklanan bir tarım yöntemidir. Bu yaklaşım, bitki nöbetleşmesi, yeşil gübre, kompost ve biyolojik zararlı kontrolü gibi uygulamaları içerir. Bütün bu unsurlar değerlendirildiğinde bu deneme şeker pancarı üreticilerinin aşırı ve bilinçsizce gübreleme yapmasının önüne geçebilmek ve organik gübre kullanımının şeker pancarı üretimindeki verim ve kaliteye etkisini istatistiksel olarak standart üretimle karşılaştırmak için kurulmuştur. **Gereç ve Yöntem:** Bu çalışmada organik şeker pancarı yetiştiriciliği için en uygun gübrenin tespit edilmesi amaçlanmıştır. Denemede PGPR grubu bakteriler içeren gübreler, çiftlik gübresi, katı solucan gübresi, vermicast çayı ve leonardit içerikli uygulamalar yapılmıştır. Her uygulama parseli için 15'er pancar rastgele seçilip kalite kontrol laboratuvarına gönderilmiştir. Betalyser cihazı ile teknolojik kalite verileri ölçülmüştür. Elde edilen sayısal verilere, deneme desenine göre varyans analizi uygulanmıştır. F testi ile farklılıkların önem düzeyi belirlenmiş ve önemli farklar tespit edilmesi durumunda, ortalamalar Asgari Önemli Fark (LSD) yöntemiyle gruplandırılmıştır. **Bulgular:** Yapılan istatistik analizlerin sonucuna göre 7,01 t/da verim açısından en iyi PGPR grubu bakterileri içeren uygulama, şeker varlığı açısından %17,16 ile leonarditle birlikte PGPR grubu bakterileri içeren uygulama olmuştur. **Sonuç ve Değerlendirme:** Bu denemede PGPR'ler bitkilerin daha iyi besin almasını sağlayarak, hormonal aktiviteyi devreye sokarak ve bir yandan da köklerde doğal olarak bulunan zararlı mikroflorayı engelleyerek çeşitli kirleticileri nötralize etme ve birçok bitki hastalığını düzenleme kapasitesine sahiptir. Daha iyi gelişme ve daha yüksek verim sağlamıştır. Belirli bir PGPR'nin verimliliği, yerel toprak koşullarına göre optimize edilerek ve uyarlanarak daha da artırılabilir. Bunların, sürdürülebilir tarım üzerinde bir dizi olumsuz etkisi olan, pestisitler ve kimyasal gübrelere yerini alması beklenmektedir.

Anahtar Kelimeler: PGPR; Şeker Pancarı; Organik Tarım

COMPARATIVE STUDY OF SOME SUGAR PROFILES OF HONEY AND FRUITS**Elsever Asadov**

Nakhchivan State University, Faculty of Medicine, Nakhchivan, Azerbaijan

orcid.org/0000-0003-2892-2974

Sevgi KolaylıFaculty of Science, Department of Chemistry, Karadeniz Technical University, Trabzon,
Türkiye

orcid.org/0000-0003-0437-6139

ABSTRACT

Carbohydrates, particularly sugars, are essential biomolecules that impart taste and flavor to foods. They are primarily sourced from plants in the form of monosaccharides, disaccharides, and oligosaccharides. Beyond serving as a vital energy source, sugars significantly influence the taste and aroma profiles of food products. In this study, the sugar profiles of certain bee products and fruits were compared. Utilizing High-Performance Liquid Chromatography with Refractive Index Detection (HPLC-RID) and eight distinct sugar standards, analyses were conducted on chestnut and pine honey, apricots, and grapes. The methanolic extracts revealed that, in honey samples, fructose and glucose were predominant, followed by sucrose and maltose, with trace amounts of melezitose and trehalose. In apricots, the sugars were primarily ordered as sucrose, glucose, melibiose, and fructose. Grape samples exhibited a sugar composition sequence of fructose, glucose, and sucrose. These findings underscore the variability in sugar compositions across different natural products, highlighting the influence of botanical origin on sugar content and profile.

Keywords: *sugar, profiles, honey, apricot, grapes***Introduction**

Natural or raw honey is a complex mixture, with sugars accounting for approximately 95% of its dry weight. The primary sugar components are fructose and glucose; however, it has been reported that honey also contains 20 to 25 different sugar types, including sucrose, maltose, arabinose, melibiose, melezitose, trehalose, and trehalose. The sugar profile of honey is considered a critical analytical parameter in distinguishing natural honey from adulterated or counterfeit honey samples (Wu et al. 2017; Raatz et al. 2015; Kolaylı et al 2012; Kolaylı et al 2024). The sugar composition of honey varies depending on its floral origin, and there are differences in both the total sugar content and sugar profiles between floral honeys and honeydew honeys (Ucurum et al. 2023).

Similar to honey, fruits naturally contain sugars, primarily fructose, glucose, and sucrose. These sugars not only define the fruit's flavor profile but also serve as a significant source of energy. Fructose is typically the predominant sugar, found in higher concentrations in ripe fruits. The sugar content in fruits varies depending on the type of fruit, its ripeness, and growing conditions. Additionally, the fiber and bioactive compounds present in fruits positively influence the metabolism of these sugars, contributing to a healthy diet (Beckles, D. M. (2012; Basson et al. 2010). Fruits with high sugar content are characterized by their intense sweet flavors and are rich in energy. Bananas, grapes, figs, mangoes, cherries, and dates are among the fruits in this

category. These fruits typically have high levels of fructose and glucose. Dried fruits, in particular, contain more concentrated sugars due to their reduced water content. Despite their high sugar content, these fruits are also rich in fiber, vitamins, and minerals, making them a valuable part of a balanced diet.

The aim of this study is to determine and compare the sugar compositions of various honey and fruit types.

Materials and Methods

Samples

In this study, Arhavi chestnut honey and Muş region floral honey were used as floral honeys, while pine honey was selected as the honeydew honey. As for fruits, apricots from the Nakhchivan region and fragrant grapes from Trabzon were included.

Sugar analysis was performed using High-Performance Liquid Chromatography with a Refractive Index Detector (HPLC-RID). For this purpose, standard sugars, including ribose, fructose, glucose, sucrose, maltose, trehalose, melezitose, and melibiose, were used. The analysis was conducted on a reverse-phase NH₂ column (EC 200/4.6 Nucleosil 100-5 NH₂) under isocratic conditions (Bogdanov & Baumann, 1998). The mobile phase consisted of acetonitrile and ultrapure water, with a flow rate set at 1.5 mL/min.

One gram of sample (honey or homogenized fruit) was homogenized with 10 mL of 70% ethanol, followed by filtration. The filtration was evaporated to dryness, and the remaining residue was re-dissolved in pure water and then injected into the HPLC system. For sample preparation, 1 g of homogenized grape extract was mixed with 10 mL of 80% ethanol, filtered, and the filtrate evaporated to dryness. The residue was reconstituted in ultrapure water, filtered, and analyzed using the HPLC system. Separation was achieved on a reverse-phase -NH₂ column (EC 250/4 Nucleosil, Macherey-Nagel) with an isocratic mobile phase of 80% acetonitrile and 20% ultrapure water. The injection volume was 20 µL, with a mobile phase flow rate of 1.5 mL/min and a column temperature of 30°C.

Results and Discussions

The study conducted using High-Performance Liquid Chromatography with a Refractive Index Detector (HPLC-RID) focused on analyzing the sugar content and sugar profiles of selected fruit and honey varieties by employing seven different sugar standards. The findings, summarized in Table 1, provide valuable insights into the variations in total sugar content and the presence of specific sugar components across different samples.

Among the honey varieties analyzed, chestnut honey exhibited the highest total sugar content, approximately 78%, followed by honey from the Muş region with 72% and pine honey with 55%. Notably, pine honey was found to contain melebiose and melezitose, while these sugars were below detectable limits in chestnut and flower honey. This finding aligns with previous research suggesting that honeydew-based honeys, such as pine honey, typically contain complex oligosaccharides that are absent or present in lower concentrations in nectar-based honeys like chestnut and flower honey. The relatively lower sugar content in pine honey compared to other varieties may be attributed to its unique composition and source of production, which involves honeydew secretions rather than direct nectar collection.

The study also examined the sugar content of two different fruit samples. White grapes, sourced from the Manisa region, were found to contain approximately 9% total sugar, whereas apricots belonging to the *Prunus armeniaca* L. genotype from the Nakhchivan region exhibited a significantly higher total sugar content of around 18%. The nearly twofold increase in sugar

content in apricots compared to grapes is particularly noteworthy, suggesting that apricots are a richer source of natural sugars and, consequently, provide higher caloric value. This finding is in line with nutritional data indicating that apricots, especially those cultivated in arid and semi-arid regions, tend to accumulate more sugars due to environmental factors such as temperature, sunlight exposure, and water availability.

Overall, the study highlights the variability in sugar composition among different honey and fruit varieties, emphasizing the influence of botanical and geographical origins on sugar profiles. The identification of specific sugars in pine honey, the relatively high sugar content in chestnut honey, and the pronounced difference in sugar levels between grapes and apricots contribute to a broader understanding of the nutritional and compositional characteristics of these natural products. Further research with a larger sample size and additional analytical techniques could provide deeper insight into the biochemical and environmental factors influencing sugar composition in honey and fruits.

Table 1. Sugar profiles of the honey and fruits samples

	Chestnut honey	Muş plateau honey	Pine honey (Muğla)	Grape (<i>Vitis vinifera</i>) Manisa	Apricot (<i>Prunus armeniaca</i> L) Nakhchivan
Fructose%	40,04	36.20	28.30	5,45	0.72
Glucose%	28.20	32.65	25.00	4.20	5.60
Sucrose%	nd	1.20	0.50	0.35	12.10
Maltose%	nd	0.80	nd	nd	nd
Trehalose%	nd	nd	nd	nd	0.04
Melibiose%	nd	nd	1.20	nd	0.95
Melezitose%	nd	nd	0.50	nd	nd
Total sugar	%78	%72	%55	%9	%18

Fructose, often known as fruit sugar, is one of the main naturally occurring sugars in most fruits. However, unlike fruits such as grapes (Eyduran et al. 2015), figs (Slatnar et al. 2011), dates (Assirey, 2015), and apples (Zupan et al. 2016), apricots generally do not contain fructose in higher concentrations than sucrose and glucose (Su et al. 2020). This distinct sugar composition sets apricots apart from many other fruit varieties. Our study aligns with previous research indicating that the sugar profiles of pine honey, chestnut honey, and flower honey are composed of similar types of sugars (Uçurum et al. 2023; Can et al. 2015).

References

- Basson, C. E., Groenewald, J. H., Kossmann, J., Cronjé, C., & Bauer, R. (2010). Sugar and acid-related quality attributes and enzyme activities in strawberry fruits: Invertase is the main sucrose hydrolysing enzyme. *Food Chemistry*, *121*(4), 1156-1162.
- Beckles, D. M. (2012). Factors affecting the postharvest soluble solids and sugar content of tomato (*Solanum lycopersicum* L.) fruit. *Postharvest biology and technology*, *63*(1), 129-140.
- Can, Z., Yildiz, O., Sahin, H., Turumtay, E. A., Silici, S., & Kolayli, S. (2015). An investigation of Turkish honeys: Their physico-chemical properties, antioxidant capacities and phenolic profiles. *Food chemistry*, *180*, 133-141.
- Kolayli, S., Boukraâ, L., Şahin, H., & Abdellah, F. (2012). Sugars in honey.

Kolaylı, S., Asadov, E., Huseynova, A., Rahimova, S., & Kara, Y. (2024). Phenolic composition and antioxidant properties of black mulberry (*Morus nigra* L.) fruits and leaves. *Journal of Wildlife and Biodiversity*, 8(2), 355-364.

Raatz, S. K., Johnson, L. K., & Picklo, M. J. (2015). Consumption of honey, sucrose, and high-fructose corn syrup produces similar metabolic effects in glucose-tolerant and-intolerant individuals. *The Journal of nutrition*, 145(10), 2265-2272.

Uçurum, H. Ö., Tepe, Ş., Yeşil, E., Güney, F., Karakuş, S., Kolaylı, S., ... & Çakici, N. (2023). Characterization of Turkish pine honey according to their geographical origin based on physicochemical parameters and chemometrics. *European Food Research and Technology*, 249(5), 1317-1327.

Wu, L., Du, B., Vander Heyden, Y., Chen, L., Zhao, L., Wang, M., & Xue, X. (2017). Recent advancements in detecting sugar-based adulterants in honey—A challenge. *TrAC Trends in Analytical Chemistry*, 86, 25-38.

SEED COATING TECHNOLOGIES FROM A SUSTAINABLE AGRICULTURE PERSPECTIVE

Emine ATALAY

Selcuk University, Faculty of Agriculture, Department of Field Crops, Konya, Türkiye

ORCID ID: <https://orcid.org/0000-0002-1911-4951>

ABSTRACT

Seeds are a key material in plant production, enabling the uninterrupted production of various products and raw materials over large areas. They have also been a primary food source due to their rich nutritional content. These characteristics make seeds an indispensable production element of strategic importance, ensuring the continuity of agriculture since the beginning of sedentary life.

The seed embryo develops into a new plant during germination. Germination and seedling development are critical for plant production. Healthy, pathogen-free seeds, high germination, synchronized emergence and rapid growth, and stress-resistant seedlings are considered key factors that can increase yield and quality by at least 25%. These methods are a priority in agriculture because of their role in production. Seed coating methods are among the most prominent applications.

Seed coating processes (three main forms: pelleting, film coating or a combo of these) involve adding physical, chemical or biological components to the existing seed coat via specific processes to improve germination, emergence and facilitate the sowing of small, irregular seeds and the loading of pesticides, plant nutrients, micro-organisms, growth regulators and seed protection agents.

Seed coating improves agricultural efficiency by using inputs more efficiently, reducing waste and seed loss. It also enables the controlled application of chemicals and the introduction of beneficial microbes into the soil, making it environmentally friendly. This makes seed coating a key technology in sustainable agriculture.

This paper will present general information about seed coating applications and their potential use within the concept of sustainable agriculture.

Key words: Film coating, Seed pelleting, Sustainable agriculture, Germination

INTRODUCTION

All living organisms are part of an ecosystem in which they interact with each other and with non-living entities. It is impossible for any living being, including humans, to exist without the environment in which they live (Yeni, 2014). One of the most significant points revealed by archaeological evidence is the fact that, throughout every stage of history, humans have obtained vital necessities such as food, clothing, shelter, and energy from the natural environments in which they lived and have been in a constant search for methods to access these needs more easily and sufficiently. For this reason, humans are living beings that are in continuous interaction with nature, influencing and shaping it while also being shaped by it.

Agriculture is a multidisciplinary field of science that addresses the production of plant and animal products, the improvement of these products in terms of yield and quality, the preservation of the obtained products without losing their quality, their processing and

utilization, and their marketing (Kökten & İnci, 2021). As can be understood from this definition, agriculture is not limited to cultivation activities alone; it operates in integration with many fields and, therefore, shapes numerous elements related to life, including the environment. The concept of sustainable agriculture refers to an agricultural structure in which all agricultural activities and the techniques and technologies used in these activities are conducted while conserving and enhancing natural resources, ensuring long-term benefits, and adopting a holistic approach to agricultural practices. Although various definitions of sustainable agriculture exist, it is generally described as a form of agricultural production that meets people's nutritional needs while preserving the natural resources necessary for both present and future generations (Eryılmaz Aydın & Kılıç, 2018).

In sustainable agriculture, activities aimed at balancing the agronomic, environmental, social, and economic aspects of agricultural production are fundamental (Camcı Çetin, 2022). The use of self-sustaining agricultural systems that maintain soil fertility, protect groundwater, develop renewable energy sources, and adapt production systems to climate change, as well as planning agricultural activities in a way that contributes to the sustainability of social communities, can be considered the two main approaches of sustainable agriculture (Paravar et al. 2023). The goal of agricultural activities is to preserve soil fertility while minimizing environmental damage, maintaining economic vitality in both the short and long term, and improving the welfare of the rural (Camcı Çetin, 2022).

In the last century, although the use of fertilizers, irrigation activities, and chemicals for pest and disease control in plant production has led to significant increases in yield, it has been determined that the unconscious implementation of these practices has seriously and negatively affected the ecological balance of the Earth in the long term (Haspolat, 2012). Due to this intense pressure, environmental problems arising from the destruction of natural resources have also led to major social and economic issues, such as dependency on foreign agricultural products, the disappearance of rural community characteristics due to migration to cities, and an increasing imbalance between rural and urban populations. As the disruption of the balance between humans and nature has exacerbated these problems, their global scale has made them a significant concern and a topic of discussion in all countries. (Eryılmaz Aydın & Kılıç, 2018). Today, just as it is necessary to take precautions against all other anthropogenic threats and plan for a sustainable life, the necessity of utilizing innovations that enhance efficiency and food security in agriculture has also been recognized. It is now widely accepted that such major problems cannot be overcome with traditional/conventional agricultural methods alone. Among the foremost efforts to reduce the environmental pressure arising from agricultural production is the reduction of input requirements in production.

The primary goal of effective alternative approaches needed for low-input production is to support the production of nutritious and healthy food while reducing the use of synthetic inputs, energy, land, water, and waste generation (Ma, 2019; Demirel, 2020). Any agricultural practice that serves this purpose has gained strategic importance.

The seed, which is the initial material of plant production, is defined as a generative structure with the potential to develop into a new plant, carrying the embryo formed as a result of fertilization, the nutrient-storing endosperm, and the seed coat surrounding them (Aras, 2019). The seed is not only a crucial input in plant production but also a strategic material that directly affects the efficiency of other inputs and the quality of the final product. It is estimated that a high-quality seed can increase crop yield by up to 30% (Afzal et al., 2020).

Due to its significance in plant production, adapting technical and technological advancements to seeds has gained importance, as it contributes to achieving the desired progress in agriculture (Anonymous, 2017). Seed treatments, which encompass various methods and processes, are considered the most effective practices when they are target-oriented and crop-specific, ensuring optimal plant cover formation and yield improvement under changing climatic

conditions (Afzal et al., 2020). Among these practices, seed coating techniques are increasingly popular and are actively used in the seed industry. Properly applied seed coating processes with suitable materials are also recognized as supportive technologies in achieving sustainable agriculture goals.

SEED COATING APPLICATIONS

Definition and purpose of seed coating

The observable characteristics of an organism, namely its phenotype, emerge as a result of the interaction between its genetic traits and the environment. Therefore, it can be easily stated that the primary goal of the techniques used in plant production is to maximize the genetic potential of the seed. Research has determined that the contribution of a quality seed to yield increase is at least 25%, and it is possible to increase this rate even further. Various pre-sowing applications are available to improve seed quality, and seed coating applications are among these methods (Gökçöl & Duman, 2018). These applications have found widespread use to ensure plant production and product efficiency under today's climate and environmental conditions (Altıkat, 2024).

Seed coating is a general term used for processes that involve adding substances to seeds for various purposes, such as imparting different qualities to the seed or protecting it. Seed coatings are considered carrier systems that allow the direct application or loading of desired substances onto seeds (Taylor & Harman, 1990; Kavak, 2024). Through these processes, substances in the appropriate amount and for the intended purpose can be used effectively and uniformly across all treated seeds (Çetin, 2019). Successful results are also achieved by integrating other post-harvest seed applications, such as priming, which maintains seed viability and quality or enhances seed quality, into coating technology (Hacıyusufoglu & Guler, 2015).

In seed coating, various materials with different properties are applied to the seed surface to facilitate sowing by altering the size, weight, or surface of the seeds and/or to enhance seed quality, as well as to impart components that provide functions such as protection against biotic and abiotic stress (<https://www.incotec.com/en-gb/seed-technologies/seed-coating>).

By coating the surface of seeds with a functional layer of varying thickness, the germination rate, speed, and vigor can be enhanced. Since tolerance to biotic and abiotic stress can be increased, seedling losses in the early stages can be reduced. The bioactive, organic, or microbial elements in the coating can effectively enrich the seedbed. Through micro-level applications, production costs can be reduced (Altıkat, 2024). The seed can be made larger, heavier, or more uniform, enabling homogeneous sowing. In short, seed applications are a technological approach that supports plant development, helps ensure production, and allows the genetic and physiological potential of plants to be realized (Bicakcı ve ark., 2024). Seed coating applications:

- a. Facilitating sowing operations by modifying the morphological structure of small, light, or amorphous seeds (Sharma et al., 2015),
- b. Application of protectants such as fungicides and insecticides (Taylor et al., 2001),
- c. Loading of plant nutrients (especially micronutrients) (Farooq et al., 2012),
- d. Loading of growth regulators, biostimulants, bioinoculants, and growth-promoting bacteria (Madsen et al., 2014; Ma et al., 2019; Paravar et al., 2023),
- e. Applying water retainers to the seed coat to provide tolerance to low water stress and non-water-retentive materials to enhance tolerance to high water stress (Schneider & Renault, 1997),
- f. Using colorants and markers to track varieties with different genetic traits, different applications, or different seed lots (Pedrini et al., 2017),

g. Exxtending the storage life of seeds by restricting moisture and gas exchange with special polymers (Kavak & Eser, 2006) (Kavak, 2024).

Historical Development of Coating

Although the exact origins of seed applications are not clearly known, it is assumed that they began through observations or by coincidence. It is known that seed applications date back to ancient times, as evidenced by the remnants of ancient civilizations. Historical records from around 470 BC indicate that olive oil residues, when sprinkled on plants, helped prevent diseases. Approximately 2,500 years after these findings, it was discovered that vegetable oils possess antifungal properties (Andrews, 1961). Around 30 BC, it was recorded in Ancient Egypt that seeds treated with onion juice were protected from diseases. The use of wine and crushed cypress leaves to protect seeds from storage insects dates back to 60 AD (Hazra & Patanjali, 2016). It has been stated that seeds treated with extracts obtained from cypress trees were also protected from birds and rodents due to the bitter taste of the coating (<https://summitseedcoatings.com/blog/history-of-seed-coatings-theyve-been-around-longer-than-you-d-think/>). In Ancient Egypt, it is known that seeds were coated with clay mud and formed into balls to protect them from weather conditions and living organisms. This process, developed to adapt to the variable conditions of the Nile River, later became widespread and spread to different countries, with other materials besides clay being added to the coating (<https://www.trtworld.com/life/the-simple-seedballs-giving-kenya-s-forests-a-helping-hand-44594>). Another record mentions that in the early 17th century, it was discovered that wheat seeds recovered from a shipwreck that had remained in salty seawater for a while were healthier against fungi compared to other seeds. (<https://www.croplife.com/special-reports/seed-care-evolves-through-the-years/>). It has been determined that fungal spores germinate less on seeds treated with compounds such as sulfur and copper sulfate (Andrews, 1961).

Over time, with the increasing interest in seed applications and the development of new chemicals, both new methods and new materials were developed and used in seed applications. While the first patent for seed coating applications was obtained in 1868, the first scientific studies related to techniques still used today began in the 1940s. The initial aim of these studies was to make seeds more suitable for mechanized agriculture. The first steps toward the commercial development of seed coating were taken in the 1960s, and this technology quickly became widespread. During these years, the invention of systemic fungicides and their addition to seed coatings were reported to reduce early seedling losses and increase yield. (<https://summitseedcoatings.com/blog/history-of-seed-coatings-theyve-been-around-longer-than-you-d-think/>). Approximately 15 years later, in the early 1970s, it was recorded that 43% of vegetable growers in the USA and Europe began applying precision agriculture techniques along with coated seeds (Kayaalp, 2014, Çetin, 2019).

It is known that seed applications, which began modestly with organic materials, quickly gained traction and evolved into a giant industry by incorporating synthetic inputs through the transfer of technologies to agriculture. Along with many products used in seed coating, such as colorants, polymers, fillers, and other additives, the market for biological seed applications, particularly those containing biological products (biofertilizers, biopesticides, bioinoculants, biostimulants), is reported to be gaining more value globally with each passing year (Afzal ve ark., 2020).

The global seed treatment market was reported to be valued at \$15.11 billion in 2024. It is expected that the market will reach approximately \$17 billion in 2025 and \$39 billion by 2032. The same report also highlights that coating technologies, particularly those suitable for sustainable agriculture, are on an upward trend (<https://www.fortunebusinessinsights.com/industry-reports/seed-treatment-market-100156>). Another report stated that the agricultural biological products market, consisting of products

derived from plant extracts, microorganisms, and other organic materials, is rapidly gaining value and that its market share is gradually increasing (<https://www.marketsandmarkets.com/Market-Reports/agricultural-biological-market-100393324.html>).

General Properties of Materials Used in Coating

A wide variety of materials with different functions, compositions, or origins can be used for seed coating purposes. Seed coating materials are characterized by their functions as active ingredients, liquids, or solid particles. Additionally, these materials are categorized by their origin into synthetic chemicals, natural products or derivatives of natural products, biological agents, and minerals (Afzal et al., 2020).

In general, according to their intended use and function: **Polymers** are the most widely used coating material for various purposes, such as adhering applied substances to the seed surface, preventing active ingredients from contacting the environment, and protecting the seed from external factors (e.g., moisture, dust, pathogens) through the layer they form. **Plant nutrients** are preferred for balancing soil pH, improving soil structure, and supporting germination and early seedling development. **Colorants** serve purposes such as distinguishing seeds from others, enhancing visibility during sowing, and deterring animals due to their vibrant colors. **Binders** are used for similar purposes as polymers and to maintain the viability of microbial agents. **Microbial agents** include symbiotic microorganisms that support plant growth, help overcome stress factors, and facilitate water and nutrient uptake (<https://summitseedcoatings.com/blog/seed-preparation/what-goes-into-seed-coatings-seed-coating-materials-and-what-they-do/>). **Protective chemicals**, including pesticide- and fungicide-effective coatings, prevent harmful microorganisms from damaging seeds (Çakmakçı et al., 2018; Kavak, 2024).

Some materials within these categories are suitable for organic use and labeling (Afzal et al., 2020). Due to the importance given to sustainable agriculture, both commercial and scientific research trends are inclined toward the use of biological materials. Therefore, factors such as the toxicity level of the coating, particle size, distribution on the surface, porosity (i.e., permeability), and water-holding capacity are important criteria in determining the coating material (Javed et al., 2022). Since healthy plant development is crucial, utmost attention is paid to ensuring that coating materials are compatible with biological systems (Dadlani, 2025). Water-holding capacity is considered a criterion that affects the better growth of seeds under arid conditions (El Bouchtaoui & Ablouh, 2025). Another important criterion is the biodegradability of the coating (Makvandi et al., 2025).

It has been found that some natural compounds present in plants, due to their antimicrobial effects, are effective in controlling seed-borne bacterial and fungal diseases. These active components can be used alone or in combination with physical treatments as an alternative to chemical disinfectants for seed disinfection. Fixed and volatile oils, as well as other antimicrobial components with antifungal and antibacterial properties, have been identified in various plants, particularly in compounds derived from tea tree, clove, mint, rosemary, bay leaf, black cumin, lavender, sage, and thyme (Kavak, 2024).

The use of beneficial microorganisms that have a symbiotic relationship with plants (bioinoculants) is another noteworthy component in seed applications. *Rhizobium* and *mycorrhiza* are the best known symbiotic microorganisms in this context (Taylor & Harman, 1990). Symbiotic microorganisms have positive effects such as nitrogen fixation, mineralization of organic phosphorus in the soil, facilitating more effective access of plant roots to water and minerals, enhancing plant resistance to biotic and abiotic stress, and increasing plant tolerance to pathogens (Korkutal & Bahar, 2024; Kavak, 2024). *Rhizobium* species have

been used for many years in seed applications of legume species specifically for nitrogen fixation (Taylor & Harman, 1990).

Biostimulants are another important component in coating and seed applications. It is known that the biostimulant effects of symbiotic microorganisms promote plant growth and development. Non-microbial biostimulants such as chitosan, humic acid, and seaweed extracts are also widely used (Korkutal & Bahar, 2024).

Seed Coating Techniques

Different coating techniques are applied based on the size, shape, weight, and intended use of the seeds (Pedrini et al., 2017). There are three main techniques that are most commonly used in seed coating. **Film Coating** refers to the application of colorants to the seed surface in a thin film along with adhesive materials. **Pelleting** can be defined as the application of a coating material containing solid particles to the seed surface in a way that changes the shape of the seed. **Encrusting** involves applying a thin layer that partially alters the shape of the seed (Kavak, 2024).



Figure 1. Seed coating (Javed et al., 2022)

Pelleting is a method in which the size and weight of seeds are increased. It is particularly applied to small seeds (Hacıyusufoğlu et al., 2015). Typically, lime, clay, or polymers are used as the main coating material (Hazra & Patanjali, 2016). Although the primary purpose is the morphological alteration of seeds, any substance needed for seed germination, seedling development, or the plant's later growth stages can be added to the coating material (Kavak, 2024). The pelleting process provides an advantage, especially in mechanized sowing systems, as it reduces sowing errors (Javed et al., 2022).

Film coating is the process of applying a thin polymer layer to the seed surface without affecting the seed's morphological structure (Hacıyusufoğlu et al., 2015). Through film coating, components such as plant nutrients, bioinoculant elements, or pesticides can be evenly distributed on the seed surface (Pedrini et al., 2017). Additionally, this technique allows for the use of very low doses of the applied substances (Ma, 2019).

In the encrusting technique, the seed surface is covered with a thicker layer compared to film coating, resulting in a partial increase in seed weight (Afzal et al., 2020). This technique helps achieve not only a partial weight increase but also a smoother surface and a more uniform shape of the seed (Ma, 2019). In the encrusting process, the seed is first pelleted, and then the pelleted seed is coated with a film. This method also prevents the loss of materials used in pelleting, especially active ingredients (Kayaalp, 2014).

SEED COATING AND SUSTAINABLE AGRICULTURE

Seed coating technologies, which predominantly use conventional inputs, have found significant application areas in sustainable agriculture in recent years with the introduction of biological materials into the sector. Through coating techniques and seed applications used in conjunction with coating, advantages can be achieved in many aspects critical to sustainable agriculture.

Reducing Seed Usage

The seed is one of the largest inputs in plant production. Therefore, a reduction in seed usage contributes to one of the goals of sustainable agriculture, which is reducing input usage. Additionally, reducing the high cost of seeds is a significant factor for achieving economical production. Research has indicated that seed coating technology can alter the physical and physiological properties of seeds. These changes improve seed quality, increase germination and emergence rates, enhance tolerance to biotic and abiotic stress, and reduce germination losses (Hacıyusufoğlu et al., 2015; Sharma et al., 2015; Pedrini et al., 2017; Afzal et al., 2020; Javed et al., 2022). Consequently, it has been stated that the desired production can be achieved with less seed usage during sowing (Javed & Afzal, 2018).

Reduction of Mechanization or Facilitation of Sowing Processes

Seed coating helps seeds better adapt to mechanized sowing processes by altering their physical properties. Mechanical equipment used in sowing may face difficulties with seeds that have surface features such as hair-like structures or that are small and irregularly shaped (Hacıyusufoğlu & Güler, 2015). Coated seeds are more uniformly shaped, aiding sowing machines in planting without errors and ensuring the homogeneous distribution of seeds. It has also been stated that coating contributes to maintaining the sowing settings of the equipment used in planting (Javed et al., 2022). The compatibility of coated seeds with different machines also contributes to reducing mechanization. It has been determined that coated seeds can be sown with many different types of sowing machines (Bicakci et al., 2020). Sowing machines with adjustable row spacing enable sowing at accurate distances and depths with minimal seed loss (Panda & Mondal, 2020). Since coated seeds can be sown precisely, the sowing process becomes more efficient, and pelleted seeds are more easily handled by mechanical sowing equipment (Pedrini et al., 2017). Due to their structural properties, seeds that were previously sown by hand or broadcasting can now be sown using a seed drill without the need for special equipment (Hacıyusufoğlu et al., 2015). Thus, seed coating allows for reduced mechanization, error-free sowing, and the use of fewer seeds during planting..

Reduction of Chemical Usage

It is known that the seed coating process affects the microenvironment of each seed (Javed et al., 2022). It has been emphasized that seed coating technology can significantly reduce the use of chemicals in disease and pest management and that effective disease and pest control can be achieved by applying active ingredients directly to the seed in smaller amounts (Panda & Mondal, 2020). At the same time, seed applications accelerate seed germination, strengthen seedling development, and increase tolerance to biotic and abiotic stress, leading to more resilient plants. This reduces the need for chemical protectants such as pesticides and fungicides (Rocha et al., 2019; Panda & Mondal, 2020).

It has been stated that using biological agents in seed coating can reduce the need for chemical protectants, enhance the plant's metabolism, improve the uptake and availability of nutrients, and thereby reduce fertilizer use (Rocha et al., 2019).

Soil and Water Conservation

Organic materials used in seed coating act as soil conditioners by improving soil structure and enhancing water-holding capacity, especially in arid regions. Additionally, they reduce environmental impacts by retaining protective agents or plant nutrients around the seed. By reducing leaching from the soil, they contribute to the efficient use of fertilizers and other inputs. It is also stated that these coatings can absorb certain harmful compounds present in the environment (Altıkat, 2024).

It is stated that bioinoculants or beneficial microorganisms used in seed coating have a soil-structuring effect, thereby providing benefits in improving the utilization of nutrient elements (Chirag & Amir, 2023).

Pest and Disease Management

Today, it is widely accepted that biopesticides have the potential to reduce dependence on chemical seed applications (Afzal et al., 2020; Bicakci et al., 2020). Biopesticides generally target specific pests, reduce the likelihood of resistance, and have minimal impact on non-target organisms and the environment (<https://vlsci.com/blog/seed-treatment-guide/>).

In sustainable agriculture, seed applications containing plant extracts, essential oils, and beneficial microorganisms are considered an alternative method for protection against diseases and pests. It has been stated that plant extracts, essential oils, and animal-derived natural components such as propolis are used to prevent or reduce many fungal diseases in seeds (Kavak, 2024).

Microbial seed coating is another application that provides protection against diseases and pathogens (Afzal et al., 2020). Inoculation with antagonist organisms is used in controlling plant seed pathogens (Bonome et al., 2020).

Better Physical and Physiological Development

It has been stated by many researchers that bioinoculants and symbiotic microorganisms can enhance plant yield, improve nutrition, and increase tolerance to biotic and abiotic stresses. These components support seedling development, increase tolerance to biotic and abiotic stress, and reduce germination losses (Sharma et al., 2015; Pedrini et al., 2017). Through coating processes, inoculation can be performed with fewer resources and greater effectiveness (Rocha et al., 2019). This method allows the application of beneficial microorganisms without significantly disturbing the microfauna.

Biostimulants can be defined as natural compounds that stimulate or regulate physiological processes in plants. When used in small amounts, these compounds can alter plant metabolism, improving crop yield and quality. Applying biostimulants through seed coating technology offers significant potential for production improvement compared to direct soil application or foliar application at later growth stages (Afzal et al., 2020).

Increasing Tolerance to Abiotic Stress

Ecological factors significantly affect plant development and yield by influencing plant metabolism. While favorable environmental conditions promote plant growth, adverse conditions create stress in plants, negatively impacting their development (Ceritoğlu et al., 2021). Major abiotic stress factors such as drought, salinity, and extreme temperatures (high or low) lead to yield losses and reduced quality in agricultural production (Bicakci et al., 2020).

It is stated that seed coating is an approach that enables adaptation to different stress conditions and improves crop performance under such conditions (Javed et al., 2022). Through coating, active ingredients with protective and supportive functions are imparted to the seed during the most sensitive stages of germination and early seedling development, providing a significant advantage in plant growth (Bicakci et al., 2020).

In addition to physical and biochemical substances, the use of appropriate biological agents is considered an application with the potential for sustainability (Sharma et al., 2015). Biological applications show diversity in use depending on the product and purpose (Javed et al., 2022). Biological seed applications have the potential to mitigate the harmful effects of temporary abiotic stress. They help plants enhance their resistance to drought and salinity (Afzal et al., 2020). Another study demonstrated that when beneficial root bacteria were introduced to the rhizosphere region through coating, they promoted growth and development in both favorable and unfavorable environmental conditions (Javed et al., 2022).

Supplementation for Plant Nutrients

It has been determined that addressing the deficiency of micronutrients in the soil through coating can increase yield by 44%. The same researchers stated that nutrient supplements provided through seed applications are more economical and effective compared to traditional methods (Farooq et al., 2012).

It has been stated that treating or coating seeds with a solution containing micronutrients before sowing is a simple and practical way to contribute to better seedling growth. Seeds enriched in this way can improve seed germination, seedling vigor, crop growth, and stress tolerance, especially in soils deficient in specific nutrients (Muhammed & Pekşen, 2020).

CONCLUSION

According to estimates by many international organizations, including the United Nations, arable land is expected to decrease to nearly half of today's values, reaching as low as 1.5 decares within the next 25 years due to the impact of various factors. Additionally, it is predicted that agricultural problems arising from the effects of climate change will lead to an approximately 25% decrease in agricultural productivity. Balancing these losses through sustainable soil and water management, sustainable agricultural practices, creating an informed society, and implementing smart environmental policies has become a necessity (Atalay, 2024).

Sustainable agriculture is the establishment of an agricultural structure that not only conserves natural resources in the long term but also utilizes agricultural technologies that do not harm the environment (Turhan, 2005). Effective alternative approaches are highly important in sustainable agriculture. Any agricultural practice that serves this purpose has gained strategic importance. Seed applications are among these practices.

The initiation of seed applications, especially using biologically-based coating materials, provides sustainable solutions to the increasing demands of global agriculture. The success of seed coating technology in this context depends on the development and use of low-cost, readily available, environmentally friendly, and biologically compatible coating materials that can serve different purposes as alternatives to chemical inputs.

Seed coating technologies may sometimes face challenges in aligning with existing systems or regulations related to sustainable agriculture. Therefore, the environmental impact, economic feasibility, and system compatibility of these technologies should be carefully evaluated from multiple perspectives.

Seed coating plays a key role in transforming agricultural production into an ecosystem-friendly model. However, it is necessary to transfer technology at accessible costs for small farmers and to develop coating formulations compatible with local conditions.

Additional research and development are needed to develop biological materials that can be used in sustainable agriculture systems. There are several issues to be researched, such as the difficulty of optimizing biologically-derived materials, R&D costs, the long duration of the biodegradation process, and the effects on non-target organisms. This technical and technological transformation can be achieved through R&D studies conducted jointly by multiple scientific disciplines. Since academic studies often work with small seed lots and quantities, the support of industry representatives who operate with much larger seed batches and quantities on an industrial scale is also a crucial aspect. To promote the widespread adoption of this system and replace conventional products, education programs, support, and incentives are needed.

REFERENCES

- Afzal, I., Javed, T., Amirkhani, M., & Taylor, A. G. (2020). Modern seed technology: Seed coating delivery systems for enhancing seed and crop performance. *Agriculture*, 10(11), 526. <https://doi.org/10.3390/agriculture10110526>
- Altıkat, S. (2024). Biyokömür ve elektrostatik kaplama yönteminin tohum performansına etkisi. *Journal of Agriculture*, 7(2), 302-309. <https://doi.org/10.46876/ja.1600751>
- Andrews, C. H. (1961). A history of seed treatment. *Proceedings of the Short Course for Seedsmen*, 51. <https://scholarsjunction.msstate.edu/seedsmen-short-course/51>
- Anonymous (2017). TÜBİTAK Türkiye Sanayi Sevk ve İdare Enstitüsü, Tohum Dağıtıcıları Alt Birliği, Tohumluk Dağıtım Sektörü Ulusal Strateji Raporu. <https://todab.org.tr/tr/sayfalar?altsayfa=23>
- Aras, B. (2019). Türkiye’de tohumculuk sektörünün mevcut durumu, sorunlar ve öneriler. *Journal of the Institute of Science and Technology*, 9(3), 1763-1775. <https://doi.org/10.21597/jist.531701>
- Atalay, E. (2024). Effective approaches for sustainable agriculture: Land degradation neutrality. *3rd International Conference on Sustainable Ecological Agriculture*. pp 147-169.
- Bicakci, T., Aksu, E., & Arslan, M. (2020). Determination of germination characteristics of covered alfalfa (*Medicago sativa* L.) seeds in drought stress conditions. *Journal of Tekirdag Agriculture Faculty*, 17(2), 124-136. <https://doi.org/10.33462/jotaf.520836>
- Bonome, L. T. S., Bittencourt, H. H., Moura, G. S., Franzener, G., & de Carvalho, J. H. (2020). Natural products for alternative seed treatment. In: Tiwari, A.K. (eds) *Advances in Seed Production and Management* (pp. 399-418). Springer. https://doi.org/10.1007/978-981-15-4198-8_18
- El Bouchtaoui, F. Z., Ablouh, E. H., Mouhib, S., Kassem, I., Kadmiri, I., Hanani, Z., & El Achaby, M. (2025). Hydrophobic Nanostructured Coatings of Colloidal Lignin Particles Reduce Nutrient Leaching and Enhance Wheat Agronomic Performance and Nutritional Quality. *ACS Applied Materials & Interfaces*. 17(8), 12578-12596. <https://doi.org/10.1021/acsami.4c19243>
- Camcı Çetin, S. (2022). Yeşil devrim: Tamam mı? Devam mı? *Uluslararası Turizm, Ekonomi ve İşletme Bilimleri Dergisi*, 6(2), 109-115.

- Ceritoğlu, M., Erman, M., Çığ, F., Şahin, S., & Acar, A. (2021). Bitki gelişimi ve stres toleransının geliştirilmesi üzerine sürdürülebilir bir strateji: Priming tekniği. *Türkiye Tarımsal Araştırmalar Dergisi*, 8(3), 374-389. <https://doi.org/10.19159/tutad.883564>
- Çakmakçı, S., Yıldırım, G. H., & Yıldırım, C. (2018). Tohum kaplama yöntemleri ve kaplamada kullanılan materyaller. *Tarımın Sesi*, 41, 64-67.
- Çetin, D. (2019). *Tohum kaplama teknolojisinin bazı küçük tohumlu bitkilerde kullanımının ekonomik yönden değerlendirilmesi* (Yüksek lisans tezi). Kahramanmaraş Sütçü İmam Üniversitesi. Fen Bilimleri Enstitüsü. Kahramanmaraş.
- Chirag, A. P., & Mohammad, A. (2023). Seed coating technology: A sustainable approach for improving seed quality and crop performance. *Pharma Innovation*, 12(10), 970-978.
- Dadlani, M. (2025). Seed enhancement technologies. In *Indian Seed Sector: Evolution, Technology, Trade and Sustainability* (pp. 129-150). Springer. https://doi.org/10.1007/978-981-96-0714-3_8
- Demirel, F. (2020). Bitki ve hayvan biyoteknolojisi; hücreli tarım ve nano-teknoloji. *Journal of Agriculture*, 3(2), 1-9. <https://doi.org/10.46876/ja.822503>
- Eryılmaz Aydın, G., & Kılıç, O. (2018). Türkiye’de sürdürülebilir tarım ve iyi tarım uygulamaları. *KSÜ Tarım ve Doğa Dergisi*, 21(4), 624-631. <https://doi.org/10.18016/ksudobil.345137>
- Farooq, M., Wahid, A., & Siddique, K. H. (2012). Micronutrient application through seed treatments: A review. *Journal of Soil Science and Plant Nutrition*, 12(1), 125-142. <http://dx.doi.org/10.4067/S0718-95162012000100011>
- Gökçöl, A., & Duman, İ. (2018). Tohum kaplama teknolojileri. *TÜRKTÖB, Türkiye Tohumcular Birliği Dergisi*, 7(26), 23-25.
- Hacıyusufoğlu, A. F., Akbaş, T., & Şimşek, E. (2015). Bazı küçük çaplı tohumlara peletle tohum kaplama yönteminin uygulanması. *Tarım makinaları bilimi dergisi*, 11(3), 257-263.
- Hacıyusufoğlu, A. F., & Guler, E. (2015). Innovative developments in the seed coating systems. *International Journal of Scientific and Technological Research*, www.iiste.org ISSN, 2422-8702.
- Haspolat, I. (2012). Genetiği değiştirilmiş organizmalar ve biyogüvenlik. *Ankara Üniversitesi Veteriner Fakültesi Dergisi*, 59, 75-80.
- Hazra, D. K., & Patanjali, P. K. (2016). Seed coating formulation technologies: An environmental biology friendly approach for sustainable agriculture. *Bioscience Methods*, 7(5): 1-10. <https://doi.org/10.5376/bm.2016.07.0005>
- <https://summitseedcoatings.com/blog/history-of-seed-coatings-theyve-been-around-longer-than-you-d-think/> (access date: 21.02.2025)
- <https://summitseedcoatings.com/blog/seed-preparation/what-goes-into-seed-coatings-seed-coating-materials-and-what-they-do/> (access date: 21.02.2025).
- <https://vlsci.com/blog/seed-treatment-guide/> (access date: 21.02.2025)
- <https://www.croplife.com/special-reports/seed-care-evolves-through-the-years/> (access date: 21.02.2025).
- <https://www.fortunebusinessinsights.com/industry-reports/seed-treatment-market-100156> (access date: 21.02.2025)
- <https://www.incotec.com/en-gb/seed-technologies/seed-coating> (access date: 21.02.2025)

<https://www.marketsandmarkets.com/Market-Reports/agricultural-biological-market-100393324.html> (access date: 21.02.2025)

<https://www.trtworld.com/life/the-simple-seedballs-giving-kenya-s-forests-a-helping-hand-44594> (erişim tarihi: 21.02.2025)

Javed, T., & Afzal, I. (2018). Impact of seed pelleting on germination potential, seedling growth and storage of tomato seed. *Acta Horticulturae*, 1273, 417-424. <https://doi.org/10.17660/ActaHortic.2020.1273.54>

Javed, T., Afzal, I., Shabbir, R., Ikram, K., Zaheer, M. S., Faheem, M., Ali, H. H. & Iqbal, J. (2022). Seed coating technology: An innovative and sustainable approach for improving seed quality and crop performance. *Journal of the Saudi Society of Agricultural Sciences*, 21(8), 536-545. <https://doi.org/10.1016/j.jssas.2022.03.003>

Kaufman, G. (1991). Seed coating: A tool for stand establishment, a stimulus to seed quality. *HortTechnology*, 1(1), 98-102. <https://doi.org/10.21273/HORTTECH.1.1.98>

Kavak, S. (2024). Bölüm 9: Sebze tohumlarında kaliteyi iyileştirici güncel uygulamalar. In A. Balkaya & L. Arın (Eds.), *Sebze Tohum Üretimi ve Teknolojisinde Güncel ve Yenilikçi Uygulamalar* (pp. 129-150). İKSAD Yayınları. <https://doi.org/10.5281/zenodo.13822740>

Kavak, S., & Eser, B. (2006). *Farklı polimer kaplama materyal ve uygulamalarının soğan tohumlarında depo ömrü ve yaşlanma üzerine etkileri* (Doktora tezi). Ege Üniversitesi. Fen Bilimleri Enstitüsü. İzmir.

Kayaalp, Ö. (2014). *Prototip tohum peletleme sisteminin geliştirilmesi* (Yüksek lisans tezi). Süleyman Demirel Üniversitesi. Fen Bilimleri Enstitüsü. Isparta

Korkutal, İ., & Bahar, E. (2024). Mikorizalar ve bağcılıkta kullanımı. *ANADOLU Ege Tarımsal Araştırma Enstitüsü Dergisi*, 34(1), 97-107. <https://doi.org/10.18615/anadolu.1442118>

Kökten, K., & İnci, H. (2021). *Tarım uygulamalarında yenilikçi yaklaşımlar*. İKSAD Yayınları. ISBN: 978-625-8007-32-9.

Ma, Y. (2019). Seed coating with beneficial microorganisms for precision agriculture. *Biotechnology Advances*, 37(7), 107423. <https://doi.org/10.1016/j.biotechadv.2019.107423>

Madsen, M. D., Davies, K. W., Mummey, D. L., & Svejcar, T. J. (2014). Improving restoration of exotic annual grass-invaded rangelands through activated carbon seed enhancement technologies. *Rangeland Ecology & Management*, 67, 61-67. <https://doi.org/10.2111/REM-D-13-00050.1>

Mohammed, M., & Pekşen, E. (2020). Influence of Zn seed priming and coating on germination and seedling growth in wheat. *Anadolu Tarım Bilimleri Dergisi*, 35(2), 259-267. <https://doi.org/10.7161/omuanajas.745891>

Makvandi, M., Bakhshandeh, A., Moshatati, A., Moradi Telavat, M. R., & Khodaei Joghhan, A. (2025). The effect of nitrogen fertilizer and sugarcane residue compost on soil physicochemical properties and wheat grain yield under terminal heat stress in Ahwaz. *Plant Productions*. <https://doi.org/10.22055/ppd.2025.48005.2209>

Panda, D., & Mondal, S. (2020). Seed enhancement for sustainable agriculture: An overview of recent trends. *Plant Archives*, 20(1), 2320-2332. e-ISSN:2581-6063.

Paravar, A., Piri, R., Balouchi, H., & Ma, Y. (2023). Microbial seed coating: An attractive tool for sustainable agriculture. *Biotechnology Reports*, 37, e00781. <https://doi.org/10.1016/j.btre.2023.e00781>

Pedrini, S., Merritt, D. J., Stevens, J., & Dixon, K. (2017). Seed coating: Science or marketing spin? *Trends in Plant Science*, 22(2), 106-116. <https://doi.org/10.1016/j.tplants.2016.11.002>

- Rocha, I., Ma, Y., Souza-Alonso, P., Vosátka, M., Freitas, H., & Oliveira, R. S. (2019). Seed coating: a tool for delivering beneficial microbes to agricultural crops. *Frontiers in plant science*, 10, 1357. <https://doi.org/10.3389/fpls.2019.01357>
- Salem, Y., Sunoqrot, S., Hammad, A., Rajha, H. N., Alzaghari, L. F., Abusulieh, S., Maroun, R. G., & Louka, N. (2025). Oxidation-driven assembly of phenolic compounds from grape seeds waste into nanoparticles as potential anti-inflammatory and wound healing therapies. *ACS Applied Bio Materials*. <https://doi.org/10.1021/acsabm.4c01800>
- Schneider, A., & Renault, P. (1997). Effects of coating on seed imbibition: I. Model estimates of water transport coefficient. *Crop Science*, 37, 1841-1849. <https://doi.org/10.2135/cropsci1997.0011183X003700060030x>
- Sharma, K. K., Singh, U. S., Sharma, P., Kumar, A., & Sharma, L. (2015). Seed treatments for sustainable agriculture-A review. *Journal of Applied and Natural Science*, 7(1), 521-539. ISSN: 2231-5209.
- Taylor, A. G., & Harman, G. E. (1990). Concepts and technologies of selected seed treatments. *Annual Review of Phytopathology*, 28, 321-339. <https://doi.org/10.1146/annurev.py.28.090190.001541>
- Taylor, A. G., & Kwiatkowski, J. (2001). Polymer film coatings decrease water uptake and water vapour movement into seeds and reduce imbibitional chilling injury. *2001 BCPC Symposium Proceedings No. 76: Seed Treatment: Challenges and Opportunities*, 215-220.
- Turhan Ş 2005. Tarımda Sürdürülebilirlik ve Organik Tarım. *Tarım Ekonomisi Dergisi*, 11(1): 13-24.
- Yeni, O. (2014). Sürdürülebilirlik ve sürdürülebilir kalkınma: Bir yazın taraması. *Gazi Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 16(3), 181-208.

**MOLECULAR MARKER TECHNIQUES FOR THE DETECTION OF DNA
CYTOSINE METHYLATION****DNA SİTOZİN METİLASYONUNUN BELİRLENMESİNDE KULLANILAN
MOLEKÜLER MARKIR TEKNİKLERİ****Adnan AYDIN**Iğdır University, Faculty of Agriculture, Department of Agricultural Biotechnology,
Iğdır/TürkiyeORCID ID: <https://orcid.org/0000-0002-8284-3751>**ABSTRACT**

DNA cytosine methylation is a crucial epigenetic modification that regulates gene expression and genome stability. The detection of DNA methylation is essential for understanding various biological processes, including development, aging, and diseases such as cancer. Several marker techniques have been developed to accurately detect and analyze DNA cytosine methylation, each offering unique advantages and limitations. Bisulfite sequencing is considered the gold standard, enabling single-base resolution by converting unmethylated cytosines to uracils while leaving methylated cytosines intact. Other methods, such as methylation-specific PCR (MSP), utilize primers that are specific to either methylated or unmethylated DNA, offering a quicker and more targeted approach. High-throughput sequencing techniques, including methylated DNA immunoprecipitation sequencing (MeDIP-seq), allow genome-wide analysis of methylation patterns with high sensitivity. Additionally, techniques like methylation-sensitive restriction enzyme digestion, array-based methods, and quantitative PCR have emerged as cost-effective alternatives. Each technique varies in terms of sensitivity, resolution, throughput, and cost, making the selection of the appropriate method dependent on the research question, sample type, and available resources. This review highlights the diverse marker techniques used in DNA cytosine methylation detection, their respective strengths, and applications in epigenetic research.

Key Words: Epigenetic, epi-Markers, DNA cytosine methylation**ÖZET**

DNA sitozin metilasyonu, gen ekspresyonunu ve genom stabilitesini düzenleyen önemli bir epigenetik modifikasyondur. DNA metilasyonunun tespiti, gelişim, yaşlanma ve kanser gibi hastalıklar dahil olmak üzere çeşitli biyolojik süreçlerin anlaşılması için gereklidir. DNA sitozin metilasyonunu doğru bir şekilde tespit etmek ve analiz etmek için her biri benzersiz avantajlar ve sınırlamalar sunan çeşitli marker teknikleri geliştirilmiştir. Bisülfid dizileme altın standart olarak kabul edilir ve metillenmemiş sitozinleri urasillere dönüştürürken metillenmiş sitozinleri sağlam bırakarak tek baz çözünürlüğü sağlar. Metilasyona özgü PCR (MSP) gibi diğer yöntemler, metillenmiş veya metillenmemiş DNA'ya özgü primerler kullanarak daha hızlı ve daha hedefe yönelik bir yaklaşım sunar. Metillenmiş DNA immünopresipitasyon sekanslama (MeDIP-seq) dahil olmak üzere yüksek verimli sekanslama teknikleri, metilasyon modellerinin yüksek hassasiyetle genom çapında analiz edilmesine olanak tanır. Ayrıca, metilasyona duyarlı restriksiyon enzim sindirimi, dizi tabanlı yöntemler ve kantitatif PCR gibi teknikler uygun maliyetli alternatifler olarak ortaya çıkmıştır. Her teknik hassasiyet, çözünürlük, verim ve

maliyet açısından farklılık gösterir ve uygun yöntemin seçimini araştırma sorusuna, örnek türüne ve mevcut kaynaklara bağlı hale getirir. Bu çalışma, DNA sitozin metilasyon tespitinde kullanılan çeşitli marker tekniklerini, bunların güçlü yönlerini ve epigenetik araştırmalardaki uygulamalarını vurgulamaktadır.

Anahtar Kelimeler: Epigenetik, Epi-Markırlar, DNA sitozin metilasyonu

GİRİŞ

Epigenetik mekanizmalar, gen ekspresyonunun düzenlenmesinde, genom stabilitesinin korunmasında, hücrel farklılaşmada ve organizmaların çevresel uyum süreçlerinde kritik bir rol oynar (Allis & Jenuwein, 2016). DNA metilasyonu, histon modifikasyonları ve kodlamayan RNA'lar gibi epigenetik değişiklikler, genetik bilginin korunmasına rağmen, genlerin nasıl ve ne zaman ifade edileceğini belirleyen dinamik süreçlerdir (Bird, 2007). Bu değişiklikler, embriyonik gelişimden doku farklılaşmasına, hastalık oluşumundan çevresel yanıt mekanizmalarına kadar geniş bir biyolojik spektrumda işlev görür. Özellikle, DNA metilasyonu genellikle gen susturulmasıyla ilişkilendirilirken, histon modifikasyonları kromatin yapısının gevşemesine veya yoğunlaşmasına neden olarak gen ekspresyon seviyelerini değiştirebilir (Jaenisch & Bird, 2003).

Epigenetik regülasyonun önemi, sadece normal fizyolojik süreçlerde değil, aynı zamanda çeşitli patolojik durumların ortaya çıkmasında da kendini göstermektedir. Kanser, nörodejeneratif hastalıklar, otoimmün bozukluklar ve metabolik sendromlar gibi birçok hastalık, DNA metilasyonundaki düzensizliklerle ilişkilendirilmiştir (Schübeler, 2015). Epigenetik değişikliklerin çevresel faktörler, yaşam tarzı ve beslenme alışkanlıkları gibi dış etmenlere duyarlı olması, bu süreçlerin dinamik ve geri dönüşümlü olduğunu göstermektedir (Feil & Fraga, 2012). Bu nedenle, epigenetik değişimlerin tespit edilmesi ve analiz edilmesi, bitkilerde biyotik stres, abiyotik stres ve bitki ıslahı açısından büyük önem taşımaktadır (Sahu et al., 2013).

Son yıllarda, DNA metilasyonunun belirlenmesi için çeşitli moleküler belirteç teknikleri geliştirilmiştir. Bu yöntemler, genellikle bisülfid dönüşümüne dayalı teknikler, restriksiyon enzimleri kullanan yaklaşımlar, immünopresipitasyon bazlı yöntemler ve yeni nesil dizileme (NGS) teknolojileri olmak üzere dört ana kategoriye ayrılır (Xie et al., 2018). Bisülfid dönüşümüne dayalı yöntemler (örneğin, bisülfid dizileme, metilasyon-spesifik PCR) sitozin bazlarını ayırt ederek, genom çapında veya spesifik lokuslarda metilasyon analizi yapmaya olanak tanır (Clark et al., 2006). Restriksiyon enzimlerine dayalı yöntemler, metilasyona duyarlı enzimler kullanarak, DNA metilasyon seviyelerini dolaylı olarak ölçer ve özellikle büyük ölçekli taramalar için uygundur (Chung et al., 2021). İmmünopresipitasyon tabanlı teknikler (örneğin, metillenmiş DNA immünopresipitasyonu [MeDIP], kromatin immünopresipitasyonu [ChIP]) belirli metilasyon ve histon modifikasyonlarını hedefleyerek, spesifik DNA bölgelerindeki epigenetik değişimleri yüksek hassasiyetle tespit edebilir (Zhou et al., 2011). Yeni nesil dizileme tabanlı yöntemler (örneğin, tam genom bisülfid dizileme [WGBS], azaltılmış temsil bisülfid dizileme [RRBS]) tek baz seviyesinde kapsamlı metilasyon haritaları oluşturulmasını sağlayarak, genom çapında epigenetik analizler için güçlü bir araç sunmaktadır (Lister et al., 2009).

Bu çalışmada, DNA sitozin metilasyonunu belirlemek için kullanılan başlıca moleküler belirteç teknikleri detaylı bir şekilde incelenecek, bu yöntemlerin avantajları, sınırlamaları ve kullanım alanları ele alınacaktır. Epigenetik regülasyon mekanizmalarının daha iyi anlaşılması, hastalık biyobelirteçlerinin keşfi, genetik mühendislik uygulamaları ve kişiselleştirilmiş tıp alanlarında büyük katkılar sağlamaktadır (Feil & Fraga, 2012). Bu nedenle, DNA metilasyon analizlerine yönelik güncel tekniklerin karşılaştırmalı değerlendirilmesi, epigenetik araştırmaların ilerlemesine önemli bir rehber niteliği taşıyacaktır.

DNA Metilasyon Analiz Tekniklerinin Detaylı Açıklaması

DNA metilasyonu, gen ifadesini ve genom stabilitesini düzenleyen önemli bir epigenetik modifikasyondur. DNA metilasyonunu analiz etmek için her biri benzersiz metodolojilere, avantajlara ve sınırlamalara sahip çeşitli teknikler geliştirilmiştir. Aşağıda dört ana DNA metilasyon analiz tekniğinin ayrıntılı bir açıklaması yer almaktadır:


1. Bisülfıt Sekanslama (BS-Seq)

Bisülfıt dizileme, metillenmemiş sitozinlerin sodyum bisülfıt işlemi kullanılarak urasile kimyasal olarak dönüştürülmesine dayanırken, metillenmiş sitozinler değişmeden kalır. İşlenmiş DNA'nın dizilenmesiyle, metilasyon modelleri tek nükleotid çözünürlüğünde tanımlanabilir.

Adımlar:

a) DNA İzolasyonu ve Bisülfıt İşlemi:

- Genomik DNA izolasyonu gerçekleştirilir ve sodyum bisülfıt ile muamele edilir.
- Metillenmemiş sitozinler (C) urasile (U) dönüştürülürken, 5-metilsitozinler (5mC) değişmeden kalır (Şekil 1).



Şekil 1. Bisülfıt dönüşüm reaksiyonu

b) PCR Amplifikasyonu ve Sekanslama:

- Bisülfıt ile muamele edilmiş DNA, PCR kullanılarak istenen gen bölgesi çoğaltılır.
- Urasiller dizilemede timin (T) olarak okunurken, metillenmiş sitozinler hala sitozin olarak okunur.

c) Metilasyon Analizi:

- Bisülfıt ile muamele edilmiş ve edilmemiş diziler karşılaştırılarak, metillenmiş sitozinler metillenmemiş olanlardan ayırt edilebilir. Bu amaçla geliştirilmiş farklı programlar bulunmaktadır.

Avantajlar:

- Metilasyon analizi için tek baz çözünürlüğü sağlar.
- Genom çapında metilasyon modellerini analiz edebilir.

Sınırlamalar:

- Bisülfıt dönüşümü DNA yapısının bozulmasına neden olabilmekte ve bundan dolayı da analiz edilecek DNA dizileri sınırlı tutulması gerekmektedir. Ayrıca dizayn edilen primerlerin hedef bölgeye bağlanması ve ampikon oluşturması normal geliştirilen primerlere göre daha zordur.
- PCR amplifikasyon önyargıları kantitatif metilasyon analizini etkileyebilir.

2. Metilasyona Duyarlı Restriksiyon Enzimi (MSRE) Analizi

Bu yöntem, DNA'yı sindirmek için metilasyona duyarlı ve metilasyona bağlı restriksiyon enzimlerini kullanır ve metillenmiş ve metillenmemiş DNA bölgeleri arasında ayırım yapılmasını sağlar.

Adımlar:

a) Enzim Sindirimi:

- İki tip restriksiyon enzimi kullanılır:
- Metilasyona duyarlı enzimler (örn. *HpaII*) sadece metillenmemiş DNA'yı keser.
- Metilasyona bağlı enzimler (örn. *MspI*) hem metillenmiş hem de metillenmemiş DNA'yı keser (Tablo 1).

b) PCR veya Elektroforez Analizi:

- Sindirilmiş DNA, PCR kullanılarak çoğaltılır veya fragmanlar jel elektroforezi kullanılarak analiz edilir.
- Spesifik parçaların varlığı veya yokluğu metilasyon durumunu gösterir.

Tablo 1. *MspI* ve *HpaII* Restriksiyon enzim metilasyon duyarlılıkları.

Hedef Bölge	<i>MspI</i>	<i>HpaII</i>	Metilasyon Durumu
5'→CCGG←3' 3'←GGCC←5'	Keser	Keser	Herhangi bir metilasyon yok.
5'→CC ^m GG←3' 3'←GGC ^m C←5'	Keser	Kesmez	Metilasyon sadece içteki sitozinlerde mevcut.
5'→C ^m C ^m GG←3' 3'←GGCC←5'	Kesmez	Keser	Sadece tek sarmalda iki sitozin de metilli.
5'→C ^m CGG←3' 3'←GGCC←5'	Kesmez	Keser	Sadece tek sarmalda dıştaki sitozinde metilasyon mevcut.
5'→C ^m C ^m GG←3' 3'←GGC ^m C ^m ←5'	Kesmez	Kesmez	Ful metilli.

Avantajlar:

- Hızlı ve uygun maliyetli.
- Belirli DNA bölgelerini analiz etmek için uygundur.

Sınırlamalar:

- Genom çapında metilasyon analizi için uygun değildir.
- Restriksiyon enzim bölgesine bağımlılık çözünürlüğü sınırlar.

3. Metillenmiş DNA İmmünopresipitasyonu (MeDIP)

MeDIP, metillenmiş DNA parçalarını immünopresipite etmek için 5-metilsitozine (5mC) özgü antikolar kullanır ve bunlar daha sonra sekanslama (MeDIP-seq) veya mikro diziler kullanılarak analiz edilebilir.

Adımlar:

a) DNA'nın kesilmesi:

- Genomik DNA, sonikasyon veya enzimatik sindirim kullanılarak daha küçük parçalara ayrılır.

b) Antikor Tabanlı Zenginleştirme:

- Metillenmiş DNA parçalarını seçici olarak yakalamak için 5mC'ye özgü antikolar kullanılır.

c) Tespit ve Analiz:

- Zenginleştirilmiş DNA, yüksek verimli dizileme (MeDIP-seq) kullanılarak analiz edilir veya mikro diziler (MeDIP-chip) üzerine hibridize edilir.

Avantajlar:

- Genom çapında metilasyon analizi için uygundur.
- Bisülfıt işlemleri gerektirmez, DNA bütünlüğünü korur.

Sınırlamalar:

- Bisülfıt dizilemeye kıyasla daha düşük çözünürlük.
- Mutlak kantifikasyon yerine sadece göreceli metilasyon seviyeleri sağlayabilir.

4. Tüm Genom Bisülfıt Sekanslama (WGBS)

Tüm genom bisülfıt dizileme (WGBS), genom çapında tek baz çözünürlüklü metilasyon haritaları sağlayan gelişmiş bir bisülfıt dizileme şeklidir.

Adımlar:

a) DNA Ekstraksiyonu ve Bisülfıt İşlemi:

- Genomik DNA, metillenmemiş sitozinleri urasile dönüştüren sodyum bisülfıt ile muamele edilir.

b) Kütüphane Hazırlama ve Yüksek Verimli Dizileme:

- Bisülfıt ile muamele edilmiş DNA, dizileme kütüphaneleri oluşturmak için kullanılır.
- Yeni nesil dizileme (NGS) genom boyunca metilasyon haritaları oluşturur.

c) Veri Analizi ve Metilasyon Haritalaması:

- Dizileme verileri referans genomla hizalanır.
- Her sitozindeki metilasyon durumunu belirlemek için biyoinformatik araçlar kullanılır.

Avantajlar:

- Tüm genom boyunca tek nükleotid çözünürlüğü sağlar.
- DNA metilasyonunun kantitatif analizine izin verir.

Sınırlamalar:

- Yüksek dizileme maliyetleri.
- Karmaşık biyoinformatik analiz gerektirir.

DNA sitozin metilasyonunun belirlenmesinde her ne kadar farklı yöntem ve yaklaşımlar olsa da en fazla kullanılan yöntemler açıklanmıştır. Ayrıca bu yöntemlerin bir biri ile karşılaştırmalı olarak anlatımı ise Tablo 2'de belirtilmiştir.

Tablo 2. Tekniklerin Karşılaştırılması

Teknik	Çözünürlük	Genom Kapsamı	Kantitatif Özellik	Maliyet	En İyi Kullanım Alanı
Bisülfid Sekanslama (BS-Seq)	Tek Baz	Gen Spesifik ya da Genom Çapında	Evet	Orta	Yüksek Çözünürlü Metilasyon Analizi
Metilasyona Duyarlı Restriksiyon Enzimi (MSRE) Analizi	Düşük	Lokus Spesifik	Hayır	Düşük	Hızlı ve Uygun Maliyetli Tarama
Metillenmiş DNA İmmünopresipitasyonu (MeDIP)	Orta	Genom Çapında	Göreceli	Orta	Genom Çapında Metilasyon Profillemesi
Tüm Genom Bisülfid Sekanslama (WGBS)	Tek Baz	Genom Çapında	Evet	Yüksek	Yüksek Çözünürlü ve Kapsamlı Metilasyon analizi

Bu tekniklerin her birinin kendine özgü güçlü yanları ve sınırlamaları vardır. Uygun yöntemin seçimi, araştırma hedeflerine, maliyet hususlarına ve çalışma için gereken çözünürlük seviyesine bağlıdır.

Bu çalışmada, DNA sitozin metilasyonunu belirlemek için kullanılan başlıca moleküler belirteç teknikleri incelenerek, epigenetik araştırmalarda kullanılan yöntemlerin avantajları, sınırlamaları ve kullanım alanları ele alınmıştır. Epigenetik regülasyonun temel mekanizmalarının anlaşılması, hastalık biyo-belirteçlerinin keşfi, genetik mühendisliği, bitki ıslahı, ve kişiselleştirilmiş tıp alanlarında önemli katkılar sağlamaktadır.

KAYNAKLAR

- Allis, C. D., & Jenuwein, T. (2016). *The molecular hallmarks of epigenetic control*. Nature Reviews Genetics, 17(8), 487-500.
- Bird, A. (2007). *Perceptions of epigenetics*. Nature, 447(7143), 396-398.
- Chung, H., Lee, J., Kim, Y., & Park, J. (2021). *Epigenetic modifications and stress tolerance in plants*. Plant Physiology and Biochemistry, 162, 120–130.
- Clark, S. J., Harrison, J., & Paul, C. L. (2006). *Bisulfite sequencing for DNA methylation analysis*. Methods in Molecular Biology, 301, 27-45.
- Feil, R., & Fraga, M. F. (2012). *Epigenetics and the environment: Emerging patterns and implications*. Nature Reviews Genetics, 13(2), 97-109.
- Jaenisch, R., & Bird, A. (2003). *Epigenetic regulation of gene expression: How the genome integrates intrinsic and environmental signals*. Nature Genetics, 33(3), 245-254.

Lister, R., Pelizzola, M., Downen, R. H., Hawkins, R. D., Hon, G., Tonti-Filippini, J., & Ecker, J. R. (2009). *Human DNA methylomes at base resolution show widespread epigenomic differences*. *Nature*, 462(7271), 315-322.

Sahu, P. P., Pandey, G., Sharma, N., Puranik, S., Muthamilarasan, M., & Prasad, M. (2013). *Epigenetic mechanisms of plant stress responses and adaptation*. *Current Genomics*, 14(2), 95-111.

Schübeler, D. (2015). *Function and mechanisms of DNA methylation*. *Nature*, 517(7534), 321–326.

Xie, H., et al. (2018). *DNA methylation in drought stress response of plants*. *Plant Molecular Biology*, 96(1-2), 85-98.

Zhou, X., Maricque, B., Xie, M., Li, D., Sundaram, V., Martin, E. A., ... & Wang, T. (2011). *The Human Epigenome Browser at Washington University*. *Nature Methods*, 8(12), 989-990.

THE IMPORTANCE OF RANGELANDS AS CARBON SINKS

Fatih KUMBASAR

Ondokuz Mayıs University, Graduate Education Institute, Department of Field Crops,
Samsun, Turkey

ORCID ID: <https://orcid.org/0000-0002-0379-3355>

Mehmet CAN

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: <https://orcid.org/0000-0003-0230-6209>

Gülcan KAYMAK BAYRAM

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: <https://orcid.org/0000-0002-0915-0529>

İlknur AYAN

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: <https://orcid.org/0000-0002-5097-9013>

Zeki ACAR

Ondokuz Mayıs University, Agricultural Faculty, Department of Field Crops, Samsun,
Turkey

ORCID ID: <https://orcid.org/0000-0002-0484-1961>

ABSTRACT

This study examines the role of grasslands in the carbon cycle and their potential in combating climate change. Grasslands, ranking second only to forests as terrestrial carbon sinks, store approximately 25-34% of global terrestrial carbon stocks. These ecosystems capture atmospheric carbon dioxide (CO₂) through photosynthesis, converting it into organic matter and storing significant amounts of carbon in the soil. Deep-rooted plant species play a crucial role in long-term carbon sequestration by distributing carbon across various soil depths. Additionally, grasslands contribute to effective water retention, acting as natural barriers against drought and soil erosion.

Grasslands also serve as a primary forage source for livestock, and proper grazing management can enhance both animal welfare and greenhouse gas (GHG) mitigation. The presence of leguminous forage species has been shown to reduce methane production from ruminants. Research indicates that controlled grazing practices improve soil organic carbon (SOC) levels,

preserving soil fertility. Studies conducted in Turkey reveal significant declines in SOC stocks when grasslands are converted into agricultural land. However, appropriate grazing management and pasture rehabilitation efforts can prevent carbon losses and enhance sequestration potential.

Beyond carbon storage, grasslands play a vital role in biodiversity conservation, soil health improvement, and sustainable water resource management. Studies conducted in the United States, Europe, and China demonstrate that well-managed grazing systems can increase soil carbon storage capacity by 20-50%. Furthermore, in certain regions, grasslands may outperform forests as carbon sinks. In fire-prone areas, grasslands provide a more stable carbon storage alternative, as carbon remains stored in the soil rather than being released into the atmosphere through biomass combustion.

In conclusion, grasslands serve as crucial carbon sinks, offering significant climate change mitigation potential while supporting livestock production, water management, and soil health. Sustainable grassland management and rehabilitation can enhance carbon sequestration, reduce greenhouse gas emissions, and strengthen ecosystem services. On a global scale, the conservation and improvement of grassland ecosystems present an effective strategy in the fight against climate change.

INTRODUCTION

Climate change is one of the most significant environmental issues of our time. The increase in global greenhouse gas emissions has led to rising temperatures, ecosystem degradation, and a reduction in biodiversity. In this context, natural carbon sinks such as forests, wetlands, and grasslands play a critical role in balancing atmospheric carbon dioxide (CO₂) levels (IPCC, 2023). The concept of sink first defined in the UN Framework Convention on Climate Change, is defined as any process, activity or mechanism that removes a greenhouse gas, an aerosol or a precursor of a greenhouse gas from the atmosphere (UNFCCC, 1992). Kyoto Protocol Land Use, Land Use Change and Forestry Guide (LULUCF) divides sink areas into six groups; It is categorized as (1) forest areas, (2) meadow and pasture areas, (3) agricultural areas, (4) wetlands, (5) residential areas and (6) other areas (IPCC, 2003).

Numerous ecosystem services in grasslands are supported by the movement of carbon (C) among plants, microorganisms, soils, and the atmosphere, which serves as the principal medium for ecosystem energy flow and biogeochemical cycle (Chapin et al., 2006). Thus, a key component of grassland conservation and sustainable use is understanding the mechanisms that regulate C fluxes and pools in grassland (Liu et al., 2022).

Although the C pool in grasslands is less than that in forests, it nevertheless contributes roughly 25–34% of the total terrestrial C supply in the world (Bai and Cotrufo, 2022; IPCC, 2007). The difference between C uptake, primarily by photosynthesis, and losses, primarily by respiration, leaching, and erosion, determines the C balance of grasslands, just like it does in forests. However, due to their lower density and shorter stature of plants compared to forests, grasslands can be subject to stronger winds, more intense sun radiation, and larger temperature swings. Therefore, the regulation of the grassland C balance is more heavily influenced by abiotic processes (Austin, 2011; Grünzweig et al., 2022).

Grasslands, by storing carbon in soil and vegetation, have great potential in mitigating the effects of climate change. Covering approximately 24% of the Earth's land area, grasslands are among the largest ecosystems for carbon storage, second only to forests (FAO, 2023).

It is estimated that grasslands retain at least 10% of the global carbon stock in soil (Anderson, 1991; Eswaran et al., 1993). It is known that grasslands are important carbon sinks with a level of 200-300 billion tons (Pg) of CO₂ (Batjes and Sombroek, 1997). Even small changes in

grassland carbon absorption inputs can cause greater and longer-lasting carbon sequestration in pastures (Scurlock and Hall, 1998).

This review will address the carbon cycle, carbon sinks, and the role of grasslands in ecosystems, and will examine the impact of sustainable grassland management on carbon absorption based on scientific data.

THE IMPORTANCE OF CARBON

Carbon is a nonmetallic chemical element found widely in nature. Carbon, which is found both naturally and in combination with other elements, constitutes approximately 0.2% of the Earth's crust by weight.

CO₂ (carbon dioxide), which constitutes approximately 0.04% of the atmosphere and is found dissolved in all natural waters, carbonate minerals such as limestone and marble, and hydrocarbons, which are the main building blocks of coal, oil and natural gas, are the most abundant compounds.

Carbon is one of the most important building blocks of living cells and the most important element of the biological system. 40-50% of the total weight of plants and microorganisms consists of carbon. The atmosphere plays the most important role in the carbon cycle. Here, carbon is found in the form of carbon dioxide (CO₂) gas. There is approximately 0.04% CO₂ gas in the atmosphere. It has been determined that the safe level of CO₂ in the atmosphere is 0.03% (Jones, 2017).

Carbon is found in organic and inorganic forms in soil. There is a significant amount of carbon in the form of calcium and magnesium carbonate compounds in rocks such as limestone and dolomite, which are carbonate-containing rocks. However, since these rocks do not decompose or dissolve easily, they do not play an important role in the carbon cycle and ecosystem services.

CARBON CYCLE

It refers to the exchange of C within and between C reservoirs connected by a network of physical, chemical and biological processes.

The C cycle consists of a large number of intertwined cycles that differ according to some reservoir processes. The C cycle includes biotic and abiotic processes and transfers C within and between four main reservoirs.

These reservoirs are;

- » lithosphere (solid outer shell),
- » hydrosphere,
- » atmosphere,
- » biosphere connected by a complex set of natural and anthropogenic processes.

The main source of C in the C cycle is CO₂ and in the absence of anthropogenic influence, the C cycle is generally stable.

Carbon exchange between reservoirs includes photosynthesis, respiration, water-atmosphere interface and gas exchange with weathering.

Carbon is found dissolved in water in the form of carbon dioxide and bicarbonate (HCO₃) in lakes, streams, seas and oceans.

The transfer of carbon from the atmosphere to water in various ways and the respiration and decomposition of consumer organisms living in water are the reasons for the presence of carbon in the hydrosphere.

Organic compounds, which are the building blocks of all living organisms, are carbon-based. Carbon is also found in the biosphere as organic molecules. This is also the reason why fossilized organisms contain high amounts of carbon.

Carbon, which is present as an organic compound in the structure of living cells, is converted into CO₂ by decomposers (decayers) when the organism dies and released into the atmosphere.

Photosynthesis, respiration and decomposition play key roles in the soil carbon cycle. Most of the carbon dioxide in the atmosphere enters the cycle through photosynthesis. Plants take in atmospheric carbon dioxide through photosynthesis and convert it into organic carbon. Some of the carbon taken in by plants is returned to the atmosphere through respiration. The remaining carbon is stored as organic compounds in plant tissues.

The global carbon cycle is closely linked to the greenhouse effect. Carbon is constantly moving between air, plants and soil, and changes in any of these three components always affect the other components.

Globally, there is twice as much carbon in soil organic matter as in the atmosphere, and as a result, relatively small changes in soil organic matter can have a large impact on CO₂ in the air.

Some of the organic carbon created by producers reaches consumer organisms through the food chain. Consumers release some of the carbon they take into the atmosphere as CO₂ through respiration. The remaining part is stored in organisms. Dead organisms and wastes are transformed by decomposers on the soil. During the decomposition process, some of the carbon mixes with the atmosphere in the form of carbon dioxide from the soil due to microbial respiration, while some of the carbon remains in the soil in the form of humus. 60% of the humus in the soil consists of organic carbon (Piccolo et al., 2018).

CARBON EMISSION

Carbon emissions are the release of carbon dioxide gas into the atmosphere as a result of the combustion of carbon-containing fuels (fossil fuels: oil, natural gas, coal, etc.).

In 2021, 73.2% of the total CO₂ equivalent greenhouse gas emissions in the world were from the energy sector, 5.2% from the industrial processes and product use sector, 3.2% from the waste sector, and 18.4% from the agricultural sector (IEA, 2022).

Table 1. Countries that emit the most CO₂

Ranking	Countries	CO ₂ (billion tonnes)	The rate in the World (%)
1	China	15.6	29.16
2	USA	6.0	11.19
3	India	3.9	7.33
4	Russia	2.5	4.80
5	Brasil	1.3	2.44
6	Indonesia	1.2	2.31
7	Japan	1.1	2.20
8	Iran	0.9	1.77
9	Mexico	0.81	1.52
13	Türkiye	0.6	1.28

(Crippa et al. 2022)

It has been reported that there will be an increase of 0.9% in the world in 2022, reaching 53.8 billion tons of carbon dioxide equivalent emissions (IEA, 2022). In Turkey, the total greenhouse gas emissions in 2021 were determined as 523.9 million tons of carbon dioxide equivalent (TUIK, 2022). The sources of emissions from agriculture in our country can be listed as animal husbandry, rice farming and other agricultural activities.

The rapid increase in atmospheric carbon (CO₂) concentrations throughout the industrial era is due to anthropogenic (human activities).

Today, the most important sectors causing greenhouse gas emissions are energy, transportation, industry and agriculture.

Rangeland areas, which are vital for livestock practices that lead in greenhouse gas emission production from the agricultural sector, are natural assets and rural-ecologically used areas that are directly affected by climate change and directly affect climate change.

CARBON FOOTPRINT

It is a measure of the damage caused to the environment by human activities in terms of the amount of greenhouse gas produced, measured in units of carbon dioxide.

Primary footprint is a measure of direct CO₂ emissions from the combustion of fossil fuels, including household energy consumption and transportation, while secondary footprint is a measure of the indirect CO₂ emissions from the entire life cycle of the products we use, related to their manufacture and eventual degradation.

Areas where more carbon is absorbed than is emitted are called carbon sinks. The most important carbon sinks are forests, meadows and pastures, and wetlands.

IMPORTANCE OF RANGELANDS

Covering approximately 24% of the world's land and ranking second after forests, meadow-pasture ecosystems provide a significant portion of the roughage required by animals and also constitute the most important and largest source of biological wealth in countries.

In countries where animal product costs are high, as in our country, pastures are given special importance by being the cheapest source of roughage.

Thanks to their high species diversity, they produce more balanced roughage for domestic animals. In our country, our pastures are an indicator of our social and economic conditions.

Due to resource use and legal problems, some of our pastures have been converted to agricultural areas or included in forest borders, and the remaining significant portion has lost its productivity and self-sustainability due to increasing grazing pressure.

In pasture ecosystems, climate, soil, vegetation and animal factors form a whole in mutual interaction. The continuity and productivity of vegetation in pasture areas can be protected by conscious and planned grazing.

Meadows and pastures are natural ecosystems that have undertaken many ecological functions such as the protection and sustainability of natural resources, the indispensable nature of wildlife, creating genetic resources with biodiversity and serving different uses.

Meadows and pastures play an active role in reducing the greenhouse effect on the warming of the earth and reduce the greenhouse effect that occurs with the increase in carbon dioxide by assimilation.

Table 2. Land use changes in Turkey over the years

Years	Forest (%)	Rangelands(%)	Cropland (%)
1945	13.55	50.19	18.85
1950	13.37	49.54	20.54
1960	13.58	36.77	32.49
1970	23.44	27.58	35.07
1980	25.91	27.16	36.15
1990	25.91	18.19	35.74
2000	26.56	18.47	33.84
2010	27.63	18.75	31.3
2017	27.81	18.75	29.99
2022	29.4	18.75	30.70

(TUIK, 2023)

The Kyoto Protocol Guidelines for Land Use, Land Use Change and Forestry (LULUCF) categorize sink areas into six groups as (1) forest areas, (2) pastures and meadows, (3) agricultural areas, (4) wetlands, (5) settlements and (6) other areas (IPCC, 2003).

Pasture areas are very important in terms of their biodiversity, high carbon absorption and erosion prevention capacity. In addition, as stated in the Intergovernmental Panels on Climate Change (IPCC, 2023), they are important terrestrial carbon sinks thanks to their plant diversity and soil structure.

THE IMPORTANCE OF RANGELANDS AS CARBON SINKS

Rising atmospheric CO₂ levels and the resulting increases in average temperatures have increased interest in developing strategies to biosequester C in order to mitigate climate change. Pastures are the areas with the highest carbon sequestration in terrestrial ecosystems in the world and in our country, after forests. Pastures contribute to approximately 25-34% of the world's terrestrial C stock (FAO, 2023).

Within the scope of the "Soil Organic Carbon Project" carried out to determine the Organic Carbon Capacity of Turkish Soils, the soil organic carbon amounts of our country's soils were determined (Anonymous, 2018).

Table 3. Land Use and SOC Amounts Found in Turkey

Land Use Status	Land (ha)	SOC (tonnes C/ha)	SOC Stock (%)
Forest	24.180.644	55.68	38.33
Rangeland	23.568.338	49.77	33.39
Cropland	26.316.375	35.96	26.94
Bare Land	1.172.581	12.78	0.43
Sow pastures	796.519	16.12	0.37
Wetlands	393.100	49.71	0.56

Organic matter, which forms the source of soil organic carbon, is also an important indicator of terrestrial ecosystem productivity and soil fertility.

According to the results of the project called Soil Organic Carbon (2018), the total amount of carbon stock in Turkey at 30 cm soil depth was determined as 3.51 billion tons. As a result of the study, the highest amount of soil organic carbon was determined as,

Black Sea Region 67.83

Mediterranean Region 52.36

Eastern Anatolia Region-46.92

Aegean Region- 43.18

Marmara Region-40.67

Central Anatolia Region- 38.50

Southeastern Anatolia Region- 29.46 tons C/ha.

In a study conducted in Eskişehir, in soil samples taken from different places, the places with the highest soil organic carbon were determined as forest, pasture and agricultural areas, respectively. Soil organic carbon in the forest was determined as 32.23 million tons, in pasture 31.37 million tons and in agricultural areas as 16.88 million tons. They reported that soil organic matter and carbon ratio increased as the distance from the settlement in pasture areas increased, and that this was related to the vegetation ratio and composition. Organic matter values decreased as the depth increased in pasture and forest areas; They determined that the values in agricultural areas were quite low and there was no significant and linear change (Evliyaoglu and Avdan, 2019).

Pastures are areas with high plant diversity. Due to these features, plants can store carbon at different depths of the soil with different root depth developments. Perennial plants in particular contribute to soil organic matter with both root biomass and deeper roots. Since carbon is accumulated at different depths in pastures, these areas are more stable carbon stores.

In France, it has been determined that the difference in C retention capacity between agricultural lands and pastures is approximately 25 tons/ha. Researchers have stated that this difference in

pastures is due to the amount of organic matter accumulated at different root depths (Soussana et al., 2004). Pasture lands accumulate carbon for 15 to 25 years after they are established, and then this accumulation tends to continue at a constant level. This is generally an indication that the soil has reached carbon saturation, which varies according to its chemical, biological and physical properties. When surface soils are saturated with carbon, the use of deep-rooted species allows carbon to be retained deeper and more in the soil (Ontl and Janowiak, 2017).

Due to the lack of tillage, there is continuous vegetation on the soil surface in pastures. This situation causes continuous carbon input from aboveground vegetation and release of large amounts of organic carbon into the subsoil through root exudates and decomposition of deep roots. In addition, the mulch layer consisting of plant residues contributes to the accumulation of organic matter in the soil and increases C accumulation in the soil.

The continuous covering of the soil surface with vegetation also reduces soil losses. In areas with low vegetation cover, soil loss occurs from the soil surface and soil organic matter decreases over time. Soil organic matter is decomposed by microorganisms, soil temperature increases and organic carbon is released. This released carbon passes into the atmosphere as CO₂. In addition, the decrease in organic matter can lead to soil losses (Lal, 1995; Lal et al., 1998).

Net C storage in pasture soils can vary between years and sites. Even when net primary productivity (organic matter remaining after respiration) is low in natural pastures as a result of drought or low temperature, they have the potential to sequester significant amounts of CO₂ and play an important role in the global C cycle (Bolinder et al., 2007).

The amount of organic matter in pasture soils varies depending on topographic conditions and grazing intensity.

In a study conducted in three different pasture sections at 1900, 2000 and 2200 meters altitude in the pastures of Aydın Village in the Ardanuç District of Artvin Province, it was determined that the highest organic matter rate was 6.27% in the pasture at 2200 m altitude, 3.10% in the pasture section at 2000 altitude, and 5.66% at 1900 altitude. In the light of these results, it was emphasized that the amount of organic matter increased depending on the altitude, and the low amount of organic matter in the pasture at 2000 altitude was attributed to the decrease in the proportion of gramineous and legume forage plants in the botanical composition due to grazing pressure (Bilgin, 2010).

In pastures in Erzurum Palandöken, soil organic matter content was found to be 2.30% at 2000 m altitude, 4.10% at 2500 m altitude, and 6.84% at 3000 m altitude. As altitude increased, the amount of organic matter in the soil also increased (Çomaklı et al., 2012).

RANGELANDS AND MITIGATION CLIMATE CHANGE EFFECTS

Plants in pastures absorb CO₂ during their growth and store it in different tissues. Aboveground plant parts are grazed by ruminant animals and C eventually returns to the soil as fertilizer or to the atmosphere through enteric fermentation. The remaining plants and roots eventually decompose and C is stored in the soil. Methane is the reduction of CO₂ and H₂ formed as a result of anaerobic fermentation of nutrients in the rumen of ruminant animals to Methane (CH₄) by methanogenic bacteria. It is stated that the total contribution of methane and nitrous oxide to global warming is between 17-27% on a global scale. The proportion of methane emissions from animals is 16% of all methane emissions (Tseten et al., 2022).

Since pastures are used for grazing, energy loss is less. In addition, since animals graze fresh plants on pastures, methane release is less. The presence of plant diversity, especially legume forage plants containing tannin, reduces methane release. Methane release from the digestive system can be reduced by improving the quality of roughage (Chung et al., 2013).

According to a study conducted in France, the gross carbon footprint for 1 liter of milk production was 1.2-1.3 kg CO₂, while the net carbon footprint was determined as 0.8-1.2 kg CO₂ in calculations considering carbon sequestration in pastures and meadows. It has been reported that carbon sequestration in pastures and meadows can cause a 5-30% reduction in carbon emissions from dairy farming (Salvador et al., 2017).

THE IMPORTANCE OF RANGELAND MANAGEMENT AS A CARBON SINK

According to a study conducted in the United States, both natural and sown pastures are strong carbon sinks, holding up to 0.9 metric tons of carbon per year (Adewopo et al., 2014). In addition, it has been determined that soil carbon increases by 50% in pastures with more productive plant species and appropriate grazing and fertilization management. Adopting more effective pasture management practices is also beneficial in terms of pasture productivity and livestock production (Adewopo et al., 2014; Xu et al., 2016). It is estimated that there can be an average increase of 2.9% in global carbon stocks with appropriate grazing management (Conant et al., 2001).

Within the scope of the Sustainable Pasture Management Project, a rotational grazing system was applied in a study called ‘‘Alternative Grazing Management Demonstration Program in Karacadağ (Şanlıurfa Section)’’. It was determined that the area covered with plants was between 23.1-72.2% at the beginning of grazing, while it was between 28.9-80% at the end of the grazing period. The proportion of legumes in the botanical composition was between 9.4-14.28% at the beginning, and 10.68-15.49% at the end. While the soil organic matter ratio was between 0.70-1.49% at the beginning, it was determined to be between 1.62-1.82% at the end of the grazing period (Ayan et al., 2022).

As a result of a study carried out in a pasture area that was ploughed and abandoned in Samsun ecological conditions, the organic matter content of the plant-covered area and soil as a result of the pasture improvement processes applied are shown in the Table (Mut and Ayan, 2009).

Table 4. Plant-covered area and organic matter contents

Years	Area Covered with Plants (%)	Organic matter (%)
2005	38.3-49.8	2.8
2006	47.8-60.6	2.5-5.9
2007	56.7-64.5	3-5.7
2008	60-73.5	3.0-5.8

In low-covered pastures, soil carbon losses can be limited by appropriate grazing management and increasing the proportion of deep-rooted forage crops.

Soil carbon recovery is generally a slow process, taking decades to centuries depending on the carbon balance of the system (Bai et al., 2008). Despite these slow changes, the global carbon sequestration potential from restoring degraded pastures is significant, with the potential to sequester approximately 3 Gt C/year, equivalent to a 50 ppm reduction in atmospheric CO₂ over 50 years (Lal, 2009).

The effects of grazing on soil carbon may depend on interactions between soil, plant species and climate.

It has been reported that when grazing frequency exceeds pasture capacity, soil organic carbon stocks decrease by an average of 0.9% per year, which is due to the inability of frequently grazed plant roots to fix C to the soil (Abdalla et al., 2016).

A 25.9% decrease in soil organic carbon was determined in a freely grazed pasture in Argentina (Abril and Bucher, 1999).

In pastures with controlled grazing, soil organic carbon increased by 48% per year in China (Guo et al., 2001), 35-45% in South Africa (Chaplot et al., 2016) and 20% in the United States (Manley et al., 1995).

Approximately 20% of the world's pastures have been converted to agricultural land (Ramankutty et al., 2008). In the central states of America, many soils have lost 30-50% of their carbon (25 to 40 metric tons of carbon per hectare) due to conversion to agricultural land (Lal, 2002).

In a review of 115 studies, it was determined that agricultural lands converted to pasture gained approximately 3-5% C. It was also determined that the C content gained by pasture soils was affected by pasture management and especially fertilization (Conant et al., 2001).

As a result of conversion of agricultural lands to pastures, the amount of organic C in the soil increases by 18%, which can provide an annual retention of 332 kg of C (Guo and Gifford, 2002; Post and Kwon, 2000).

Considering the studies conducted within the scope of the "Soil Organic Carbon Project" of Turkey, as a result of the conversion of 1 million hectares of pasture land to agricultural land; It is stated that there may be a decrease of approximately 13.8 million tons C in the amount of soil organic carbon, and this decrease will lead to a decrease of approximately 0.39% in the total soil organic carbon stock of Türkiye (Anonymous, 2018).

CONCLUSION

We are faced with a global warming problem that has been accelerating over the last century. The reason for this is the increase in the density of various heat-trapping gases in the atmosphere, primarily carbon dioxide, methane, nitrogen oxide and water vapor.

In pasture ecosystems, climate, soil, vegetation and animal factors interact to form a whole. Pastures are of great importance in terms of animal production, as well as human, environmental, animal and soil health.

Pastures are important terrestrial carbon sinks thanks to their plant diversity and soil structure.

Since carbon is accumulated at different depths in pastures, these areas are more stable carbon storage areas.

The fact that the soil surface is constantly covered with vegetation in pastures and the amount of organic matter increases also reduces soil losses through erosion.

Methane emissions are lower since animals graze fresh plants in pastures. The presence of plant diversity, especially legume forage plants containing tannin, reduces animal-based methane emissions.

In order for pastures to make the expected contribution to reducing greenhouse gas emissions;

Pastures should be improved and developed to the extent of their capacity.

With correct pasture management,

Biodiversity should be protected

Underground and aboveground plant density should be kept high

Grazing should be regulated in terms of time and intensity, taking into account surface forms and erosion dynamics

Pasture ecosystems should be managed with a holistic approach in terms of protection and sustainability of the environment, wildlife, soil and water resources.

As a result, pastures regulate greenhouse gas emissions with good pasture management, while simultaneously providing many ecosystem services such as

- providing quality feed for animals,
- protecting soil and water, and
- regulating microclimate.

REFERENCES

- Abdalla, K., Chivenge, P., Everson, C., Mathieu, O., Thevenot, M., & Chaplot, V. (2016). Long-term annual burning of grassland increases CO₂ emissions from soils. *Geoderma*, 282, 80-86.
- Abril, A., & Bucher, E. H. (1999). The effects of overgrazing on soil microbial community and fertility in the Chaco dry savannas of Argentina. *Applied Soil Ecology*, 12(2), 159-167.
- Adewopo, J. B., Silveira, M. L., Xu, S., Gerber, S., Sollenberger, L. E., & Martin, T. A. (2014). Management intensification impacts on soil and ecosystem carbon stocks in subtropical grasslands. *Soil Science Society of America Journal*, 78(3), 977-986.
- Anderson, J.M., (1991). The effects of climate change on decomposition processes in grassland and coniferous forests. *Ecological Applications*, 1, 326-347.
- Anonymous, 2018. Soil Organic Carbon Project – ÇEM-TUBITAK-BILGEM, 2018, Ankara
- Austin, A. T. (2011). Has water limited our imagination for aridland biogeochemistry?. *Trends in ecology & evolution*, 26(5), 229-235.
- Ayan, İ., Acar, Z. Töngel, M.Ö. and Can, M. (2022). Alternative Grazing Management Demonstration Program Final Report In Karacadağ (Şanlıurfa Section), Protection And Sustainable Management Of Türkiye Steppe Ecosystems Project, GCP/TUR/061/GFF, 24 p.
- Bai, Y., Cotrufo, M. F., (2022). Grassland soil carbon sequestration: Current understanding, challenges, and solutions. *Science*, 377(6606), 603-608.
- Bai, Z.G.; Dent, D.L.; Olsson, L; Schaepman, M.E. (2008). Proxy global assessment of land degradation. *Soil Use and Management* 24: 233–234.
- Batjes, N. H., Sombroek, W. G., (1997). Possibilities for carbon sequestration in tropical and subtropical soils. *Global Change Biology*, 3, 161-173.
- Bilgin, F. (2010). Artvin Ardanuç-Aydın Köyü yaylası mera vejetasyonu ile bazı toprak özelliklerinin yükseltiye göre değişiminin irdelenmesi. (Yüksek Lisans Tezi.), Artvin Çoruh Üniversitesi, Fen Bilimleri Enstitüsü, Orman Mühendisliği Ana Bilim Dalı, 91s.
- Bolinder, M. A., Janzen, H. H., Gregorich, E. G., Angers, D. A., & VandenBygaart, A. J. (2007). An approach for estimating net primary productivity and annual carbon inputs to soil for common agricultural crops in Canada. *Agriculture, Ecosystems & Environment*, 118(1-4), 29-42.
- Chapin, F. S., Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi, D. D., ... & Schulze, E. D. (2006). Reconciling carbon-cycle concepts, terminology, and methods. *Ecosystems*, 9, 1041-1050.
- Chaplot V, Dlamini P, Chivenge P. (2016). Potential of grassland rehabilitation through high density-short duration grazing to sequester atmospheric carbon. *Geoderma* 271, 10-17.

- Chung M L, Shilton A N, Guieysse B, Pratt C (2013). Questioning the accuracy of greenhouse gas accounting from agricultural waste: A case study. *Journal of Environmental Quality* 42, 654-659.
- Çomaklı, B., Fayetörbay, D., Daşçı, M. (2012). Changing of botanical composition and canopy coverage ratio in rangelands at different altitudes. *Atatürk Üniversitesi Ziraat Fakültesi Dergisi*, 43(1): 17-21.
- Conant RT, Paustian K, Elliott ET (2001) Grassland management and conversion into grassland: effects on soil carbon. *Ecological Applications*, 11, 343–355.
- Crippa, M., Guizzardi, D., Banja, M., Solazzo, E., Muntean, M., Schaaf, E., Pagani, F., Monforti-Ferrario, F., Olivier, J., Quadrelli, R., Risquez Martin, A., Taghavi-Moharamli, P., Grassi, G., Rossi, S., Jacome Felix Oom, D., Branco, A., San-Miguel-Ayanz, J. and Vignati, E., CO2 emissions of all world countries - JRC/IEA/PBL (2022). Report, EUR 31182 EN, Publications Office of the European Union, Luxembourg, 2022, doi:10.2760/730164, JRC130363.
- Eswaran, H., van den Berg, E., Reich, P. (1993). Organic carbon in soils of the world. *Soil Science Society of America Journal*, 57, 192-4.
- Evliyaoğlu, G. ve Avdan, Z. (2019). Farklı Alan Kullanımlarında Toprak Organik Karbon Dinamiğinin Belirlenmesi, Yüksek Lisans Tezi.
- FAO (2023). Global assessment of soil carbon in grasslands. – From current stock estimates to sequestration potential. FAO Animal Production and Health Paper No. 187. Rome, FAO. <https://doi.org/10.4060/cc3981en>.
- Grünzweig, J. M., De Boeck, H. J., Rey, A., Santos, M. J., Adam, O., Bahn, M., Yakir, D. (2022). Dryland mechanisms could widely control ecosystem functioning in a drier and warmer world. *Nature ecology & evolution*, 6(8), 1064-1076.
- Guo, L. B., Gifford, R. M. (2002): Soil carbon stocks and land usechange: a meta analysis. *Global Change Biol.*8, 345–360.
- IEA (2022). CO₂ Emissions from Fuel Combustion 2022: Highlights (Paris: International Energy Agency, 2022) p.98.
- IPCC (2003). Good Practice Guidance for Land Use, Land-Use Change and Forestry. IPCC National Greenhouse Gas Inventories Programme Technical Support Unit, Printed in Japan.
- IPCC (2007). The physical science basis. Contribution of working group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. *Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA*, 996(2007), 113-119.
- IPCC (2023). *Climate Change 2023: Synthesis Report*. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J.Romero (eds.)]. IPCC, Geneva, Switzerland, pp. 35-115, doi: 10.59327/IPCC/AR6-9789291691647.
- Jones, N. (2017). How the world passed a carbon threshold and why it matters. *Yale Environment*, 360, 26.
- Lal, R. (1995). Global soil erosion by water and carbon dynamics. In: Lal R., Kimble J.M., Levine E. and Stewart B.A. (eds), *Soils and Global Change*. CRC/Lewis Publishers, Boca Raton, FL, pp. 1–34.
- Lal, R. (2009). Challenges and opportunities in soil organic matter research. *European Journal of Soil Science* 60(2): 158–169.

- Lal, R. 2002. Soil carbon dynamics in cropland and rangeland. *Environmental Pollution* 116 (3): 353–362 [https://doi.org/10.1016/S0269-7491\(01\)00211-1](https://doi.org/10.1016/S0269-7491(01)00211-1).
- Lal, R., Kimble, J. M., Follett, R. F., & Cole, C. V. (1998). *The potential of US cropland to sequester carbon and mitigate the greenhouse effect*. CRC Press.
- Liu, L., Sayer, E. J., Deng, M., Li, P., Liu, W., Wang, X., Piao, S. (2022). The grassland carbon cycle: Mechanisms, responses to global changes, and potential contribution to carbon neutrality. *Fundamental Research*.
- Manley, J. T., Schuman, G. E., Reeder, J. D., & Hart, R. H. (1995). Rangeland soil carbon and nitrogen responses to grazing. *Journal of soil and water conservation*, 50(3), 294-298.
- Mut H. and Ayan, İ. (2009). *Determination of effectiveness of different improvement methods on a plowed and abandoned pasture*. Phd Tesis. Ondokuz Mayıs University.
- Ontl, T., & Janowiak, M. (2017). Grassland and Carbon Management. *US Department of Agriculture, Forest Service, Climate Change Resource Center*.
- Piccolo, A., Spaccini, R., Drosos, M., Vinci, G., & Cozzolino, V. (2018). The molecular composition of humus carbon: recalcitrance and reactivity in soils. In *The future of soil carbon* (pp. 87-124). Academic Press.
- Post, W. M., Kwon, K. C. (2000): Soil carbon sequestration and land-use change: processes and potential. *Global Change Biol.*6,317–327.
- Ramankutty, N., Evan, A. T., Monfreda, C., & Foley, J. A. (2008). Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global biogeochemical cycles*, 22(1).
- Salvador, S., Corazzin, M., Romanzin, A., & Bovolenta, S. (2017). Greenhouse gas balance of mountain dairy farms as affected by grassland carbon sequestration. *Journal of environmental management*, 196, 644-650.
- Scurlock, J. M. O., Hall, D. O. (1998). The global carbon sink: a grassland perspective. *Global Change Biology*, 4(2), 229-233.
- Soussana, J. F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T., Arrouays, D. (2004): Carbon cycling and sequestration opportunities in temperate grasslands. *Soil Use Manage.*20, 219–230.
- Tseten, T., Sanjorjo, R. A., Kwon, M., & Kim, S. W. (2022). Strategies to mitigate enteric methane emissions from ruminant animals.
- TUIK (2022). Turkish Statistical Institute. <https://data.tuik.gov.tr/Bulten/Index?p=Greenhouse-Gas-Emissions-Statistics-1990-202253701&dil=2#:~:text=TURKSTAT%20Corporate&text=The%20greenhouse%20gas%20inventory%20results,4.1%20tonnes%20CO2%20eq>.
- TUIK, (2023). <https://data.tuik.gov.tr/Kategori/GetKategori?p=tarim-111&dil=1>
- UNFCCC (1992). <https://unfccc.int/resource/docs/convkp/conveng.pdf> (Access date: 20.02.2025).
- Xu, S., Silveira, M. L., Inglett, K. S., Sollenberger, L. E., & Gerber, S. (2016). Effect of land-use conversion on ecosystem C stock and distribution in subtropical grazing lands. *Plant and soil*, 399, 233-245.

PHYTOCHEMICAL CONTENT OF LAURUS NOBILIS LEAVES AND EXTRACT; POTENTIAL FOR USE IN VALUE-ADDED COSMETICS, HEALTH AND FOOD PRODUCTS

Öğr. Gör. Musa KARADAĞ

Iğdır Üniversitesi, Teknik Bilimler Meslek Yüksekokulu, Kimya Ve Kimyasal İşleme
Teknolojileri Bölümü, Iğdır, Türkiye

Dr. Yunus BAŞAR

Iğdır Üniversitesi, Araştırma Laboratuvarı Uygulama ve Araştırma Merkezi, Iğdır, Türkiye

Prof. Dr. Mehmet Hakkı ALMA

Iğdır Üniversitesi, Araştırma Laboratuvarı Uygulama ve Araştırma Merkezi, Iğdır, Türkiye

ABSTRACT

Medicinal and aromatic plants have been used for centuries in traditional medicine, nutrition and cosmetics. *Laurus nobilis* (laurel), a medicinal plant cultivated in Europe, Asia and the tropics, is an evergreen plant that grows all year round in the Mediterranean climate. It plays an important role in the preservative, cooking, flavoring, cosmetic, chemical and pharmaceutical industries in various culinary cultures around the world. The aim of this study was to investigate the potential of the bioactive compounds obtained for use in value-added products in the cosmetic, health and food sectors by analyzing the phytochemical content of *L. nobilis* leaves (headspace-GC-MS/MS) and methanol-chloroform extract (HPLC). When the phenolic content was analyzed by HPLC, gallic acid (48.77 ng/μL), alizarin (15.36 ng/μL) and naringin (14.11 ng/μL) were determined as the main components. In addition, eucalyptol (54.33%), sabinene (13.15%) and α -pinene (8.61%) were detected in high amounts of volatile compounds by headspace GC-MS/MS. These compounds are known for their numerous biological activities, such as antioxidant, antimicrobial and anti-inflammatory properties. The biological activities of these compounds increase the usability of the leaf and extract in various industrial applications such as wound healing creams, acne treatment products and natural flavor additives. They also show that the leaf and extract of *L. nobilis* can be used as an effective ingredient in cosmetic products that prevent skin aging and in food preservation systems.

Keywords: *Laurus nobilis*, laurel, phytochemical content, volatile compounds, phenolic compounds.

INTRODUCTION

Laurus nobilis (laurel) is an aromatic plant that is widespread in the Mediterranean region and is used in both gastronomy and medicine (Patrakar et al., 2012). In addition to its use as a spice in food, *L. nobilis* leaves is used in traditional medicine for its antiseptic, antibacterial, anti-inflammatory and digestive properties and is also being intensively studied by modern science for its various bioactive compounds (Siriken et al., 2018). The rich presence of phytochemicals such as phenolic acids, flavonoids and terpenes has been reported to increase the antioxidant and antimicrobial potential of *L. nobilis* leaves. *L. nobilis* leaves, thanks to its antioxidant properties, it supports the use of food to extend shelf life and in cosmetics, anti-ageing creams

and serums (Kıvrak et al., 2017). *L. nobilis* leaves are known for their antibacterial effect and can be used in cosmetics and products to treat acne. Thanks to its anti-inflammatory effect, it is also used in pharmacology for wound healing ointments and topical treatment (Chahal et al., 2017).

In recent years, the increasing demand for natural and bioactive compounds has brought the use of plant sources such as *L. nobilis* leaves to the fore as raw materials in food, cosmetics and health products (Awada et al., 2023). However, the effective use of these bioactive compounds requires accurate analytical characterization. Chromatographic methods such as HPLC, GC-MS, GC-MS/MS are very popular for both quantitative and qualitative analysis of phytochemical compounds (Braithwaite & Smith, 2012).

This study aims to determine the bioactive compounds in *L. nobilis* leaves and its extract by analyzing the content of phytochemicals with headspace GC-MS/MS (volatile components) and HPLC (phenols and flavonoids), and to discuss the potential applications of these compounds in different industries.

MATERIALS and METHODS

L. nobilis leaves were supplied by a local producer in the Defne district of Hatay in Türkiye, dried and ground into powder. 20 g of the powdered *L. nobilis* leaves (50%) was extracted with methyl alcohol (250 mL) and chloroform (250 mL) in a ratio of one to one. At the end of the extraction, the solvents were evaporated using a rotary evaporator. A portion of the *L. nobilis* leaf extract obtained was analyzed by high performance liquid chromatography (HPLC) for the profiles of phenolic and flavonoid compounds. The other extract sample was stored in a refrigerator at +4 °C to be used for the production of value-added products.

Analysis of phenolic compounds by HPLC

High performance liquid chromatography (HPLC) was used to determine the content of phenolic flavonoids in the methanol-chloroform extract of *L. nobilis* leaves. The instrument was equipped with a DAD (300/200 nm wavelength) detection and an 8 micron reversed phase Hi-plex (300x7.7) analytical column. The column temperature was 30 °C, eluent A; 83% water (0.1 formic acid) and eluent B; 17% acetonitrile (0.1 formic acid), the solvent flow rate was set to 0.8 mL/min. The injection volume was set to 10 µL. Sample preparation: 20 mg of the extract was weighed on a precision balance and dissolved in methanol. 1 mL of the sample was withdrawn using an automatic pipette and filtered through a 0.45 micron filter. It was diluted 1:1 with pure water and injected into the device (Başar et al., 2024b).

Analysis of the volatile components by GC-MS/MS

Analysis of volatile components in *L. nobilis* leaves was performed using an Agilent 7000 A GC/MS Triple Quad with 7890 GC, 7693 autosampler and 7697A headspace sampler. The instrument was equipped with an Agilent J&W HP-5 ms UltraInert 5% (phenyl)-methylpolysiloxane (30 m × 0.25 mm × 0.25 µm) GC column. The instrument oven temperature was set to 130 °C, the loop temperature to 140 °C, the transfer line temperature to 145 °C, and the vial equilibration time to 30 min. GC conditions: The initial column oven temperature was set to 60 °C and then ramped to 220 °C at a rate of 4 °C/min and held for 10 min. Components were qualitatively determined using the NIST library and NIST MS Search 2.2 libraries (Başar et al., 2024a; Gül et al., 2024).

RESULT and DISCUSSION

L. nobilis leaves are a rich source of volatile constituents, phenolic and flavonoid compounds and thanks to these properties it is becoming a very important ingredient for value-added products in the cosmetic, health and food industries. In our study, the bioavailability and mechanisms of

action of the bioactive compounds in *L. nobilis* leaves were analyzed by HPLC and GC-MS/MS to investigate them in detail.

HPLC analysis results

In the HPLC analysis of the *L. nobilis* leaves methanol-chloroform extract, screening was performed using 20 standards. According to the result of the analysis, 13 compounds were determined (Figure 1). Accordingly, gallic acid (48.7703 ng/uL), ascorbic acid (19.6933 ng/uL), alizarin (15.3591 ng/uL) and naringin (14.1149 ng/uL) were detected in the highest amounts (Table 1). The phenolic content of the ethanol extract of *L. nobilis* leaves was analyzed by UPLC-MS/MS.

Table 1. Results of the HPLC analysis of the methanol-chloroform extract of the leaves of *L. nobilis*

No	R.T. [min]	m/z (g/mol)	Formula	Compound Name	Amount[ng/ul]
1	3.310	176.12	C ₆ H ₈ O ₆	Ascorbic acid	19.6933
2	4.267	170,12	C ₇ H ₆ O ₅	Gallic acid	48.7703
3	5.811	154.12	C ₇ H ₆ O ₄	Protocatechuic acid	0.5394
4	6.318	290.26	C ₁₅ H ₁₄ O ₆	Catechin	-
5	7.953	138.12	C ₇ H ₆ O ₃	Hydroxybenzoic acid	2.2179
6	10.000	168.14	C ₈ H ₈ O ₄	Vanillic Acid	-
7	10.455	154.12	C ₇ H ₆ O ₄	Gentisic Acid	-
8	16.796	164,04	C ₉ H ₈ O ₃	p-Coumaric acid	0.1647
9	17.889	610.5	C ₂₇ H ₃₀ O ₁₆	Rutin	1.9394
10	20.223	194.18	C ₁₀ H ₁₀ O ₄	Ferulic Acid	-
11	27.165	580.5	C ₂₇ H ₃₂ O ₁₄	Naringin	14.1149
12	28.810	164.16	C ₉ H ₈ O ₃	o-Coumaric acid	0.3357
13	29.581	610.6	C ₂₈ H ₃₄ O ₁₅	Neohesperidin	-
14	30.805	146.14	C ₉ H ₆ O ₂	Coumarin	-
15	32.238	228.24	C ₁₄ H ₁₂ O ₃	Resveratrol	0.4490
16	35.017	302,236	C ₁₅ H ₁₀ O ₇	Quarceetin	0.3013
17	35.480	148.16	C ₉ H ₈ O ₂	Trans-sinamic acid	-
18	36.616	610.6	C ₂₈ H ₃₄ O ₁₅	Hesperidin	1.0634
19	38.523	240.21	C ₁₄ H ₈ O ₄	Alizarin	15.3591
20	39.686	222.24	C ₁₅ H ₁₀ O ₂	Flavone	3.9411

According to the analysis results, rosmarinic acid, syringic acid, gallic acid, protocatechuic acid, chlorogenic acid, caffeic acid, p-coumaric acid, ferulic acid and gallic acid were detected (Bachir-bey et al., 2023; Dobrosłavić et al., 2021). Our results are consistent with the literature. In the analyses, the contents may vary depending on the harvest time, altitude, soil structure and the standards used in the analyses.




Figure 1. HPLC chromatogram of the methanol-chloroform extract of the leaves of *L. nobilis*

Determination of the content of volatile components of the leaves of *L. nobilis* by GC-MS/MS

The content of volatile compounds in the leaves of *L. nobilis* was analyzed by headspace GC-MS/MS (Figure 2). During the analysis, 30 volatile compounds were detected. According to the analytical result, eucalyptol (1,8-cineole) (54.33%), sabinene (13.15%), α -pinene (8.61%), β -pinene (5.35%), α -terpineol acetate (3.20%) and α -terpineol (3.16%) were detected in the highest amounts (Table 2).

Table 2. Content of volatile compounds in *L. nobilis* leaves by headspace GC-MS/MS

No	RT (min.)	RI (NIST lib.)	Compound Name	%
1	8,79	925	Tricyclene	1,44
2	9,03	929	α -Pinene	8,61
3	10,40	974	Sabinene	13,15
4	10,51	979	β -Pinene	5,35
5	10,95	991	β -Myrcene	1,29
6	11,45	1005	α -Phellandrene	0,34
7	11,89	1017	α -Terpinene	0,63
8	12,22	1022	o-Cymene	0,98
9	12,58	1032	Eucalyptol	54,33
10	13,42	1060	γ -Terpinene	1,09
11	13,73	1070	cis-Sabinene hydrate	0,26
12	14,50	1086	Isoterpinolene	0,32
13	14,89	1099	Linalool	1,40
14	16,37	1139	trans-Pinocarveol	0,24
15	17,36	1166	δ -Terpineol	0,73
16	17,74	1177	4-Terpineol	1,55
17	18,23	1189	α -Terpineol	3,16
18	18,42	1193	Myrtenal	0,14
19	19,47	1228	Nerol	0,12
20	20,38	1257	Bergamol	0,22
21	21,47	1286	Isobornyl acetate	0,08
22	21,66	1294	2-Undecanone	0,15
23	22,49	1315	δ -Terpineol, acetate	0,18
24	23,57	1350	α -Terpineol acetate	3,20
25	23,82	1357	Eugenol	0,22
26	24,96	1391	β -Elemene	0,27
27	25,27	1402	Methyleugenol	0,35
28	25,88	1419	Caryophyllene	0,10
29	30,61	1576	Spathulenol	0,08
30	30,79	1581	Caryophyllene oxide	0,04

Composition of the essential oil of *L. nobilis* In the GC-MS/MS analysis, eucalyptus (36.8%), α -terpinyl acetate (14.6%), α -pinene (4.6%), methyl eugenol (4.2%), (Z)-3-hexanol (3.8%), β -pinene (3.6%), sabinene (3.1%), terpinen-4-ol (6.4%) were determined as the main components (Kasabri et al., 2013). The results of our study appear to be generally consistent with the literature.




Figure 2. Headspace GC-MS/MS chromatogram of volatile compounds from *L. nobilis* leaves

CONCLUSION

In this study, conducted on the leaves of *L. nobilis*, the rich phytochemical profile of the plant was analyzed in detail and the results showed that laurel extract has great potential as a valuable ingredient in the cosmetic, health and food industries. The results of the HPLC analysis showed that bay leaf extract contains high concentrations of phenolic compounds, especially gallic acid and alizarin, in significant amounts. The strong antioxidant properties of these compounds suggest that they can be used in cosmetic products and food preservatives that prevent skin aging. In addition, it is believed that the extract has the potential to be used in the treatment of metabolic disorders associated with diabetes due to its insulin sensitivity enhancing effects. GC-MS/MS analysis showed that the laurel extract has a high content of terpenoids such as eucalyptol, sabinene and α -pinene in the form of volatile compounds. These volatile compounds can be used in topical treatments, wound healing ointments and anti-acne products thanks to their antimicrobial and anti-inflammatory properties. In addition, these components can be evaluated as a natural additive in the food industry due to their aromatic and flavor-enhancing effects.

The rich content of bioactive components in *L. nobilis* extract therefore makes it an extremely valuable natural resource for use in modern health and cosmetic products. This study underlines that bay leaf should be supported by optimized extraction methods and clinical studies to expand its industrial applications. With its rich phenolic and volatile constituents, *L. nobilis* leaves extract is attracting attention as a natural and bioactive raw material. While this study highlights the potential use of bay laurel in modern health, cosmetic and food products, it shows the need for more advanced extraction methods and clinical studies. The results of the analysis suggest that laurel extract should be developed for use in high value-added products.

Acknowledgments

The authors would like to acknowledge their contributions to the project numbered YiP0723i08 entitled "Production of functional food, health and cosmetic products from plant-based raw materials" conducted at Iğdır University.

REFERENCES

Awada, F., Hamade, K., Kassir, M., Hammoud, Z., Mesnard, F., Rammal, H., & Fliniaux, O. (2023). *Laurus nobilis* leaves and fruits: a review of metabolite composition and interest in human health. *Applied Sciences*, 13(7), 4606.

- Bachir-bey, M., Khaled Khodja, Y., Belmouhoub, M., Ladjouzi, R., Dahmoune, F., & Khettal, B. (2023). The botanical study, phytochemical composition, and biological activities of *Laurus nobilis* L. leaves: A review [The botanical study, phytochemical composition, and biological activities of *Laurus nobilis* L. leaves: A review]. *International Journal of Secondary Metabolite*, 10(2), 269-296. <https://doi.org/10.21448/ijsm.1171836>
- Başar, Y., Gül, F., Karadağ, M., Alma, M. H., Demirtas, İ., & Tel, A. Z. (2024a). Determination of the volatile, phenolic and fatty acid contents of *Helichrysum plicatum* by chromatographic methods. *Turkish Journal of Biodiversity*, 7(2), 83-94.
- Başar, Y., Gül, F., Nas, M. S., Alma, M. H., & Çalimli, M. H. (2024b). Investigation of value-added compounds derived from oak wood using hydrothermal processing techniques and comprehensive analytical approaches (HPLC, GC-MS, FT-IR, and NMR). *International Journal of Chemistry and Technology*, 8(1), 53-61.
- Braithwaite, A., & Smith, J. (2012). *Chromatographic methods*. Springer Science & Business Media.
- Chahal, K., Kaur, M., Bhardwaj, U., Singla, N., & Kaur, A. (2017). A review on chemistry and biological activities of *Laurus nobilis* L. essential oil. *Journal of Pharmacognosy and Phytochemistry*, 6(4), 1153-1161.
- Dobrosłavić, E., Elez Garofulić, I., Zorić, Z., Pedisić, S., & Dragović-Uzelac, V. (2021). Polyphenolic Characterization and Antioxidant Capacity of *Laurus nobilis* L. Leaf Extracts Obtained by Green and Conventional Extraction Techniques. *Processes*, 9(10), 1840. <https://www.mdpi.com/2227-9717/9/10/1840>
- Gül, F., Başar, Y., & Demirtas, İ. (2024). Phytochemical Content and In Silico Molecular Docking Studies of *Achillea biebersteinii* and *A. millefolium* Plants. *Bütünleyici ve Anadolu Tıbbi Dergisi*, 5(2), 10-24.
- Kasabri, V., Abu-Dahab, R., & Afifi, F. (2013). "Evaluation of the volatile oil composition and antiproliferative activity of *Laurus nobilis* L. (Lauraceae) on breast cancer cell line models". *Records of Natural Products*.
- Kıvrak, Ş., Göktürk, T., & Kıvrak, İ. (2017). Assessment of volatile oil composition, phenolics and antioxidant activity of bay (*Laurus nobilis*) leaf and usage in cosmetic applications. *International Journal of Secondary Metabolite*, 4(2), 148-161.
- Patrakar, R., Mansuriya, M., & Patil, P. (2012). Phytochemical and pharmacological review on *Laurus nobilis*. *International journal of pharmaceutical and chemical sciences*, 1(2), 595-602.
- Sırıken, B., Yavuz, C., & Güler, A. (2018). Antibacterial Activity of *Laurus nobilis*: A review of literature. *Medical Science and Discovery*, 5(11), 374-379.

BIOLOGICAL CONTROL OF STORED PRODUCT DAMAGE

DEPOLANMIŞ ÜRÜN ZARARLILARIYLA BİYOLOJİK MÜCADELE

Gönül USTA GEBEŞ

University of Iğdır, Faculty of Agriculture, Department of Plant Protection, Iğdır, Turkey

ORCID ID: <https://orcid.org/0000-0002-4961-4658>

Celalettin GÖZÜAÇIK

University of Iğdır, Faculty of Agriculture, Department of Plant Protection, Iğdır, Turkey

ORCID ID: <https://orcid.org/0000-0002-6543-7663>

ABSTRACT

Food safety and security is one of the biggest concerns in today's world. Preservation and safe storage of grains are essential for timely delivery to consumers. Different food commodities such as harvested legumes and grains, processed plant and animal food products and semi-perishable products need to be stored safely in order to be used for domestic or commercial purposes. During long-term storage, stored products are prone to contamination, increased temperature, mold and damage by biotic and abiotic factors. Among the biotic factors, insects, mites, rodents, birds and microorganisms cause tremendous losses in storage. Our aim is to protect these stored product pests by using biological agents (such as predators and parasitoids) with the help of a natural method called biological control. Scientific studies conducted in our country and worldwide within the scope of biological control on stored product pests were evaluated. Stored product pests are insects, mites and other organisms that are released during the storage of agricultural products and cause product losses. These harmful organisms significantly reduce the quality of agricultural products and cause economic losses. Traditional control methods, chemical control drugs, have negative effects on the environment, animal and human health. Storage pests also develop resistance to chemical drugs. Therefore, the importance of biological control methods continues to increase. In this report, biological control methods applied against storage pests and the advantages of these methods were evaluated.

Key Words: Biological Control, Store, Pest, Predator

ÖZET

Gıda güvenliği ve emniyeti günümüz dünyasının en büyük endişelerinden biridir. Tahılların muhafazası ve güvenli bir şekilde depolanması, tüketicilere zamanında ulaştırılması için elzemdir. Hasat edilmiş baklagiller ve tahıllar, işlenmiş bitki ve hayvan gıda ürünleri ve yarı bozulabilir ürünler gibi farklı gıda emtialarının, evsel veya ticari amaçlarla kullanılabilmesi için güvenli bir şekilde depolanması gerekir. Uzun süreli depolama süresince biyotik ve abiyotik faktörler tarafından depolanmış ürünler kirlenmeye ve hasara eğilimlidir. Biyotik etkenler arasında böcekler, akarlar, kemirgenler, kuşlar ve mikroorganizmalar depolamada muazzam bir kayba neden olur. Depolanmış ürün zararlıları, tarım ürünlerinin depolanması sırasında açığa çıkan ve ürün kayıplarına sebep olan böcekler, akarlar ve diğer organizmalardır. Bu zararlı organizmalar, tarım ürünlerinin kalitesini ciddi oranda düşürerek ekonomik kayıplara

yol açmaktadır. Geleneksel mücadele yöntemi olan kimyasal mücadele ilaçlarının çevre, hayvan ve insan sağlığı üzerinde olumsuz etkileri bulunmaktadır. Depo zararlıları da kimyasal ilaçlara karşı direnç geliştirmektedir. Bu nedenle, biyolojik mücadele yöntemlerinin önemi artarak devam etmektedir. Bu bildiride, depo zararlılarına karşı uygulaması yapılan biyolojik mücadele yöntemleri ve bu yöntemlerin avantajları değerlendirilmiştir.

Anahtar kelimeler: Biyolojik Mücadele, Depo, Zararlı, Predatör

GİRİŞ

Depolanmış ürün zararlıları, başta tahıl, baklagil, kurutulmuş sebze ve meyveler gibi tarım ürünlerinin depolanması esnasında ürünlerin besin değerini düşürmekte, bakteri ve küf oluşumuna neden olmakta ve ticari değerini azaltmaktadır. Bu zararlılar geleneksel olarak kullanılan kimyasal pestisitler ile kontrol edilmeye çalışılsa da bu mücadelenin çevremize ve insan sağlığı üzerinde görülen olumsuz etkileri nedeniyle alternatif çözüm yolları aranmaktadır.

Uygun iklim koşulları nedeniyle ülkemiz fazla sayıda depo zararlısının gelişimi için elverişli ortamlar oluşturmaktadır. Depolara koyulmuş ürünlerde görülen zararlılar yaklaşık %10 ağırlık kayıplarına neden olmaktadır (Prewet, 1975; Esther et al., 2014). Depo zararlısı olan böceklerin en temel sınıflandırması beslenme alışkanlıklarına, yani "birincil zararlılar" ve "ikincil zararlılar"a dayanmaktadır (Srivastava & Subramanian, 2016). Tahıllarda bulunan birincil zararlılar arasında *Sitophilus* spp., *Trogoderma granarium* Everts, *Sitotroga cerealella* Olivier and *Rhyzopertha dominica* Fabricius görülmekte ve sağlam tahıl danelerinde zarara neden olmaktadır (Reichmuth, 2000a; Donnelly & Phillips, 2001a; Sarker et al., 2019). Sekonder zararlılar ise kırıklı veya daha öncesinde zarar görmüş danelerde beslenebilmektedirler.

Depo zararlıları ile mücadelede dünyada ve ülkemizde genel olarak insektisitler kullanılmaktadır. Depolama alanlarında kullanılan birçok pestisite karşı zararlı böceklerin direnç geliştirdiği görülmektedir (Arthur 1996). Kimyasal mücadele kapsamında en sık kullanılan yöntemlerden biri hızlı, az maliyetli ve etkili çözüm sağlayan fumigasyondur (Zettler and Arthur, 2000).

Dünyada, kısmi bir şekilde başarılı olan fumigasyon uygulamalarından sonra canlı kalan böcekler, fosfine karşı direnç kazanarak popülasyonlarını giderek arttırmışlardır (Zettler, 1996; Tyler et al., 1983; Bell, 2000).

Havalandırma ve hermetik depolama ülkemizde kimyasal mücadele dışında uygulanan en yaygın yöntemdir. Hermetik depolama da depo içerisine giren oksijen miktarı ile zararlı böcekler tarafından tüketilen oksijenin eşit olduğu koşullar oluşturulur ve bu depo ortamı için tam gaz geçirmezlik olması gerekmektedir (Ferizli & Emekçi, 1999).

Havalandırma, tahılların genel konveksiyonel havalanmasının tersi yönünde olacak şekilde bir güç yardımı ile yapılan bir uygulama türüdür. Bu yöntem ile tahılın ısısı düşürülerek zararlı böceğin gelişimi engellenmektedir ve entegre mücadele kapsamında uygulanmaktadır. Isı 20 °C den düşük olursa zararlı gelişimi yavaşlar, 10 °C' de durur. Organik ürünlerin depolanmasında ülkemizde uygulanan diğer bir yöntem, -10 ila -18 °C'de ürünlerin tutularak zararlı böceklerin öldürüldüğü düşük sıcaklık uygulamasıdır. (Sarıyörük & Köseoğlu, 1987; Bülbül, 1993; Ferizli et al. 2004).

Biyolojik mücadele, zararlı organizmaların popülasyonlarını doğal düşmanları (parazitoitler, predatörler ve patojenler) kullanılarak kontrol altına alma ve düzenlemedir. Bu yöntem, zararlıların popülasyon yoğunluklarını doğal yöntemler ile düşürerek, kimyasal pestisit kullanımını en aza indirmeyi hedefler. Bu mücadele çeşidi, sürdürülebilir tarım uygulamalarının önemli bir parçası olarak yer almaktadır.

Depolanmış ürün zararlılarına karşı biyolojik mücadele çalışmaları

Depolanmış ürün böcek zararlılarına karşı böcek parazitoitlerinin ve predatörlerinin kullanımı zaman zaman çeşitli yazarlar tarafından incelenmiştir (Matthias, 2010; Flinn ve Scholler, 2012). Entegre zararlı yönetimi yaklaşımındaki ilerlemeden sonra, biyolojik kontrol en sürdürülebilir ve önemli bileşenlerden biri haline gelmiştir. Depolanmış tahıllardaki böcek ilacı kalıntıları, organik tarıma yönelik yaklaşım, gıda güvenliği biyolojik kontrole olan ilgiyi artırmıştır (Flinn ve Scholler, 2012). Entegre mücadele kapsamında, biyolojik mücadele için çok sayıda parazitoit ve predatör depolanmış ürün zararlılarında kullanılmaktadır (Schöller & Flinn, 2000; Schöller et al., 2006). Solà (2018) tarafından depolanmış ürünlerde veya gıda endüstrisinde biyolojik kontrol, sırasıyla %51, %21 ve %18'i parazitoit, entomopatojen ve avcıydı. Akarlar (%56) ve heteropteranlar (%24) başlıca avcılardı. Parazitoitlerin, pteromalidler ve braconidler (%70) ve ardından bethylidler, ichneumonidler ve trichogrammatidler (%24) ve chalcididler ve eulopidler (%3) yoğunlukta olduğunu belirtmiştir

Predatör

Predatörler, depolanmış ürün zararlılarını avlayarak popülasyonlarını kontrol altına alır. Depolarda kullanılan bazı predatörler, *Xylocoris flavipes* (Reuter) (Hemiptera: Anthocoridae), *Lyctocoris* spp (Hemiptera: Anthocoridae), *Amphibolus wenator* (Klug) (Hemiptera: Reduviidae) (LeCato et al., 1977; Brower & Mullen, 1990; Ahmed et al., 1991; Brower & Press, 1992; Sing & Agrobast, 2008; Imamura et al., 2008; Rahman et al., 2009; Upadhyay, & Ahmad, 2011; Saha et al., 2012; Ballal et al., 2013; Schöller, 2014; Gözüaçık et al., 2019; Saikia & Borkakati, 2020). Reichmuth (2000b) tarafından *Stegobium paniceum* Linnaeus, *Acanthoscelides obtectus* Say, *Sitophilus zeamais* Motschulsky, *Sitophilus granarius* Linnaeus, *Tribolium confusum*, *Tribolium castaneum*, *Lasioderma serricorne* Fabricius, *Plodia interpunctella* Hubner, *S. cerealella*, *Ephestia cautella* Walker, *Corcyra cephalonica* Stainton depolanmış ürün zararlılarının yer aldığı ortamlarda *X. flavipes* predatöründe tespit edilmiştir. Donnelly and Phillips (2001b), yaptıkları çalışmalarında buğday taneleri ilave edilmiş cam kaplar ve boş cam kaplardan oluşan farklı iki yaşam ortamında, *X. flavipes*'in farklı oranlarda olan *O. surinamensis*, *T. castaneum*, *R. dominica* ve *P. interpunctella* zararlılarının fonksiyonel tepkilerini incelemiştir. *X. flavipes*'in biyolojisi, preovipozisyon, ovipozisyon, postovipozisyon süreleri ve *T. confusum*, *E. kuehniella*, *T. granarium* depo zararlılarında av tercihlerini belirleyen çalışma yürütülmüştür (Usta Gebeş, & Gözüaçık, 2024).

Parazitoit

Parazitoitler, zararlıların yumurtalarına veya larvalarına kendi yumurtalarını bırakarak zararlının ölmesine neden olan organizmalardır. *Bracon hebetor* Say (Hymenoptera: Braconidae), *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae), *Anisopteromalus claudrae* (Howard) (Hymenoptera: Pteromalidae), *Antrocephalus* spp (Hymenoptera: Chalcididae), *Trichogramma deion* Riley (Hymenoptera: Trichogrammatidae), parazitoitleri depo zararlılarına karşı kullanılmaktadır (Grieshop et al., 2006; Upadhyay & Ahmad, 2011; Press et al., 1974). El-Gawad et al. (2009), *Anisopteromalus calandrae* (Howard) parazitoitinin doğrudan salınmasının (torbaların içinde) böcek popülasyonları üzerindeki etkisini incelerken, *Callosobruchus maculatus* Fabricius, *Callosobruchus chinensis* Linnaeus'ın bakla fasulyesi tohumları, *R. dominica* ve *Sitophilus oryzae* Linnaeus'ın buğday taneleri üzerindeki etkisini araştırdı. Sonuçlar, *A. calandrae*'nin salınmasının bakla fasulyesi üzerindeki test edilen böceklerin aylık artış oranını %35,20 ile %42,14 arasında azalttığını gösterdi. Bu oranlar, 6 aylık depolama süresinin sonunda popülasyonda %31,24 ile %36,7 arasında azalmaya yol açtı. Reppchen et al. (2003), *Lariophagus distinguendus* (Forster) (Hymenoptera: Pteromalidae) biyolojik ajanını *S. granaries* popülasyon büyümesini baskılama yeteneğini yedi aylık bir süre boyunca incelemiştir. Serbest bırakılan parazitoit sayısından bağımsız olarak, *S. granarius*'un popülasyon büyüklüğü tüm süre boyunca %79-94 oranında azalmıştır.

Entomopatojenler

Entomopatojen bakteriler, funguslar ve nematodlar, zararlı olan böcekleri enfekte edip öldüren mikroorganizmalardır. *Beauveria bassiana* (Balsamo), *Lecanicillium lecanii* (Zimm.), *Metarhizium anisopliae* (Metchnikoff) and *Paecilomyces farinosus* (Holmsk.) fungusları depo zararlılarına karşı kullanılmaktadır (Büda & Pečiulytė, 2008). Ticari olarak temin edilebilen entomopatojenik funguslardan bazıları *B. bassiana*, *M. anisopliae* ve bakteri *Bacillus thuringiensis* (Berliner) olup, esas olarak depolanmış tahıl zararlılarına, özellikle böceklere karşı test edilmiştir (Batta, 2016). *B. thuringiensis* var. *kurstaki*'nin depolanmış tahıl zararlıları üzerinde etkili olduğu bildirilmiştir ((Díaz-Gomez et al., 2000; Shapiro-Ilan et al., 2007). Özellikle *P. interpunctella* ve diğer güveler de etkili olduğunu bildirmişlerdir (Lord et al., 2007). Entomopatojenlerin, özellikle fungusların kullanımı depolama koşullarında umut verici olarak değerlendirilmiştir. Rumbos & Athanassiou (2017a), depolanmış zararlı yönetiminde kullanılan entomopatojenik fungusları vurgulamıştır.

Diğer patojenler, örneğin protozoalar ve virüsler de depolanmış tahıl böcek zararlılarına karşı test edilmiştir. Protozoalar, yani *Nosema whitei* (Milner, 1973) ve *Nosema plodiae* (Kellen & Lindegren, 1974), un böceklerini istila ettiğini uzun zamandır bildirilmişlerdir. Benzer şekilde, virüsler (çoğunlukla bakulovirüsler) de *P. interpunctella* ve *E. cautella* gibi lepidoptera zararlılarında büyük ölçüde incelenmiştir (Vail et al., 1991).

Nematod bilimi ilerledikçe, böcekleri kontrol etmek için entomopatojenik nematodların kullanımı zararlı yönetiminde yeni bir yaklaşım haline gelmektedir (Arthurs et al., 2004). *Steinernema* ve *Heterorhabditis* isimli iki önemli cinsin tarla mahsulü zararlılarına karşı etkili olduğu kanıtlanmış ve ayrıca incelenerek depolanmış tahıl böceklerine karşı da etkili olduğu bulunmuştur (Lacey et al., 2000; Vashisth et al., 2013; Saleh et al., 2015; Yurt et al., 2015; Gözel, 2019; Yan et al., 2020 Rumbos ve Athanassiou, 2017b; Gözel, 2019; Yan et al., 2020). Athanassiou et al. (2010), laboratuvar koşulları altında depolanmış buğdayda bulunan dört zararlı böceğe (*E. kuehniella*, *R. dominica*, *S. oryzae*, *T. confusum*) karşı üç nematod türünün (*Heterorhabditis bacteriophora* Poinar, *Steinernema carpocapsae* Weiser ve *Steinernema feltiae* Filipjev) etkinliğini test etmiştir. Ülkemizde izole edilen entomopatojen nematodların (EPN) laboratuvarında patates güvesi *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae) son dönem larvalarına karşı etkinlikleri çalışılmıştır. Uygulamada *H. bacteriophora*, *Steinernema affine* Bovien, *S. feltiae* ve *S. carpocapsae* EPN türleri kullanılmıştır. Çalışmada kullanılan EPN türleri zamana bağlı olarak *P. operculella* larvalarında yüksek düzeyde ölüm sağlamıştır (Gözel et al., 2020). Az yoğunluktaki nematod uygulamaları yapılan başka bir çalışmada ise, birinci larva döneminde cm² 'ye 5 ve 10 IJ yoğunluğundaki *H. bacteriophora* ve *S. carpocapsae*'nin %98-100 ölüme sebep olduğu tespit edilmiş (Moawad et al., 2018).

Biyolojik kontrol kullanılarak depolanmış tahılın korunması, çevre sağlığına zarar vermeden potansiyel bir araç/tekniktir (Matthias, 2010). Bir dizi böcek patojeni, laboratuvar koşullarında depolanmış tahıl böceklerine karşı test edilmiştir (Batta & Kavallieratos, 2018; Kumar et al., 2019).

SONUÇ

Ham ürünlerden tüketiciye kadar tüm depolama zinciri boyunca ürünleri istiladan uzak tutmak için entegre bir yönetim stratejisine ihtiyaç vardır. Biyolojik kontrol bu stratejinin bir parçasıdır (Schöller, 2010). Birçok zararlı türünün böcek ilaçlarına karşı direncinin artması, kayıtlı aktif bileşik sayısının azalması ve çevre üzerindeki olumsuz etkiler de depolanmış ürün zararlı kontrolünde toksik bileşiklerin kullanımına alternatifler geliştirmek için önemli nedenlerdir (Riudavets, 2018).

Depo zararlılarına karşı biyolojik mücadele yöntemleri, insan sağlığına zararı olmayan, çevre dostu ve sürdürülebilir bir çözüm sunmaktadır. Bu yöntemler, kimyasal kullanımını azaltarak

tarımsal ürünlerin kalitesini korumakta ve ekonomik kayıpları engellemektedir. Biyolojik mücadele, önümüzdeki yıllarda depo zararlılarıyla mücadelede önemini arttırarak devam ettirecektir.

REFERENCES

- AI Now Institute, (2017). *AI Now Report 2017*. Erişim adresi https://ainowinstitute.org/AI_Now_2017_Report. Son Erişim 26 Ekim 2021.
- Ahmed, K. N., Khatun, M., & Rahman, M. M. (1991). Biological notes on *Xylocoris flavipes* (Reuter)(Hemiptera: Anthocoridae). *Journal Asiatic Society of Bangladesh Sci*, 17, 65-67.
- Arthurs, S., Heinz, K. M., & Prasifka, J. R. (2004). An analysis of using entomopathogenic nematodes against above-ground pests. *Bulletin of Entomological Research*, 94(4), 297-306. doi: 10.1079/BER2003309
- Arthur, F. H. (1996). Grain protectants: current status and prospects for the future. *Journal of Stored Products Research*, 32(4), 293-302.
- Athanassiou, C. G., Kavallieratos, N. G., Menti, H., & Karanastasi, E. (2010). Mortality of four stored product pests in stored wheat when exposed to doses of three entomopathogenic nematodes. *Journal of Economic Entomology*, 103(3), 977-984.
- Ballal, C. R., Gupta, T., & Joshi, S. (2013). Production and evaluation of the warehouse pirate bug *Xylocoris flavipes* (Reuter)(Hemiptera: Anthocoridae). In *Presented during the International Congress on Insect Science*. 2013, 14-17 February, GKVK, Bangalore.
- Batta, Y. A. (2016). Recent advances in formulation and application of entomopathogenic fungi for biocontrol of stored-grain insects. *Biocontrol Science and Technology*, 26(9), 1171-1183. doi: 10.1080/09583157.2016.1201458
- Batta, Y. A., & Kavallieratos, N. G. (2018). The use of entomopathogenic fungi for the control of stored-grain insects. *International Journal of Pest Management*, 64(1), 77-87. doi: 10.1080/09670874.2017.1329565
- Bell, C. H. (2000). Fumigation in the 21st century. *Crop protection*, 19(8-10), 563-569.
- Brower, J. H., & Mullen, M. A. (1990). Effects of *Xylocoris flavipes* (Hemiptera: Anthocoridae) releases on moth populations in experimental peanut storages. *Journal of Entomological Science*, 25(2), 268-276.
- Brower, J. H., & Press, J. W. (1992). Suppression of residual populations of stored-product pests in empty corn bins by releasing the predator *Xylocoris flavipes* (Reuter). *Biological Control*, 2(1), 66-72.
- Būda, V., & Pečiulytė, D. (2008). Pathogenicity of four fungal species to Indian meal moth *Plodia interpunctella* (Hübner)(Lepidoptera: Pyralidae). *Ekologija*, 54(4).
- Bülbül, S. 1(993). Researchs on the possibilities of the use of cooling technique for the controlling of the dried fig moth, (*Cadra cautella* Wlk. (Lepidoptera:Pyralidae) (in Turkish). Ph.D thesis. Ege University, Graduate School. 114 p.
- Díaz-Gomez, O., Rodríguez, J. C., Shelton, A. M., Lagunes-T, A., & Bujanos-M, R. (2000). Susceptibility of *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) populations in Mexico to commercial formulations of *Bacillus thuringiensis*. *Journal of Economic Entomology*, 93(3), 963-970.
- Donnelly, B. E., & Phillips, T. W. (2001a,b). Functional response of *Xylocoris flavipes* (Hemiptera: Anthocoridae)-effects of prey species and habitat. *Environmental Entomology*, 30(3), 617-624.

- El-Gawad, A., Hany, A. S., El-Aziz, A., & Sayed, A. M. (2009). Effect of releasing the parasitoid *Anisopteromalus calandrae* (Hymenoptera: Pteromalidae) on certain coleopteran stored products beetles in Egypt. *Egyptian Academic Journal of Biological Sciences. A, Entomology*, 2(2), 211-219.
- Emekci, M., & Ferizli, A. G. (2000). Current status of stored products protection in Turkey. *IOBC wprs Bulletin*, 23(10), 39-46.
- Esther, M., Sharon, M., Abirami, C. V., & Alagusundaram, K. (2014). Grain storage management in India. *Journal of Postharvest Technology*, 2, 12–14. Available online at: <http://jph.t.in/MenuScriptFile/03b17047-fa10-48ed-b0cb-95fb9222798f.pdf>
- Ferizli, A.G., & Emekçi, M. (1999). An alternative and environmentally safe storage method: hermetic storage (in Turkish). Orta Anadolu’ da Hububat Tarımının Problemleri ve Çözüm Yolları Sempozyumu, 8-11 Haziran 1999.
- Ferizli, A.G., Emekci, M., Tütüncü, S., & Navarro, S. (2004). The efficacy of phosphine fumigation against dried fruit pests in Turkey. *IOBC WPRS (OILB SROP) Integrated Protection of Stored Products*, 27(9) 265-269, Turkey, September 16-19, 2003.
- Gözel, Ç. (2019). Yerel entomopatojen nematodların Amerikan beyaz kelebeği *Hyphantria cunea* (Drury)(Lepidoptera: Arctiidae) üzerindeki etkinliği. *Türkiye Biyolojik Mücadele Dergisi*, 10(1), 17-28.
- Gözel, Ç., Hürkan, A., & Gözel, U. (2020). Bazı entomopatojen nematodların etkinliğinin patates güvesi *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae) ’nin mücadelesinde değerlendirilmesi *Türkiye Biyolojik Mücadele Dergisi*, 11: 165-173.
- Gözüaçık, C., Konuksal, A., Karaca, C. and Çerçi, B., (2019). A Preliminary Study on the Predator, *Xylocoris flavipes* (Reuter) (Hemiptera: Anthocoridae) of Insect Pests of Stored in Northern Cyprus. International Agricultural Congress of Muş Plain, (24-27 September 2019), Bildiriler kitabı, sayfa 217-222.
- Grieshop, M. J., Flinn, P. W., & Nechols, J. R. (2006). Biological control of Indianmeal moth (Lepidoptera: Pyralidae) on finished stored products using egg and larval parasitoids. *Journal of Economic Entomology*, 99(4), 1080-1084.
- Imamura, T., Murata, M., & Miyanoshita, A. (2008). Biological aspects and predatory abilities of hemipterans attacking stored-product insects. *Japan Agricultural Research Quarterly: JARQ*, 42(1), 1-6.
- Kellen, W. R., & Lindegren, J. E. (1974). Comparative virulence of *Nosema plodiae* and *Nosema heterosporum* in the indian meal moth, *Plodia interpunctella*. *J. Invertebr. Pathol.* 23, 242–245. doi: 10.1016/0022-2011(74)9 0191-8
- Kumar, K. K., Sridhar, J., Murali-Baskaran, R. K., Senthil-Nathan, S., Kaushal, P., Dara, S. K., et al. (2019). Microbial biopesticides for insect pest management in India: current status and future prospects. *J. Invertebr. Pathol.* 165, 74–81. doi: 10.1016/j.jip.2018.1 0.008
- Lacey, L. A., Knight, A., & Huber, J. (2000). Microbial control of lepidopteran pests of apple orchards. In *Field Manual of Techniques in Invertebrate Pathology: Application and evaluation of pathogens for control of insects and other invertebrate pests* (pp. 557-576). Dordrecht: Springer Netherlands.
- LeCato, G. L., Collins, J. M., & Arbogast, R. T. (1977). Reduction of residual populations of stored-product insects by *Xylocoris flavipes* (Hemiptera: Anthocoridae). *Journal of the Kansas Entomological Society*, 84-88.
- Lord, J. C., Campbell, J. F., Sedlacek, J. D., & Vail*, P. V. (2007). Application and evaluation of entomopathogens for managing insects in stored products. ,” in *Field Manual of Techniques*

- in *Invertebrate Pathology*, eds L. A. Lacey and H. K. Kaya (Frankfort: Springer), 677–693. doi: 10.1007/978-1-4020-5933-9_33
- Matthias, S. (2010). “Prospects for biological control of stored-product pests,” in *International European Symposium on Stored Product Protection “Stress on Chemical Products,”* (January) (Germany), 25–31.
- Moawad S.S., M.M.E. Salah, H.M. Metwally, I.M. Ebadah & Y.A. Mahmoud 2018. Protective and curative treatments of entomopathogenic nematodes against the potato tuber moth, *Phthorimaea operculella* (Zell.). *Bioscience Research*, 15(3): 2602-2610
- Press, J. W., Flaherty, B. R., & Arbogast, R. T. (1974). Interactions among *Plodia interpunctella*, *Bracon hebetor*, and *Xylocoris flavipes*. *Environmental Entomology*, 3(1), 183-184.
- Prevett, P. F. (1975). Stored product pests causing losses of stored food. *FAO Plant Protection Bulletin*, 23, 115-117.
- Rahman, M. M., Islam, W., & Ahmed, K. N. (2009). Functional response of the predator *Xylocoris flavipes* to three stored product insect pests. *International Journal of Agriculture and Biology*, 11(3), 316-320.
- Reichmuth, C. (2000a,b). Biological control in stored product protection. *International Organisation for Biological and Integrated Control Bulletin*, 23(10), 11-24.
- Reppchen, A., Schöller, M., Prozell, S., Adler, C., Reichmuth, C., & Steidle, J. (2003). The granary weevil *Sitophilus granarius* is suppressed by the parasitoid *Lariophagus distinguendus* Förster (Hymenoptera: Pteromalidae). *Proceedings of the 8th International Working Conference on Stored Product Protection*, York. UK: 230-232.
- Riudavets, J. (2018). Biological control of stored products pests.
- Rumbos, C. I., & Athanassiou, C. G. (2017). Use of entomopathogenic fungi for the control of stored-product insects: can fungi protect durable commodities?. *Journal of pest science*, 90, 839-854. doi: 10.1007/s10340-017-0849-9
- Saha, S. R., Islam, W., & Parween, S. (2013). Influence of humidity and *Tribolium* beetle food source on the life history characteristics of predator, *Xylocoris flavipes* (Hemiptera: Anthocoridae). *Tropical Agricultural Research and Extension*, 15(1).
- Saikia, D. K., & Borkakati, R. N. (2020). Evaluation of Anthocorid Predators against Storage Pests of Rice in Assam Situation. *International Journal of Current Microbiology and Applied Sciences*, 9(8), 3180-3185.
- Saleh, M. M. E., Hussein, M. A., Hafez, G. A., Hussein, M. A., Salem, H. A., & Metwally, H. M. (2015). Foliar application of entomopathogenic nematodes for controlling *Spodoptera littoralis* and *Agrotis ipsilon* (Lepidoptera: Noctuidae) on corn plants. *Advances in Applied Agricultural Science*, 3: 51-61.
- Sariyörük, N., & Köseoğlu, A. (1987). *Effect of freezing methods to the fig insects in the natural dried figs* (Doctoral dissertation, B. Sc. Thesis. Ege Univ., Food Engineering Dept. 37 p).
- Sarker, A. C., Islam, W., & Parween, S. (2019). Developmental parameters of *Xylochoris flavipes* (Reuter)(Hemiptera: Anthocoridae) fed on life-stages of *Rhyzopertha dominica* (f.). *Journal of Bio-Science*, 27, 11-21.
- Schöller, M., & Flinn, P. W. (2000). Parasitoids and predators. *Alternatives to pesticides in stored-product IPM*, 229-271.
- Schöller, M., Flinn, W.P., Grieshop, M.J. and Zdarkova, E., (2006). *Biological Control of Stored Product Pests*. *Insect Management for Food Storage and Processing*, 67-87.

- Schöller, M. (2010). Biological control of stored-product insects in commodities, food processing facilities and museums. *Julius-Kühn-Archiv*, (425), 596.
- Schöller, M. (2014). Cryptocephaline Egg Case Provides Incomplete Protection from Generalist Predators (Coleoptera: Chrysomelidae). *Psyche: A Journal of Entomology*, 2014(1), 176539.
- Shapiro-Ilan, D. I., Lacey, L. A., & Siegel, J. P. (2007). Microbial control of insect pests of stone fruit and nut crops. In *Field Manual of Techniques in Invertebrate Pathology: Application and Evaluation of Pathogens for Control of Insects and Other Invertebrate Pests* (pp. 547-565). Dordrecht: Springer Netherlands. doi: 10.1007/978-1-4020-5933-9_26
- Sing, S. E., & Arbogast, R. T. (2008). Predatory response of *Xylocoris flavipes* to bruchid pests of stored food legumes. *Entomologia Experimentalis et Applicata*, 126(2), 107-114.
- Srivastava, C., & Subramanian, S. (2016). Storage insect pests and their damage symptoms: an overview. *Indian Journal of Entomology*, 78(special), 53-58. doi: 10.5958/0974-8172.2016.00025.0
- Solà Cassi, M. (2018). *Approaches for the biological control of stored product pests*.
- Tyler, P. S., Taylor, R. W., & Rees, D. P. (1983). Insect resistance to phosphine fumigation in food warehouses in Bangladesh. *International Pest Control*, 25, 10-13.
- Upadhyay, R. K., & Ahmad, S. (2011). Management strategies for control of stored grain insect pests in farmer stores and public ware houses. *World Journal of Agricultural Sciences*, 7(5), 527-549
- Usta Gebeş, G., & Gözüaçık, C. (2024). Determination of Biology and Prey Preference of the Predator Insect, *Xylocoris flavipes* (Reuter) (Heteroptera: Anthocoridae) Against Storage Pests. *Kahramanmaraş Sütçü İmam Üniversitesi Tarım Ve Doğa Dergisi*, 27(Ek Sayı 1 (Suppl 1), 114-123. <https://doi.org/10.18016/ksutarimdogavi.1425154>
- Vail, P. V., Tebbets, J. S., Cowan, D. C., & Jenner, K. E. (1991). Efficacy and persistence of a granulosis virus against infestations of *Plodia interpunctella* (Hübner)(Lepidoptera: Pyralidae) on raisins. *Journal of Stored Products Research*, 27(2), 103-107. doi: 10.1016/0022-474X(91)90019-9
- Vashisth, S., Chandel, Y. S., & Sharma, P. K. (2013). Entomopathogenic nematodes-A review. *Agricultural Reviews*, 34(3), 163-175.
- Yan, J. J., Sarkar, S. C., Meng, R. X., Reitz, S., & Gao, Y. L. (2020). Potential of *Steinernema carpocapsae* (Weiser) as a biological control agent against potato tuber moth, *Phthorimaea operculella* (Zeller)(Lepidoptera: Gelechiidae). *Journal of Integrative Agriculture*, 19(2), 389-393.
- Yurt, Ç., Gözel, Ç., & Gözel, U. (2015). Bazı entomopatojen nematod türlerinin *Pieris brassicae* (Linnaeus)(Lepidoptera: Pieridae) üzerindeki etkinlikleri. *Türkiye Biyolojik Mücadele Dergisi*, 6(2), 77-84.
- Zettler, J. L. (1996, April). Influence of resistance of future fumigation technology. In *Proceedings of the International Conference on Controlled Atmosphere and Fumigation in Stored Products* (pp. 21-26).
- Zettler, J. L., & Arthur, F. H. (2000). Chemical control of stored product insects with fumigants and residual treatments. *Crop protection*, 19(8-10), 577-582.

**EVALUATION OF THE EFFECTS OF SHEEP MANURE, WORM COMPOSITION
AND MELASSES APPLICATIONS ON BLACK ARONIA (*Aronia Melanocarpa*)
YIELD**

**KOYUN GÜBRESİ, SOLUCAN GÜBRESİ VE MELAS UYGULAMALARININ
SİYAH ARONYA (*Aronia Melanocarpa*) VERİMİ ÜZERİNDEKİ ETKİSİNİN
DEĞERLENDİRİLMESİ**

Dr. Öğr. Üyesi Ruhşen AYDIN KARAAĞAÇ

Kastamonu Üniversitesi, (İhsangazi Meslek Yüksekokulu), (Veterinerlik Bölümü),
Kastamonu, Türkiye

ORCID ID: <https://orcid.org/0000-0001-7009-5314>

Ahmet Taner KALE

Kastamonu Üniversitesi, (İhsangazi Meslek Yüksekokulu), (Veterinerlik Bölümü),
Kastamonu, Türkiye

ORCID ID: <https://orcid.org/0009-0007-9379-9905>

ÖZET

Giriş ve Amaç: Siyah aronya (*Aronia melanocarpa*), Kuzey Amerika ve Kanada'nın Güneydoğusunda yetişen, Rosacea familyasına ait, kışın yapraklarını döken çok yıllık çalı formunda, uzun ömürlü, üzüksü meyveler grubuna ait bir bitkidir. Meyvesi yüksek miktarda antioksidan içermektedir. İnsan sağlığına olumlu etkisinden dolayı 'süper meyve' olarak bilinmektedir. Yetiştirmesi kolay olup, her toprak tipinde ve yükseklikte yetişmektedir. Bu araştırmanın temel amacı koyun gübresi, solucan gübresi ve melas uygulamalarının Siyah aronya (*Aronia melanocarpa*) verimliliği üzerindeki etkisini değerlendirmektir. Fidan gelişme dönemi sonunda bitkilerde ortalama sürgün uzunluğu, ortalama sürgün çapı, ortalama gövde çapı, ortalama yaprak alanı, ortalama kök uzunluğu ve meyve vermesi durumunda verim özelliklerinin belirlenmesidir. Ayrıca çalışma verimliliği artırmak için kullanılan organik gübrelerin potansiyel etkilerini vurgulayarak, sürdürülebilir tarım uygulamalarına değerli bilgiler katmaktadır. **Gereç ve Yöntem:** Çalışma Kastamonu İli İhsangazi ilçesi Çay mahallesi yerleşkesinde yer alan açık arazide yürütülmüş olup, materyal olarak Siyah Aronya (*Aronia melanocarpa*) çeşidine ait fidanlar kullanılmıştır. Gübreleme (%10) Melas 10 fidan, koyun gübresi 10 fidan ve solucan gübresi 10 fidana uygulanacak şekilde gruplandırılmıştır. Kontrol grubu 10 fidana herhangi bir uygulama yapılmamıştır. Denemede fidanlarda düzenli olarak sulama, gübreleme ve yabancı ot temizliği yapılmıştır. **Bulgular:** Uygulama sonucunda her gruptaki bitkilerin morfolojik ölçümleri yapılmıştır. Bitkinin boyu, kalınlığı, yaprak alanı, sürgün sayısı, sürgün boyu, sürgün kalınlığı değerleri her bir grup için ayrı ayrı belirlenerek karşılaştırılmıştır. Fidanların toprak adaptasyon süresi sebebiyle çalışmamız süresince (1 yıl) meyve vermemiştir. **Tartışma ve Sonuç:** Kontrol grubuna kıyasla gübreleme işlemi yapılan tüm bitkiler önemli ölçüde gelişme göstermiştir. Gübreler arasında en iyi etkiyi koyun ve solucan gübresi göstermiştir. Bu araştırmanın sonuçları, mahsul verimini artırmak, yüksek kaliteli, besin açısından zengin aronia ürünleri üretmek için çevre dostu, ekonomik ve organik yaklaşımlar arayan çiftçileri teşvik edecektir.

Anahtar Kelimeler: Siyah Aronya (*Aronia melanocarpa*), Melas, Solucan Gübresi, Koyun gübresi

ABSTRACT

Introduction and Purpose: Black Aronia (*Aronia melanocarpa*) is a perennial shrub-shaped, perennial, berry-like fruit plant belonging to the Rosacea family, which grows in the Southeast of North America and Canada and sheds its leaves in winter. Its fruit contains high amounts of antioxidants. It is known as a ‘super fruit’ due to its positive effects on human health. It is easy to grow and grows in all soil types and altitudes. The main purpose of this study is to evaluate the impact of sheep manure, worm manure, and molasses applications on the productivity of Black Aronia (*Aronia melanocarpa*). The determination of average shoot length, average shoot diameter, average stem diameter, average leaf area, average root length, and yield characteristics in the case of fruiting at the end of the seedling development period. In addition, the study adds valuable information to sustainable agricultural practices by emphasizing the potential effects of organic fertilizers used to increase productivity. **Materials and Methods:** The study was conducted in an open field in the Çay neighborhood of the İhsangazi district of Kastamonu province. The saplings of the Black Aronia (*Aronia melanocarpa*) variety were used as material. Fertilization was grouped as (10%) Molasses to be applied to 10 saplings, sheep manure to 10 saplings, and worm manure to 10 saplings. No application was made to the control group of 10 saplings. In the experiment, regular irrigation, fertilization, and weed cleaning were carried out on the saplings. **Results:** As a result of the application, morphological measurements of the plants in each group were made. Plant height, thickness, leaf area, shoot number, shoot length, and shoot thickness values were determined separately for each group and compared. Due to the soil adaptation period of the seedlings, they did not bear fruit during our study (1 year). **Discussion and Conclusion:** All fertilized plants showed significant growth compared to the control group. Among the fertilizers, sheep and worm manure showed the best effect. The results of this study will encourage farmers looking for environmentally friendly, economical, and organic approaches to increase crop yields and produce high-quality, nutrient-rich aronia products.

Key Words: Black Aronia (*Aronia melanocarpa*), Molasses, Worm Castings, Sheep Manure

GİRİŞ

Aronya Meloideae sınıfının Rosales takımının Rosaceae (Gülgiller) familyasının *Maleae* cinsinden Aronya şeklinde tanımlanmıştır. (Lupascu, 2019). Üzüksü meyveleri sebebiyle bu cinsin İngilizce genel adı ‘Chokeberry’ olarak tanımlanmıştır. Bu cins içerisinde *Aronia melanocarpa* (Michx) Elliot (Black chokeberry, siyah aronya), *Aronia arbutifolia* (L.) Elliot (Red chokeberry, kırmızı aronya) ve *Aronia prunifolia* (Marsh) (Purple chokeberry, mor aronya) olmak üzere bilinen 3 tür mevcuttur (Kulling ve Rawel, 2008; Šnebergrová vd., 2014).

Anavatanı Kanada ve Kuzey Amerika olan aronya, sağlık üzerine olumlu etkilerinin keşfedilmesinden sonra Rusya’da 1990’lı yılların ilk çeyreğinde ekilmeye başlanmıştır. Güçlü terapotik potansiyele sahip bu bitki, 20. yüzyılın ikinci yarısından itibaren Doğu Avrupa devletlerinde de ticari amaçlı yetiştirilmeye başlanmıştır. Günümüzde üretimi en çok Rusya’da görülürken Almanya ve Doğu Avrupa ülkelerinde de yetiştiriciliği halen devam etmektedir (Fidancı, 2015; Šnebergrová vd., 2014).

Türkiye’de aronya yetiştiriciliği ile ilgili ilk çalışmalar 2012 yılında Atatürk Bahçe Kùltürleri Merkez Araştırma Enstitüsü’nde (ABKMAE) fidan üretimi ile başlamıştır. 2014 yılında Yalova ve Kırklareli’nde küçük bahçeler kurulmuş, 2017 yılında ticari bir şekilde yetiştiricilik çalışmaları başlayıp ilk büyük aronya bahçeleri kurulmuştur. Hatta yine aynı yılda ilk hasat şenliği düzenlenmiş, meyvenin tanıtım ve yayım çalışmalarına başlanmıştır. Bunların dışında yine 2017 yılında başta Yalova olmak üzere Çanakkale, Samsun, İstanbul, Antalya ve Bursa’da küçük bahçeler kurulmaya başlanmıştır (Kulling vd.,2008, Nas vd.,2012, Vidalie, 1986).

Siyah meyveli *Aronia melanocarpa* yetiştirilmesi kolay olup, her toprak tipinde ve yükseklikte yetişmektedir. Kendine verimli olup tozlayıcıya gerek duymaz. Çoğaltılması çoğunlukla dal çeliklerinden olmakla birlikte tohumdan ebeveyninin birebir aynı özelliklerini taşıyan bitkiler çoğaltılabilmektedir. Aronyada, ilaçlama ve bakım yapmaya gerek yoktur. Sadece gübreleme, yabancı otların ayırt edilmesi ve sulama yeterlidir. Meyveler bu türde Ağustos ayı sonunda tümüyle olgunlaşmakta olup, sulama yapılmadığı takdirde dal üzerinde kurumaya başlamaktadır. Yağış alan bölgelerde meyveler daha uzun süre bozulmadan bitki üzerinde kalabilmektedir (Brand 2010).

Kırmızı meyveli aronya (*A. arbutifolia*) türü içerisinde yer alan bitkiler diğer iki türe göre daha dik büyümektedir. Yapraklar ters yumurta şeklinde, üst kısmı parlak yeşil, alt kısmı ise gri renkli ve tüylüdür. Sonbaharda parlak kırmızı rengini almaktadır. Çiçeklenme bu türde de Mayıs ayında gerçekleşmektedir. Kırmızı renkli meyvelerin olgunlaşma tarihi Eylül ayı sonu ve Ekim ayı başlarıdır. Aralık ayına kadar bozulmadan bitki üzerinde kalabilmektedir (Brand 2010).

Kırmızı meyveli aronya türüne daha sıcak alanlarda rastlanırken, siyah meyveli aronya daha çok soğuk bölgelerde bulunmaktadır. Mor meyveli aronya ise (*A. prunifolia*) *A. arbutifolia* ve *A. melanocarpa* türlerinin melezi olup, meyveleri mor-siyah renklidir. Yaprak ve gövde üzerindeki tüy yoğunlukları diğer iki türün ortalamasıdır (Brand 2010, Poyraz Engin ve Boz 2019).

Aronia melanocarpa , yaklaşık -30°C 'ye kadar soğuğa dayanıklı olup, çok geç çiçek açtıklarından ilkbahar donlarından etkilenmezler (Mckay, 2001; Ochmain vd., 2012). Aronya meyveleri kalın dış kabuğu sebebiyle diğer üzümü meyvelere göre çevre koşullarına daha dayanıklıdır. Meyveler bitki üzerinde 7 haftaya kadar kalabilmekte ve mekanik hasadı yapılabilmektedir. Hasat sonrası birkaç hafta soğuk depoda kalabildiği ve taşıma sırasındaki zarar görmeye karşıda oldukça dayanıklı olduğu bildirilmiştir (Kulling ve Rawel, 2008; Mckay, 2001).

Aronya günümüzde taze olarak tüketilebildiği gibi buruk bir tadı sebebiyle gıda sanayisinde meyve suyu, konsantre, reçel, kuru meyve, çay, şarap, likör ve diğer alkollü içeceklerle de işlenmektedir (Kapçı, 2013; Wilkes, 2014). Aronya meyveleri koyu renklerinden dolayı zengin bir doğal gıda boyası hammaddesidir (Strigl vd., 1995). Aronya kek, pasta, çörek ve ekmek gibi fırıncılık ürünlerinde çikolata, yoğurt ve tatlılarda hem çeşitliliği hem de gıdanın fonksiyonel özelliğini artırmak için taze veya dondurulmuş olarak da kullanılmaktadır. Üzümsü meyvelerin taze ve işlenmiş ürün olarak pazarlanabilmesi satış potansiyellerini giderek artırmaktadır. İçerik olarak fazlaca flavanoid ve antosiyanin bulundurmaktadır. Bu durum aronyanın pazarlanabilirliğini olumlu etkileyen bir faktör olarak görülmektedir. Diğer taraftan fenol, vitamin, mineral ve antioksidan bakımından da oldukça zengin olup, insanların birçok kalp ve kanser hastalıklarına karşı korunmasına yardımcı olmaktadır (Tolic vd., 2015). İçeriğindeki mineraller, antioksidanlar ve C vitamini bakımından da zengin olması nedeniyle bağışıklık sisteminin güçlendirilmesine katkıda bulunmaktadır (Cuşmenco ve Bulgaru, 2020).

Denev vd., (2019) siyah kuş kirazı (*Aronia melanocarpa*) meyvelerinden elde edilen farklı polifenolik preparasyonların antioksidan, antimikrobiyal ve nötrofil modüle edici aktiviteleri üzerine bir çalışma yapmışlardır. *A. melanocarpa* meyvelerinin birçok polifenol içerdiği ve özellikle içerdikleri proantosiyanidinlerin vücutta sindirim sistemi içinde vücudu patojenlere karşı koruyan fagosit hücrelerine etki ederek antimikrobiyal ve antioksidan işlevlerine doğrudan katkı sağladığı bildirilmiştir.

Organik gübrelerin; topraktaki mikroorganizmalara karbon ve enerji kaynağı sağladıkları, toprağı ıslah etmelerinden dolayı verim ve kaliteyi arttırdıkları bilinmektedir. Çiftlik gübreleri etkisini çok yönlü olarak göstermektedir. Kimyasal gübrelerde olduğu gibi etkileri tek yönlü değildir (Erdal, 2018; Ali vd.,2019).

Çiftlik gübresinin kalitesi ve içeriği; hayvanın cinsi, yaşı, beslenme durumu, yataklık materyali, saklama koşulları ve araziye uygulama şekli olmak üzere pek çok faktöre bağlı olarak değişiklik göstermektedir (Lampkin, 2002; Çıtak vd.,2011).

Yapılan araştırmalarda çiftlik gübreleri arasında koyun gübresinin inek gübresinden daha fazla kuru madde içermesi nedeniyle daha yararlı olduğu bildirilmiştir (Ali vd., 2019).

Koyun gübresi, katı, sıcak bir gübre olup aerobik ayrışmaya uygundur. Azot, potasyum ve fosfor oranları çok yüksektir. Toprağın işlenmesini kolaylaştırır, sıkışmasını engelleyerek köklerin daha kolay gelişmesine yardımcı olur (Url1).

Toprak solucanları bitkisel ve hayvansal atıklarla beslenmektedir. Hayvan dışkıları ile bitkisel üretim artıkları ve evsel atıklar (kağıt, talaş, sebze ve meyve kabukları) gibi organik maddeler solucanlara besin olarak verilip, "vermikompost" denilen organik gübre elde edilmektedir. Hümik asit ve büyüme hormonları içeren Vermikompost, bitki gelişimini düzenler, toprak yapısını iyileştirir ve toprağa besin elementi kazandırarak bitkisel üretimi artırır (Ay, 2016). Vermikompost bitkilerde verimi artırdığı gibi çimlenmeyi de hızlandırmaktadır (Sezgin ve Şimşek, 2017).Solucan gübresi bitki beslemede önemli bir rol üstlenmekte ve organik tarımın geleceğinde de ciddi katkılar sağlayabileceği düşünülmektedir. Ülkemiz şartlarında solucan gübresiyle ilgili bilimsel çalışmalar henüz istenilen düzeyde olmaması sebebiyle bu konuyla ilgili çalışmalar artırılmalıdır. Vermikompostun bu özelliklerinden ve en önemlisi de doğal ekosistemlere zarar vermemesi nedeniyle hem ülkemizde hem de dünyada kullanımı giderek artmaktadır (Abacıoğlu vd.,2020).

Melas, bünyesinde bol miktarda organik madde, makro ve mikro elementler bulunduran şekerpancarının şekere dönüşmesi için şeker fabrikasında yapılan işlemlerden yan ürün olarak açığa çıkan bir preparattır. Melas uygulamaların bitkilerin vejetatif gelişimleri üzerine olumlu etkiler yaptığı bilinmektedir (Ulus ve Yavuzaslanoğlu, 2017).

İhsangazi, Batı Karadeniz Bölgesi'nde, Kastamonu İl'ine bağlı, yüzölçümü 384 km² olan bir ilçedir. Kastamonu İl alanı içinde % 2,9'luk bir yer kaplayan İhsangazi İlçesi, Ilgaz Dağları'nın kuzey eteklerinde kurulmuştur. İhsangazi İlçesi, Karadeniz iklim kuşağında bulunmakla beraber, sıcaklık ve yağış rejimi açısından daha sert bir yapıya sahiptir. İlçe tarihinde en düşük sıcaklık -26,9 C°, en yüksek sıcaklık 37,7 C° olarak kaydedilmiştir. Yıllar itibariyle ortalama yağış miktarı ise 450 mm'dir. Mayıs ve Haziran aylarında yağışlar artarken, Temmuz ve Ağustos aylarında azalmaktadır (Url2).

Bu çalışma sayesinde organik gübreleme yöntemlerinin *Aronya melanocarpa* bitki verimine etkilerinin araştırılmasına katkı sağlamanın yanı sıra son yıllarda artan ticari gübreleme yöntemlerinde gerçekleşen toprak kirliliği sorununun da ortadan kaldırılması sağlanacaktır.

Ayrıca aronya meyvesinin biyo-kimyasal içeriğinin çeşit, uygunluk, iklim, hava, hasat zamanı ve gübreleme gibi faktörlere bağlı olarak değiştiği birçok çalışma sonuçlarının yer aldığı bir derleme tarafından rapor edilmiştir (Kulling ve Rawel, 2008).

Kurutma firmaları, meyve suyu firmaları, gıda takviyesi ürün firmaları üretilen ürüne talep göstermektedir. Türkiye'de üretimin artması ile sanayisinin daha da gelişeceği düşünülmektedir (Poyraz Engin ve Boz, 2019).

Koyun gübresi, solucan gübresi ve melasın Siyah aronia (*Aronia melanocarpa*) bitkilerine uygulanması, bitkinin genel sağlık, üretkenlik ve besin içeriğinde artış ile sonuçlanacaktır. Koyun gübresindeki zengin Azot ve diğer temel besinler, toprağın verimliliğini artırabilir ve bitkiler için dengeli bir besin profili sağlayabilir. Solucan gübresi, içerdiği organik maddelerle faydalı mikroorganizmaları ile toprak yapısının iyileştirilmesine ve besin bulunabilirliğine katkıda bulunabilir. Bir şeker kaynağı olan melas eklenmesi de, toprakta besin döngüsünü teşvik ederek mikrobiyal aktiviteyi teşvik edebilir. Sonuç olarak, bu uygulamalar, siyah aronia

bitkileri için daha elverişli, besin açısından zengin bir ortam yaratacağı, sonuçta hasat edilen meyvenin kalitesini ve miktarını olumlu yönde etkileyeceği varsayımı yapılabilir.

Bu araştırma, *A. melanocarpa* yetiştiriciliğinde verimliliği artırmak için kullanılan organik gübrelerin potansiyel etkilerini vurgulayarak, sürdürülebilir tarım uygulamalarına değerli bilgiler katmaktadır. Organik değişikliklerin etkisinin anlaşılması, sürdürülebilir tarım uygulamalarına artan ilgiyle paralellik göstermektedir. Koyun gübresi, solucan gübresi ve melas gibi organik girdiler, sentetik gübrelelere olan bağımlılığı en aza indiren çevre dostu yaklaşımların bir parçası olabilir. Ayrıca bu girdilerin etkilerini incelemek, bu değişikliklerin toprak sağlığının kritik bileşenleri olan toprak yapısının, verimliliğinin ve mikrobiyal çeşitliliğin geliştirilmesine nasıl katkıda bulunduğunun değerlendirilmesine de yardımcı olacaktır.

MATERYAL ve YÖNTEM

Materyal Temini

Çalışma Kastamonu Üniversitesi İhsangazi Çay mahallesi yerleşkesinde yer alan açık arazide yürütülecektir. Araştırmada materyal olarak My botanik sitesinden (www.mybotanik.com) temin edilen Siyah Aronya (*Aronia melanocarpa*) çeşidine ait fidanlar kullanılmıştır.

Solucan gübresi tarım ürünleri marketlerinden, koyun gübresi İhsangazi Meslek Yüksekokulu Koyunculuk Ünitesinden, melas da Erzurum Şeker Fabrikasından temin edilmiştir.

Dikim Öncesi Uygulamalar

Arazide fidanların ekileceği alanlar belirlenmiş, topraktaki oksijen seviyelerinin iyileştirilmesi amacıyla toprak sürülerek havalandırılmıştır. Kök bölgesinde yeterli oksijen, kök gelişimi ve genel bitki sağlığı için gereklidir. Toprağın havalanması aynı zamanda faydalı mikrobiyal aktiviteyi teşvik etmesi açısından da gereklidir. Çalışmada, melas, koyun gübresi ve solucan gübreleri uygulanmıştır. Kontrol grubu fidanlara herhangi bir uygulama yapılmamıştır.

Deneme tesadüf parselleri 1 m x 4 m mesafe olacak şekilde 3 tekerrür ve her tekerrürde 4 fidan olacak şekilde dikilmiştir.

Dikim Sonrası Yapılan Uygulamalar

Denemede fidanlarda düzenli olarak sulama ve gübreleme yapılmıştır. Ayrıca siyah aronya bitkileriyle besin, su ve güneş ışığı gibi temel kaynaklar için rekabet edebilen, büyümelerini ve gelişmelerini engelleme potansiyeline sahip olan yabancı otların da temizliği düzenli olarak yapılmıştır. Çalışmada, melas, koyun gübresi ve solucan gübrelerinin %10'luk konsantrasyonları uygulanmıştır. Kontrol grubu fidanlara herhangi bir uygulama yapılmamıştır.

SONUÇ ve TARTIŞMA

Deneme sonunda fidanların boyu, gövde çapı, yıllık sürgün uzunlukları, sürgün çapları, sürgün sayıları (adet) ve yaprak alanları belirlenmiştir. Sürgün uzunlukları ve bitkinin boyu cetvelle, sürgün çapı ile bitkinin gövde çapları ise kumpas kullanılarak ölçülmüştür. Büyüme mevsimi sonunda, fidanların olgunlaşmış yapraklarından alınan dörder numunenin alan hesaplamaları, AutoCAD programı kullanılarak yapılmıştır. Her deneme alanındaki bitkilerin değerlerinin ortalaması alınmış ve karşılaştırılmıştır. Çalışma süresi boyunca aronya bitkileri toprağa adapte olmuş ve başarılı bir gelişim göstermiştir. Mevsim geçişi ve adaptasyon süreci nedeniyle belirlenen çalışma döneminde meyve gelişimi gözlemlenmemiş olduğundan, meyve ile ilgili değerlendirme yapılamamıştır.

Uygulamaların bitki boyu uzunluğuna, bitki gövde çapı, sürgün boyu, sürgün çapı, sürgün sayısı ve yaprak alanı etkilerine ait sonuçlar Çizelge 1.1. , 1.2. , 1.3. , 1.4. , 1.5.ve 1.6. 'da verilmiştir.

Çizelge 1.1. Uygulamaların Ortalama Bitki Boyuna Etkileri

Uygulama (%10)	Ortalama Bitki Boyu (cm)			
	1.Deneme (4 bitki ort.)	2.Deneme (4 bitki ort.)	3.Deneme (4 bitki ort.)	Ortalama (cm)
Toprak Solucanı Gübresi	110	96	104,75	103,58
Koyun Gübresi	85	90	82,5	85,83
Melas	95	92,25	93,12	93,45
Kontrol	67	60	70	65,66

Çizelge 1.2. Uygulamaların Ortalama Bitki Gövde Çapına Etkileri

Uygulama (%10)	Ortalama Bitki Gövde Çapı (mm)			
	1.Deneme (4 bitki ort.)	2.Deneme (4 bitki ort.)	3.Deneme (4 bitki ort.)	Ortalama (mm)
Toprak Solucanı Gübresi	1,37	1,1	1,32	1,26
Koyun Gübresi	1,1	1	1	1,03
Melas	1,62	1,59	1,57	1,59
Kontrol	0,7	0,6	0,9	0,73

Çizelge 1.3. Uygulamaların Ortalama Sürgün Boyuna Etkileri

Uygulama (%10)	Ortalama Sürgün Boyu (cm)			
	1.Deneme (4 bitki ort.)	2.Deneme (4 bitki ort.)	3.Deneme (4 bitki ort.)	Ortalama (cm)
Toprak Solucanı Gübresi	53,25	50	55	52,75
Koyun Gübresi	50	55	50	51,66
Melas	45	30	38,25	37,75
Kontrol	29	25	26,5	26,83

Çizelge 1.4. Uygulamaların Ortalama Sürgün Çapına Etkileri


Uygulama (%10)	Ortalama Sürgün Çapı (mm)			
	1.Deneme (4 bitki ort.)	2.Deneme (4 bitki ort.)	3.Deneme (4 bitki ort.)	Ortalama (mm)
Toprak Solucanı Gübresi	0,27	0,32	0,22	0,27
Koyun Gübresi	0,37	0,31	0,35	0,34
Melas	0,3	0,27	0,31	0,29
Kontrol	0,2	0,1	0,15	0,15

Çizelge 1.5. Uygulamaların Ortalama Sürgün Sayısına Etkileri

Uygulama (%10)	Ortalama sürgün sayısı (adet)			
	1.Deneme (4 bitki ort.)	2.Deneme (4 bitki ort.)	3.Deneme (4 bitki ort.)	Ortalama (adet)
Toprak Solucanı Gübresi	10	12	10	10,66
Koyun Gübresi	20	19	22	20,33
Melas	23	18	14	18,33
Kontrol	7	5	10	7,33

Çizelge 1.6. Uygulamaların Ortalama Yaprak Alanına Etkileri

Uygulama (%10)	Numune No	En	Boy	Yaprak Alanı (cm ²)	Ortalama (cm ²)
Toprak Solucanı Gübresi	1	5	9,5	37,30641	38,4845075
	2	6	10,5	49,48008	
	3	5	9	35,34292	
	4	4,5	9	31,80862	
Koyun Gübresi	1	5	8	31,41592	33,72303
	2	4,5	8,5	30,04147	
	3	5,5	9	38,87721	
	4	5,5	8	34,55752	
Melas	1	5	8,8	34,55752	33,673945
	2	5	9	35,34292	
	3	5	8	31,41592	
	4	5	8,5	33,37942	
Kontrol	1	4	7,5	23,53194	18,9402275
	2	3	6,5	15,31526	
	3	4	6,5	20,42035	
	4	3	7	16,49336	

**Şekil 1.** Çeşitli Gübre Uygulamalarının (%10) Siyah aronia (*Aronia melanocarpa*) Verimi Üzerine Etkisi

Aronya fidanlarının gelişimi üzerinde solucan gübresi, koyun gübresi ve melas uygulamalarının etkileri genel olarak olumlu yönde olmuştur. Bitki boyu gelişiminde en iyi etkiyi solucan gübresi göstermiş, bunu sırasıyla melas ve koyun gübresi takip etmiştir. Bitki çapı gelişiminde ise melas uygulamaları, diğer uygulamalara göre daha olumlu sonuçlar vermiştir. Sürgün boyu gelişiminde, solucan ve koyun gübresi birbirine çok yakın değerlere sahip olup, melasa kıyasla daha olumlu sonuçlar elde edilmiştir. Sürgün çaplarında ise, kontrole göre olumlu sonuçlar gözlemlenmiş olmasına rağmen, uygulamalar arasında belirgin bir fark bulunmamıştır. Sürgün sayısı gelişiminde en iyi etkiyi koyun ve melas uygulamaları göstermiştir. Yaprak gelişiminde,

uygulanan tüm preparatlar, kontrol grubuna göre olumlu sonuçlar elde etmiştir; yaprak alanını kontrole göre en fazla artıran uygulama solucan gübresi olmuş, bunu koyun gübresi ve melas takip etmiştir.

Genel olarak, aronya fidan yetiştiriciliğinde solucan gübresi, koyun gübresi ve melas uygulamaları tavsiye edilebilir. Bu araştırma, *Aronia melanocarpa* yetiştiriciliğinde verimliliği artırmak amacıyla kullanılan organik gübrelerin potansiyel etkilerini vurgulayarak, sürdürülebilir tarım uygulamalarına önemli bilgiler sunmaktadır. Organik değişikliklerin etkisinin anlaşılması, sürdürülebilir tarım uygulamalarına duyulan artan ilgiyle paralellik göstermektedir. Koyun gübresi, solucan gübresi ve melas gibi organik girdiler, sentetik gübrelere olan bağımlılığı en aza indirerek çevre dostu yaklaşımlar geliştirmeye olanak tanıyabilir. Ayrıca, bu girdilerin etkilerini incelemek, toprak sağlığının temel bileşenleri olan toprak yapısı, verimlilik ve mikrobiyal çeşitliliğin geliştirilmesine nasıl katkı sağladığının değerlendirilmesine de yardımcı olacaktır.

Bitki yetiştirme sahası olarak soğuk iklime kolay adapte olabilen *A. melanocarpa*'nın Rusya'da yoğun olarak yetiştirilmesinden ilham alınarak, soğuk ve bol yağışlı iklime sahip Kastamonu İhsangazi ilçesindeki adaptasyonu ve yetiştirme performansları değerlendirilmiş ve olumlu sonuçlar elde edilmiştir.

TEŞEKKÜR

Bu çalışma Türkiye Bilimsel ve Teknolojik Araştırma Kurumu tarafından maddi olarak desteklenmiştir (proje no: 1919B012314796). Desteklerinden dolayı TÜBİTAK'a teşekkür ederiz.

KAYNAKLAR

- Abacıoğlu, E., Yatgın, S., Tokel, E., Yücesoy, P. 2020. Vermikompostun (solucan gübresi) üretimi ve bitki beslemesindeki önemi. Bartın University International Journal of Natural and Applied Sciences, 3(1): 1-10.
- Ali, B., Üstündağ, E., Alatürk, F. 2019. Farklı gübre ortamlarında yem bitkisi *Pennisetum benthium*'un yetiştirilmesi. Türkiye 13. Ulusal, 1. Uluslararası Tarla Bitkileri Kongresi, Antalya.
- Ay, T., 2016, Çevre Odaklı Üretim Ve Tarımsal Girişimcilik Bağlamında: Vermikültür, Journal of Life Economics, 3 (2), 1-18.
- Brand, M.H. (2010). *Aronia*: Native shrubs with untapped potential. *Arnoldia*, 67: 14-25.
- Cuşmenco, T., Bulgaru, V., 2020. Quality Characteristics and Antioxidant Activity of Goat Milk Yogurt with Fruits. *Ukrainian Food Journal*, 9(1): 86-98.
- Çıtak, S., Sönmez, S., Koçak, F., Yaşın, S. 2011. Vermikompost ve ahır gübresi uygulamalarının ıspanak (*Spinacia oleracea* var. l.) bitkisinin gelişimi ve toprak verimliliği üzerine etkileri. *Batı Akdeniz Tarımsal Araştırma Enstitüsü Derim Dergisi*, 28 (1): 56-69.
- Denev, P., Číž, M., Kratchanova, M., Blazheva, D., 2019. Black Chokeberry (*Aronia melanocarpa*) Polyphenols Reveal Different Antioxidant, Antimicrobial and Neutrophil-Modulating Activities. *Food Chemistry*, 284: 108-117.
- Erdal, İ., Küçükyumuk, Z., Şimşek, K., Basır, M., Baysal, G.D. 2018. Farklı hayvan gübrelerinin domatesin gelişimi ve mineral beslenmesine etkisi. *Süleyman Demirel Üniversitesi, Zir. Fak. Derg., Özel Sayısı*, 295-302.

- Fidancı, A. (2015). Türkiye İçin Yeni Bir Minör Meyve: Aronia Bitkisi ve Yetiştirme Teknikleri, VII. Ulusal Bahçe Bitkileri Kongresi, 25-29 Ağustos Çanakkale, 1177-1180.
- Kapçı, B. (2013). Characteristic components and antioxidant potential of black chokeberry (*Aronia melanocarpa*) products. Istanbul Technical University, Graduate School of Science Engineering and Technology, Department of Food Engineering.
- Kulling, S.E., Rawel, H.M. 2008. Chokeberry (*Aronia melanocarpa*)—A review on the characteristic components and potential health effects. *Planta medica*, 74(13), 1625-1634.
- Lampkin, N. 2002. Organic Farming. Old Pond Publishing, 104 Valley Road Ipswich, IPI 4PA, U.K.
- Lupascu, N. Candar, E., Cherim, M., Erimina, C. L., Sirbu, R. (2019). Research concerning the efficiency of *Aronia melanocarpa* for pharmaceutical purpose. *European Journal of Medicine and Natural Sciences*, 2(2): 1-7.
- Mckay, S. A. (2001). Demand increasing for aronia and elderberry in North America. *New York Fruit Quarterly*, 9(3): 2-3.
- Nas, M. N., Gokbunar, L., Sevgin, N., Aydemir, M., Dagli, M., Susluoglu, Z. 2012. Micropropagation of mature *Crataegus aronia* L., a medicinal and ornamental plant with rootstock potential for pome fruit. *Plant Growth Regulation*, 67(1), 57-63.
- Ochmian, I. Grajkowski, J., Smolik, M. (2012). Comparison of some morphological features, quality and chemical content of four cultivars of choke berry fruits (*Aronia melanocarpa*). *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 40: 253–260.
- Poyraz Engin, S., Boz, Y. 2019. Türkiye ve dünyada aronya (*Aronia melanocarpa* (Michx) Elliot) yetiştiriciliği. *Bahçe* 48 (Özel Sayı 1): 247–252.
- Sezgin, M. ve Şimşek, E., 2017, Bazı orman ağacı ve çalı türleri tohumlarının çimlendirilmesinde vermikompost ürünlerinin etkileri.
- Šnebergrová, J. Čížková, H., Neradová, E., Kapcı, B., Rajchl, A., Voldrich, M. (2014). Variability of characteristic components of aronia. *Czech Journal of Food Science*, 32(1): 25–30.
- Strigl, A. W. Leitner, E., Pfannhauser, W. (1995). Qualitative und quantitative analyse der anthocyane in schwarzen apfelbeeren (*Aronia melanocarpa* Michx. Ell.) mittels TLC, HPLC. *Z Lebensm Unters Forsch* 201:266–268.
- Tolić, M.T. Jurcevic, I. L., Krbavcic, I.P., Markovic, K., Vahcic, N. (2015). Phenolic content, antioxidant capacity and quality of chokeberry (*Aronia melanocarpa*) products. *Food Tech. Biotech.*, 53(2): 171-179.
- Ulus, F. ve Yavuzaslanoğlu, E., 2017, The effect of different fertilizer applications on plant and fruit yield in greenhouse organic tomato growing, *Turkish Journal of Agriculture-Food Science and Technology*, 5 (13), 1757-1761.
- Url1 https://samsun.tarimorman.gov.tr/Belgeler/Yayinlar/Kitaplarimiz/organik_gubreler_ve_onemi.pdf [Erişim Tarihi 26.10.2023]
- Url2 <https://www.kuzka.gov.tr/> [Erişim Tarihi 23.10.2023]
- Vidalie H., 1986. In vitro culture and its applications in horticulture, *Econ. Bot.*, 51(1), 92–93.
- Wilkes, K., Howard, L. R., Brownmiller, C., Prior, R. L. (2014). Changes in chokeberry (*Aronia melanocarpa* L.) polyphenols during juice processing and storage. *J. Agric. Food Chem.*, 62: 4018–4025. doi.org/10.1021/jf404281n

**DETECTION AND ANALYSIS OF BURNED AREAS WITH SENTINEL-2
SATELLITE IMAGERY AND FIRE SEVERITY MAP: 2025 LOS ANGELES FIRE**

Rüştü ÇALLI

Iğdır University, Vocational School of Technical Sciences, Department of Architecture and
Urban Planning, Iğdir, Türkiye

Emirhan ÖZDEMİR

Iğdır University, Vocational School of Technical Sciences, Department of Architecture and
Urban Planning, Iğdir, Türkiye

Mitat Can YILDIZ

Iğdır University, Vocational School of Technical Sciences, Department of Architecture and
Urban Planning, Iğdir, Türkiye

Alihsan ŞEKERTEKİN

Iğdır University, Vocational School of Technical Sciences, Department of Architecture and
Urban Planning, Iğdir, Türkiye

ABSTRACT

Wildfires pose a significant threat to both natural ecosystems and human communities, with increasing frequency and intensity due to climate change. The 2025 Los Angeles fire, particularly the Palisades Fire, caused extensive damage and highlighted the urgent need for effective monitoring and analysis of burned areas. This study aims to utilize Sentinel-2 satellite imagery to detect and analyze the extent and severity of the burned areas, providing valuable insights for wildfire management and recovery efforts. Sentinel-2 images were acquired on 26 August 2024 (pre-fire) and 12 January 2025 (post-fire) to capture the conditions before and after the fire. Normalized Burn Ratio (NBR) images were generated for both prefire and postfire conditions using the near-infrared (NIR) and shortwave infrared (SWIR) bands. The differenced NBR (dNBR) was calculated by subtracting the postfire NBR from the prefire NBR, highlighting the burned areas. Based on the dNBR values, a fire severity map was created, classifying the area into seven categories: Enhanced Regrowth High, Enhanced Regrowth Low, Unburned, Low Severity, Moderate-Low Severity, Moderate-High Severity, and High Severity. The area analysis revealed that the classes covered the following areas: Enhanced Regrowth High (0.307 km²), Enhanced Regrowth Low (1.616 km²), Unburned (270.053 km²), Low Severity (75.307 km²), Moderate-Low Severity (49.142 km²), Moderate-High Severity (39.268 km²), and High Severity (1.107 km²). The results indicate that there were significant areas with varying levels of fire severity. The analysis identified regions with enhanced regrowth, suggesting that some areas are beginning to recover. The findings highlight the importance of using satellite imagery for timely and accurate wildfire assessment, which can aid in effective management and recovery planning. This study demonstrates the effectiveness of using Sentinel-2 imagery for detecting and analyzing burned areas.

Key Words: Remote Sensing; Fire Severity Map; Normalized Burn Ratio Index

INTRODUCTION

Forest fires cause damage to vegetation, disrupt the habitats of living beings in these areas, and lead to economic losses. Monitoring these fires, determining the extent of the destruction, and analyzing the changes occurring in the burned areas over time are of great importance both ecologically and economically. Thanks to innovations and advancements in satellite technology, remote sensing techniques have become an effective tool for monitoring forested areas and analyzing forest fires (Çiçekli, 2024).

In the literature, numerous studies have been conducted using remote sensing techniques to detect the damages caused by forest fires (Abdikan et al., 2022; Kurbanov et al., 2022; Ghali and Akhloufi, 2023). After forest fires, satellite images play a crucial role in accurately mapping the affected areas (Chen et al., 2015). Various remote sensing datasets with medium spatial resolution, such as Sentinel-2 MSI and Landsat-8 OLI, have been used to detect burned areas during or after the fire (Gülci et al., 2021; Howe et al., 2022). Navarro et al. (2017) analyzed images obtained from the Sentinel-2A satellite to assess the wildfires that occurred in Madeira in August 2016. In that study, Normalized Difference Vegetation Index (NDVI), Green Normalized Difference Vegetation Index (GNDVI), Normalized Burn Ratio (NBR), and red-edge bands were used to examine the post-fire conditions. Additionally, the NDVI was calculated using red-edge bands (NDVI_{reXn}) to evaluate post-fire conditions. The research confirmed previous findings, demonstrating that the NDVI_{re1n} index provided the best results in detecting burned areas using Sentinel-2A MSI images.

Quintano et al. (2018) compared Sentinel-2 MSI and Landsat-8 OLI satellites to assess the damage caused by a large forest fire in the Sierra Gata region of central-western Spain on August 11, 2015, which burned approximately 8,000 hectares. As part of this assessment, pre- and post-fire NBR images were calculated. The results indicated that the fire severity maps generated with Landsat-8 OLI provided an adequate evaluation of damage severity (κ statistic = 0.80) and the spatial distribution of emergency response interventions related to the wildfire.

García-Llamas et al. (2019) analyzed a forest fire that occurred in Spain in 2017 using Landsat-8 OLI and Sentinel-2 MSI satellite data. They employed dNDVI and differenced Normalized Burn Ratio Index (dNBRI) indices to identify burned areas. The study's findings revealed that Sentinel-2 MSI satellite data, due to its higher resolution, offered better accuracy compared to Landsat-8 OLI. Additionally, it was determined that the NBRI index exhibited strong correlation values with burned areas, regardless of the sensor used.

Kesgin Atak and Tonyaloğlu (2020) examined a wildfire that occurred in 2019 in the districts of Gaziemir, Buca, and Karabağlar in İzmir. In their study, they utilized Landsat-8 OLI and Sentinel-2A satellite data to detect burned areas. They applied NDVI, Atmospherically Resistant Vegetation Index (ARVI), NBR, NBR2, and Burn Area Index (BAI) indices to identify the burned areas. Furthermore, they employed an object-based image classification approach using multi-resolution segmentation and the nearest neighbor supervised classification technique. The study results demonstrated that NDVI, NBR2, and ARVI indices produced sufficiently accurate outputs for detecting burned areas. Yılmaz et al. (2022) investigated a forest fire that occurred in the Gelibolu district of Çanakkale in 2020. In their study, they used Sentinel-2 MSI and Landsat-8 OLI satellite images to detect burned areas. By calculating BAI, NDVI, NBRI, and Normalized Difference Moisture Index (NDMI) indices before and after the fire, they generated difference images and identified the burned areas. Accuracy assessments based on the obtained results indicated that the NDVI index provided the most accurate detection.

The objective of this study is to determine the areas affected by the forest fire that occurred on January 7, 2025, in Los Angeles, California, which resulted in significant destruction. Based on the given literature, this study uses Sentinel-2 satellite images by calculating the dNBR index

based on pre- and post-fire images. On the final stage, fire severity map was generated using the dNBR image.

MATERIAL AND METHODOLOGY

Study Area and Materials

The selected study area is the recent forest fire zone in Los Angeles, California, USA, in which the fire occurred on January 7, 2025 (Figure 1). The fire caused significant damage to cultivated agricultural lands, large farms, and a substantial portion of the forested area surrounding the region.




Figure 1: Map of the Study Area and Pre- and Post-Fire Sentinel-2 Images

In this study, Sentinel-2, including 13 spectral bands with spatial resolutions of 10 m, 20 m, and 60 m, was utilized. These bands cover the electromagnetic spectrum in the range of 443–2190 nm. The bands consist of four visible and one near-infrared (NIR) bands with a spatial resolution of 10 m, red-edge and shortwave infrared (SWIR) bands with a spatial resolution of 20 m, and three atmospheric correction bands with a spatial resolution of 60 m. One of the primary missions of this satellite is mapping land cover and land use (Dereli, 2019).

For Sentinel-2, two sets of images were analyzed: a pre-fire image from August 26, 2024, and a post-fire image from January 12, 2025. The characteristics of Sentinel-2 MSI are presented in Table 1.

Table 1: Characteristics of Sentinel-2 Bands Used

Spectral Bands	Wavelength (nm)	Resolution (m)
Band 1 (Coastal aerosol)	0.443	60
Band 2 (Blue)	0.490	10
Band 3 (Green)	0.560	10
Band 4 (Red)	0.665	10
Band 5 (Vegetation red edge)	0.705	20
Band 6 (Vegetation red edge)	0.740	20
Band 7 (Vegetation red edge)	0.783	20
Band 8 (NIR)	0.842	10
Band 8A (Vegetation red)	0.865	20
Band 9 (Water vapour)	0.945	60
Band 10 (SWIR-Cirrus)	1.375	60
Band 11 (SWIR1)	1.610	20
Band 12 (SWIR2)	2.190	20

Methodology

Sentinel-2 images were obtained from the Copernicus Browser (ESA) website (URL-1). Since the used images belong to the Level 2A series, they were already atmospherically corrected. Therefore, in the preprocessing stage, only image clipping was performed, and due to the difference in spatial resolution between the bands used in the NBR calculation—where the NIR band has a 10 m resolution and the SWIR band has a 20 m resolution—the bands were resampled to 20 m resolution. The clipping and resampling processes were conducted using ESA's SNAP software (Figure 2).




Figure 2: Workflow diagram.

- **Normalized Burn Ratio (NBR)**

The NBR is a widely used remote sensing index for determining the effects of forest fires. This index is calculated by using the ratio of near-infrared (NIR) and shortwave infrared (SWIR2) bands. The calculation of NBR is based on the formula shown in Equation (1). The NBR value indicates burned areas when it is low after a fire, while higher values represent healthy vegetation. This index serves as an important tool for assessing fire severity and its impacts (Key & Benson, 2006).

$$NBR = \frac{NIR - SWIR_2}{NIR + SWIR_2} \quad (1)$$

In Sentinel-2 satellite images, Band 8 represents the NIR (Near-Infrared) and Band 12 represents the SWIR2 (Shortwave Infrared 2) (Table 1). For calculating NBR, Band 8 and Band 12 are used. The NBR value ranges between -1 and +1. An NBR value close to -1 indicates burned vegetation.

- **Differenced Normalized Burn Ratio (ΔNBR)**








The Differenced (ΔNBR or $dNBR$) is a remote sensing index used to identify burned areas and assess fire severity after forest fires. ΔNBR is calculated by taking the difference between the pre-fire and post-fire NBR indices. The calculation of ΔNBR is based on the formula shown in

Equation (2). This difference better highlights the impact of the fire and accentuates the distinction between burned areas and healthy vegetation. Areas with high ΔNBR values indicate more intense burning and damage, while low ΔNBR values signify areas with minimal burning or healthy vegetation. This method is widely used to map the effects of forest fires and monitor post-fire recovery processes (Key & Benson, 2006).

$$\Delta\text{NBR or } \Delta\text{NBR} = \text{PrefireNBR} - \text{PostfireNBR} \quad (2)$$

The ΔNBR values, derived by taking the difference between pre-fire and post-fire NBR images, were classified into seven categories based on threshold values in Table 2. This process was carried out to generate fire severity map (Keeley, J. E. 2009).

Table 2: ΔNBR ranges according to fire severity levels.

	SEVERITY LEVEL	ΔNBR RANGE
	Enhanced Regrowth, high	(-0.500) – (-0.251)
	Enhanced Regrowth, low	(-0.251) – (-0.101)
	Unburned	(-0.100) – (+0.099)
	Low Severity	(+0.100) – (+0.269)
	Moderate-Low Severity	(+0.270) – (+0.439)
	Moderate-High Severity	(+0.440) – (+0.659)
	High Severity	(+0.660) – (+1.300)

RESULTS

Post-fire chemical changes in plants affect spectral reflectance, creating noticeable variations in different bands. The NBR index, calculated using the mathematical relationship between the NIR band (sensitive to chlorophyll in healthy plants) and the SWIR-2 band (sensitive to water content in plants, as well as oxides and minerals like clay in the soil), provides a sensitive indicator of chemical changes in burned areas (Lutes et al., 2006; Yılmaz et al., 2021). In the images obtained by applying the NBR index, each pixel takes a value between -1 and +1. A value of +1 represents healthy vegetation, while a value of -1 indicates bare ground or recently burned areas. To determine the level of damage to vegetation, the NBR index was applied to satellite images on a GIS platform, and pre-fire and post-fire conditions were compared (Figure 3). Based on the pre- and post-fire NBR images, it is clear from Figure 3 that a very large area is burned.




Figure 3. Application of the NBR index for pre-fire (a) and post-fire (b) conditions.

The Δ NBR image was created on the GIS platform by taking the difference between the pre-fire and post-fire NBR analysis results. The resulting Δ NBR image was classified on the GIS platform based on the value of each pixel. The classification process was performed using the Δ NBR threshold values according to the fire severity ranges provided in Table 2, and fire severity map was generated as shown in Figure 4. The fire severity map reveals the burned areas obviously, especially from moderate-low severity to high severity areas represents the burned areas.




Figure 4. Fire severity map of the study area.

The area values for each class were calculated from the classified images. The total burned area was calculated as the sum of moderate-low severity, moderate-high severity, and high-severity burned areas based on the ΔNBR results. Based on this analysis, total burned area was calculated as 89.517 km², which represents 54.3% of the considered test site. The data for all calculated classes are provided in Table 3.

Table 2: Total burned areas.

CLASS	ΔNBR (km ²)	Alan (%)
Enhanced Regrowth, high	0.307	
Enhanced Regrowth, low	1.616	
Unburned	270.053	
Low Severity	75.307	
Moderate-Low Severity	49.142	29.81
Moderate-High Severity	39.268	23.82
High Severity	1.107	0.67
Total Burned Area	89.517 km²	54.3

CONCLUSION

In this study, the area affected by the forest fire that broke out on January 7 in Los Angeles, California, USA, was selected as the study area. Analyses were conducted using Sentinel-2 satellite images and GIS platforms. In this study, fire severity map was created using ΔNBR . Sentinel-2 satellite images are a highly effective tool for obtaining information after forest fires. Thanks to ΔNBR , the identification of burned forest areas and the analysis of fire severity were

carried out quickly and reliably. In this study, vegetation damaged by the fire exhibited negative values post-fire, which were represented in red on the maps. Based on the Δ NBR image, it was determined that the fire area predominantly experienced moderate-low and moderate-high severity burns. According to the fire severity maps generated using the Δ NBR index, the total burned area in the study area was found to be 89.517 km². Based on the Δ NBR results, 45.69% of the burned area was classified as low severity, 29.81% as moderate-low severity, 23.82% as moderate-high severity, and 0.67% as high severity.

REFERENCES

- Abdikan, S., Bayik, C., Sekertekin, A., Bektas Balcik, F., Karimzadeh, S., Matsuoka, M., & Balik Sanli, F. (2022). Burned area detection using multi-sensor SAR, optical, and thermal data in Mediterranean pine forest. *Forests*, 13(2), 347.
- Chen G., Metz M.R., Rizzo D.M., Meentemeyer R.K., (2015), Mapping fire severity in a disease-impacted forest landscape using Landsat and ASTER imagery, *International Journal of Applied Earth Observation and Geoinformation*, 40(2015), 91-99.
- Çiçekli, S. Y. (2024). Yanmış Orman Alanlarının Uzaktan Algılama Teknikleri ile Araştırılması: Kozan Orman Yangını Örneği. *Çukurova Üniversitesi Mühendislik Fakültesi Dergisi*, 39(2), 503-514.
- Dereli M., (2019), Sentinel-2A uydu görüntüleri ile Giresun il merkezi için kısa dönem arazi örtüsü değişiminin belirlenmesi, *Afyon Kocatepe Üniversitesi Fen ve Mühendislik Bilimleri Dergisi*, 19(2), 361-368.
- García-Llamas, P., Suárez-Seoane, S., Fernández-Guisuraga, J. M., Fernández-García, V., Fernández-Manso, A., Quintano, C., ... & Calvo, L. (2019). Evaluation and comparison of Landsat 8, Sentinel-2 and Deimos-1 remote sensing indices for assessing burn severity in Mediterranean fire-prone ecosystems. *International Journal of Applied Earth Observation and Geoinformation*, 80, 137-144.
- Ghali, R., & Akhloufi, M. A. (2023). Deep learning approaches for wildland fires using satellite remote sensing data: Detection, mapping, and prediction. *Fire*, 6(5), 192.
- Gülci, S., Yüksel, K., Gümüş, S., & Wing, M. (2021). Mapping wildfires using Sentinel 2 MSI and Landsat 8 imagery: spatial data generation for forestry. *European Journal of Forest Engineering*, 7(2), 57-66.
- Howe, A. A., Parks, S. A., Harvey, B. J., Saberi, S. J., Lutz, J. A., & Yocom, L. L. (2022). Comparing Sentinel-2 and Landsat 8 for burn severity mapping in Western North America. *Remote Sensing*, 14(20), 5249.
- Keeley, J. E. (2009). Fire intensity, fire severity and burn severity: a brief review and suggested usage. *International journal of wildland fire*, 18(1), 116-126.
- Kesgin Atak B., & Tonyaloğlu E., (2020). Evaluating spectral indices for estimating burned areas in the case of Izmir/Turkey. *Eurasian Journal of Forest Science*, 8(1), 49-59. <https://doi.org/10.31195/ejejfs.657253>
- Key, C. H., & Benson, N. C. (2006). Landscape Assessment: Ground measure of severity, the composite burn index; and remote sensing of severity, the normalized burn ratio. Pp. LA1-LA51. FIREMON: Fire Effects Monitoring and Inventory System. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- Kurbanov, E., Vorobev, O., Lezhnin, S., Sha, J., Wang, J., Li, X., ... & Wang, Y. (2022). Remote sensing of forest burnt area, burn severity, and post-fire recovery: A review. *Remote Sensing*, 14(19), 4714.

Lutes DC., Keane RE., Caratti JF., Key CH., Benson NC., Sutherland S., Gangi LJ. Firemon: Fire effects monitoring and inventory system. technical report RMRS-GTR-164-CD. US Department of Agriculture, Forest Service, Rocky Mountain Research Station: Fort Collins, CO, USA, 2006.

Navarro, G., Caballero, I., Silva, G., Parra, P. C., Vázquez, Á., & Caldeira, R. (2017). Evaluation of forest fire on Madeira Island using Sentinel-2A MSI imagery. *International Journal of Applied Earth Observation and Geoinformation*, 58, 97-106.

Quintano C., Fernández-Manso A., Fernández-Manso O., (2018), Combination of Landsat and Sentinel-2 MSI data for initial assessing of burn severity, *International Journal of Applied Earth Observation and Geoinformation*, 64(2018), 221-225.

URL-1 <https://browser.dataspace.copernicus.eu/>

Yılmaz OS., Oruç MS., Ateş AM., Gülgen F. Orman yangın şiddetinin Google Earth Engine ve Coğrafi Bilgi Sistemleri kullanarak analizi: Hatay-Belen örneği. *Journal of the Institute of Science and Technology* 2021; 11(2): 1519-1532.

Yılmaz, B., Demirel, M., & Balçık, F. (2022). Yanmış alanların sentinel-2 msı ve landsat-8 olı ile tespiti ve analizi: Çanakkale/Gelibolu orman yangını. *Doğal Afetler ve Çevre Dergisi*, 8 (1). <https://doi.org/76-8.10.21324/dacd.941456>

BIBLIOMETRIC ANALYSIS OF THE USE OF SATELLITE-BASED REMOTE SENSING TECHNOLOGIES IN CROP MAPPING (2018-2025)

Mitat Can YILDIZ

Iğdir University, Vocational School of Technical Sciences, Department of Architecture and Urban Planning, Iğdir, Türkiye

Emirhan ÖZDEMİR

Iğdir University, Vocational School of Technical Sciences, Department of Architecture and Urban Planning, Iğdir, Türkiye

Rüştü ÇALLI

Iğdir University, Vocational School of Technical Sciences, Department of Architecture and Urban Planning, Iğdir, Türkiye

Alihsan ŞEKERTEKİN

Iğdir University, Vocational School of Technical Sciences, Department of Architecture and Urban Planning, Iğdir, Türkiye

ABSTRACT

Remote sensing technologies have revolutionized agricultural applications thanks to the broad perspective and detailed analytical capabilities provided by satellite data. Crop mapping, in particular, is critical for agricultural land management and productivity analysis. This study examines research conducted between 2018 and 2025 using the keywords "remote sensing", "satellite", and "crop mapping" through bibliometric analysis. The objective is to identify academic trends in this field, analyze collaborations between authors, and assess international relationships, thereby highlighting global contributions to crop mapping with remote sensing technologies. A total of 175 studies were identified in the Scopus database, including 132 research articles, 35 conference papers, 5 review articles, and 2 book chapters, and 1 editorial comment related to the specified keywords. Bibliometric analysis was conducted using the VOSviewer platform. The analysis focused on author collaborations, international relations, highly cited studies, and keyword distributions. The results indicate that the journal Earth and Planetary Sciences contributed the most, with 97 studies. Zheng Niu was identified as the most cited author with 290 citations. China emerged as the leading country in terms of both the number of publications and citations. Approximately 92% of the studies were written in English. Co-authorship analyses revealed strong collaborations between authors and countries. Keyword analyses showed that "remote sensing", "satellite", and "crop mapping" were the primary focus of studies during this period. The impact of remote sensing technologies on crop mapping holds great potential for modern agricultural practices. The results of the bibliometric analysis reveal that studies in this field have made significant contributions both academically and practically. In particular, crop mapping with satellite data is crucial for the development of sustainable agricultural management strategies. China's leadership in this field presents an

opportunity to enhance international collaboration. The findings of this study provide a valuable guide for future research in this domain.

Keywords: Remote Sensing, Crop Mapping, Bibliometric Analysis, Scopus, VOSviewer

INTRODUCTION

With the rapid advancement of modern technologies, significant progress has been made in data collection and analysis processes. The development of satellite technologies, the widespread adoption of artificial intelligence-based analysis methods, and the increase in high-resolution data sources have brought about major transformations in various fields, including the agricultural sector. Agriculture is a strategic sector in terms of global food security, economic development, and ecosystem balance, making it crucial to enhance the efficiency and sustainability of production processes. In this context, remote sensing techniques are increasingly preferred for monitoring agricultural areas, assessing crop productivity, and optimizing land use (Bégué et al., 2018).

Crop mapping is a vital application aimed at improving agricultural productivity. It involves identifying the types, health status, growth processes, and spatial distribution of crops in agricultural fields. Traditional crop mapping methods generally rely on field surveys, sampling, and direct observation, which are time-consuming and costly. However, with technological advancements, satellite-based remote sensing techniques have become a valuable tool in crop mapping studies, offering fast and cost-effective solutions that cover large areas (Khanal et al., 2020; Sishodia et al., 2020).

Remote sensing is a technological field that enables the detection and analysis of the physical characteristics of a specific area through electromagnetic waves without direct contact. In agricultural applications, remote sensing provides significant advantages in monitoring crop health, detecting water stress, identifying diseases, and forecasting yields. Satellite imagery, which is the most commonly used data source in these techniques, offers extensive spatial coverage and enables continuous monitoring of agricultural areas over time (Mulla, 2013).

Van Tricht et al. (2018) demonstrated that the combined evaluation of Sentinel-1 and Sentinel-2 satellite images significantly improved the accuracy of crop mapping in Belgium. Tian et al. (2019) utilized multi-satellite data and an algorithm considering crop growth stages for winter crop mapping in China, significantly enhancing classification accuracy. Xu et al. (2020) developed DeepCropMapping, a multi-temporal deep learning approach for dynamic corn and soybean mapping, increasing spatial generalization. Sun et al. (2019) proved that multi-source and multi-temporal remote sensing data significantly improved crop type mapping accuracy in subtropical agricultural regions. Zhang et al. (2018) introduced a 10-meter resolution method integrating optical and synthetic aperture radar (SAR) imagery for up-to-date rice field mapping in China. Hao et al. (2020) enhanced crop classification accuracy using transfer learning with Cropland Data Layer (CDL) data as training samples. Mazzia et al. (2020) improved crop monitoring efficiency in precision agriculture by enhancing satellite-based vegetation indices using UAVs and machine learning. Song et al. (2021) compared the performance of Landsat, Sentinel, and MODIS data in crop type mapping, finding that Sentinel data provided the highest accuracy. Tian et al. (2021) demonstrated the effectiveness of time-series analysis in agricultural monitoring by integrating Sentinel-1A images according to crop growth cycles for summer maize mapping. Csillik et al. (2019) improved crop classification accuracy using Sentinel-2 data and an object-based dynamic time warping (DTW) method with time constraints.

As evidenced in above given literature, in recent years, the use of remote sensing methods in crop mapping has rapidly increased, overcoming the limitations of traditional techniques and bringing groundbreaking advancements to agricultural management processes. In particular,

multispectral and hyperspectral satellite images play a crucial role in distinguishing crop types and assessing crop health by utilizing the spectral signatures of vegetation. Spectral indices such as the Normalized Difference Vegetation Index (NDVI), Soil-Adjusted Vegetation Index (SAVI), and Enhanced Vegetation Index (EVI) allow for monitoring crop growth and making yield predictions using satellite data (Wu et al., 2022). This study aims to examine the use of satellite-based remote sensing technologies in crop mapping between 2018 and 2025 through a bibliometric analysis, identifying trends, research gaps, and international collaborations in the literature. The findings will provide valuable guidance for developing agricultural management strategies and future research directions.

MATERIAL AND METHOD

In this study, a bibliometric analysis method was applied to determine scientific trends in the field of remote sensing and satellite technologies for crop mapping. Bibliometric analysis is a quantitative method used to evaluate publication trends, academic collaborations, and scientific impacts within a specific research area. The study utilized the Scopus database, which includes international academic publications, and a systematic search was conducted based on predefined keywords. Scopus is a multidisciplinary database that covers various sources such as scientific journals, conference proceedings, and books. Researchers use this database to access the latest studies in their fields and conduct comprehensive literature reviews. Additionally, Scopus is frequently preferred for citation analysis and research performance evaluation (Falagas et al., 2008). The workflow followed in this study is illustrated in Figure 1.




Figure 1. Workflow diagram

During the data collection process, only academic publications from 2018 to 2025 were included using the keywords "remote sensing," "satellite," and "crop mapping" (Figure 2). VOSviewer software was used to perform bibliometric analyses (Zupic and Čater, 2015). Van Eck and Waltman (2010) stated that VOSviewer is a user-friendly software developed for bibliometric mapping and visualizing relationships within scientific literature. Using this software, detailed examinations were conducted by visualizing author collaborations, academic connections between countries, citation analyses, and keyword network structures.

The screenshot displays the Scopus search interface. At the top left is the Scopus logo. To the right is a search bar with a magnifying glass icon and the word 'Search'. Below this is a light blue banner with the text: 'Welcome to a more intuitive and efficient search experience. [See what is new](#)'. To the right of the banner is a toggle for 'Advanced query'. Below the banner are two search input fields. The first is labeled 'Search within' and has a dropdown menu showing 'Article title, Abstract, Keywords'. The second is labeled 'Search documents *' and contains the query: '"remote sensing" AND "satellite" AND "crop mapping"'. To the right of the second field is a close button (X). Below these fields are buttons for '+ Add search field', 'Reset', and 'Search' (with a magnifying glass icon). At the bottom of the search area are navigation tabs: 'Documents', 'Preprints', 'Patents', 'Secondary documents', and 'Research data'. A 'Beta' badge is visible above the 'Documents' tab. Below the search area, it states '175 documents found'.

Figure 2. Selection of search criteria in Scopus database interface

Within the scope of the bibliometric evaluation, the methodology focused on examining the distribution of academic publications over time, scientific productivity across different countries, citation analyses, and research trends. Based on the obtained data, this study aims to reveal the current state of academic research in remote sensing and crop mapping and provide a scientific framework to guide future research. Accordingly, this bibliometric analysis process seeks to answer the following questions:

1. What trends are observed in the distribution of studies on remote sensing, satellites, and crop mapping between 2018 and 2025? Has the research volume in this field increased or decreased?
2. How is academic productivity geographically distributed in this field? Which countries have published the most studies on remote sensing and crop mapping?
3. From a scientific impact perspective, which countries have received the highest number of citations, and how is the academic collaboration network structured among these countries?
4. When analyzing scientific impact at the author level, who are the top 10 most cited authors, and how are their total document and citation counts distributed?
5. When examining the thematic distribution of publications, which topics are most frequently addressed in the field of remote sensing and crop mapping?
6. Based on keyword analyses, which concepts are prominent in the field of remote sensing and crop mapping, and how do these concepts form a network structure?

RESULTS AND DISCUSSION

Between 2018 and 2025, a bibliometric analysis conducted on Scopus regarding crop mapping, remote sensing, and satellite technologies identified 175 publications that met the specified criteria. When examining the distribution of publication types, research articles constituted the largest share (132, 75.4%), followed by conference papers (35, 20%), review papers (5, 2.9%), book chapters (2, 1.1%), and editorial comment (1, 0.6%).

The predominance of research articles highlights the critical role of peer-reviewed journal publications in advancing the field. The 20% share of conference papers indicates that researchers frequently present their findings at scientific meetings, demonstrating the dynamic development of the field. However, the limited number of review studies (2.9%) suggests a need for comprehensive studies that summarize existing knowledge and identify research gaps. The low representation of book chapters and editorial comments further indicates that this topic is primarily addressed through scientific journal articles and conference papers. The nature of

remote sensing and crop mapping research, which relies on technological innovations, emphasizes the rapid dissemination of findings through academic journals and conferences. The yearly distribution of publications over the last ten years is presented in Figure 3.




Figure 3. Number of publications in the last 10 years

The highest number of publications occurred in 2024 (46 publications), while the lowest was in 2016 (3 publications), with an annual average of approximately 19 publications. These findings indicate a significant increase in publication numbers in 2024. A gradual rise has been observed since 2016, except for 2023, with 2024 marking the peak. Since 2025 is not yet complete, the current four publications for this year may increase by the end of the year.

Since 2018, the majority of academic studies have been published in English (161 publications). Chinese is the second most common language, with 12 publications. French and Russian have only one publication. This distribution reflects the dominance of English in global academic literature, ensuring broad international accessibility. The relatively high number of Chinese publications may indicate the increasing impact of research from China. The limited number of French and Russian publications suggests that studies in these languages remain scarce or focus on specific regional contexts. This language distribution confirms that the international academic community prefers English for accessibility and impact, while publications in other languages may contribute significantly at regional or national levels.

Analyzing the geographical distribution of publications, contributions from 58 different countries have been identified in Figure 4. China leads with 58 publications (30.7%), likely due to state-supported projects and investments in remote sensing technologies. The United States follows with 27 publications (14.3%), and India, with 19 publications (10.1%), stands out for its scientific productivity. In Europe, Italy, Germany, and Canada each contributed 11 publications, while France (7), Spain (5), and the United Kingdom (4) also conducted significant research in this field. In Africa and South America, countries like Nigeria, Morocco, Brazil, and Egypt show limited but growing academic interest. Turkey has contributed two publications, indicating a potential increase in regional interest.




Figure 4. Distribution of publications related to remote sensing, satellite and crop mapping terms by country.

These findings demonstrate the global distribution of crop mapping and remote sensing research. While some countries exhibit high scientific productivity, the field is still developing in other regions. Future initiatives should focus on increasing international collaborations, promoting open-access publications, and supporting scientific productivity in developing countries to advance research from a broader perspective.

According to citation analysis, China leads with 1,686 citations, highlighting its global impact in remote sensing, crop mapping, and satellite technologies. The United States follows with 923 citations, reinforcing the dominance of these two countries in scientific research. Among European nations, Germany (288), Italy (254), Belgium (350), France (122), and the Netherlands (132) have notable citation counts. Belgium's strong position with 350 citations suggests that regional academic research holds significant international influence. Iran ranks high with 289 citations, indicating growing recognition of Middle Eastern research in this field. Ukraine, with 172 citations, stands out as a country with increasing international scientific contributions. Canada, with 157 citations, remains an influential research hub in North America, while the Netherlands and France have relatively lower citation counts. The distribution of the top 10 most-cited countries is shown in Figure 5.




Figure 5. Top 10 most cited countries

These findings highlight the continued leadership of China and the United States in scientific research while demonstrating the significant contributions from Europe and the Middle East. Citation counts reveal that scientific productivity should be evaluated not only by the number of publications but also by the global academic impact of the studies.

A co-authorship analysis conducted using VOSviewer identified 46 countries with at least one publication and one citation. Among these, 35 countries had direct collaboration links. When evaluating the total strength of co-authorship connections for each of these 35 countries, China had the highest number of connections. This result is attributed not only to China's dominance in publication and citation counts but also to its central role in international research collaborations. Figure 6 presents the visualization of the co-authorship analysis along with its density analysis.




Figure 6. (a) Cross country co-authorship network visualization (b) cross country co-authorship density network visualization

China's emergence as a key player in the global research network is linked to its large-scale projects and extensive academic collaborations. The findings indicate that scientific productivity is influenced not only by publication and citation numbers but also by international academic partnerships. Countries engaged in widespread collaborative research tend to hold stronger positions in the scientific ecosystem (Figure 6).

The top 10 most-cited researchers in crop mapping using remote sensing and satellite imagery, along with their publication and citation counts, are presented in Table 1.

Table 1. Number of documents and citations of the first 10 authors in the citation ranking

Authors	Documents	Citations
Zheng Niu	3	290
Haifeng Tian	2	276
Kristof Van Tricht	1	262
Anne Gobin	1	262
Sven Gilliams	1	262
Isabelle Piccard	1	262
Liping Di	3	229
Jingfeng Huang	5	226
Ni Huang	1	191
Jie Pei	1	191

According to the table, Zheng Niu stands out as the researcher with the highest total citation count, with 3 publications and a total of 290 citations. He is followed by Haifeng Tian, who has 2 publications and 276 citations. Notably, Kristof Van Tricht and his colleagues (Anne Gobin, Sven Gilliams, and Isabelle Piccard) have produced one of the most highly cited individual studies, receiving 262 citations from a single publication. This indicates that their research on the synergistic use of Sentinel-1 and Sentinel-2 data has made a significant impact in the academic field. Other researchers, such as Liping Di (3 publications, 229 citations) and Jingfeng Huang (5 publications, 226 citations), stand out in terms of productivity. Particularly, Jingfeng Huang has the highest number of documents (5), but his total citation count of 226 is relatively lower. This suggests that Huang has conducted extensive research on various topics, but none of his studies have received exceptionally high citations individually. Lastly, Ni Huang and Jie Pei are listed with a single publication receiving 191 citations. These authors contributed to Tian et al.'s (2019) study on winter crop mapping.

The analysis results contribute to identifying prominent studies in the literature and understanding their academic interactions. The findings indicate that authors such as Van Tricht (2018), Tian (2019), Sun (2019), and Xu (2020) are represented by large nodes and are among the most highly cited works. This demonstrates that these authors have made significant contributions to the field and are frequently referenced in the literature. Additionally, clusters displayed in different colors on the network indicate research groups concentrating on specific thematic areas (Figure 7).




Figure 7. Network visualization of authors' citation density

In general, citation analyses show that studies on the use of satellite data in agricultural mapping have resonated widely within the academic community and that certain researchers have emerged as key figures in this field. Particularly, topics such as multi-source data utilization and deep learning methods appear to be common factors in highly cited studies.

Table 2. Number of documents and citations of the first 10 authors in the citation ranking

No	Authors	Title	Year	Citations
1	Van Tricht et al.,	Synergistic use of radar sentinel-1 and optical sentinel-2 imagery for crop mapping: A case study for Belgium	2018	262
2	Tian et al.,	Mapping winter crops in China with multi-source satellite imagery and phenology-based algorithm	2019	191
3	Xu et al.,	DeepCropMapping: A multi-temporal deep learning approach with improved spatial generalizability for dynamic corn and soybean mapping	2020	169
4	Sun et al.,	Using of multi-source and multi-temporal remote sensing data improves crop-type mapping in the subtropical agriculture region	2019	119
5	Zhang et al.,	Mapping up-to-date paddy rice extent at 10 M resolution in China through the integration of optical and synthetic aperture radar images	2018	110
6	Hao et al.,	Transfer Learning for Crop classification with Cropland Data Layer data (CDL) as training samples	2020	105
7	Mazzia et al.,	UAV and machine learning based refinement of a satellite-driven vegetation index for precision agriculture	2020	97
8	Song et al.,	An evaluation of Landsat, Sentinel-2, Sentinel-1 and MODIS data for crop type mapping	2021	96
9	Tian et al.,	Summer Maize Mapping by Compositing Time Series Sentinel-1A Imagery Based on Crop Growth Cycles	2021	85
10	Csillik et al.,	Object-based time-constrained dynamic time warping classification of crops using Sentinel-2	2019	81

Table 2 presents the authors, titles, publication years, citation counts, and reference numbers of the most highly cited studies. The most cited study, conducted by Van Tricht et al. (2018), examines the impact of integrating Sentinel-1 and Sentinel-2 data for agricultural mapping and has received 262 citations. This research highlights the significant contributions of combining radar and optical data for precision agriculture and crop identification.

Tian et al. (2019) rank second with 191 citations for their study on mapping winter crops in China using multi-source satellite data and algorithms that consider plant growth stages. Xu et al. (2020) rank third with their study, which employs a deep learning-based multi-temporal model for mapping maize and soybeans, receiving 169 citations. Additionally, studies by Sun et al. (2019), Zhang et al. (2018), Hao et al. (2020), Mazzia et al. (2020), Song et al. (2021), Tian et al. (2021), and Csillik et al. (2019) focus on mapping different plant species, applying deep learning approaches, conducting precision agriculture analyses using Unmanned Aerial Vehicles (UAVs), and performing comparative analyses of various satellite data sources.

These findings suggest that the most highly cited studies in agricultural mapping predominantly utilize Sentinel and Landsat data, conduct multi-temporal analyses, and integrate machine

learning techniques. Particularly, the combination of radar and optical data and the adoption of deep learning approaches have significantly advanced precision agriculture applications.

The bibliometric analysis revealed that publications are largely concentrated in specific subject areas. Earth and Planetary Sciences (26.9%) and Computer Science (18.6%) are the most productive disciplines in terms of publication output, followed by Agriculture and Biological Sciences (13.9%), Engineering (10.3%), and Environmental Science (7.5%). These five disciplines together account for 77.2% of the total publications. In contrast, fields such as Neuroscience, Medicine, Business, Chemistry, and Decision Sciences have very limited publications. The particularly low representation of multidisciplinary research (only 0.8%) suggests that interdisciplinary studies have not yet developed sufficiently. However, the increasing integration of data analytics and modeling methods from disciplines such as Computer Science and Engineering with other scientific fields has been observed. The distribution of publications by research areas is illustrated in Figure 8.




Figure 8. Distribution of publications in terms of study subject

Overall, a significant disparity exists between fundamental sciences (Physics, Chemistry, Mathematics) and applied sciences (Engineering, Environmental Sciences, Earth Sciences) in terms of publication volume. Additionally, the concentration of publications in certain disciplines highlights the need for a more balanced approach to research direction. A co-occurrence author keyword analysis using VOSviewer identified 95 keywords appearing at least twice, forming 12 distinct clusters. This analysis provides a crucial framework for understanding the key topics in crop mapping and remote sensing research, as well as academic trends among researchers. The findings show that the most frequently used keywords are "crop mapping" (54 occurrences, total link strength: 148) and "remote sensing" (53 occurrences, total link strength: 147), indicating that these two terms form the foundation of the field. This suggests a strong relationship between crop mapping and remote sensing methods, with researchers focusing heavily on agricultural mapping through satellite data. The prominence of the keyword "Sentinel-2" (28 occurrences, total link strength: 94) highlights the widespread use of Sentinel-2 data, provided by the European Space Agency, in crop mapping studies. The frequent occurrence of "random forest" (25 occurrences, total link strength: 79) among the keywords demonstrates the significant role of machine learning methods in agricultural

Encouraging multidisciplinary studies could lead to a more balanced distribution of publications across different fields. Combining environmental and engineering sciences with data science and artificial intelligence applications would be a significant step toward boosting academic productivity.

An analysis at the country level reveals that China has the highest number of publications, positioning it as the leader in academic productivity. The United States and India follow China in terms of publication count. While European countries contribute significantly with their technological infrastructure, regions such as Africa and South America exhibit limited but growing research interest. The relatively low number of publications from Turkey suggests that more scientific studies should be conducted in this field. Citation analysis indicates that the most highly cited publications predominantly originate from China, with the United States and European countries also producing highly cited works. Scientific collaboration analysis shows an increasing trend in multi-author studies and highlights that international co-authorship enhances the impact of scientific publications. Academic network analyses point to China as the country with the most research collaborations. Among the most cited researchers, Zheng Niu leads in total citations, while Haifeng Tian and Kristof Van Tricht also achieve high citation counts. Notably, Van Tricht's single publication receiving 262 citations underscores its significant impact in the field.

Based on these findings, increasing international collaborations, diversifying publication types, and supporting the scientific productivity of developing countries could further strengthen the research field. Expanding inter-country collaborations and promoting open-access publications would facilitate more effective knowledge sharing within the global scientific community. These results indicate that the field of vegetation mapping, remote sensing, and satellite technologies is rapidly evolving and innovative. Future studies incorporating more systematic reviews and interdisciplinary approaches could provide a more holistic perspective on advancements in the field. Additionally, the increasing use of machine learning, artificial intelligence, and high-resolution satellite imagery could enable future research to integrate automated data processing and real-time monitoring techniques. Overall, scientific activities in vegetation mapping, remote sensing, and satellite technologies have increased in recent years, yet they remain dominated by certain countries and research groups. In the future, promoting international partnerships and supporting open-access policies will be critical for expanding the scientific influence of this field across a broader geographical scope.

REFERENCES

- Bégué, A., Arvor, D., Bellon, B., Betbeder, J., De Aballeyra, D., PD Ferraz, R., ... & R. Verón, S. (2018). Remote sensing and cropping practices: A review. *Remote Sensing*, 10(1), 99.
- Csillik, O., Belgiu, M., Asner, G. P., & Kelly, M. (2019). Object-based time-constrained dynamic time warping classification of crops using Sentinel-2. *Remote sensing*, 11(10), 1257.
- Falagas, M. E., Pitsouni, E. I., Malietzis, G. A., & Pappas, G. (2008). Comparison of PubMed, Scopus, web of science, and Google scholar: strengths and weaknesses. *The FASEB journal*, 22(2), 338-342.
- Hao, P., Di, L., Zhang, C., & Guo, L. (2020). Transfer Learning for Crop classification with Cropland Data Layer data (CDL) as training samples. *Science of The Total Environment*, 733, 138869.
- Khanal, S., Kc, K., Fulton, J. P., Shearer, S., & Ozkan, E. (2020). Remote sensing in agriculture—accomplishments, limitations, and opportunities. *Remote Sensing*, 12(22), 3783.

- Mazzia, V., Comba, L., Khaliq, A., Chiaberge, M., & Gay, P. (2020). UAV and machine learning based refinement of a satellite-driven vegetation index for precision agriculture. *Sensors*, 20(9), 2530.
- Mulla, D. J. (2013). Twenty five years of remote sensing in precision agriculture: Key advances and remaining knowledge gaps. *Biosystems engineering*, 114(4), 358-371.
- Sishodia, R. P., Ray, R. L., & Singh, S. K. (2020). Applications of remote sensing in precision agriculture: A review. *Remote sensing*, 12(19), 3136.
- Song, X. P., Huang, W., Hansen, M. C., & Potapov, P. (2021). An evaluation of Landsat, Sentinel-2, Sentinel-1 and MODIS data for crop type mapping. *Science of Remote Sensing*, 3, 100018.
- Sun, C., Bian, Y., Zhou, T., & Pan, J. (2019). Using of multi-source and multi-temporal remote sensing data improves crop-type mapping in the subtropical agriculture region. *Sensors*, 19(10), 2401.
- Tian, H., Huang, N., Niu, Z., Qin, Y., Pei, J., & Wang, J. (2019). Mapping winter crops in China with multi-source satellite imagery and phenology-based algorithm. *Remote sensing*, 11(7), 820.
- Tian, H., Qin, Y., Niu, Z., Wang, L., & Ge, S. (2021). Summer maize mapping by compositing time series sentinel-1A imagery based on crop growth cycles. *Journal of the Indian Society of Remote Sensing*, 49, 2863-2874.
- Van Eck, N., & Waltman, L. (2010). Software survey: VOSviewer, a computer program for bibliometric mapping. *scientometrics*, 84(2), 523-538.
- Van Tricht, K., Gobin, A., Gilliams, S., & Piccard, I. (2018). Synergistic use of radar Sentinel-1 and optical Sentinel-2 imagery for crop mapping: A case study for Belgium. *Remote Sensing*, 10(10), 1642.
- Wu, B., Zhang, M., Zeng, H., Tian, F., Potgieter, A. B., Qin, X., ... & Loupian, E. (2022). Challenges and opportunities in remote sensing-based crop monitoring: a review. *Natl Sci Rev* 10 (4): nwac290.
- Xu, J., Zhu, Y., Zhong, R., Lin, Z., Xu, J., Jiang, H., ... & Lin, T. (2020). DeepCropMapping: A multi-temporal deep learning approach with improved spatial generalizability for dynamic corn and soybean mapping. *Remote Sensing of Environment*, 247, 111946.
- Zhang, X., Wu, B., Ponce-Campos, G. E., Zhang, M., Chang, S., & Tian, F. (2018). Mapping up-to-date paddy rice extent at 10 m resolution in china through the integration of optical and synthetic aperture radar images. *Remote Sensing*, 10(8), 1200.
- Zupic, I., & Čater, T. (2015). Bibliometric methods in management and organization. *Organizational research methods*, 18(3), 429-472.

TISSUE CULTURE STUDIES ON *CYCLAMEN* SPP.

Aslihan KULOGLU

Dicle University, Diyarbakır Agricultural Vocational School, Department of Seed Technology, 21280 Diyarbakır, Türkiye

Kocaeli University, Institute of Natural and Applied Sciences, Department of Horticultural Sciences, 41001 Kocaeli, Türkiye

ORCID ID: <https://orcid.org/0000-0002-7176-0838>

Aysun CAVUSOGLU

Kocaeli University, Faculty of Agriculture, Department of Plant Protection, 41285 Kocaeli, Türkiye

Kocaeli University, Institute of Natural and Applied Sciences, Department of Horticultural Sciences, 41001 Kocaeli, Türkiye

ORCID ID: <https://orcid.org/0000-0001-6921-7097>

ABSTRACT

Over the recent years, a lot of research has been studied on the popular ornamental plant cyclamen, especially in the area of tissue culture. Enhancing the growth, development, and genetic diversity of *Cyclamen* species has been made possible via tissue culture, which also provides an effective means of mass multiplication and the preservation of valuable genotypes. New developments have also looked into the possibility of using *in vitro* methods to improve genetics through somaclonal variety and conserve endangered *Cyclamen* species. According to certain research, tissue culture methods might cause somatic mutations, which can result in the emergence of plants with distinctive characteristics like improved flower color or resistance to disease. To guarantee survival and healthy growth, researchers stress the significance of maximizing ambient conditions and introducing plants to soil conditions gradually. Another difficulty that must be addressed for tissue culture applications to be successful is the occurrence of physiological diseases, such as hyperhydricity. This review discusses the possibility of tissue culture as a method to improve cyclamen propagation and conservation by combining the results of multiple investigations. In order to further improve the quality and diversity of cyclamen plants, it draws attention to current issues and potential research avenues. These include developing disease-resistant cultivars, optimizing growth conditions, and incorporating molecular techniques. Tissue culture can contribute significantly to the long-term viability of cyclamen production in both ornamental and conservation contexts by increasing the effectiveness of propagation techniques and promoting genetic variety.

Key Words: Cyclamen, Tissue Culture, Sustainability

INTRODUCTION

Cyclamen is a common geophyte cultivated throughout temperate zones (Takamura, 2007), with species growing in Turkey, Lebanon, Syria, Cyprus, Greece, Israel, and Iran, as well as enclaves in North Africa (Grey-Wilson, 2003). Cyclamen, known for its exquisite blossoms, became popular in the nineteenth century. Although wild *Cyclamen persicum* plants typically

flower from winter to spring, the production time of commercial flowering cyclamen plants varies by nation, depending on demand and climatic conditions (Takamura, 2007).

Cyclamen tissue culture studies have looked into a variety of methods for effective propagation. While vegetative propagation, including micropropagation, is not strictly a breeding technique, it is extremely important in plant breeding programs since it allows for the cloning of bred cyclamen plants. Nakayama (1980) attempted vegetative propagation by notching, but *in vivo* techniques proved difficult. Thus, researchers have attempted to develop dependable tissue culture procedures.

Mayer-Hörster (1956) was the first to report *in vitro* proliferation via organogenesis from tuber explants, but systemic bacteria in these explants slowed the development of tissue culture techniques. To address this issue, alternate explants containing fewer microorganisms were identified, including etiolated petiole sections (Ando & Murasaki, 1983) and aseptic seedling tissue (Wainwright & Harwood, 1985). In addition, leaf blade tissue has been employed in rare circumstances.

Cyclamen plants are a major topic of debate in these works. The agronomic and biotechnological developments in cyclamen cultivation are the main topic of this review paper, which also highlights tissue culture methods and how they are used in plant improvement and propagation. The research also highlights the significance of sustainability in tissue culture procedures, emphasizing how *in vitro* propagation can lessen environmental effect by encouraging the conservation of endangered species and eliminating the need for collecting wild plants.

In this review article, the tissue culture studies on cyclamen are examined in recent years and the studies conducted are as follows.

Direct and Indirect Organogenesis in Tissue Culture for *Cyclamen* spp.

The *in vitro* production of tuberous roots from the root ends of rooted tuberous stems devoid of shoots in *Cyclamen persicum* was investigated as a vegetative propagation technique in the 2004 study by Lim and Junzo. Cyclamen is frequently propagated from seedlings in Japan, however this method is expensive because of the high cost of seeds and their sluggish, uneven germination. The researchers looked for a technique for genetically homogeneous propagation in order to solve this problem. As explants, the researchers cultivated rooted tuberous stems devoid of shoots on conditions containing sucrose and benzylaminopurine (BA). The root tips connected to the tuberous stem developed into tuberous roots after 30 days of culture. The medium's composition determined how these roots developed. The 1/3 MS medium with 0,05 mg/L NAA, 0,5 mg/L BA, and 5% sucrose had the highest percentage of tuberous root development. The study also discovered that the location of the root cut in the explants had a substantial impact on the tuberous roots' growth rates. Cutting off a tiny section of the root end at the beginning of tissue culture produced the best development.

Bürün & Şahin (2009) investigated *Cyclamen alpinum* (formerly *C. trochopteranthum*) germination *in vitro* and *in vivo*. The seeds were provided from red pine forest around the Gökbel-Dalyan regions on May. The seeds were then stored at 20°C with and without capsules. Different temperatures were used to examine the germination. The findings indicated that after 8 months of storage, germination peaked in December (95%) and that seeds kept for 5–6 months had germination rates ranging from 35% to 87%. While keeping seeds in capsules increased *in vivo* germination rates on filter paper and in peat-perlite-sand combinations, the storage method had no effect on *in vitro* germination in Murashige-Skoog medium. With 95% germination *in vitro* at both temperatures and 93% *in vivo* at 20°C, germination was best at 15°C and 20°C in the dark. After 8 to 9 months of storage, the maximum germination rates 94% *in vitro* and 83%

in vivo occurred. Furthermore, it was discovered that the best media for *in vivo* germination was a peat-perlite-sand mixture, which produced 93% germination for *C. alpinum*.

The effects of various tissue culture mediums and plant hormones on gynogenesis in *Cyclamen* species specifically, *Cyclamen persicum* and *Cyclamen pseudibericum* were examined by Tütüncü et al. (2019). The study's objective was to maximize *in vitro* haploidization conditions in cyclamen, a genus that has breeding difficulties because of things like ploidy variances and self-incompatibility. Ovule explants grown on ½MS and B5 medium supplemented with different quantities of 6-(γ,γ -dimethylallylamino) purine (2iP) and 2,4-dichlorophenoxyacetic acid (2,4-D) were evaluated by the researchers. According to the results, *Cyclamen persicum* produced the most callus (70%) on B5 medium when it contained 1,5 mg/L 2,4-D and 0,5 mg/L 2iP, whereas *Cyclamen pseudibericum* produced 48% on B5 medium when it contained 1 mg/L 2,4-D and 0,3 mg/L 2iP. *Cyclamen persicum* had a 55% success rate for embryo induction when callus was regenerated on ½MS medium using 1,5 mg/L 2,4-D and 0,8 mg/L 2iP. *Cyclamen pseudibericum* did not exhibit embryo induction on any of the tested media, nevertheless. The study found that B5 medium was better for callus formation, but the ½MS medium with high auxin and low cytokinin concentrations was more successful for embryo induction. 42 plantlets were eventually recovered from callus that had been regenerated on ½MS medium with 0,5 mg/L 2iP and 2,0 mg/L 2,4-D.

Shahabi et al. (2023) aimed to develop an efficient protocol for microtuber formation and direct shoot regeneration in *Cyclamen persicum* Mill. using *in vitro* leaves and tubers from the cultivars 'Pure White', 'Neon Pink', and 'Dark Violet'. In this study, explants were cultured on ½ Murashige and Skoog (MS) basal medium supplemented with various concentrations of benzyladenine (BA) or thidiazuron (TDZ) for direct shoot regeneration or microtuber formation, and the number of shoots and microtubers per explant was recorded. The highest direct shoot regeneration (27.1 shoots per explant) was obtained from leaf explants of the 'Pure White' cultivar on medium supplemented with 0,25 mg/L TDZ and 0.1 mg/L α -naphthaleneacetic acid (NAA), while the highest microtuber proliferation (17.1 microtubers per explant) was observed in tuber cut cultures of the 'Neon Pink' and 'Pure White' cultivars grown in darkness with 1.0 mg/L BA and 0.1 mg/L NAA. Additionally, the best rooting results were achieved with 0.5–1,0 mg/L NAA, and 58% of the rooted shoots successfully acclimated to greenhouse conditions.

Using *in vitro* anther and ovule culture, Sevindik et al. (2023) sought to create an effective haploid plant regeneration strategy for both the commercial Melody F1 cultivar and wild *Cyclamen persicum*. This was accomplished by using DAPI staining to identify microspores at the uninuclear stage and by cold pre-treating flower buds prior to culture. The best conditions for anther and ovule development were determined by testing several culture mediums. For *C. persicum*, the optimum ovule culture procedures differed by genotype, however the highest haploid embryo production (100%) was achieved using B5 medium with 1 mg/L NAA. Within four months, the regenerated plants had effectively adapted to the soil and flow cytometry had shown spontaneous double haploidization. This work offers a very efficient technique for producing haploid and double-haploid.

Al-Khafaji et al. (2024) sought to determine the ideal microspore developmental stage and bud size in order to create haploid *Cyclamen coum* plants using anther culture. Using 2% acetocarmine staining, the late uninuclear stage was identified, and the associated bud size was measured to be between 7.64 and 8.23 mm. After a 4-day cold pre-treatment at 4°C, anthers were cultivated on B5 medium enriched with NAA, 2,4-D, and kinetin. In the first year of the study, twelve various media were investigated; in the second year, that number was reduced to seven. In combination with NAA (1 or 2 mg/L). These findings imply that embryo development was successfully induced by cold pre-treated anthers collected at the proper flower bud size, and that the best medium for *Cyclamen coum* embryo regeneration was B5 enriched with NAA and kinetin.

Somatic Embryogenesis in Tissue Culture for *Cyclamen* spp.

Furukawa et al. (2002) work used aseptic seedlings from 18 wild *Cyclamen* species to create somatic embryos. The seeds were first cultivated at 15-20°C in the dark until they germinated, after which they were moved to a 16/8 light/dark photoperiod at 24°C. 3.0 g/L gellan gum and 30.0 g/L sucrose were present in the modified MS medium that was used to cultivate the aseptic seedlings. Aseptic seedlings were successfully obtained from 14 wild *Cyclamen* species, including *C. africanum*, *C. cilicium*, *C. coum*, *C. cyprium*, *C. graecum*, *C. hederifolium* var. *album*, *C. libanoticum*, *C. mirabile*, *C. parviflorum*, *C. persicum*, *C. pseudibericum*, *C. purpurascens*, *C. rohlfsianum*, and *C. trochopteranthum*, but seeds from *Cyclamen balearicum*, *C. creticum*, *C. intaminatum*, *C. balearicum*, *C. creticum*, *C. intaminatum*, and *C. repandum* did not germinate. These seedlings' sectioned petioles and leaves were inoculated into two different kinds of induction media at 24°C in the dark. Calli were absent from, *C. cyprium*, *C. libanoticum*, and *C. parviflorum*, three of the fourteen species. After subculturing, the calli were moved to a somatic induction medium devoid of plant growth regulators. Species differences in somatic embryogenesis potential, callus type, and formation frequency were significant. Shoot production was accomplished for the first time in *C. africanum* and *C. hederifolium* var. *album*, while somatic embryogenesis was accomplished effectively in *C. africanum*, *C. cilicium*, *C. persicum*, *C. rohlfsianum*, and *C. trochopteranthum*.

Prange et al. (2010) published one of the first research on the regeneration of *Cyclamen graecum*, *Cyclamen mirabile*, and *Cyclamen trochopteranthum* via somatic embryogenesis. Their research built embryogenic callus cultures from seedling tissue and created a regeneration system that goes from protoplast to plant. The ability to develop embryogenic callus and regenerate through somatic embryogenesis was discovered to be highly genotype-dependent. *C. mirabile* produced up to 1461 somatic embryos from 0.5 g of callus, which were impacted by culture media composition, plant growth regulators, CaCl₂, and activated charcoal. Furthermore, up to 1.4×10^6 protoplasts were recovered from 1 g of *C. graecum* cell suspension, exhibiting varied growth responses in agarose and alginate embedding. Somatic embryos were effectively regenerated from both protoplast culture systems, grown into plantlets, and adapted to greenhouse conditions. These findings provided light on the regeneration potential of wild *Cyclamen* species and could serve as a foundation for future fusion research.

In order to improve the clonal propagation of *Cyclamen persicum*, Hoenemann et al. (2010) studied somatic embryogenesis. Developmental abnormalities and non-embryogenic callus formation were among the difficulties in the process that were examined in the study. In order to compare zygote and somatic embryos, as well as embryogenic and non-embryogenic cultures, gene expression analysis was carried out at different developmental stages of embryogenic cell cultures. 417 genes were identified as differentially expressed using cDNA microarrays and real-time PCR, with a significant overrepresentation of genes encoding apoplast-related enzymes. These findings were validated by histological analyses. The study highlighted the apoplast's important function in somatic embryogenesis and proposed that focusing on this element could enhance the procedure. In order to improve molecular control and increase the effectiveness of micropropagation in *C. persicum*, the authors also suggested physiological indicators.

Winkelmann et al. (2015) sought to improve *in vitro* somatic embryogenesis in *Cyclamen persicum* by comparing its metabolite profile to that of zygotic embryos in order to solve issues such as deformities, asynchronous development, and defects in maturation and germination. Rather of assessing different culture conditions and plant growth regulators, a metabolomic analysis was performed with zygotic embryos as a reference. Two sets of extracts from two to four biological replicates were examined, revealing and measuring 52 metabolites from various tissues, including endosperm and testa. The investigation indicated considerable differences, with zygotic embryos containing almost 40 times more proline than somatic embryos. Epicatechin, which is renowned for its ability to scavenge reactive oxygen species, was most

plentiful in the testa, but sucrose was substantially more concentrated in zygotic embryos. Furthermore, an unidentified trisaccharide was discovered to be greatly enriched in zygotic embryos, providing insights that may help improve *in vitro* somatic embryogenesis systems.

DISCUSSION/ FUTURE ASPECTS

Because of their varied cultivars, adaptability to different growing circumstances, and aesthetic appeal, cyclamen species are highly valued ornamental plants. But maintaining their sustainable production is still difficult, especially in light of commercial demands and environmental pressures. In this regard, tissue culture methods are essential for the conservation, genetic enhancement, and large-scale reproduction of *Cyclamen* species. Optimizing culture conditions, enhancing somaclonal variation control, and incorporating biotechnological developments like genome editing and metabolomic analyses should be the main goals of future study. Furthermore, improving commercial output will require the creation of economical and ecologically beneficial micropropagation techniques. Collaboration between researchers, breeders, and industry stakeholders can further support the adoption of tissue culture methods, ensuring both the conservation and economic viability of cyclamen cultivation in a sustainable manner. Furthermore, the implementation of sustainable production strategies will safeguard the survival of valuable *Cyclamen* species while satisfying the expanding demand in the ornamental plant market. To ensure that these advancements align with sustainable agricultural practices, future research should focus on optimizing *in vitro* culture conditions for various *Cyclamen* species, experimenting with new media formulations, and investigating the long-term ecological and economic impacts of large-scale tissue culture propagation.

REFERENCES

- Al-Khafaji, M., Yıldız, M., Koçak, M., & Tütüncü, M. (2024). Production of microspore-derived plants by anther culture of *Cyclamen coum*. *Black Sea Journal of Agriculture*, 7(6), 720–728. doi: [10.471115/bsagriculture.1554595](https://doi.org/10.471115/bsagriculture.1554595)
- Ando, T., & Murasaki, K. (1983). *In vitro* propagation of cyclamen by the use of etiolated petioles. *Technological Bulletin of the Faculty of Horticulture Chiba University*, 32, 1-5.
- Bürün, B., & Şahin, O. (2009). *In vitro* and *in vivo* germination of *Cyclamen alpinum* seeds. *Turkish Journal of Botany*, 33(4), 277–283. doi: [10.3906/bot-0709-8](https://doi.org/10.3906/bot-0709-8)
- Furukawa, K., Kakihara, F., & Kato, M. (2002). Somatic embryos produced from aseptic seedlings of wild *Cyclamen* species. *Journal of Society of High Technology in Agriculture*, 14 (2), 71–80.
- Grey-Wilson, C. (2003). *Cyclamen: a guide for gardeners, horticulturists and botanists*, New edition. London Batsford.
- Hoenemann, C., Richardt, S., Krüger, K., Zimmer, A. D., Hohe, A., & Rensing, S. A. (2010). Large impact of the apoplast on somatic embryogenesis in *Cyclamen persicum* offers possibilities for improved developmental control *in vitro*. *BMC Plant Biology*, 10, 77. doi: [10.1186/1471-2229-10-77](https://doi.org/10.1186/1471-2229-10-77)
- Lim, J. G., & Junzo, F. (2004). *In vitro* formation of tuberous roots from root ends in the rooted tuberous stem without shoots in *Cyclamen persicum* Mill. *Korean Journal of Plant Resources*, 7(3), 222-225.
- Mayer-Hörster, L. (1956). Wachstum and organbildung an *in vitro* kultivierten segmenten von *Pelargonium zonale* und *Cyclamen persicum*. *Planta*, 47, 401-446. doi: [10.1007/BF01911864](https://doi.org/10.1007/BF01911864)

- Nakayama, M. (1980). Vegetative propagation of cyclamen by notching of tuber II. Effects of scooping site and notching size on the regeneration of cyclamen tuber. *J. Japan. Soc. Hort. Sci.* 49 (2), 228-234.
- Prange, A. N. S., Bartsch, M., Serek, M., & Winkelmann, T. (2010). Regeneration of different *Cyclamen* species via somatic embryogenesis from callus, suspension cultures and protoplasts. *Scientia Horticulturae*, 125, 442-450. doi: 10.1016/j.scienta.2010.04.018
- Sevindik, B., İzgü, T., Tütüncü, M., Çürük, P., Sarı, N., & Mendi, Y. Y. (2023). Double-haploid plant production through anther and ovule culture of wild *Cyclamen persicum* Mill. and Melody F1 cyclamen cultivar. *In Vitro Cellular & Developmental Biology - Plant*, 59, 711–723. doi: [10.1007/s11627-023-10364-2](https://doi.org/10.1007/s11627-023-10364-2)
- Shahabi, M., Emadpour, M. & Moieni, A. (2023). Highly efficient microtuber formation, direct shoot regeneration, and root induction in *Cyclamen persicum* Mill. from *in vitro* seedling-derived tuber and leaf segments. *In Vitro Cellular & Developmental Biology-Plant*, 59, 475–482. doi: [10.1007/s11627-023-10353-5](https://doi.org/10.1007/s11627-023-10353-5)
- Takamura, T. (2007). Cyclamen. In: Anderson, N.O. (eds) Flower Breeding and Genetics. Springer, Dordrecht. 459-489. doi.org/[10.1007/978-1-4020-4428-1_16](https://doi.org/10.1007/978-1-4020-4428-1_16)
- Tütüncü, M., Özcan, M., & Mendi, Y. Y. (2019). Effects of different tissue culture media on gynogenesis in some cyclamen species. *Anadolu Journal of Agricultural Sciences*, 34, 239-249. doi: [10.7161/omuanajas.529247](https://doi.org/10.7161/omuanajas.529247)
- Wainwright, H., & Harwood, A.C. (1985) *In vitro* organogenesis and plant regeneration of *Cyclamen persicum* Mill. using seedling tissue. *J. Hort. Sci.* 60(3), 397-403. doi: 10.1080/14620316.1985.11515644
- Winkelmann, T., Ratjens, S., Bartsch, M., Rode, C., Niehaus, K., & Bednarz, H. (2015). Metabolite profiling of somatic embryos of *Cyclamen persicum* in comparison to zygotic embryos, endosperm, and testa. *Frontiers in Plant Science*, 6, 597. doi: 10.3389/fpls.2015.00597

AGRO-PRODUCTION CHARACTERISTICS OF SOILS IN THE NAKHCHIVAN AUTONOMOUS REPUBLIC (ON THE EXAMPLE OF SHAHBUZ AND ORDUBAD DISTRICTS)

Leyla IBRAHIMOVA

Nakhchivan State University, Department of Reclamation Engineering, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0005-3955-1074>

ABSTRACT

Introduction and Purpose: Our study area is one of the most productive agricultural regions of the country. Fruit growing is especially developed here. Shahbuz and Ordubad regions, which are also distinguished by their grain crops, are distinguished by the export of the fruits they produce. The productivity of lands used for gardening and grain cultivation for a long time decreases over time. For this reason, it is important to study the soils in these regions. Therefore, our study is dedicated to the current issue. **Materials and Methods:** During the study, the results of the studies conducted in the area, the field research method, and the processing of satellite images were used. ArcGIS software was used during the processing of satellite images. When evaluating the soils, both the parameters of the soils as a whole and their economic indicators were taken into account. The zoning and agro-production grouping carried out were based on the economic indicators of 1989, 2009, and 2019, namely, productivity per hectare and the degree of soil nutrient supply, soil layer thickness, slope inclination, mechanical composition of soils, and salinity degree. **Results:** High-quality soils, when used intensively under agricultural crops, ensure high and stable yields. **Discussion and Conclusion:** The lands used in agriculture in the Shahbuz and Ordubad districts are grouped as follows: high-quality, good-quality, medium-quality, low-quality and conditionally unsuitable lands. High-quality lands include: Mountain-meadow, mountain-meadow-steppe, mountain-brown soils. Such a group of soils is characterized by high natural fertility and profile. In such soils, the thickness of the humus layer reaches 50-60 cm, and the amount of humus reaches 4-9%. Such quality soils are evaluated with 100-81 points. The total area in the Ordubad district is 0.5 thousand ha, of which 0.2 thousand hectares or 40% are used for sowing and gardening, and 70% are used as high-quality hayfields and pastures. In the Shahbuz district, such lands account for 0.8 thousand ha, of which 50% are used for sowing and gardening, and 50% are used as high-quality pastures.

Keywords. agriculture, productivity, Nakhchivan AR, soils

INTRODUCTION

Agro-production groups of soils, types and varieties of soils, combined into groups by common environmental conditions, agronomic properties, similarity of qualitative features and fertility level (Amanova, Hajiyeva and Jafarova, 2024). Agro-production groups of soils are characterized by the uniformity of the necessary agrotechnical and melioration measures. For soils combined into one agro-production group, their suitability for the cultivation of certain agricultural crops is noted; soil properties that limit crop yield are identified; a set of necessary agrotechnical, melioration and anti-erosion measures is developed (Bhatia, Jha, Sarkar and Sarangi, 2023).

According to scientists, there are 3 categories of agro-production groups of soils:

In the 1st category, soils are grouped in accordance with the requirements of a specific agricultural crop;

In the 2nd category - in accordance with the requirements of individual groups of crops (row crops, cereal and cereal grains, technical non-row crops, etc.);

in the 3rd category – for all groups of crops.

The first 2 categories were developed only for crops that are demanding of soil conditions. The most widespread are the groups of the 3rd category (Amanova and Hajiyeva, 2023).

To be combined into one agricultural production group, soils must be characterized by similarity according to the following criteria (Payen, Basset-Mens and Perret, 2015):

- water-air and thermal properties and regimes, identified based on the assessment of the granulometric composition, composition and thickness of the humus horizon, as well as taking into account the geomorphological and hydrological conditions of the soil location (Urbano, Barquero, González-Andrés, 2022);
- nutrient regime and conditions for the application of fertilizers (content of mobile forms of nitrogen, phosphorus, potassium, content of organic matter and gross reserve of nutrients, reaction of the environment, oxidation-reduction conditions, content of microelements, etc.);
- the attitude of soils to cultivation (physical and mechanical properties of soils - plasticity, stickiness, hardness, shear resistance, cohesion; the possibility of crust formation and waterlogging; proximity of the dates of physical maturity of soils; features of deepening the arable layer, etc.);
- need for melioration measures (drainage, irrigation, gypsum application, liming, etc.) taking into account the characteristics of the hydrological regime and relief conditions;

METHODOLOGY

During the research, we used historical, mathematical-statistical, and field research methods. We conducted field research in Shahbuz and Ordubad districts. We also used materials from researchers who conducted research on the lands of the area. According to international experience, we divided the lands of the area into 5 areas according to agro-production groupings.

RESULT

When assessing the soils, the parameters of the soils as a whole, as well as their economic indicators, were taken into account.

The zoning and agro-production grouping carried out were based on the economic indicators of 1987, 2009, 2019, namely, productivity per hectare and the degree of soil nutrient supply, soil layer thickness, slope inclination, mechanical composition of soils, and salinization degree.

The soils used in agriculture in the Shahbuz and Ordubad districts are grouped as follows: high-quality, good-quality, medium-quality, low-quality and conditionally unsuitable soils.

Group I. High-quality soils:

These include mountain-meadow, mountain-meadow-steppe, mountain-brown soils. Such a group of soils is characterized by high natural fertility and profile.

In such soils, the thickness of the humus layer reaches 50-60 cm, and the amount of humus reaches 4-9%.

Such quality soils are evaluated with 100-81 points. In Shahbuz district, such soils are 0.8 thousand ha, of which 50% is arable and fallow land, and 50% is used as high-quality pasture.

Thus, high-quality soils provide high and stable yields with intensive use under agricultural crops. (Table 1).

Group II. Good quality soils:

These soils are evaluated with 80-61 points. They cover an area of 1.3 thousand ha in Shahbuz district. This group of soils is distributed on gently sloping slopes and in areas subject to weak erosion and occupies brown mountain-forest, mountain-brown, primitive mountain-meadow, and washed mountain-brown soils. On good quality soils, the specific weight of arable, fallow, and perennial crops is 45.7%, and hayfields and pastures is 54.3%.

The mechanical composition of this group of soils is clayey and medium clayey. The thickness of the humus layer varies between 20-30 cm, and its amount varies between 3-4.5%. It is moderately supplied with nutrients. In order to obtain high and stable yields from this group of soils, it is necessary to correctly use modern advanced agrotechnical rules.

Group III. Medium-quality soils

These soils were evaluated with 60-41 points. These soils are weakly and moderately exposed to erosion. In addition to the types and varieties we indicated in the previous group, irrigated gray-brown, alluvial, gray-brown soils are also included.

The thickness of the humus layer varies between 25-30 cm, and its amount varies between 2-3.0%. It is poorly supplied with nutrients. In particular, a lack of nitrogen is observed.

The total area in Ordubad district is 7.7 thousand ha, and in Shahbuz district - 4.3 thousand ha. The main land areas are used for cultivation and fallow, 22.1% in Ordubad, and 60.5% in Shahbuz.

In addition to improving agricultural culture on these high-quality lands, modern advanced agrotechnical rules should be used, rotational crops (especially soil conservation) should be given wide space, comprehensive measures to combat erosion should be implemented on a large scale, etc.

Table 1. Agro-production groups of soils in Ordubad region

Agro-production groups	Quality of soils		Total area Th.ha	Including								
	Bonitet score	Bonitet class		planted and laid to rest	perennial crops	Mowed areas	pastures	frest	Other soils			
High quality	100-81	X-IX	0,5	Th.ha %	60	-	20	20	20	20	20	70,5
				Th.ha	0,3	-	0,1	0,1	0,1	0,1	0,1	
Good quality	80-61	VIII-VII	3,8	Th.ha %	36,8	10,5	50	50	50	50	50	70,5
				Th.ha	1,4	0,4	1,9	1,9	1,3	4,4	0,3	
Medium quality	60-41	VI-V	7,7	Th.ha %	22,1	15,6	57,1	57,1	57,1	57,1	57,1	70,5
				Th.ha	1,7	1,2	4,4	4,4	1,3	2,5	0,1	
Low quality	40-21	IV-III	3,9	Th.ha %	25,6	10,3	64,1	64,1	64,1	64,1	64,1	70,5
				Th.ha	1,0	0,4	2,5	2,5	-	0,1	0,1	
Conditionally useless lands	20-10	II-I	0,8	Th.ha %	50	37,5	-	-	-	-	-	70,5
	low			Th.ha	0,4	0,3	-	-	-	-	-	
Other soils			87,2	Th.ha %								70,5
				Th.ha								

Group IV. Low-quality soils

These soils are evaluated with 40-20 points. The group of such quality soils includes chestnut, gray and gray-brown soils of plain areas. These soils occupy 3.9 thousand ha in Ordubad district, and 10.8 thousand ha in Shahbuz district.

In Shahbuz, 27.8% of these lands are under cultivation and fallow, and 71.3% are under hayfields and pastures. In Ordubad district, 64% are used as pastures, and the remaining areas are used as orchards.

Such soils are notable for their low nutrient content and lack of moisture.

Such soils are suitable for planting drought-resistant tree species such as almond, pistachio and. In order to restore fertility in such quality soils, a number of control measures should be implemented. The quality of such soils can be improved by using the achievements of modern agricultural technology and applying organic and mineral fertilizers.

Group V. Conditionally unsuitable soils

These soils are assessed with a score of 20-10 and lower.

Such soils occupy 0.8 thousand ha in the Ordubad region, and 0.2 thousand ha in Shahbuz.

The productivity of agricultural crops grown on such soils is very low. Conditionally unsuitable soils require agro-ameliorative work based on large investments.

Table.

CONCLUSION AND DISCUSSION

The total area of Ordubad district is 0.5 thousand ha, of which 0.2 thousand hectares or 40% are suitable for sowing. 70% of this is arable and fallow land, and the rest is used as high-quality hay and pasture.

In the territory of Ordubad district, the specific weight of arable, peaceful and perennial crops is 47.3%, and hayfields and pastures are 52.7%.

REFERENCES

- Amanova Sh.S., Hajiyeva G.N. (2023) Investigation of Natural Condition in Urban Landscapes of Plain Areas Based on GIS, C. R. Acad. Bulg. Sci., 76(11), 1679
- Amanova Sh.S., Hajiyeva A.Z., & Jafarova F.M. (2024). Preparation of future development scenarios of urban landscapes in accordance with natural and socio-economic conditions (on the example of the cities of the Kura-Araz lowland). *Visnyk of V. N. Karazin Kharkiv National University, Series Geology. Geography. Ecology*, (60), 305-322. <https://doi.org/10.26565/2410-7360-2024-60-22>
- Bhatia L., Jha H., Sarkar T., Sarangi P.K. (2023) Food Waste Utilization for Reducing Carbon Footprints Towards Sustainable and Cleaner Environment: A Review. *Int. J. Environ. Res. Public Health*, 20, 2318
- Payen, S., Basset-Mens, C., Perret, S. (2015) LCA of Local and Imported Tomato: An Energy and Water Trade-Off. *J. Clean. Prod.*, 87, 139–148
- Urbano B., Barquero M., González-Andrés F. (2022) The Environmental Impact of Fresh Tomatoes Consumed in Cities: A Comparative LCA of Long-Distance Transportation and Local Production. *Sci. Hortic.*, 301, 111126

PESTICIDE USE IN İĞDIR PROVINCE**Muhammet ÖTER**

Ağrı Ziraat Karantina Müdürlüğü, Ağrı, Türkiye

ORCID ID: <https://orcid.org/0009-0005-8037-4567>**Celalettin GÖZÜAÇIK**

İğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, 76000, İğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6543-7663>**ÖZET**

Pestisit, zararlı organizmaların engellemesi, kontrol altında tutulması ya da zararlarını azaltması için kullanılan madde ya da maddelerden oluşan kimyasal karışımlardır. Günümüzde tarımsal üretimde yaşanan sorunlardan birisi olan verim ve kaliteyi etkileyen hastalık, zararlı ve yabancı otlarla mücadelede en çok tercih edilen yöntem kimyasal mücadeledir. Türkiye’de 2023 yılında 57.766 ton, İğdır ilinde ise toplam 164.693 kg-lt pestisit kullanılmıştır. İğdır ilinde kullanılan bu pestisitlerin 108.000 kg/lt insektisit, 29.293 kg/lt fungusit, 21.250 kg/lt herbisit ve 5970 kg/lt akarisitir. İğdır ili mikroklima özelliklerinden dolayı birçok kültür bitkisinin yetişmesi için uygun vejetasyona sahiptir. İlde meyve, sebze, tahıl ve yem bitkileri yetiştiriciliği yapılmaktadır. Bununla birlikte ürünlerde görülen birçok zararlı (hastalık, böcek, akar, yabancı ot vb.) türe karşı, farklı bitki gelişim dönemlerinde uygun ya da uygun olmayan zaman ve dozlarda pestisit uygulamaları söz konusudur. Bu, zararlılara karşı kullanılan pestisitlerin etkisini kısa sürede göstermesi, kullanımının kolay olması, ürünün pazar değerinin artırılması düşüncesi gibi nedenlerle tercih edilmektedir. Yoğun pestisit kullanımının çevreye ve insan sağlığına kısa ve uzun vadede zararlarının da olduğu bilinmektedir. Günümüzde insan, çevre sağlığı ile biyolojik çeşitliliğin korunması ön plana çıkmıştır. Bu nedenlerle, tarımsal zararlılarla mücadelede pestisit tüketiminin azaltılması ve sürdürülebilir agro-ekosistemlerin oluşturulması için, daha bilinçli ve kontrollü kimyasal kullanılması önemlidir. Aynı zamanda zararlılarda direnç geliştirme riskleri bulunduğunu ve kalıntı sorunu, doğal dengenin tahribi gibi olumsuz etkilerde bu fikri desteklemektedir. Bu çalışmada İğdır ilindeki zararlılarla mücadelede pestisit kullanımına dikkat çekmek amacıyla ele alınmıştır.

Anahtar Kelimeler: İğdır, Pestisit, Tarım**ABSTRACT**

Pesticides are chemical mixtures used to prevent, control, or reduce the damage caused by harmful organisms. One of the key issues in modern agricultural production is the management of diseases, pests, and weeds that affect both yield and quality. Among the various methods for addressing these challenges, chemical control is the most commonly preferred approach. In 2023, a total of 57,766 tons of pesticides were used in Turkey, while in İğdır province, 164,693 kg/liters of pesticides were applied. The pesticides used in İğdır consisted of 108,000 kg/liters of insecticides, 29,293 kg/liters of fungicides, 21,250 kg/liters of herbicides, and 5,970 kg/liters of acaricides. Due to its microclimatic characteristics, İğdır province has favorable vegetation for the cultivation of a variety of crops, including fruits, vegetables, cereals, and forage plants.

However, many pests (such as diseases, insects, mites, weeds, etc.) are present in these crops, leading to pesticide applications at different stages of plant development, sometimes at inappropriate times or in incorrect doses. This approach is preferred due to the rapid effectiveness of pesticides, their ease of use, and the perception that they increase the market value of the crops. However, it is well-established that the intensive use of pesticides can have both short-term and long-term negative impacts on the environment and human health. Today, the protection of human and environmental health, as well as the conservation of biodiversity, has become a priority. Therefore, reducing pesticide consumption and promoting sustainable agro-ecosystems by using chemicals in a more controlled and informed manner is essential. Additionally, the risks of resistance development in pests, along with concerns about pesticide residues and the disruption of natural ecological balances, further support the need for more responsible pesticide use. This study aims to highlight the use of pesticides in pest management in Iğdır province.

Keywords: Iğdır, Pesticides, Agriculture

GİRİŞ

Tarımsal faaliyetler, tarih boyunca insanoğlunun yaşamını sürdürebilmesi için temel bir rol oynamış ve her dönemde büyük bir öneme sahip olmuştur. İnsanlık tarihi boyunca, tarım toplumların ekonomik yapısının ve kültürel gelişiminin şekillenmesinde belirleyici bir faktör olmuştur. Tarım, sadece insanların besin ihtiyaçlarını karşılamakla kalmaz, aynı zamanda hayvancılık, tekstil, ilaç, enerji ve sanayi gibi birçok sektöre de hammadde sağlar.

Günümüzde, dünya nüfusunun hızla artması ve çevresel değişiklikler, tarımın önemini bir kez daha gözler önüne sermektedir. 2050 yılına kadar dünya nüfusunun 9,7 milyara ulaşması beklenmektedir, bu da gıda üretiminin %70 oranında artırılmasını gerektirmektedir (Anonim, 2024). Ayrıca, yaşlanan nüfusun gıda talebindeki değişiklikler de dikkate alınmaktadır; 2050 yılına kadar yaşlı nüfusun %135 oranında artması beklenmekte ve bu durum, gıda pazarında önemli değişikliklere yol açacağı öngörülmektedir. Dünya Sağlık Örgütü (WHO), bu öngörüler ışığında, sürdürülebilir tarım uygulamalarının benimsenmesinin, gıda israfının azaltılmasının ve sağlıklı beslenme alışkanlıklarının teşvik edilmesinin önemini vurgulamaktadır. Bunun yanı sıra, iklim değişikliği ve çevresel faktörlerin tarım ve gıda güvenliği üzerindeki olumsuz etkilerinin azaltılabilmesi için küresel iş birliğinin gerekliliği de belirtilmektedir. Sınırlı doğal kaynaklar ve hızla artan dünya nüfusu göz önünde bulundurulduğunda, sürdürülebilir tarım ve sağlıklı gıda üretiminin önemi giderek daha da artmaktadır. Birim alandan elde edilen verimin artırılması ve kaliteli ürünlerin üretilmesi, tarımsal üretimde hastalıklar, zararlılar ve yabancı otlarla mücadeleyi gerektirmektedir. Bu mücadele, yalnızca verimliliği artırmaya değil, aynı zamanda sağlıklı ve güvenli gıda temin etmeye de yardımcı olmaktadır. Ancak, bu mücadelede kullanılan pestisitlerin etkili ve dikkatli bir şekilde yönetilmesi gerekmektedir. Pestisitler, ürünlerin korunmasını sağlarken, çevresel etkiler ve insan sağlığı üzerindeki olumsuz sonuçlar da göz önünde bulundurulmalıdır.

Pestisitlerin kullanımı çok eski tarihlere dayanmaktadır. M.Ö. 1500'lere ait bir papirüs üzerinde bit, pire ve eşek arılarına karşı insektisitlerin hazırlanışına dair kayıtlar bulunmuştur. Bilinen ilk pestisit Mezopotamya'da yaklaşık 4500 yıl önce antik Sümer'de kullanılan elemental kükürt tozudur. Onbeşinci yüzyılda arsenik, civa ve kurşun gibi toksik kimyasallar tarım ürünlerindeki zararlıların öldürülmesinde kullanılmışlardır. Onyedinci yüzyılda nikotin sülfat, insektisit olarak kullanılmak üzere tütünden ekstrakte edilmiştir. Ondokuzuncu yüzyılda iki doğal pestisit kullanılmaya başlanmıştır. Bunlardan biri krizantemden elde edilen pyrethrum (pire otu) ve diğeri de tropik bitki köklerinden elde edilen rotenon'dur (Miller, 2002). Pestisitler, ürünlerde arzu edilmeyen organizmaları yok etmede kullanılan sentetik organik bileşiklerdir. Bitki koruma amacıyla kullanılan her türlü ilaç ile preparatlar ve bunların imalinde kullanılan maddeler bu gruba girmektedir. Pestisitlerin yararlarının yanı sıra uzun süreli

kullanımları sonucunda ekosisteme ve insan sađlıđına zarar verdiđi saptanmıřtır, bundan dolayı kimyasalların tarım amaçlı kullanımları bazı kurallarla sınırlandırılmıřtır. Pestisit kalıntılarının önemi ilk kez 1948 yılında insan vücudunda organik klorlu pestisit kalıntılarının bulunmasıyla anlaşılmıřtır. Pestisitlerin bazıları toksikolojik açıdan zarar oluřturmazken, bazılarının kanserojen olduđu ve sinir sistemini etkilediđi saptanmıřtır. Pestisit kalıntılarının en önemli kaynađı gıdalardır. Bu nedenle, 1960 yılında FAO ve WHO tarafından “pestisit kalıntıları kodex komitesi” kurulmuřtur. Bu komitenin çalıřmaları sonucu konuyla ilgili tanımlamalar yapılmıř ve bilimsel arařtırma verilerine dayanarak gıdalarda bulunmasına izin verilen maksimum kalıntı deđerleri saptanmıřtır (Akman vd., 2004).

Uzun süreli ve bilinçsiz pestisit kullanımı, çevresel bozulmalara ve insan sađlıđına ciddi zararlar verebilmektedir. Bu nedenle, pestisitlerin kullanımı ve kalıntıları, dünya genelinde sıkı düzenlemelere tabi tutulmaktadır. FAO (Gıda ve Tarım Örgütü) ve WHO (Dünya Sađlık Örgütü), pestisitlerin güvenli kullanımını sađlamak için Codex Alimentarius (Gıda Kodu) aracılıđıyla Maksimum Kalıntı Seviyeleri (MRL) belirleyerek, dünya genelinde tarımda kullanılan kimyasalların güvenli seviyelerde kalmasını sađlamayı hedeflemektedir (FAO, 2022; WHO, 2023).

Sonuç olarak, pestisitlerin kullanımı yalnızca tarımsal verimliliđi sađlamakla kalmayıp, aynı zamanda çevresel sürdürülebilirliđi ve insan sađlığını koruma açısından büyük bir sorumluluk tařımaktadır. Bu nedenle, pestisit kullanımının yönetilmesi, bilimsel arařtırmalar ve etkili düzenlemelerle, güvenli ve sađlıklı gıda üretimi için elzemdir.

Pestisitlerin Çevreye Etkileri

Tarımsal alanlara uygulanan pestisitler; hava, su ve toprađa, oradan da bu ortamlarda yařayan diđer canlılara geçmekte ve dönüşüme uğramaktadır. Bir pestisitinin çevredeki hareketlerini, onun kimyasal yapısı, fiziksel özellikleri, formülasyon tipi, uygulama şekli, iklim ve tarımsal kořullar gibi faktörler etkilemektedir. Tarım ürünlerini zararlı, hastalık ve yabancı otlardan korumak ve kaliteli ürün elde etmek için bütün dünyada pestisitler kullanılmakta, bu kullanım sonucunda insan, hayvan ve çevre sađlığı bakımından bazı problemler ortaya çıkmaktadır (Yıldırım, 2008). Tarımsal alanlara uygulanan pestisitler; hava, su ve toprađa, oradan da bu ortamlarda yařayan diđer canlılara geçmekte ve dönüşüme uğramaktadır. Bir pestisitinin çevredeki hareketlerini, onun kimyasal yapısı, fiziksel özellikleri, formülasyon tipi, uygulama şekli, iklim ve tarımsal kořullar gibi faktörler etkilemektedir. Tarım ürünlerini zararlı, hastalık ve yabancı otlardan korumak ve kaliteli ürün elde etmek için bütün dünyada pestisitler kullanılmakta, bu kullanım sonucunda insan, hayvan ve çevre sađlığı bakımından bazı problemler ortaya çıkmaktadır (Yıldırım, 2008)

Pestisitlerin Toprađa Etkileri

Tarımsal üretimde bilinçsiz ve aşırı miktarda kullanılan kimyasal gübre ve pestisitler çevre üzerinde olumsuz etkiler yaratmaktadır. Genelde, bütün pestisitler kullanıldıklarında az veya çok toprađa karıřtırılır ve toprađı kirletirler. Pestisitler, toprakta bir taraftan yer altı suyuna dođru ilerlerken, diđer taraftan karbondioksit, metan ve su gibi daha az zehirli yada zehirsiz bileřenlere indirgenmektedirler. Pestisitinin yarılanma ömrü kısaltıldıkça, parçalanması hızlanır ve önemli bir kısmını yer altı suyuna ulařmadan daha yolda kaybeder. Bunun tersi olduđunda ise, pestisitinin önemli bir kısmının yer altı suyuna ulařması ve ulařtıđı yerdeki kaynađı kirletmesi muhtemeldir. Diđer taraftan, katyonik özelliklere sahip pestisitler toprakta kuvvetlice sorbe edilerek bu kimyasalların topraktaki hareketleri azaltılır. Dolayısıyla, katyon özelliđi tařıyan pestisitlerin yer altı suyunu kirletme riski nötr veya anyon özelliđi tařıyan pestisitlere göre daha düşüktür. Zirai mücadele kullanılan teknikler insan ve çevre sađlığı açısından özel bir öneme sahiptir. Genel olarak bitki hastalık ve zararlılarıyla, zamanında ve dođru mücadele yapılmadıđında ürün kaybı olduđu bilinmektedir. Bu kaybı önlemek için en fazla kullanılan

yöntemler arasında, kimyasal mücadele gelmektedir. Ülkemizde polikültür tarımın yapıldığı özellikle Akdeniz ve Ege Bölgeleri'nde yoğun düzeyde pestisit kullanılmaktadır (Karaca & Turgay, 2012).

Kalıntı Problemi ve Zehirlenmeler

Pestisitler öneriler doğrultusunda kullanılmadığı zaman kalıntıları ile insan sağlığı ve çevrede olumsuz etkilere yol açmaktadır. Bu kalıntılar, tarım ürünü dış pazarını ve iç tüketimi de olumsuz etkilemektedir. Ayrıca üretim, formülasyon hazırlama, taşıma, yükleme ve uygulama sırasında deri ve solunum yoluyla maruz kalma (akut zehirlenme) şeklinde mesleki zehirlenmelere sebep olabilirler. Genellikle organik fosforlar ve karbamatlılar bu tip zehirlenmeye neden olurlar. Bunlar vücutta kolinesteraz enzimini engelleyerek asetil kolin birikimine yol açarlar. Kaza ile meydana gelen zehirlenmelerde pestisitlerin yaprak ve topraktaki kalıntıları veya onların toksik dönüşüm ürünleriyle temas sonucu hastalıklar meydana gelebilmektedir. Aşırı dozlarda alınmadıkça organik klorlu pestisitlerin insanlara akut zehirlilikleri enderdir. Bu bileşikler daha çok kronik zehirlenmelere neden olmakta, sinir sistemini etkilemekte ve karaciğere zarar vermektedirler (Tiryaki ve ark., 2010).

Pestisitlerin Suya Etkileri

Su en büyük doğal kaynaklarımızdan biri ve yaşamın temelidir. İnsanoğlu içmek, pişirmek ve yıkanmak için temiz suya ihtiyaç duyar. Temiz su çiftçilerin ürünlerini sulamak ve çiftlik hayvanlarını beslemek için esastır. Yeraltı suları dünyanın temel taze su kaynağıdır. Toprak ve bitki uygulamalarından sonra toprak yüzeyinde kalan pestisitler, yağmur suları ile yüzey akışı şeklinde veya toprak içerisinde aşağıya doğru yıkanmak suretiyle taban suyu ve diğer su kaynaklarına ulaşabilirler. Eğim, bitki örtüsü, formülasyon, toprak tipi ve yağış miktarına bağlı olarak taşınan pestisitler yeraltı sularına geçebilirler. Pestisitler yeraltı suyuna ulaştıktan sonra da parçalanmaya devam ederler. Fakat daha az ışık, sıcaklık ve oksijen nedeniyle daha düşük oranda parçalanırlar. Yeraltı suyu kirlendiği zaman; kirli su akıntıları, nehirler ve göllerde de bulaşma görülebilir. Bulaşma kaynakları durdurulmuş olsa bile bir aküferin doğal işlemlerle kendi kendisini saflaştırması zaman gerektirir. Yeraltı suyu kirlendiğinde temizliği çok pahalı ve zordur. Yeraltı su kirliliğine karşı en iyi koruma kirliliğin önlenmesidir (Yücel, 2007).

Iğdır ilinde Pestisit Kullanımı

Türkiye kara kütlelerinin büyük yer kapladığı Kuzey Yarım Kürenin orta kuşağında, 36-42 derece kuzey paralelleri ile 26-45 derece doğu meridyenleri arasında yer almaktadır. Kuzey Yarım Kürede yer almak coğrafi konum açısından çok önemlidir. Bilindiği gibi, karaların büyük bir kısmı Kuzey Yarım Kürede yer almakta, bu yarım kürede kara parçaları birbirine daha yakın bulunmakta, yeraltı ve yerüstü kaynakları daha zengin bir potansiyele sahip olmaktadır (Değirmenci, 2019). Türkiye, özellikle Anadolu tarih boyunca çeşitli uygarlıklara ev sahipliği yapmış bir coğrafyadır. Bu uygarlıklarda tarımsal faaliyetler, bölgenin iklim çeşitliliği nedeniyle çok çeşitli alanlarda yoğun bir şekilde yapılmıştır. Bu uygarlıkların en önemlileri, Asurlar, Hititler, Selçuklular, Anadolu Selçukluları ve Osmanlılar olmuştur. (Yavuz, 2005).

Iğdır İli, Doğu Anadolu Bölgesinin Erzurum 39056' Kuzey Kars Bölümünde yer alır. 44048' doğu boylamı ve enlemi Iğdır İli toprakları üzerinden geçmektedir. Kuzey ve kuzeydoğu sınırını Aras Nehri ve bu nehrin yatağı boyunca geçen Ermenistan sınırını teşkil eder. Güneydoğusunda ve doğusunda Nahçıvan ve İran, güneyde Ağrı İli, batı ve kuzeybatısında Kars İli yer almaktadır.



Şekil 1. Iğdır İli Harita Görünümü


Iğdır ovası ve çevresi Türkiye ve Doğu Anadolu ölçüsünde kendine özgü iklim özellikleri ile yöresel klima alanı içine girmektedir. Iğdır, yazları sıcak ve kurak, kışları ılıman bir iklime sahiptir. Iğdır İli, çevresindeki yüksek alanlardan tamamen farklı bir iklime sahiptir. Bu farklılıkların sebebi, sıcaklığın yüksek ve yağışların az oluşudur. Yarı kurak bir iklime sahip olan Iğdır ili, Doğu Anadolu ölçüsünde kendine özgü iklim koşullarıyla bir yöresel klima alanı oluşturmaktadır. Iğdır Ovası Doğu Anadolu Bölgesi'nin mikro klima özelliği gösteren, en alçak ve yüzölçümü en geniş olan ovalarından biridir. Doğu Anadolu gibi yüksek platolar ve dağlık bölgelerin geniş yer kapladığı bir bölgede bulunan ve sahip olduğu bağlı yüksekliği ile havza olarak belirlenen Iğdır ili çevresine göre gerek iklim gerekse toprak ve bitki örtüsü gibi doğal çevre özellikleri bakımından oldukça değişik özellikler gösterir.

Iğdır ilinin tarıma elverişli alanı oldukça geniştir. Toplamda 118.525 hektarlık bir tarım alanına sahip olan il, şeker pancarı, pamuk, karpuz, domates ve birçok meyve-sebze yetiştirmek için uygun bir ortama sahiptir. İl genelinde 97.041 hektarlık sulanabilir tarım arazisi, 2.233 hektar ormanlık alan ve 146.571 hektar çayır-mera alanı bulunmaktadır. Bölgede şeker pancarı üretimi, Erzurum Şeker Fabrikası'nın 1956 yılında kurulmasıyla başlamıştır ve şeker pancarı üretimi halen önemli bir tarım faaliyeti olarak devam etmektedir. Meyve bahçeleri ise toplamda 1.525 hektar alanı kaplayarak büyük bir yer kaplamaktadır. Bu alanların %74'ü kayısı, %16,1'i elma, %6,2'si şeftali, %1,7'si armut, %0,5'i ise diğer meyveler ve bağ tarımı alanlarından oluşmaktadır. Yöre halkının temel ihtiyaçlarını karşılamak için yetiştirilen meyveler, bölge ekonomisinde önemli bir yer tutmaktadır. Iğdır ilinin tarımsal alanları oldukça geniş bir yelpazede yer almaktadır. Bölgede sulanabilir tarım arazisi, orman alanları, çayır-mera ve tarım alanları bulunmaktadır. Tarıma elverişli alanlarda, özellikle şeker pancarı, pamuk, karpuz, domates gibi sebzeler ve çeşitli meyveler yetiştirilmektedir. Bu ürünlerin bir kısmı sadece Iğdır'da tüketilirken, bir kısmı da çevre illere gönderilmektedir. Ayrıca, sebze ve meyveler dışında da soğan, patlıcan, biber, lahana, turp, fasulye gibi ürünlerin yetiştiriciliği yapılmaktadır (Aksoy ve ark., 2012).

Iğdır ilinin kendine özgü iklim özellikleri olması ve klima alanı olmasından ötürü tarımı yapılan ürün grupları çeşitlenmiştir. Buna bağlı olarak Iğdır ilinde yoğun bir tarım yapılmakta ve bazı sorunlarla karşılaşmaktadır. Iğdır ilinde tarım alanlarında başlıca yaşanan sorunlar şunlardır; su kaynaklarının yetersizliği ve sulama sorunları, toprak erozyonu ve bozulması, pestisit ve kimyasal gübre kullanımı, hava koşullarının belirsizliği (iklim değişikliği), pazar erişimi ve ekonomik sorunlar, tarım işçiliği sorunları, tarım politikalarının yetersizliği gibi sorunlar yaşanmaktadır. Pestisit ve kimyasal gübre kullanımı tarım üretiminin ekstantif şartlardan entansif şartlara taşınmasında önemli rol oynamaktadır. Entansif tarım, birim alandaki verimi artırmak için kullanılan girdi miktarının artırılmasını ve modern tekniklerin uygulanmasını içermektedir. Birim alandaki verimin artırılmasında doğrudan verimi etkileyen hastalık ve zararlılarla mücadele edilmesi önemli bir yer tutmaktadır. Ziraî mücadele, zararlı etmenlerin ülkeye veya bir bölgeye girmesini önleyici karantina tedbirlerinden, canlı unsurların kullanıldığı biyolojik mücadeleye ve yoğun girdinin kullanıldığı kimyasal mücadeleye kadar değişik yöntemleri kullanır. Bu yöntemlerden kimyasal mücadele en fazla kullanılanıdır. Çünkü yüksek etkililiğe sahiptir, hızlı sonuç verir, bilinçli ve kontrollü kullanıldığında ekonomiktir ve ürünü toksin salgılayan organizmalardan da koruyabilir (Delen ve ark., 2005). Iğdır ilinde pestisit kullanımı çevre illere göre fazladır.

Ziraî mücadelede kimyevî yöntemlerin yoğun kullanımı insan sağlığı ve çevreye olumsuz etkiler gibi birçok meseleyi de beraberinde getirmektedir (Peker, 2012). Türkiye’de tarımda girdi kullanımındaki dengesizlik ve üreticinin bilinçsizliği iki önemli mesele olarak görülmektedir (Olhan, 1997). Yoğun ve bilinçsiz bir şekilde kullanılmaları sonucunda gıdalarda, toprak, su ve havada kimyasalların kendisi ya da parçalanma ürünleri kalabilmektedir (Oğuz, 1996). Dünya genelinde tarım sisteminin ayrılmaz bir parçası olarak kimyasal kullanımının, tarım ürünlerinde kalıntı riski ve çevreye olumsuz etki yapması dikkatle üzerinde durulması gereken bir konudur (Peker, 2012). Çünkü sürdürülebilir bir tarım faaliyeti ya da gıda üretimi, zararlılarla mücadelenin çevreye ve insana en az seviyede zarar verecek şekilde yürütülmesini gerektirmektedir.

Iğdır ilinde 124693,51 hektar tarım alanı vardır ve 164,693 kg-lt pestisit kullanılmıştır. Bu pestisitler gruplarına göre incelendiğinde 2023 yılında 107,980 kg-lt insektisit, 29,293 kg-lt fungusit, 21,250 kg-lt herbisit, 5,970 kg-lt akarisit ve diğer gruplar olmak üzere kullanıldığı bilinmektedir (Şekil 2).




Şekil 2. Iğdır ilinde Pestisit gruplarına göre kullanım miktarı (kg-lt)

Iğdır ilinde Ertürk ve ark. (2012) tarafından yapılan bir araştırmaya göre, Iğdır ilinde ve ilçelerinde bitki üretimi yapan tarım işletmesi sahiplerinin bitki korumayla ilgili uygulamalarını gerçekleştirirken, tabiata ve insan sağlığına verebilecekleri zarar hakkında bilgili olup olmadıkları araştırılmıştır. Çalışmanın verileri Gıda, Tarım ve Hayvancılık Bakanlığı Iğdır İl


Müdürlüğünün Çiftçi Kayıt Sisteminde kayıtlı toplam 104 tarım işletmesinin sahipleriyle yüz yüze yapılan anket çalışmalarıyla elde edilmiştir. Tarım ilâçlarının çevreye zararı konusunda çiftçilerin yarısına yakın kısmının zarar yoktur demesi, dörtte birinin de kalıntılar konusunda zararsız olacağını ifade etmesi çiftçinin bilinç seviyesinde yetersizliği ifade etmektedir. Bilhassa ilâçlama âletlerinin temizleme sularının atılmasında yeterli dikkatin gösterilmemesi, ilâç ambalajları ile kalan ilâçların muhafaza ve imhasındaki cevaplar çiftçilerin hem çevre hem de ziraî mücadele konusunda ki bilinçsizliğini göstermektedir (Ertürk ve ark., 2012).

İğdır ve diğer serhat illerinin tarımsal üretimi kıyaslaması

İğdır iline komşu olan Kars ilinde 598357,92 hektar tarım alanı, Ağrı ilinde 466650,6 hektar tarım alanı bulunmaktadır (Anonim, 2025). Kars ve Ağrı illerinde pestisit kullanımı İğdır iline nazaran çok düşüktür (Şekil 3, Şekil 4).



Şekil 3. Kars ili pestisit kullanımı (kg-lt)




Şekil 4. Ağrı ili pestisit kullanımı (kg-lt)

İğdır'ın tarımsal üretimi, coğrafi konumu ve iklimi sayesinde diğer serhat illerine kıyasla daha verimli ve çeşitlidir. Özellikle meyve üretimi (özellikle elma, kayısı, kavun ve şeftali) konusunda İğdır, diğer illere göre daha fazla çeşitliliğe ve üretim kapasitesine sahiptir. Seracılık faaliyetleri de İğdır'da yaygındır, bu da tarımsal çeşitliliği artıran önemli bir faktördür. İğdır'ın verimli toprakları ve sıcak iklimi, daha kısa sürede hızlı ürün almayı sağlar.

Kars ve Ardahan gibi iller, soğuk iklim nedeniyle genellikle hayvancılığa daha fazla dayalı tarımsal faaliyetler yürütürken, bu illerde buğday ve arpa gibi tahıllar daha yaygın, sebze ve meyve üretimi ise genellikle sınırlıdır. Erzurum, Van ve Ağrı illeri de benzer şekilde daha az çeşitliliğe sahip olup, soğuk iklim koşulları nedeniyle tarımsal verimlilikleri Iğdır'a göre daha düşüktür. Sonuç olarak, Iğdır'ın iklimsel avantajları ve mikroklima özellikleri, bu ili serhat illeri arasında tarımsal çeşitlilik ve verimlilik açısından öne çıkarmaktadır. Bu avantajlar, özellikle meyve ve sebze üretiminde önemli bir fark yaratmaktadır.

Iğdır'da tarım ve zararlıların genel durumu

Iğdır ilinde tarımsal faaliyetler oldukça gelişmiştir. Gerek mikroklima özelliği gerekse coğrafi yapısı, bu bölgedeki tarımsal çeşitliliği desteklemiş ve farklı iklim koşullarına uygun birçok ürünün yetiştirilmesine olanak sağlamıştır. Iğdır, diğer serhat illerine oranla daha geniş bir tarımsal ürün yelpazesi ile dikkat çekmektedir. Bölgenin sahip olduğu verimli alüvyal topraklar, sulama olanakları ve uygun iklim koşulları, tarımsal üretimin çeşitlenmesine katkı sağlamaktadır. Iğdır'da yetiştirilen başlıca tarım ürünleri arasında elma, kayısı, kavun, karpuz, domates, şeftali ve nektarin gibi ürünler yer almaktadır. Elma ve kayısı, Iğdır'ın meyvecilik sektöründe en önemli ve yaygın üretilen ürünlerindendir. Özellikle elma, bölgeye özgü farklı çeşitleriyle hem iç piyasada hem de dışarıya satılmak üzere üretilmektedir. Kavun ve karpuz da Iğdır'ın yaz aylarında en çok yetiştirilen sebze meyvelerindendir ve sıcak iklim koşullarına uygun bu ürünler hem yerel pazarlarda hem de diğer bölgelere satılmaktadır. Domates, Iğdır'da geniş alanlarda yetiştirilen önemli bir sebze olup, hem taze tüketim hem de sanayiye girdi olarak kullanılmaktadır. Ayrıca, şeftali ve nektarin gibi meyveler de Iğdır'ın tarımsal üretiminde yer almakta ve bölgenin meyve üretimi çeşitliliğini artırmaktadır.



Şekil 5. Iğdır İlinde Meyve ve Sebze Üretim Miktarları

Iğdır'da kültür bitkilerindeki başlıca önemli zararlılar şunlardır:

- **Elma İç Kurdu (*Cydia pomonella* L.):** Elma yetiştiriciliği, Iğdır'daki tarımsal üretimde önemli bir yer tutmaktadır. Elma iç kurdu, elma ağaçlarının meyvelerine zarar vererek meyve kalitesini bozan bir zararlıdır.
- **Akdeniz Meyve Sineği (*Ceratitis capitata* (Wiedemann)):** Iğdır ilinde geçici şeftalimeyvelerine zarar verir. Dişi sinekler meyvelerin içine yumurtalarını bırakır ve larvalar meyveyi bozar.

- **Kırmızı Örümcek (*Tetranychus urticae* Koch):** Özellikle elma ve şeftali ağaçlarında büyük zararlara yol açar, meyve gelişimini engeller.
- **Yonca hortumlu böceği (*Hypera postica* (Gyllenhal)):** Yoncada görülen zararlıdır.
- **Yaprak Bitleri (Aphididae):** Çeşitli sebze ve meyve türlerinde yaygın olup, bitkilerin zayıflamasına ve virüslerin yayılmasına neden olur.
- **San jose kabuklu biti (*Quadraspidiotus perniciosus* Comst.):** Iğdır ilinde sert çekirdekli meyve ağaçlarında zarar oluşturur.
- **Domates güvesi (*Tuta absoluta*. Meyrick):** Domates Iğdır için önemli bir tarım ürünüdür. Domates güvesinin zararını önlemek için kimyasal mücadele tercih edilmektedir.
- Ek olarak tarımsal alanlarda birçok fungal ve bakteriyel hastalıklarla birlikte yabancı ot içinde kimyasal mücadele yoğun olarak tercih edilmektedir.

Iğdır'da bu zararlılarla yoğun bir kimyasal mücadele yapılmaktadır. Kimyasal mücadele kısa vadede hızlı sonuçlar vererek ürünlerin verim ve kalitesini artırmakta olsa da, aşırı pestisit kullanımı çevre üzerinde kalıcı tahribatlar bırakabilmektedir. Özellikle Iğdır ilinde hava kirliliği önemli bir sorundur. Pestisitlerin hava yoluyla yayılması, hava kalitesinin daha da bozulmasına neden olmaktadır. Bu kirlilik, solunum yolu hastalıklarını artırabilir ve ekosistemler üzerinde olumsuz etkiler yaratabilir.

Bir diğer önemli çevresel sorun ise su kirliliğidir. Pestisitler, sulama suyu veya yağmur suları aracılığıyla yer altı ve yüzey su kaynaklarına karışarak suyun kimyasal maddelerle kirlenmesine yol açmaktadır. Bu durum, hem içme suyu hem de tarımsal sulama suyu kalitesini tehlikeye atmaktadır. Su kaynaklarında biriken pestisit kalıntıları, insan sağlığı için tehdit oluşturmakla kalmaz, aynı zamanda sucul yaşam üzerinde yıkıcı etkiler yaratır. Suda bulunan kimyasal maddeler, sucul organizmaların yaşamını engelleyebilir ve biyolojik çeşitliliği olumsuz etkileyebilir.

SONUÇ

Alternatif mücadele yöntemleri, pestisit kullanımının çevre üzerindeki olumsuz etkilerini azaltmada büyük önem taşımaktadır. Biyolojik mücadele, entegre zararlı yönetimi (IPM) ve doğal düşmanların kullanımı gibi çevre dostu yöntemlerle zararlılarla mücadele edilmesi, ekosistemlerin korunmasına yardımcı olur ve tarımsal verimliliği sürdürülebilir kılar. Bu yöntemler, pestisitlerin yerine geçerek çevresel sürdürülebilirliği sağlar ve insan sağlığı üzerindeki riskleri en aza indirir.

Sonuç olarak, Iğdır'daki pestisit kullanımının çevreye olan olumsuz etkileri, hava kirliliği, toprak erozyonu ve su kirliliği gibi önemli çevresel sorunlara yol açmaktadır. Bu durum, yalnızca ekosistemler ve biyolojik çeşitlilik açısından değil, insan sağlığı açısından da büyük bir tehdit oluşturmaktadır. Bu nedenle, zararlılarla mücadelede alternatif ve çevre dostu yöntemlerin benimsenmesi büyük bir önem taşımaktadır.

KAYNAKÇA

- Akman, Y., Ketenoğlu, O., Kurt, L., Düzenli, S., Güney, K., & Kurt, F. (2004). *Çevre Kirliliği (Çevre Biyolojisi)*. Palme Yayıncılık.
- Aksoy, Y., Savaş, D., & Dursun, S. (2012). Iğdır İlinin Mekânsal Analiz Raporu. Iğdır Çevre ve Şehircilik İl Müdürlüğü. https://webdosya.csb.gov.tr/db/kentges/editordosya/kiu_igdir_mekansal_analiz.pdf (Erişim Tarihi: 23.05.2023).

- Anonim. (2024). World Population Growth Rate 1950-2024. <https://www.macro Trends.net/global-metrics/countries/WLD/world/population-growth-rate> (Erişim Tarihi: 21.02.2025).
- Anonim. (2025). <https://corinecbs.tarimorman.gov.tr/> (Erişim Tarihi: 19.02.2025).
- Değirmenci, Y. (2019). Öğretmen Adaylarının Bakış Açısıyla Türkiye'nin Coğrafi Konumunun İncelenmesi. *Ağrı İbrahim Çeçen Üniversitesi Sosyal Bilimler Enstitüsü Dergisi*, 5(2), 91-110.
- Delen, N., Durmuşoğlu, E., Güncan, A., Güngör, N., Turgut, C., & Burçak, A. (2005). Türkiye'de Pestisit Kullanımı, Kalıntı ve Organizmalarda Duyarlılık Azalışı Sorunları. *Türkiye Ziraat Mühendisliği 6ncı Teknik Kongresi*, 21 sayfa.
- Ertürk, Y. E., Bulak, Y., & Uludağ, A. (2012). Iğdır ili tarım işletmelerinin zirai mücadele uygulamalarında çevreye duyarlılıkları.
- FAO. (2022). *Codex Alimentarius - Pesticide Residues*. Food and Agriculture Organization. FAO.org
- FAO. (2021). *Synthesis Report*.
- <https://openknowledge.fao.org/server/api/core/bitstreams/ecb51a59-ac4d-407a-80de-c7d6c3e15fcc/content> (Erişim Tarihi: 23.02.2025).
- Karaca, A., & Turgay, O. C. (2012). Toprak Kirliliği. *Toprak Bilimi ve Bitki Besleme Dergisi*, 1(1), 13-19.
- Miller, G.T. (2002). *Living in the Environment* (12th ed.). Belmont: Wadsworth/Thomson Learning.
- Olhan, E. (1997). *Türkiye'de Bitkisel Üretimde Girdi Kullanımının Yarattığı Çevre Sorunları ve Organik Tarım Uygulaması-Manisa Örneği* (Doktora tezi). Ankara Üniversitesi Fen Bilimleri Enstitüsü.
- Oğuz, C. (1996). Konya İli Çumra İlçesinde Domates Yetiştiriciliği Yapan Tarım İşletmelerinde Verimlilik Analizi Üzerine Bir Çalışma. *Türkiye 2. Tarım Kongresi*, Çukurova Üniversitesi Basımevi, Adana.
- Peker, A.E. (2012). Konya İli Domates Üretiminde Tarımsal İlaç Kullanımına Yönelik Çevresel Duyarlılık Analizi. *Iğdır Üniversitesi Fen Bilimleri Enstitüsü Dergisi*, 2(1), 47-54.
- Tiryaki, O., Canhilal, R., & Horuz, S. (2010). Tarım İlaçları Kullanımı ve Riskleri. *Erciyes Üniversitesi Fen Bilimleri Enstitüsü Fen Bilimleri Dergisi*, 26(2), 154-169.
- WHO. (2023). *Pesticides and Health*. World Health Organization. WHO.int
- Yavuz, F. (2005). *Türkiye'de Tarım*. Tarım ve Köyişleri Bakanlığı Yayınları, 1, 252.
- Yıldırım, E. (2008). Tarımsal Zararlılarla Mücadele Yöntemleri ve Kullanılan İlaçlar. *Atatürk Üniversitesi Ziraat Fakültesi Yayınları*, No:219, Erzurum.
- Yücel, Ü. (2007). Pestisitlerin İnsan ve Çevre Üzerine Etkileri. *Nükleer Araştırma ve Eğitim Merkezi, Nükleer Kimya Bölümü*. Ankara.
- TÜİK. (2024). Hayvansal Üretim İstatistikleri. <https://data.tuik.gov.tr/Bulten/Index?p=Hayvansal-Uretim-Istatistikleri-2024-53935> (Erişim Tarihi: 08.03.2025).
- <https://www.tarimorman.gov.tr/GKGM/Menu/115/Resmi-Tarimsal-Ilac-Istatistikleri> (Erişim Tarihi: 13.02.2025).

CLINICAL USE OF HORMONES

HORMONLARIN KLİNİK KULLANIMI

Ağalarova Səma

Nakhchivan State University Faculty of Natural Sciences and Agriculture department of
veterinary medicine

ORCID ID: <https://orcid.org/0009-0009-9756-1691>

ABSTRACT

Introduction and Purpose

Hormones are chemical substances that regulate various

physiological processes in the animal organism. These substances manage different functions of the organism, processes such as reproduction, growth, metabolism and stress responses. In the veterinary field, hormones are widely used both to treat reproductive problems and to support the general health of animals. Proper use and regulation of hormones helps maintain the health of animals and increase their productivity. This article examines the role of hormones in the animal organism, their uses in veterinary medicine, and the risks of hormone use

Hormones are chemical compounds that regulate many physiological processes in the animal organism and ensure the continuity of vital functions. These biological substances help the organism adapt to environmental changes by controlling important processes such as growth, metabolism, reproduction and stress. The functions of hormones not only regulate natural biological processes, but also play a major role in the treatment of various diseases, especially in solving reproductive problems. The correct use of hormones in veterinary practice is considered an effective method to improve the health of animals and increase their productivity.

Hormone therapies are widely used to regulate the reproductive health of animals, accelerate their developmental processes and treat certain diseases. However, there are also some risks associated with hormone use. Improper use or imbalances can threaten animal health. Therefore, the correct administration of hormones and the expertise of veterinarians in this field are of great importance.

This article will discuss the role of hormones in the animal organism, their uses in veterinary medicine and the potential risks of hormone use. In-depth information about how hormones work, in which situations they are used and their possible side effects will be given.

Hormones and their role in the animal organism

Key Words: Hormones; Hormones in animals ; Veterinary Medicine ; Hormone therapy ; Animal health

Özet

Giriş ve Amaç

Hormonlar, hayvan organizmasında çeşitli fizyolojik süreçleri düzenleyen kimyasal maddelerdir. Bu maddeler, organizmanın farklı fonksiyonlarını, üreme, büyüme, metabolizma

ve stres tepkileri gibi süreçleri yönetir. Veterinerlik alanında hormonlar, hem üreme problemlerinin tedavisinde hem de hayvanların genel sağlığını desteklemek amacıyla yaygın olarak kullanılmaktadır. Hormonların doğru kullanımı ve düzenlenmesi, hayvanların sağlığını korumaya ve verimliliğini artırmaya yardımcı olur. Bu makale, hormonların hayvan organizmasındaki rolünü, veterinerlikte kullanım alanlarını ve hormon kullanımının risklerini incelemektedir.

Hormonlar, hayvan organizmasındaki birçok fizyolojik süreci düzenleyen ve hayati işlevlerin devamlılığını sağlayan kimyasal bileşiklerdir. Bu biyolojik maddeler, büyüme, metabolizma, üreme ve stres gibi önemli süreçleri kontrol ederek organizmanın çevresel değişikliklere uyum sağlamasına yardımcı olur. Hormonların işlevleri sadece doğal biyolojik süreçleri düzenlemekle

kalmaz, aynı zamanda çeşitli hastalıkların tedavisinde, özellikle üreme problemlerinin çözülmesinde de büyük rol oynar. Veterinerlik pratiğinde hormonların doğru bir şekilde kullanılması, hayvanların sağlığını iyileştirmekte ve onların verimliliğini artırmakta etkili bir yöntem olarak kabul edilmektedir.

Hayvanların üreme sağlığını düzenlemek, gelişimsel süreçlerini hızlandırmak ve bazı hastalıkları tedavi etmek amacıyla hormon tedavileri yaygın olarak kullanılmaktadır. Bununla birlikte, hormon kullanımının bazı riskleri de bulunmaktadır. Yanlış kullanım veya dengesizlikler, hayvan sağlığını tehdit edebilir. Bu nedenle, hormonların doğru şekilde uygulanması ve veteriner hekimlerin bu alandaki uzmanlığı büyük önem taşır.

Bu makale, hormonların hayvan organizmasındaki rolünü, veterinerlikteki kullanım alanlarını ve hormon kullanımının potansiyel risklerini ele alacaktır. Hormonların nasıl çalıştığı, hangi durumlarda kullanıldığı ve olası yan etkileri hakkında derinlemesine bilgi verilecektir.

Hormonlar ve Hayvan Organizmasındaki Roller

Hormonlar, organizmamızda çeşitli fonksiyonları düzenleyen biyolojik olarak aktif maddelerdir. Genellikle, kan yoluyla taşınan ve farklı organ ve dokularda etki gösteren maddelerdir. Hormonlar, hayvan organizmasında çok önemli bir rol oynar ve bunların doğru dengesi sağlık için büyük önem taşır.

Hormonlar, esas olarak endokrin bezler tarafından üretilir. Bu bezler arasında hipofiz bezi, tiroid bezi, adrenal bezler, yumurtalıklar ve testisler yer alır. Bu hormonlar, organizmanın büyümesi, gelişmesi, sindirim ve sinir sistemi gibi temel işlevleri üzerinde ciddi etkiler yapar, ayrıca üreme fonksiyonlarını da düzenler.

Üreme Hormonları

Üreme hormonları, hayvanların cinsel gelişimini ve çoğalmalarını düzenleyen önemli maddelerdir. Bunlar arasında şunlar yer alır:

- Estrojen: Dişi hayvanlarda ovülasyonu ve üreme süreçlerini uyarır. Estrojen seviyeleri, dişi hayvanların cinsel döngüsünü ve üreme fonksiyonlarını düzenler.
- Progesteron: Gebelik sürecini koruyan ve yavrunun gelişimini destekleyen bir hormondur. Gebelik başladığında progesteron seviyesi artar ve doğum başlayana kadar sabit kalır.
- Testosteron: Erkek hayvanlarda cinsel özelliklerin gelişmesini ve sperm üretimini uyarır. Ayrıca erkek hayvanların cinsel davranışlarını da düzenler.
- Oksitosin: Doğum sırasında kasların kasılmasını teşvik eden bir hormondur. Ayrıca süt üretimini, doğumun uyarılmasını ve uterusu boşaltmak için kullanılır.

Büyüme Hormonları

Büyüme hormonları, hayvanların gelişimini ve büyümesini düzenleyen önemli maddelerdir. Bunlar arasında Büyüme Hormonu (GH) ve İnsülin Benzeri Büyüme Faktörü (IGF) bulunur.

- Büyüme hormonu (GH), hayvanların iskelet, kas ve diğer dokularının gelişmesini teşvik eder.
- IGF, hücre bölünmesini uyarır ve hayvanların büyümesini destekler.

Stres ve Basınç Hormonları: Hayvanların yaşamında stres anlarında basınç ve stres hormonları önemli rol oynar. Bu hormonlar, organizmanın stres durumlarına uyum sağlamasını sağlar. En önemli stres hormonları adrenalin ve *kortizol*dur.

- Adrenalin: Tehdit ve stres durumlarında organizmaya enerji verir ve hayvanı mücadele veya kaçma durumuna getirir.
- Kortizol: Uzun süreli stres reaksiyonlarını düzenleyen hormon olup, kan şekeri...

2. Süt üretiminin artırılması: Mamalık sırasında süt üretimini artırmak için oksitosin ve diğer hormonlar kullanılabilir.

3. Hormonal yetiştirme: Hayvanların büyümesini ve gelişmesini hızlandırmak için büyüme hormonları kullanılabilir.

Hormon Kullanımının Riskleri ve Düzenlenmesi

Hormon tedavileri, doğru düzenlenmediği takdirde bazı risklere yol açabilir. Bunlar arasında hormon dengesizlikleri, üreme sorunları ve bazı endokrin hastalıklar yer alır. Hormonların doğru kullanımı oldukça önemlidir ve veteriner hekimlerin uzmanlık bilgisine dayanmalıdır. Aynı zamanda, hormon kullanımı ile ilgili etik sorunlar da gündeme gelebilir. Hormon tedavileri yalnızca veterinerlik alanında etik kurallara uygun şekilde uygulanmalıdır.

Sonuç bu makalede, hormonların hayvan organizmasındaki rolü, veterinerlikteki kullanım alanları ve hormon kullanımının potansiyel riskleri detaylı bir şekilde ele alınmıştır. Hormonlar, hayvanların büyüme, üreme ve metabolik süreçlerini düzenleyen önemli biyolojik bileşiklerdir. Veterinerlik alanında doğru bir şekilde kullanıldığında, hormonlar hem üreme problemlerinin tedavisinde hem de genel sağlık sorunlarının çözülmesinde etkili bir araçtır. Bununla birlikte, hormonların yanlış kullanımı veya dengesizlikleri, hayvan sağlığını tehdit edebilir.

Hayvanların sağlığını korumak ve verimliliğini artırmak amacıyla hormon tedavilerinin dikkatli bir şekilde uygulanması, veteriner hekimlerin uzmanlığı ve deneyimi ile mümkün olmaktadır. Bu nedenle, hormon tedavisinin her aşamasında veterinerlerin bilimsel bilgiye dayalı kararlar alması büyük önem taşımaktadır. Ayrıca, hormon kullanımının hayvan sağlığı üzerindeki olası yan etkilerini minimize etmek için düzenli izleme ve kontrol süreçleri de ihmal edilmemelidir.

Anahtar Kelimeler: Hormonlar; Hayvanlarda hormonlar ; Veterinerlik Hormon tedavisi ; Hayvan sağlığı

IMPACT OF CLIMATE CHANGE ON INTEGRATED PEST MANAGEMENT (IPM) IN TURKEY

İKLİM DEĞİŞİKLİĞİNİN TÜRKİYE'DE ENTEGRE ZARARLI YÖNETİMİNE (IPM) ETKİSİ

Öğr. Gör. Nagihan KILIÇ

Iğdır Üniversitesi, Teknik Bilimler Meslek Yüksekokulu Bitkisel ve Hayvansal Üretim
Bölümü Seracılık Programı

ORCID ID: <https://orcid.org/0000-0002-7746-8125>

Prof. Dr. Celalettin GÖZÜAÇIK

Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü

ORCID ID: <https://orcid.org/0000-0002-6543-7663>

ABSTRACT

The rapid growth of the global population and technological advancements have significantly impacted agriculture and food production. However, climate change brings various challenges that threaten these advancements. While climate change creates numerous effects on agriculture and ecosystems, it also directly impacts integrated pest management (IPM) methods. Understanding the effects of climate change on pest control is crucial for sustainable agricultural practices.

Climate change factors, such as rising temperatures, increased atmospheric CO₂ levels, irregular rainfall patterns, unpredictable weather changes, and shifting climate patterns, can influence pest populations. These changes may create more favorable living conditions for pests and increase their spread. For example, rising temperatures can drive certain pests into regions where they were previously not found. Additionally, climate events like increased rainfall or drought can reduce the effectiveness of natural enemies against pests.

Integrated Pest Management (IPM) practices aim to combat pests in sustainable agriculture through a combination of chemical, biological, cultural, and mechanical methods. However, climate change can affect the effectiveness of these integrated methods. Climatic changes may alter pest behaviors, reproductive cycles, and population densities, making it more challenging to determine the timing and choice of species for IPM interventions. Furthermore, climate stress factors may also negatively affect the effectiveness of natural enemies against pests.

In conclusion, climate change requires greater flexibility and adaptation in the application of integrated pest control methods. In this context, it is crucial to monitor pest populations and continuously update integrated control strategies, taking into account the impacts of climate change.

Keywords: Climate change, Integrated pest management, Agricultural pests

ÖZET

Dünya nüfusunun hızla artması ve teknolojik ilerlemeler, tarım ve gıda üretimini önemli ölçüde etkilemiştir. Ancak, iklim değişikliği bu ilerlemeleri tehdit eden çeşitli sorunları da beraberinde getirmektedir. İklim değişikliği, tarım ve ekosistemlerde birçok etki yaratırken, zararlılarla

entegre mücadele yöntemlerini de doğrudan etkilemektedir. İklim değişikliğinin zararlılarla mücadele üzerindeki etkilerini anlamak, sürdürülebilir tarım uygulamaları açısından önemlidir.

İklim değişikliği, sıcaklık artışı, atmosferik CO₂ seviyelerinin yükselmesi, düzensiz yağış rejimleri, düzensiz hava değişimleri ve iklimin değişen kalıpları gibi faktörler, zararlıların popülasyonlarını etkileyebilir. Bu değişiklikler, zararlılar için daha uygun yaşam koşulları yaratırken, aynı zamanda zararlıların yayılma hızını da artırabilir. Örneğin, sıcaklık artışı bazı zararlıları daha önce görülmedikleri bölgelere yönlendirebilmektedir. Ayrıca, yağışların artması veya kuraklık gibi iklim olayları, zararlılara karşı doğal düşmanların etkinliğini azaltabilir.

Entegre zararlı yönetimi (IPM) uygulamaları, kimyasal, biyolojik, kültürel ve mekanik yöntemlerin birleştirilmesiyle sürdürülebilir tarımda zararlılarla mücadeleyi hedefler. Ancak, iklim değişikliği bu entegre yöntemlerin etkinliğini etkileyebilir. İklimsel değişiklikler, zararlıların davranışlarını, üreme döngülerini ve popülasyon yoğunluklarını değiştirebilir, bu da IPM uygulamalarının zamanlaması ve tür seçimini zorlaştırabilir. Ayrıca, iklimsel stres faktörleri, zararlılara karşı doğal düşmanların etkinliğini de olumsuz etkileyebilir.

Sonuç olarak, iklim değişikliği, zararlılarla entegre mücadele yöntemlerinin uygulanmasında daha fazla esneklik ve adaptasyon gerektirir. Bu bağlamda, iklim değişikliğinin etkilerini göz önünde bulundurarak, zararlı popülasyonlarını izlemek ve entegre mücadele stratejilerini sürekli olarak güncellemek büyük önem taşımaktadır.

Anahtar kelimeler: İklim değişimi, Entegre mücadele, Tarımsal zararlı

GİRİŞ

İklim değişikliği, dünya genelinde olduğu gibi, Türkiye’de de ekolojik, ekonomik ve sosyal boyutlarıyla önemli bir sorun haline gelmiştir (Soltekin ve ark., 2021). Küresel sıcaklık artışları, yağış rejimlerindeki değişiklikler, kuraklık, sel, orman yangınları ve tarımsal üretimde azalma gibi olumsuz etkiler, Türkiye’nin doğal kaynaklarını, ekonomik istikrarını ve toplum sağlığını tehdit etmektedir (Şen, 2022). Bilimsel araştırmalar, özellikle son yıllarda iklim değişikliğinin etkilerinin giderek arttığını ve tarımsal üretimden biyoçeşitliliğe kadar geniş bir yelpazede ciddi değişimlere neden olduğunu göstermektedir (Şen ve ark., 2013)

Türkiye’nin iklim değişikliğine karşı duyarlılığı, ülkenin coğrafi konumu, tarıma dayalı ekonomisi ve su kaynaklarının sınırlı olması nedeniyle oldukça yüksektir. Özellikle Akdeniz havzasında yer alan Türkiye, sıcaklık artışları, yağış düzensizlikleri ve su kıtlığı gibi problemlerin etkilerini daha fazla hissetmektedir. Tarımsal üretimde verim kayıpları, artan zararlı popülasyonları, su kaynaklarının azalması ve hava olaylarındaki aşırı değişimler, ekosistemler ve insan sağlığı üzerinde doğrudan etkiler yaratmaktadır (Bayraç & Doğan, 2016).

İklim değişikliği, yalnızca çevresel bir sorun olmanın ötesinde, ekonomik kalkınma, enerji güvenliği ve ulusal güvenlik gibi birçok alanı da doğrudan etkilemektedir. Türkiye’nin ekonomik yapısının büyük bir kısmı tarım, sanayi ve turizm sektörlerine dayanmaktadır. Bu sektörlerde yaşanan iklim kaynaklı değişimler, ekonomik büyümeyi ve istihdamı olumsuz yönde etkileyebilir. Ayrıca, küresel gıda güvenliği açısından Türkiye’nin tarımsal üretimindeki değişimler bölgesel ve uluslararası piyasaları da etkileyebilir (Şen ve ark., 2013).

Türkiye, iklim değişikliğinin olumsuz etkilerini azaltmak ve bu değişime uyum sağlamak amacıyla ulusal ve uluslararası düzeyde çeşitli politikalar geliştirmektedir. “Paris İklim Anlaşması” kapsamında belirlenen karbon nötr hedefi, Türkiye’nin çevresel sürdürülebilirliği sağlama konusundaki kararlılığını ortaya koymaktadır. Ayrıca, yenilenebilir enerji kullanımının teşvik edilmesi, su kaynaklarının daha verimli yönetilmesi, entegre zararlı yönetimi uygulamalarının geliştirilmesi ve sürdürülebilir kentleşme stratejilerinin

benimsenmesi gibi birçok politika, iklim değişikliğine uyum sağlamada önemli adımlar olarak görülmektedir (Görgülü ve Görgülü, 2021)

Bu noktada, entegre mücadele uygulamaları iklim değişikliğiyle başa çıkmada önemli bir rol oynamaktadır. Tarım, enerji, su yönetimi ve şehircilik gibi farklı sektörleri kapsayan entegre mücadele yöntemleri, çok yönlü bir yaklaşımla hem azaltım hem de uyum politikalarını içermektedir. Özellikle biyolojik mücadele yöntemleri, doğal kaynakların korunması ve sürdürülebilir üretim teknikleri gibi uygulamalar, ekosistem dengelerini gözeterek iklim değişikliğinin olumsuz etkilerini en aza indirmeyi amaçlamaktadır. Bu kapsamda, Türkiye'nin iklim değişikliğiyle mücadelede uyguladığı entegre yöntemlerin değerlendirilmesi, etkinliklerinin belirlenmesi ve geliştirilebilecek yeni yaklaşımlar üzerinde durulacaktır (Küçük & Tunca, 2024).

Bu derlemede, Türkiye'de iklim değişikliğinin entegre mücadele yöntemlerinde etkileri incelenecek, etkinliği değerlendirilecek ve sürdürülebilir bir çevre yönetimi için öneriler sunulacaktır. İklim değişikliğiyle mücadelenin yalnızca çevresel bir mesele değil, aynı zamanda ekonomik büyüme, toplumsal refah ve ulusal güvenlik açısından da kritik bir konu olduğu vurgulanacaktır.

İklim Değişikliğinin Tarımsal Zararlılar Üzerindeki Etkileri

Küresel ısınma ile birlikte sıcaklıkların artması, zararlılar için daha uygun iklim koşulları oluşturarak onların daha önce görülmedikleri bölgelere yayılmalarına neden olmaktadır. Ilıman bölgelerde yaşayan zararlılar, yüksek rakımlara ve kuzeye doğru göç etmeye başlamaktadır. Bu durum, bu bölgelerde daha önce karşılaşmamış zararlılarla mücadeleyi gerektirmektedir (Yaşar, Kök & Kasap, 2021). Örneğin, Akdeniz ikliminde yaygın olan zeytin sineği (*Bactrocera oleae*) (Rossi) daha önce serin bölgelerde görülmezken, artık bu alanlarda da etkili hale gelmektedir. İklim değişikliğine bağlı olarak sıcaklık artışı, bu zararlının yaşam döngüsünü hızlandırmış ve yayılım alanını genişletmiştir (Petacchi ve ark., 2015).

Kış aylarının daha ılıman geçmesi, zararlı türlerin ölüm oranlarını düşürmekte ve daha fazla bireyin hayatta kalarak üreme şansını artırmaktadır. Artan sıcaklıklar, zararlı böceklerin üreme döngülerini hızlandırarak yılda daha fazla nesil vermelerine yol açmaktadır. Bu durum, tarımsal ürünlere zarar verme potansiyellerini ciddi şekilde artırmaktadır (Bale ve ark., 2002). Lahana kelebeği (*Pieris brassicae* L.) (Lepidoptera: Pieridae) ve yaprak bitleri sıcaklık artışına bağlı olarak eskisinden çok daha hızlı üremekte ve ek döller vererek bitkisel üretimi tehdit etmektedir (Bale ve ark., 2002).

İklim değişikliği yalnızca zararlı popülasyonlarını değil, aynı zamanda onların doğal düşmanlarını da etkilemektedir. Sıcaklık değişimleri, faydalı böceklerin, kuşların ve diğer predatör türlerinin yaşam döngülerini, göç süreçlerini ve üreme döngülerini olumsuz yönde etkileyerek biyolojik çeşitliliğin azalmasına neden olmaktadır. Özellikle böcek bazlı biyolojik mücadele ajanları, yüksek sıcaklıklar, değişen nem oranlarına ve artan CO₂ karşı hassasiyet göstermekte, bu da zararlıların doğal kontrol mekanizmalarının zayıflamasına yol açmaktadır. Bunun yanı sıra, habitat kaybı ve değişen iklim koşulları, bazı doğal düşman türlerinin popülasyonlarını azaltarak, entegre mücadele uygulamalarının etkinliğini tehdit etmektedir (Subedi ve ark., 2023). Özellikle biyolojik mücadelede kullanılan parazitoitler ve predatörler, sıcaklık değişimlerine karşı daha hassas olduğundan, zararlı türlerin baskılanması daha zor hale gelmektedir. Özellikle biyolojik mücadelede kullanılan parazitoitler ve avcı böcekler, sıcaklık değişimlerine karşı daha hassas olduğundan, zararlıların baskılanması daha zor hale gelmektedir (Küçük & Tunca, 2024). *Cotesia glomerata* (Linnaeus, 1758) parazitoiti, yüksek sıcaklıklara karşı hassasiyet göstererek konukçular üzerindeki baskıyı azaltabilir. Benzer şekilde, *Harmonia axyridis* (Pallas) gibi bazı uğur böceği türleri, aşırı sıcaklık koşullarında

popülasyon kaybı yaşayarak yaprak bitleri gibi zararlıları baskılamayı zorlaştırabilir (Gu ve ark., 2003).

Zararlılar böcek türleri iklim değişikliğine bağlı olarak daha uzun süre aktif kaldıkları için üreticiler tarafından sık ilaçlama yapılmaktadır. Bu durum, pestisitlerin aşırı kullanımına ve zararlılarda direnç gelişimine yol açmaktadır. Özellikle kimyasal mücadeleye karşı bağımsızlık kazanan zararlı türleri, tarımsal üretimde büyük kayıplara neden olmaktadır. Sıcaklık artışına bağlı olarak bazı tarım zararlıları mevcut kimyasal ilaçlara karşı daha dirençli hale gelmiştir (Hekimoğlu ve Altındağ, 2006). Bunun en belirgin örneklerinden biri, mısır kurdu (*Ostrinia nubilalis*) (Hübner 1796) (Lepidoptera: Pyralidae) gibi zararlılardır. Bu türler, artan sıcaklık ve tarımda yoğun pestisit kullanımını nedeniyle kimyasal ilaçlara karşı adaptasyon geliştirmiştir. Aynı şekilde, yeşil kurt (*Helicoverpa armigera*) (Hübner), pestisitlere karşı direnç kazanarak mücadeleyi zorlaştırmaktadır (Pareek ve ark., 2017). Yeni pestisit formülasyonları geliştirilse bile, zararlıların adaptasyon sürecine bağlı olarak bu mücadele yöntemlerinin etkinliği zamanla azalmaktadır.

Yüksek sıcaklıklar, uzun kuraklık dönemleri ve düzensiz yağışlar, tarımsal zararlılar üzerinde doğrudan etkiler yaratmaktadır. Bazı zararlılar sıcak ve kuru koşullarda daha hızlı yayılırken, bazıları ise nemli ve yağışlı hava koşullarında daha aktif hale gelmektedir. Mısır kurdu (*O. nubilalis*) sıcak ve kurak hava koşullarında daha fazla zarar oluştururken (Musser & Shelton, 2005) kırmızı örümcekler (*Tetranychus urticae* Koch) kurak dönemlerde hızla çoğalarak bitkiler üzerinde ciddi tahribata yol açmaktadır (Meynard ve ark., 2013).

CO₂ artışı ile birlikte bitkilerdeki ikincil (sekonder) metabolitler gibi kimyasal olaylar biyolojik etmenlerin konukçu bulma davranışlarını etkileyebilmekte ve üreme potansiyellerini artırabilmektedir (Özgen ve Karsavuran, 2009). Bazı yaprak biti türlerinin alarm feromonlarına verdikleri tepkilerin CO₂ artışı ile azaldığı ve bu nedenle doğal düşman saldırılarına karşı daha duyarlı hale geldikleri belirlenmiştir (Awmack et al., 1997). Artan CO₂ seviyeleri bitkilerin besin değerini değiştirebilir ve bu da zararlıların daha fazla beslenmesine neden olabilir. CO₂ seviyeleri arttığında bazı böcekler daha hızlı büyüyebilir ve daha uzun süre hayatta kalabilir. Bazı zararlılar için, besin değeri azalmış bitkiler daha az çekici hale gelebilir ve bu da zararlıların farklı bitkilere yönelmesine neden olabilmektedir.

Fırtına ve şiddetli rüzgarlar, zararlı türlerin dispersiyonunu kolaylaştırarak yeni habitatlara ulaşmalarına ve popülasyonlarını genişletmelerine olanak tanımaktadır. Özellikle yaprak bitleri (Aphididae) ve beyaz sinekler (*Bemisia tabaci*) (Gennadius 1889), düşük vücut kütleleri ve aerodinamik yapıları sayesinde rüzgar akımlarıyla uzun mesafeler kat edebilmekte ve yeni ekosistemlerde hızla adapte olarak tarımsal üretimi tehdit etmektedirler (Pathania ve ark., 2020).

Şiddetli yağışlar, zararlıların toprakta veya bitki yüzeylerinde barınma koşullarını etkileyerek popülasyon dinamiklerinde değişimlere yol açmaktadır. Özellikle aşırı yağışlar, bazı toprak kökenli zararlıların yaşam alanlarını bozarak popülasyonlarının azalmasına neden olabilmektedir (Subedi ve ark., 2023). Bununla birlikte, yüksek nem oranı, bazı fungal ve bakteriyel kaynaklı hastalıkların yayılımını artırarak zararlı böcekler için uygun bir ortam oluşturabilir. Örneğin, nemli koşullarda un kurdu (*Tribolium confusum* Duval) ve patates böceği (*Leptinotarsa decemlineata* Say) gibi türler daha aktif hale gelerek bitkisel üretimi tehdit edebilmektedir.

İklim değişikliğine bağlı olarak tarımsal zararlılar, konukçu bitkilerini değiştirebilir veya yeni beslenme alışkanlıkları geliştirebilir. Bu da klasik mücadele yöntemlerinin yetersiz kalmasına yol açmaktadır.

İklim Değişikliğinin Entegre Zararlı Yönetimine (IPM) Etkisi

Entegre zararlı yönetimi (IPM), zararlılarla mücadele etmek için çeşitli yöntemlerin birleştirildiği bir mücadele şeklidir. Bu yöntemler arasında kültürel, biyolojik, fiziksel, mekanik, biyoteknik ve kimyasal mücadele yöntemleri yer alır. Entegre zararlı yönetiminin temel amacı, zararlı popülasyonlarını ekonomik zarar seviyelerine indirerek sürdürülebilir tarım uygulamaları sağlamaktır. Ancak, iklim değişikliği, bu yöntemlerin etkinliğini ve uygulanabilirliğini ciddi şekilde etkileyebilmektedir.

İklim Değişikliğinin Kültürel Mücadeleye Etkisi

Kültürel mücadele yöntemleri, tarımsal zararlılarla mücadelede kimyasal uygulamalara alternatif olarak çevre dostu ve sürdürülebilir çözümler sunmaktadır. Ancak iklim değişikliği, bu yöntemlerin etkinliğini doğrudan etkilemektedir.

- Ekim nöbeti ve çeşitlilik uygulamaları: Değişen iklim koşulları, bazı bitki türlerinin gelişimini olumsuz etkileyerek ekim nöbeti programlarının yeniden planlanmasını gerektirmektedir. Özellikle sıcaklık artışı ve değişen yağış rejimleri, bazı kültür bitkilerinin de zararlı türlere karşı çeşitliliği artırmaya yönelik yeni yöntemler geliştirilmesini zorunlu kılmaktadır. (Gvozdenac ve ark., 2022).
- Toprak işleme teknikleri: Aşırı yağışlar ve kuraklık dönemleri, toprak yapısını ve nem dengesini değiştirerek zararlıların üreme ve barınma alanlarını etkileyebilir. Aşırı yağışlar topraktaki ve bitki üzerindeki bazı zararlıları öldürebilirken, aynı zamanda diğer türler için uygun mikro habitatlar oluşturabilir. Kuraklık dönemlerinde ise, bazı zararlı böcek türleri bitkilerin su stresinden faydalanarak zarar verme potansiyellerini artırır (Gvozdenac ve ark., 2022)
- Sulama yönetimi: İklim değişikliğine bağlı olarak su kaynaklarının azalması, sulama yöntemlerinde değişiklik yapılmasını gerektirir. Yanlış sulama yöntemleri, zararlılara elverişli nemli ortamlar sağlayabilirken, doğru uygulamalar zararlı popülasyonlarını kontrol altında tutmaya yardımcı olabilir. Damla sulama gibi yöntemler, fazla nemin önüne geçerek bazı zararlı türlerinin çoğalmasını engelleyebilmektedir (Radwan ve ark., 2024).
- Bitki örtüsünün ve ekosistem yönetiminin önemi: Doğal ekosistemlerin korunması ve biyolojik çeşitliliğin desteklenmesi, zararlıların doğal düşmanlarının hayatta kalmasını sağlayarak kültürel mücadelede önemli bir rol oynar. Ancak iklim değişikliği, bazı doğal bitki örtülerinin değişmesine neden olarak zararlıların ve onların doğal düşmanlarının etkileşim dengesini bozabilmektedir.

Buna ek olarak, sıcaklık, nem ve yağış değişiklikleri gibi iklimsel faktörler, bitkilerin bağışıklık sistemlerini de etkileyebilir. Stresli koşullar altında bitkiler, zararlılara karşı daha savunmasız hale gelebilir. Bu durum, kültürel mücadele yöntemlerinin etkinliğini, yani zararlılara karşı dirençli bitki çeşitlerinin yetiştirilmesi gibi stratejilerin başarısını olumsuz yönde etkileyebilir (Mohamed, 2023). Ayrıca, değişen iklim koşullarıyla birlikte toprak sağlığı da değişebilir. Toprak neminin artması veya kuraklık gibi durumlar, zararlıların daha elverişli koşullarda yaşamasını sağlayabilir ve bu durum mekanik ve kültürel mücadele yöntemlerinin etkinliğini sınırlayabilmektedir.

İklim Değişikliğinin Biyolojik Mücadeleye Etkisi

Sıcaklık artışı, zararlıların hayatta kalma sürelerini uzatabilir ve üreme döngülerini hızlandırabilir. Bu durum, zararlıların normalden daha erken, daha uzun süre aktif olmasına neden olabilir. Bu, zararlıların daha fazla ve daha geniş alanlarda zarar vermesine yol açarak, mücadele stratejilerinin planlamasında zorluklar yaratabilir. Örneğin, bazı zararlı türleri sıcak

iklimlerde hızla çoğalırken, bu artan popülasyonlara karşı biyolojik mücadelede kullanılan faydalı böcekler, artan sıcaklık ve değişen iklim koşulları nedeniyle yaşam alanlarında daralmalar yaşayabilir. Bu durum, doğal düşmanların zararlı popülasyonlarını kontrol etme becerisini azaltabilmektedir (Gvozdenac ve ark., 2022). Bazı doğal düşmanlar, sıcaklık artışıyla birlikte, bitki çeşitliliğinin ve su kaynaklarının azalması ya da CO₂ seviyelerinin artması gibi nedenlerden dolayı buldukları habitatları terk edebilmektedir. Bu durumda, biyolojik mücadele yöntemlerinin etkinliği önemli ölçüde azalır ve üreticilerin kimyasal mücadele yöntemine daha fazla yönelmesine neden olabilmektedir (Rubenstein, 1992). Entomopatojen fungus olan *Beauveria bassiana* (Balsamo) (Ascomycetes: Clavicipitaceae)'nın biyolojik aktivitesi, 25°C sıcaklıklarda azalmaktadır (Amarasekare ve Edelson, 2004). Değişen ve artan sıcaklıklar, zararlı yönetiminde kullanılan ürünlerin birçoğunun daha az etkili olmasına veya hiç etkili olmamasına neden olabileceği için daha sık insektisit uygulamaları gerektirebilir (Musser ve Shelton, 2005).

İklim Değişiminin Kimyasal Mücadeleye Etkisi

Kimyasal mücadele, tarımsal zararlıları kontrol altına almak için en yaygın kullanılan yöntemlerden biridir. Ancak iklim değişikliği, pestisitlerin etkinliği, kullanım sıklığı ve çevresel etkileri üzerinde önemli değişikliklere yol açmaktadır. Artan sıcaklık ve düzensiz yağış, kullanılan pestisitlerin etkinliğini azaltabilir. Özellikle yüksek sıcaklıklar, kimyasal bileşiklerin daha hızlı parçalanmasına neden olarak etki sürelerini kısaltabilir. Aynı şekilde şiddetli yağışlarda, pestisitlerin bitkilerden yıkanmasına ve hedef dışı alanlara taşınmasına sebep olabilir. Bu durum hem kimyasal mücadelenin başarısını düşürmekte hem de su kaynaklarında kirlilik sorunlarına yol açmaktadır. Piretroidler, organofosfatlar ve özellikle biyopestisitler gibi belirli pestisitler, yüksek sıcaklıklarda daha hızlı bozulmaktadır (Musser ve Shelton, 2005). İklim değişikliği nedeniyle değişen zararlı türleri ve göç hareketleri, mevcut kimyasal mücadele yöntemlerinin yetersiz kalmasına neden olabilmektedir. Bu nedenle, farklı etki mekanizmalarına sahip yeni nesil pestisitlerin geliştirilmesi gerekmektedir. Biyoteknoloji destekli kimyasal mücadele yöntemleri, hedef zararlıya özgü etkili çözümler sunarak çevre üzerindeki olumsuz etkileri azaltabilir (Gvozdenac ve ark., 2022).

İklim değişikliğine bağlı olarak kimyasal mücadele yöntemlerinin etkinliğinin azalması, alternatif mücadele stratejilerinin geliştirilmesini zorunlu kılmaktadır. Sürdürülebilir tarım uygulamalarına yönelik entegre mücadele yaklaşımlarının önemi her geçen gün daha da artmaktadır.

İklim Değişikliğinin Biyoteknik Mücadeleye Etkisi

Artan sıcaklıklar ve değişen hava koşulları, zararlı böceklerin davranışlarını ve biyolojik döngülerini etkileyerek feromonların etkinliğini değiştirebilir. Nem oranlarının değişmesi, feromon bileşiklerinin havadaki stabilitesini ve dağılımını etkileyebilir, bu da zararlıları cezbetme potansiyelini azaltabilir. Bunun dışında repellentlerin volatilizasyon oranlarını ve etki sürelerini değiştirebilir. Bazı bitki bazlı repellent bileşiklerinin etkinliği, sıcaklık ve UV ışınlarına bağlı olarak azalabilir, bu da zararlılar üzerinde beklenen caydırıcı etkinin düşmesine neden olabilir. Yağış rejimlerinin düzensiz olması feromon ve repellentlerin etkinliği üzerinde önemli etkilere sahiptir. Öncelikle, aşırı yağış ve yüksek nem, feromonların havada yayılma sürecini değiştirerek zararlı böceklerin bu kimyasalları algılama yetisini azaltabilir. Feromon bileşiklerinin hidrofilik özellikleri nedeniyle yoğun yağış, bu bileşiklerin hızla yıkanmasına veya seyrelemesine neden olabilir, bu da tuzakların çekiciliğini düşürerek biyoteknik mücadelenin etkinliğini azaltabilir (Heuskin ve ark., 2011).

Benzer şekilde, repellentlerin etkisi de yağış rejimlerindeki değişikliklerden doğrudan etkilenmektedir. Bitki bazlı repellentler genellikle uçucu bileşiklerden oluştuğu için aşırı yağış

ve yüksek nem, bu bileşiklerin yüzeylerden hızla uzaklaşmasına ve etkili olduğu süreyi kısaltmasına yol açabilir. Ayrıca, kurak dönemlerin ardından gelen ani yağışlar, repellentlerin yeniden uygulanmasını gerektirerek tarımsal üretim süreçlerinde ek maliyet ve iş gücü gereksinimi yaratabilir (Gvozdenac ve ark., 2022). Bu nedenle, biyoteknik mücadelede kullanılan feromon ve repellent formülasyonlarının iklim değişikliğine uyum sağlayacak şekilde geliştirilmesi önem arz etmektedir.

SONUÇ

İklim değişikliği, zararlılarla mücadelede kullanılan yöntemlerin etkinliğini azaltırken, yeni ve sürdürülebilir entegre mücadele yöntemlerinin geliştirilmesini zorunlu hale getirmektedir. Küresel ısınma, yağış rejimlerindeki değişiklikler ve ekstrem hava olayları, zararlı böcek türlerinin biyolojisini ve ekolojik dağılımını değiştirerek tarımsal üretimde ciddi tehditler oluşturmaktadır. Bu bağlamda, biyoteknik, biyolojik ve kültürel mücadele yöntemlerinin entegre edilmesi, kimyasal mücadeleye olan bağımlılığı azaltarak çevresel sürdürülebilirliği destekleyen yöntemlerin oluşturulmasını gerektirmektedir. Ayrıca, iklim değişikliğinin etkilerini minimize etmek için tarım politikalarının iklim dostu uygulamalarla uyumlu hale getirilmesi büyük önem taşımaktadır. Gelecekte yapılacak araştırmalar, iklim değişikliğine dayanıklı ve çevre dostu mücadele yöntemlerinin geliştirilmesine odaklanmalı, teknolojik yenilikler ve akıllı tarım uygulamalarıyla entegre edilerek zararlı yönetimi daha verimli hale getirilmelidir. Bu süreçte, sürdürülebilir tarım uygulamalarının benimsenmesi ve yaygınlaştırılması hem ekosistem dengesinin korunmasını hem de tarımsal verimliliğin devamlılığını sağlayacak temel unsurlar arasında yer almaktadır.

KAYNAKÇA

- Amarasekare, K. G. ve Edelson, J., 2004. Effect of temperature on efficacy of insecticides to differential grasshopper (Orthoptera: Acrididae). *Journal of Economic Entomology*, 97 (5), 1595-1602.
- Awmack, C. S., Woodcock, C.M. ve Harrington, R., 1997. Climate change may increase vulnerability of aphids to natural enemies. *Ecological Entomology*, 22, 366-368.
- Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K., Butterfield, J., Buse, A., Coulson, J.C., Farrar, J., Good, J.E.G., Harrington, R., Hartley, S., Jones, T.H., Lindroth, R.L., Press, M.C., Symrnioudis, I., Watt, A.D. and Whittaker, J.B. 2002. Herbivory in global climate change research: Direct effects of rising temperature on insect herbivores. *Global Change Biology*, 8(1): 1–16.
- Bayraç, H. N., & Doğan, E. (2016). Türkiye’de iklim değişikliğinin tarım sektörü üzerine etkileri. *Eskişehir Osmangazi Üniversitesi İİBF Dergisi*, 11(1), 23-48.
- Görgülü, Ç., & Görgülü, L. (2021). İklim değişikliğine eko-morfolojik yaklaşım: Kentsel çeper kuşak alanları. *JENAS Journal of Environmental and Natural Studies*, 3(1), 72-99.
- Gu, H., Wang, Q., & Dorn, S. (2003). Superparasitism in *Cotesia glomerata*: response of hosts and consequences for parasitoids. *Ecological Entomology*, 28(4), 422-431.
- Gvozdenac, S., Dedić, B., Mikić, S., Ovuka, J., & Miladinović, D. (2022). Impact of climate change on integrated pest management strategies. *Climate Change and Agriculture: Perspectives, Sustainability and Resilience*, 311-372.
- Gvozdenac, S., Dedić, B., Mikić, S., Ovuka, J., & Miladinović, D. (2022). Impact of climate change on integrated pest management strategies. *Climate Change and Agriculture: Perspectives, Sustainability and Resilience*, 311-372.

- Hekimoğlu, B., & Altındeğer, M. (2006). Organik Tarım ve Bitki Koruma Açısından Organik Tarımda Kullanılacak Yöntemler. *Samsun Valiliği Gıda Tarım Ve Hayvancılık İl Müdürlüğü*, 1-200.
- Heuskin, S., Verheggen, F.J., Haubruge, E., Wathélet, J.P. and Lognay, G. 2011. The use of semiochemical slow-release devices in integrated pest management strategies. *Biotechnol. Agron. Soc. Environ*, 15(3): 459- 470.
- Küçük, M. E., & Tunca, H. (2024). İklim Değişikliğinin Tarım Ürünleri ve Böcekler Üzerindeki Etkisi. *Bursa Uludağ Üniversitesi Ziraat Fakültesi Dergisi*, 38(2), 535-564.
- Meynard, C.N., Migeon, A. and Navajas, M. 2013. Uncertainties in predicting species distributions under climate change: A case study using *Tetranychus evansi* (Acari: Tetranychidae), a widespread agricultural pest. *PLoS One*, 8(6): e66445.
- Mohamed, S. (2023). Sürdürülebilir Tarım İçin Entegre Zararlı Yönetimi Stratejileri: Mevcut Uygulamaların ve Gelecekteki Yönlerin İncelenmesi. *Asya Araştırma İlerlemeleri Dergisi*, 6 (1), 475-483.
- Musser F. R., Shelton A. M., 2005. The influence of post-exposure temperature on the toxicity of insecticides to *Ostrinia nubilalis* (Lepidoptera: Crambidae). *Pest Management Science*. 61:508–510.
- Özgen, İ., Karsavuran, Y., 2009. Küresel iklim değişikliklerinin böcekler açısından değerlendirilmesi. *Harran Üniversitesi Ziraat Fakültesi Dergisi*. 13, 51-61.
- Pareek, A., Meena, B., Sharma, S., Tatarwal, M., Kalyan, R. and Meena, B. 2017. Impact of climate change on insect pests and their management strategies. *Climate change and sustainable agriculture*. New India Publishing Agency, p:253-286.
- Pathania, M., Verma, A., Singh, M., Arora, P.K. and Kaur, N. 2020. Influence of abiotic factors on the infestation dynamics of whitefly, *Bemisia tabaci* (Gennadius 1889) in cotton and its management strategies in North-Western India. *International Journal of Tropical Insect Science*, 40(4): 969–981.
- Petacchi, R., Marchi, S., Federici, S., & Ragagnini, G. (2015). Large-scale simulation of temperature-dependent phenology in wintering populations of *Bactrocera oleae* (Rossi). *Journal of Applied Entomology*, 139(7), 496-509.
- Radwan, M. H., Abdel-Hameed, I. M., & Nouredine, A. Z. (2024). The impact of various climate change factors on agricultural insect pests, crop production, and their management strategy in india-a review. *Zagazig Journal of Agricultural Research*, 51(4), 697-714.
- Rubenstein, D. I. 1992. The greenhouse effect and changes in animal behavior: effects on social structure and lifehistory strategies. In *Global Warming and Biological Diversity*. 14, 180-192.
- Soltekin, O., Altındışlı, A., & İşçi, B. (2021). İklim değişikliğinin Türkiye’de bağcılık üzerine etkileri. *Ege Üniversitesi Ziraat Fakültesi Dergisi*, 58(3), 457-467.
- Subedi, B., Poudel, A., & Aryal, S. (2023). The impact of climate change on insect pest biology and ecology: Implications for pest management strategies, crop production, and food security. *Journal of Agriculture and Food Research*, 14, 100733.
- Şen, Ö. L., Bozkurt, D., Göktürk, O. M., DüNDAR, B., & Altürk, B. (2013). Türkiye’de iklim değişikliği ve olası etkileri. *Taşkın Sempozyumu*, 29, 30.
- Şen, Z. (2022). İklim değişikliği ve Türkiye. *Çevre Şehir ve İklim Dergisi*, 1(1), 1-19.
- Yaşar, İ., Kök, Ş., & Kasap, İ. (2021). Küresel Isınma ve İklim Değişikliğinin Böcekler Üzerindeki Olası Etkileri. *Lapseki Meslek Yüksekokulu Uygulamalı Araştırmalar Dergisi*, 2(4), 67-75.

TOMATO YELLOW LEAF CURL VIRUS (TYLCV) AND CONTROL METHODS IN SOLANACEAE

Ayşe Altinkaya

Agricultural Genetic Engineer, Ayşe, Altinkaya, Multi Tohum Tar. San. Tic. A.Ş.
Antalya, Türkiye

ORCID ID: <https://orcid.org/0009-0000-7235-2249>

Melahat Özge Özen

Agricultural Engineer MSc., Melahat Özge, Özen, Multi Tohum Tar. San. Tic. A.Ş. Antalya,
Türkiye

ORCID ID: <https://orcid.org/0000-0002-0529-3258>

Nur Ülger

Agricultural Engineer MSc. R&D Assistant Manager, Nur, Ülger, Multi Tohum Tar. San.
Tic. A.Ş. Antalya, Türkiye

ORCID ID: <https://orcid.org/0000-0003-3222-8037>

Büşra Ülger

MSc. Plant Biotechnologist, R&D Engineer, Büşra, Ülger, Wageningen University, Multi
Tohum Tar. San. Tic. A.Ş. Antalya, Türkiye

ORCID ID: <https://orcid.org/0009-0007-7728-6640>

Abstract

The Solanaceae family contains important crops including tomatoes, peppers, and eggplants, that are vegetables produced and exported worldwide. These crops can be grown in tropical and subtropical climates in greenhouses, open fields, and low tunnels. However, several plant diseases and pests adversely affect their cultivation. Viruses are the most destructive plant disease agents. Plant virus diseases affect plant health and fruit quality, resulting in reduced yields and, in the case of severe diseases, loss of the entire crop. The loss of production also causes high economic damage. Tomato yellow leaf curl virus (TYLCV) is one of the most common viral diseases in Solanaceae (nightshade) crops. The virus belongs to the genus Begomovirus that affects tomato (*Solanum lycopersicum*), pepper (*Capsicum spp.*), eggplant (*Solanum melongena*), and potato (*Solanum tuberosum*) plants. TYLCV can be transmitted through infected plants, its host weeds, and its vector whitefly (*Bemisia tabaci*), as the most important factor in its widespread. TYLCV-infected juvenile plants show symptoms such as curled leaves, yellowing due to lack of chlorophyll, and severe growth retardation. In addition to phenotypic observation, serological methods such as PCR and ELISA tests are used to identify the virus. Like other viral diseases, no effective control method has been proposed so far. Therefore, essential disease management is to prevent the occurrence and spread of the pathogen. This review will provide information on the transmission routes and symptoms of TYLCV, one of the most important viral diseases in the Solanaceae family, and the control methods that can be applied, aiming to prevent high yield losses that may occur.

Keywords: Solanaceae, Begomovirus, Tomato yellow leaf curl virus, *Bemisia tabaci*

GİRİŞ

Sebzeler antik çağlardan bu yana insan beslenmesi ve sağlığı için önemli bir yere sahip olmuştur. Dünyada yaşanan nüfus artışıyla beraber insanların gıdaya olan ihtiyacı giderek artmaktadır. Bu durum insan beslenmesinde önemli bir yere sahip olan sebzelere talebin artmasına, dolayısıyla bitkisel üretim miktarında artışa gidilmesine ihtiyaç doğurmuştur. Sebze yetiştiriciliği dünyada ve ülkemizde örtü altı, açık arazi gibi farklı alanlarda yapılmaktadır (Gönül ve ark., 2023). Dünya sebze yetiştiriciliğinde ilk sırada Çin ardından, Hindistan ve ABD gelmektedir. Türkiye ise yıllık yaklaşık olarak 33 600 000 ton sebze üretimiyle dördüncü sırada yer almaktadır (TÜİK, 2024).

Ülkemizde sebze yetiştiriciliğinde ilk sırada meyveleri tüketilen sebzeler domates, biber, patlıcan, kavun, karpuz, hıyar, kabak yer almaktadır (Düden ve Daşgan,2021). Sebze üretiminde Türkiye’de en çok yetiştiriciliği yapılan domates (*Solanum lycopersicum*) 13 000 000 ton üretim miktarı ile 4. sırada yer almaktadır. Bunu patates (*Solanum tuberosum*) 5 700 000 ton üretim miktarıyla 7. sırada takip etmektedir. *Solanaceae* (patlıcangiller) familyasının diğer üyelerinden biber (*Capsicum spp.*) 3 018 775 ton ve patlıcan (*Solanum melongena*) 781 242 ton üretilmektedir (TÜİK, 2022-23).

Yeryüzünde kültüre alınan ve tarımı yapılan birçok önemli tür *Solanaceae* familyasına aittir. Bitkiler aleminin en büyük familyalarından biri olan *Solanaceae* familyası besin değeri yüksek sebzelerden, zehirli bitkilere kadar yaklaşık 100 cins ve 2500’den fazla türü kapsamaktadır. Antarktika hariç tüm kıtalara yayılmış olup en fazla çeşitliliği Orta ve Güney Amerika’da göstermiştir (Motti, 2021). Dünya genelinde yoğun olarak tropikal ve subtropikal bölgelerde yetiştirilmektedir. *Solanaceae* familyası annual, biennial veya perennial olmak üzere genellikle otsu fakat bazı türleri çalı veya küçük ağaç formunda büyüyen bitkilerden oluşmaktadır (Gebhardt, 2016). Kökler kazık kök veya saçak kök yapısına sahip olabilirler. Yapraklar, genellikle kısa saplı, basit yapılıdır. Çiçekler çoğunlukla erselik yapıda, beş adet çanak ve taç yaprakla karakterizedir. Meyve şekli türlerine göre değişkenlik göstermekte; kapsül, bakla, yumru gibi farklı tiplerde olmaktadır (Britannica, 2015). Farklı iklim koşullarına ve çevresel faktörlere kısa sürede uyum sağlayabilen *Solanaceae* familyası türleri yetiştirme sezonu süresince yüksek sıcaklığa ve kuraklığa karşı hassastır (Şahin ve Taşlıgil, 2014; Reddy ve ark., 2018; Kahraman, 2017). Yetiştiriciliğinde su teminine ve toprak havalandırılmasına dikkat edilmelidir (Katsoulas ve Kittas, 2008; Lal ve ark., 2022). Ülkemiz de ve dünyada kültüre alınan, ekonomik değere sahip başlıca önemli *Solanaceae* türleri içerisinde domates, patates, patlıcan, tütün ve biber bulunur (Afroz ve ark., 2020).

Solanaceae familyasında yer alan patates, *Solanum* cinsine ait yumruları tüketilen agronomik değere sahip önemli bir gıda ve endüstri bitkisidir. Orijini Güney Amerika olarak bilinmekte ve 16.yy’da İspanyollar aracılığıyla Avrupa’ya yayıldığı literatüre geçmiştir (Abed ve Demirhan, 2018). *Solanum* cinsine ait yetiştiriciliği yapılan çoğu patates çeşidi autotetraploiddir. Temelde 12 kromozom ($n = 12$) setine sahip *Solanum tuberosum* L. $4n = 48$ kromozom sayısına sahiptir (Reddy ve ark., 2018). Dünyada tarımı yapılan temel bitkiler arasında çeltik, buğday ve mısırın ardından dördüncü sırada yer alır (Dumanoglu ve Öztürk, 2021). Güncel verilerine bakıldığında 6 900 000 ton patates üretildiği görülmektedir (TÜİK, 2024).

Dünya genelinde yaygın olarak yetiştirilen *Solanaceae* familyasında yer alan diğer bir tarım ürünü biberdir. Biber, içerdiği zengin bileşiklerle kendine özgü aroma ve lezzet kazanarak mutfaklarda önemli bir yer edinmiştir. M.Ö. 7500’lü yıllardan bu yana yetiştiriciliği yapılan biberin anavatanı Amerika kıtası olarak bilinmektedir (Kahraman, 2017; Karim ve ark., 2021). *Capsicum* türlerinin birçoğu diploid, kromozom sayısı $2n = 24$ ’tür. 25 türü kapsayan *Capsicum* cinsi içerisinde en çok tüketimi yapılan biber türü *Capsicum annuum*’dur (Atasoy, 2020). Dünya genelinde geniş alanlarda üretimi yapılan *C. annuum* sıcak ve ılıman iklim sebzesidir. Ülkemizde hemen hemen her bölgede yetiştiriciliği yapılmaktadır (Basım ve ark., 2024). 2024

yılında 3 018 775 ton üretimiyle biber, sebze üretiminde, domates ve salatalıktan sonra üçüncü sırada yer aldığı Türkiye İstatistik Kurumu (TÜİK) tarafından rapor edilmiştir.

Patlıcan, *Solanaceae* familyası içerisinde *Solanum* cinsine üye, ılıman ve tropik iklim koşullarında yetişen ekonomik öneme sahip bir sebze türüdür. Kökeni Hindistan olarak bilinmekte ve kromozom sayısı $2n = 24$ 'dür (Kalloo, 1993). Son veriler incelendiğinde toplam 59 346 227 ton patlıcan üretildiği ve bu üretimin %64,6'sının Çin'e ait olduğu görülmektedir (FAO, 2022). Ülke ekonomisine katkı sağlayan patlıcan üretimi açık tarla ve örtü altı şartlarında yapılmaktadır (Altınok ve ark., 2012). 2023 yılında ülkemizde toplam 781 242 ton patlıcan üretilmiştir. En fazla patlıcan üretilen bölge 450 140 ton ile Akdeniz Bölgesi olmuştur (TÜİK, 2023).

Kökeninin Orta Amerika olduğu bilinen tütün *Solanaceae* familyasına mensup *Nicotiana* cinsine ait ve kromozom sayısı $2n=48$ 'dir. Tütün tarımının tarihi M.Ö. 6000'li yıllara kadar uzanmaktadır (Kenton ve ark., 1993; Şahin ve Taşlıgil, 2014). Tütün, içerdiği fenolikler nedeniyle birçok sektörde ham madde olarak kullanılan önemli endüstri bitkisidir. Tütün kökünde organik azotlu nikotin maddesini üretilip yapraklarında bu maddeyi biriktirmesiyle, yaprakları kullanılan tek yıllık bir bitkidir. (Parlak ve Karalezli, 2020). Gıda dışında kullanılan bitkiler arasında dünyada en fazla üretimi yapılan endüstri bitkilerinden biri olan tütün, dünya genelinde 120'den fazla ülkede yaklaşık 3 600 000 hektarlık alanda üretimi yapılmaktadır (Yarar, 2023).

Domates, *Solanaceae* familyasına ait ekonomik değeri en yüksek bitkidir (Binbir ve Duman, 2024). Meyvesi tüketilen domates (*S. Lycopersicum L.*), $2n=24$ kromozoma sahip otsu bir bitkidir (Kızmaz, 2024). 2022 yılı FAO verileri incelendiğinde dünya genelinde toplam sebze üretiminin %16'luk kısmını 86 107 972 ton ile domatesin karşıladığı görülmektedir. Türkiye domates (*Solanum lycopersicum*) üretiminde Çin ve Hindistan'dan sonra 3. sıraya yerleşmiştir (FAO, 2023).

Kültür bitkileri yetiştiriciliğinde çeşitli biyotik ve abiyotik kaynaklı sorunlarla karşılaşmaktadır. Ekonomik değere sahip *Solanaceae* familyasının üretiminde yüksek oranda verim kayıplarına sebep olan virüs hastalıkları bulunmaktadır (Czosnek ve Laterrot, 1997; Erdoğan ve ark., 2022). *Solanaceae* familyasında yüksek oranda verim kayıplarına Begomovirüslerden domates sarı yaprak kıvrılma virüs (TYLCV); domates lekeli solgunluk virüsü (TSWV); Tobamovirüslerden domates kahverengi buruşukluk meyve virüsü (ToBRFV); Potyviruslerden potato virus Y (PVY) sebep olan bazı virüs hastalıklarıdır (Picó ve ark., 1996; Hanson, 2022). *Solanaceae* familyası yetiştiriciliğinde sık rastlanan virüs hastalıklarından biri olan TYLCV, ortaya çıktığı alanlarda yıkıcı etkilere sahip olduğu ve enfeksiyonun yoğun olduğu durumlarda tüm mahsulün kaybına sebep olduğu bilinmektedir (Picó ve ark., 1996; Czosnek ve Laterrot, 1997; Polston ve Anderson, 1997). Türkiye'de ilk defa 1978 yılında domates üretilen bölgelerde TYLCV'nin varlığı tespit edilmiş ve kayıt altına alınmıştır (Demir, 2024). *Solanaceae* familyasından özellikle domates üzerinde yıkıcı etkisi, vektör yoluyla yayılımının hızlı olması ve diğer virüs hastalıklarında olduğu gibi etkin bir mücadele yönteminin bulunmaması nedeniyle TYLCV günümüzde değerini yitirmeyen bir konudur (Kızmaz, 2024). Bu derlemede *Solanaceae* familyasında en önemli viral hastalıklardan biri olan TYLCV'nin taşınma yolları ve oluşturduğu semptomlar, uygulanabilecek mücadele yöntemleri hakkında bilgiler sunulmakta ve oluşabilecek verim kayıplarının önlenmesine katkı sağlaması amaçlanmıştır.


KEŞFİ VE KÖKENİ

Günümüze kadar Dünya üzerine orijininin farklı baz-nükleotit dizilim varyasyonları ile TYLCV'nin yayılım gösterdiği belirlenmiştir. ICTV (2023) verilerine göre, Shotokuvirae

aleminin; Cressdnavirico şubesinde, Geplafuvirales takımında yer alan *Geminiviridae* familyası 520 *Begomovirus* türü içermekte olup ve bilinen 19 farklı ırkı bulunmaktadır.


İlk olarak 1939 yılında Ürdün tarımsal üretim alanlarında rapor edilen TYLCV (Avidov, 1944), 1959 yılında İsrail’de beyaz sinek popülasyonu sonrasında hastalık yaptığı belirlenmiştir (Cohen ve Nitzany, 1960). Çalışma indekslerine baktığımızda 1980’li yıllarda virüsün izolasyonu TYLCV’nin tanım ve mücadelesinde kilit noktası olmuştur (Czosnek ve ark., 1988). TYLCV orijini olarak kabul edilen TYLCV-IL (İsrail)’in 2.787 bp nükleotit uzunluğunda monomerik tek iplikçikten (ssDNA) oluştuğu belirlenmiştir (Zhang ve ark., 2009; Kamal ve ark., 2024). Genom üzerinde 314 bazdan oluşan intergenik bölgeye (IR) sahiptir. Şekil 1’ de gösterildiği gibi TYLCV genomu altı açık okuma çerçevesinden dört ORF (C1, C2, C3 ve C4) ve diğer iki ORF (V1 ve V2) viral iplik üzerinde yer almaktadır (Fondong, 2013; Hanson, 2022).

Şekil 1: TYLCV genomu (Medina ve ark.,2018; Hanson, 2022).



ORF	Locus	Nükleoit/amino asit
LIR	2718-223	379-
V2	294-650	357-119
V1	562-1302	741-247
C3	2605-2126	480-160
C3:C2	1303-1567	265-
C1	2717-1924	774-258
C1:C2	1569-2718	999-333
C4	2473-2177	296-
V3	224-523	300-100

Yapılan çalışmalarla gen bölgelerinde okuma farklılıkları belirlenen TYLCV-Mild, TYLCV-Sa, TYLCV-Sic, TYLCV-Mo, TYLCV-Ch ve TYLCV-Th en yaygın olarak karşılaşılan suşlarıdır (Abhary ve ark., 2006; Fauquet ve ark., 2008). Bu hastalığın başlangıç noktası olan Arap yarımadasındaki İran’ dan yapılan bir çalışmada buna ek olarak iki farklı bölgeden gelen TYLCV suşlarının tam genom dizileri kullanılarak filogenetik evrimleri incelenmiştir. Ülkenin güneyindeki TYLCV-IL ve TYLCV-OM suşlarının kuzeydoğusundaki suşlarında C1 okuma bölgesinde daha farklı dizilime barındırdığını ortaya koymuştur (Hosseinzadeh ve ark., 2014).



Şekil 2: TYLCV'nin dağılımı (EPPO, 2024).


Ülkemizde 1978 yılında sebze yetiştiriciliği yapılan alanlarda verim kayıplarında önemli azalmalarla karşılaştığı rapor edilmiştir. Coğrafik konum gereği tarımsal alanlarda ilk görülen ırkları TYLCV-IL ve TYLCV- Mild ırkları olduğu bildirilmiştir (Yılmaz, 1978; Morris, 1997).

Aynı bölgedeki tarım alanlarından toplanan değişik serolojik örneklerde dahi TYLCV yapılarının virolojik etkileri benzemediği görülmüştür (Sertkaya ve ark., 2017; Fidan ve ark., 2019; Erdoğan ve ark., 2022).

Akbar ve ark., (2024) yaptıkları bir çalışmada tam genom sekansı tamamlanmış Antarktika hariç tüm kıtadaki 39 farklı ülkeden TYLCV örneklerini (Çin 53; Kuveyt 34; İran 21; G. Kore 20; Umman 17; USA 14; Meksika 9; Japonya 9; Suudi Arabistan 6; Pakistan 5; Ürdün 4; İtalya 4; İsrail 3; İspanya 3; Dominik Cumhuriyet 2; Küba 2; Lübnan 2; Trinidad ve Tobago 2; Kosta Rika 2; Hindistan 2 ; Avusturalya 1;Portekiz 1; Fas 1; Hollanda 1; Kamerun 1; Hawaii 1; Guatemala 1; Irak 1; Türkiye 1, Yeni Kaledonya 1; İsviçre 1; Estonya 1; Mauritius 1; Sudan 1; Porto Riko 1; Mısır 1; Azerbaycan 1; Fransız Polinezyası 1) kullanarak rekombinasyon analizi ile genetik varyasyonları ortaya koymayı amaçlamışlardır. Şekil 2'de TYLCV'nin dağılımı, etki alanı ve farklılık açısından değerlendirildiğinde önemli Asya ülkeleri arasında dahi gen bölgesindeki okuma farklılığın fazla olduğu bildirilmiştir (EPPO, 2024).

KONUKÇU ARALIĞI VE TAŞINMA YOLLARI

TYLCV, *Geminiviridae* familyasından olup etkilediği host-vektör ilişkisi nedeniyle *Mastrevirüs*, *Curtovirüs* ve *Begomovirüs* olarak 3 gruba ayrılır. Begomovirüsler monokotil *Eleusine indica* (Kil ve ark., 2021) ve çoğunlukla diketiledonlar arasında etkin olmakla beraber beyaz sinek *Bemisia tabaci* ile taşınmaktadır (Zamir ve ark., 1994; Ghanim ve Czosnek, 2000). *B. tabaci*, Dünya üzerinde ekonomik öneme sahip *Solanaceae* familyası başta olmak üzere *Malvaceae*, *Fabaceae*, *Cucurbitaceae* familyasına ait ürünleri etkilemektedir (Fidan ve ark., 2019).



Şekil 3: *Bemisia tabaci*'nin dağılımı (EPPO, 2024).

Hemiptera grubundan polifag *B. tabaci* (Şekil 3) küresel ısınma etkileriyle artan sıcaklıklar, host- vektör ağındaki genişlik ile popülasyonundaki hızlı artışla TYLCV'nin taşınmasında dünya üzerindeki etki alanını genişletmiştir (Sseruwagi ve ark., 2004; Kriticos ve ark., 2020). Bilindiği gibi *B. tabaci* ondan fazla farklı esteraz fenotipi tanımlanmış olup MEAM1 (eski adıyla B biyotipi) ve MED (eski adıyla Q biyotipi) türleri TYLCV'nin yayılımında kilit bir faktör olduğu kabul edilmiştir (Cervera ve ark., 2000; Gilbertson ve ark., 2015). Ülkemizde ise yapılan çalışmalarda Göçmen ve Devran (2002), sekiz farklı *B.tabaci* biyotipinin bulunduğu kaydedilip, popülasyonun yapıları incelendiğinde 2 ana grup popülasyonun oluştuğunu; bunlarında B biyotipinin ve M biyotipinin olduğu bildirilmiştir (Cervera ve ark., 2000; Ulusoy ve ark., 2002). Satar ve ark. (2018), *B.tabaci* B biyotipi popülasyonlarının duyarlılık düzeyleri üzerine yaptıkları bir çalışmada Akdeniz bölgesinde *Solanaceae* familyasında domates, biber, bitkilerinin yanı sıra hıyar ve pamuk bitkisi üzerinden örnek aldıklarını belirtmişlerdir. Bu çalışmalar dışında ülkemizde *B. tabaci* Q biyotipini Karut ve Tok, (2014) topladıkları popülasyon örnekleri ile yapılan çalışmada kanıtlamışlardır.

Begomovirüs, Bromovirüs, Carmovirüs, Comovirüs, Curtovirüs, Enamovirüs, Luteovirüs, Mastrevirüs, Nanovirüs, Polerovirüs, Rymovirüs, Sobemovirüs, Tymovirüs ve Umbravirüs familyası üyeleri sirkülatif- nonpropagatif biçimde taşınmaktadır (Çandar ve Gümüş, 2013). Bu taşınmada virüs, vektör bünyesine en az 10 dakika beslendikten sonra, en fazla 48 saat içerisinde gerçekleşmektedir. Virüs etkisini ilerleyen generasyonlarda da göstermekte olduğu ancak vektör içerisinde çoğalmadığı bilinmektedir (Blancard ve ark., 2009).

Solanaceae familyasının ekonomik öneme sahip domates bitkisi TYLCV'nin ana konukçusu olup, yabani domates türlerinin semptom göstermeden virüs taşıyıcısı olduğu bilinmektedir (Zakay ve ark., 1991; Vidavsky ve Czosnek, 1998). İsrail'de domates (*Lycopersicon esculentum* cv. Daniella) ve pamuk (*Gossypium hirsutum* cv. Akala) bitkilerinde yapılan bir çalışmada TYLCV-Is'in hiçbir bitki ya da virüs kaynağı bulunmadan sadece virüs taşıyan beyaz sineklerin çiftleşmesi yoluyla bulaşmanın %28-56 arasında değiştiğini bildirmişlerdir. Bu sebeple enfekte olmayan bitkilerin %35'ine kadarının TYLCV ile enfekte olduğunu göstermiştir (Ghanim ve Czosnek, 2000). Günümüze kadar yapılan pek çok çalışmada bunun aksine, *B.tabaci* TYLCV yayılımında beslenme dışı enfeksiyonun etkin olmadığı, yabancı ot ve uygun sıcaklık koşullarında yaşam döngüsünü sürdürüp ekonomik zarar vermeye devam etmektedir (Marchant ve ark., 2024).

Değişen CO₂ seviyelerinin *B.tabaci* yayılımını arttırdığı ve bununla birlikte TYLCV'nin enfekte etkisinin araştırıldığı bir deneme kurulmuştur (Peñalver-Cruz ve ark., 2020; Roy ve ark., 2021). Ancak CO₂ seviyelerinden ayırım gözetmeksizin virüsün yayıldığı raporlanmıştır.

Bu çalışmada ayrıca *B.tabaci* biyotiplerinin yüksek sıcaklıkların adaptasyon-üreme yetenek azalmalarına sebep olduğu, oda sıcaklığı seviyelerinde ise bu popülasyonlarda daha fazla yayılım olduğu bildirilmiştir (Guo ve ark., 2013; Aregbesola ve ark., 2019).

Doğada yabancı otlar yetişen kökleriyle gerekli mineral ve su içeriğini sömürerek kültür bitkileriyle ilişkilerinde ürün kalite, randıman kaybının yanı sıra TYLCV'nin bitkiden bitkiye taşınmasında *B.tabaci* için konukçu olmaktadır (Prasad ve ark., 2020). Kashina ve ark., (2002) 6 farklı bölgede yapılan survey çalışmasında: *Aizoaceae* familyasından *Trianthema portulacastrum*; *Amaranthaceae* familyasından *Alternanthera sessilis*, *Amaranthus spinosus*, *Amaranthus viridis*, *Celosia trigyna*; *Asteraceae* familyasından *Acanthospermum hispidum*, *Ageratum conyzoides*, *Bidens pilosa*, *Chromolaena odorata*, *Eclipta prostrata*, *Erigeron floribundus*, *Synedrella nodiflora*; *Commelinaceae* familyasından *Commelina diffusa*, *Commelina erecta*; *Convolvulaceae* familyasından *Ipomea batatas*; *Solanaceae* familyasından *Crotalaria retusa*; *Euphorbiaceae* familyasından *Euphorbia heterophylla*, *Euphorbia hirta*; *Fabaceae* familyasından *Calopogonium mucunoides*, *Cassia obtusifolia*, *Cassia occidentalis*; *Loganiaceae* familyasından *Spigelia anthelmia*; *Malvaceae* familyasından *Sida acuta*; *Nyctaginaceae* familyasından *Boerhavia diffusa*; *Portulacaceae* familyasından *Portulaca oleracea* türlerinin TYLCV'nin konukçusu olduğu belirtilmiştir. Kıbrıs'ta yapılan 122 tür dikottan bir diğer kapsamlı çalışmada 4.000 örnekte TYLCV içlerinde *Chenopodiaceae*, *Compositae*, *Cruciferae*, *Geraniaceae*, *Leguminosae*, *Orobanchaceae*, *Plantaginaceae*, *Primulaceae*, *Umbellifera* ve *Urticaceae* familyalarında dahil olmak üzere 49 farklı türde tespit edilmiştir (Papayiannis ve ark., 2011). Begomovirüs taşınımının epidemiyolojik sonuçlarını anlamak için yapılan çalışmalarda, kontrollü koşullar altında vektör popülasyonu ile konukçusu olan kültür ürünlerini ve bununla birlikte çevre koşullarındaki özellikle yabancı ot türleri üzerinde yakın gözlemlerinin de yapıldığı kapsamlı deneysel çalışmalar yürütülmelidir. Güney Kore'de bu konuda güncelde Qiao ve ark., (2025) virüs taşınım ve yayılımı hakkında kapsamlı bir çalışma yürütmüşlerdir. Bu çalışmada TYLCV ve TSWV ile enfekteli domates örneklerine ek olarak çalışma yapılan sera etrafındaki 16 farklı yabancı ot türüne yaptıkları virüs testlerinde *Polygonacea* familyasından *Rumex crispus* her iki virüse enfekteli olduğu tespit edilmiştir (Şekil 4).



Şekil 4: TYLCV veya TSWV tespiti için domates seralarından toplanan yabancı otlar: (A) *Cerastium glomeratum*, (B) *Rorippa palustris*, (C) *Rumex crispus*, (D) *Senecio vulgaris*, (E) *Sonchus asper* ve (F) *Stellaria aquatica* (Qiao ve ark., 2025).

Kültür bitkilerinde olduğu gibi gözle kolaylıkla ayırt edilebilen semptomları yabancı otlar göstermese de İran’da yapılan bir çalışma *Solanaceae* familyasından *P. Divaricata*, *Malvaceae* familyasından *H. Trionum*, *Apiaceae* familyasından *A. Majus*, *Euphorbiaceae* familyasından *R. Communis* türlerinde TYLCV simptomatik etkiler rapor edilmiştir (Shamshiri ve ark., 2019). Mısır ve Doğu Hindistan bölgelerinde yapılan bir çalışmada *R. communis* bitkileri üzerinde *Aleyrodidae* familyasından *T. ricini* yoğun TYLCV enfeksiyon ile büyüme ve gelişmeyi etkilediği başka bir çalışmada da bildirilmiştir (Hussein ve ark., 2025).

Solanaceae familyasından ekonomik öneme sahip türlerde domates (Qiu ve ark., 2013; Baldin ve ark., 2015); patlıcan (Leite ve ark., 2003); patates (Cabanillas ve ark., 2009); tütün (Javaid ve ark., 2016) etkisini milyonlarca dolar zarara neden olmaktadır. Hastalık kaynağı olarak geniş konukçu aralığı, üreme ve adaptasyon yetenekleriyle tek taşınım yolunun *B.tabaci* bilinen TYLCV’nin yapılan ilginç bir çalışmayla tohumla da taşındığı büyük yankı yaratmıştır (Kil ve ark., 2016). TYLCV-IL izolatu ile domates bitkisinde yaptıkları çalışmada tohumla taşındığı ve domates fideleri arasında %80-84 enfeksiyon görüldüğü arasında bildirilmiştir. TYLCV çalışmalarına devam eden ekip domates dışında *Solanaceae* familyasından 4 farklı *Capsicum annuum* L. üzerinde tohum enfeksiyonu rapor etmişlerdir (Kil ve ark., 2018).

Bu çalışmalara karşılık *Solanaceae* familyasından *N. benthamiana* bitkisinde (Rosas-Diaz ve ark., 2017; Pérez-Padilla ve ark., 2020) *S. lycopersicum* ve/veya *N. benthamiana* bitkilerinde TYLCV-IL’in tohum kaynaklı olduğu olabileceğine ancak ile tohumdan bulaşmadığı sonucuna ulaştıklarını bildirmişlerdir. Uzun yıllar boyunca kabul görmüş begomovirüslerin tohumla taşınmadığı çalışmalarına karşıt yapılan çalışmalar dünya üzerinde TYLCV dağılımı açısından önem arz etmektedir. Günümüzdeki teknolojik gelişmeler göz önünde bulundurulduğunda taşınımı hakkında daha fazla çalışma yapılmasının gerektiğini düşünmekteyiz.

BELİRTİLERİ

Hastalık özellikle ana konukçusu *Solanaceae* familyasından domates başta olmak üzere enfekte olduğu tüm bitkilerin genç yapraklarında sararma ve fazla yan dal oluşumu nedeniyle çalimsı gibi bir forma neden olmaya başlanmasıyla kolaylıkla ayırt edilmektedir (Şekil 5A).



Şekil 5: TYLCV’nin Solonecea familyasına ait farklı türlerdeki belirtileri A; Hastalıkla enfekte olduktan sonra domates bitkisinin yapraklarında kıvrılma ve çalılışma belirtileri (Murray ve ark.,2023), B; Yaprakların damar aralarında ve kenarlarında kloroz (Montasser, 2012).

TYLVCV domates bitkilerinde genel bir sararmanın yanı sıra farklı belirtiler de göstermektedir (Şekil 5B). Yıkıcı semptomlara neden olan Begomovirüsler bitkide ilk olarak doğrudan floemi etkileyerek bitki taşınımını engeller ve solgunluk şeklinde görülür (Levy ve Lapidot, 2008). Meyvelerde şekil bozukluğu ve kahverengileşmeler şeklindeki renk değişimleri, yaprak ve meyvelerde konsantrik halkalı lekeler, kabarcık şeklinde pürüzlenmeler görülür (Manacorda ve ark., 2021; Bupi ve ark., 2024).

Enfeksiyon yoğunluğuna bitkinin generatif aşamasında yakalanıldığında ise meyve tutumunda olgunlaşmadan dökülerek verimde azalmalara neden olduğu bilinmektedir (Şekil 6H). Ancak TYLVCV ile meyve olgunluk aşamasında karşılaşıldığında ise alt dölllerinde semptom göstermeden gelişimin devam ettiği bildirilmiştir (Thongrit ve ark., 1986; Blancard ve ark., 2009) (Şekil 6I).



Şekil 6: H; Dayanıklı domates bitkileri ve meyveleri, I; Virüsle enfekte dayanıksız domates bitkileri ve pazarlanabilir meyvelerinde kalite farklılıkları (Marchant ve ark.,2020).

Prasad ve ark., (2020) yapraklarda mozaik şeklinde renk değişimleri, kenarlarından aşağıya veya yukarıya doğru kıvrılmaları (Şekil 7C), yaprak ayalarının daralması (Şekil 7D) gibi tipik virüs hastalıkları belirtileri gözlemlenmiştir.



Şekil 7: TYLCV'nin *Solonecea* familyasına ait farklı türlerdeki belirtileri C; Biber bitkisindeki çalılışma belirtileri, D; Biber bitkisindeki genç yapraklarında sararma ve kıvrılma.

Diğer virüs semptomlarından farklı olarak TYLCV ile enfekte olduğu tüm türlerde yaprakların negatif gravitropizm büyümesinin yanı sıra içe doğru kıvrılması domates, biber, patlıcan, kabak, fasulye gibi türlerde de sararma yaptığı pek çok çalışmada rapor edilmiştir (El-Rahmany ve ark.,2025).



Şekil 8: TYLCV'nin *Solonecea* familyasına ait farklı türlerdeki belirtileri E; TYLCV enfekte edilmemiş tütün bitkisi, F; TYLCV enfekte edilen tütün bitkisindeki semptomları (Hassan ve ark., 2016). G; Begomovirüsle enfekte olan patlıcan bitkilerinin yapraklarında yukarıya doğru kıvrılma, küçülme ve mozaikleşme semptomları (Venkataravanappa ve ark., 2014).

TEŞHİS

Begomovirüs hastalıklarının teşhisinde farklı test yöntemleri kullanılmaktadır. Virüs enfektesi sonrasında hastalık semptomlarını etkili gözlemlerle belirlemek ve erken teşhis etmek mücadelede son derece önemlidir. TYLCV'nin mücadelesinde yol haritası DNA ekstraksiyonu ve elektroforezi çalışmalarıyla başlayıp teknolojik gelişmelerle günümüze kadar artması sağlanmıştır (Yılmaz ve ark., 1991). Moleküler markör destekli seleksiyon (MAS) ıslah programlarında resesif genlerin belirlenmesinde kolaylık sağlaması ve hipostatik genlerin başarılı analizleri nedeniyle tercih edilmektedir (Güleç ve ark., 2010). Polimeraz zincir reaksiyonu (PCR) ile virüslere daha spesifik dizileme yapılırken; Revers-Transkriptaz Polimeraz Zincir Reaksiyonu (RT-PCR) yardımı ile az miktarda RNA ile gen düzeyinde okumalarla prob olarak kullanılmasına olanak sağlanmaktadır (Okutucu ve Pehlivan,2003). Gözle ayırt edilebilmesi zor karakteristik özelliklerin gen düzeyinde bilgilerini kontrollü bir şekilde aktarım takibinde kodominant SCAR (Sequence Characterized Amplified Regions) ve CAPS (Cleaved Amplified Polymorphic Sequence) markör sistemleri büyük ilgi görmektedir (Collard ve Mackill, 2008). TYLCV çalışmalarında kullanılan hastalık ve zararlıların farklı ırklarını genom düzeyinde analiz etmede DAS-ELISA (Double antibody sandwich enzyme linked immunosorbent assay) kullanılmaktadır (LecoQ, 1997; Ülger ve ark., 2023).

Avrupa' da yer alan TYLCV ırklarının izolasyonunda Dot-blot hibridizasyon, RFLP ve PAGE (polyacrylamide gel electrophoresis) yöntemleri denenmiştir. TYLCV-Is ve TYLCV-Sar ırklarının yoğun olduğu yaptıkları araştırmalarla belirlenmiştir. Denedikleri yöntemler arasından hızlı ve etkili sonuç için RFLP (Restriction endonuclease fragment length polymorphism) önerilirken, örnek başına maliyetin Dot-blot hibridizasyonda daha az olduğunu fakat PAGE yöntemi ile gen bölgesi analizinin en hızlısı olduğunu yayınlamışlardır (Accotto ve ark., 2000).

Domates bitkisinde yapılan bir çalışmada Lafrance ve ark., (2023) TYLCV, TSWV ve *Fusarium oxysporum* f. sp. *lycopersici* (Fol) dayanımlı hat belirlemede kullanılan 2 tip MAS analizini karşılaştırmışlardır. Türe ve hedef bölgeye yönelik çalışmalarda protokol 2 başarılı sonuç elde etmede daha yaygın kullanıldığını bildirmişlerdir. Dominant moleküler At-2 marker (protokol 1) ve SSR-67 (protokol 2) yöntemleriyle hassas dominant ve heterozigot bireylerin seçiminde kodominant SCAR moleküler yöntemlerin kullanılması gerektiğini düşünmektedirler.

Farklı hastalık ve zararlılara dayanıklılık için domates genotiplerinde SCAR VE CAPS markerları kullanarak tarama yapan Kıymacı ve ark., (2023), TYLCV dayanımında Ty-2 veya Ty-3 homozigot dayanıklılık genotipler dayanıklı olarak skorlamışlardır. Ekip çalışma sonunda, *Verticillium wilt*, *Meloidogyne incognita*, *Fusarium oxysporum* f. *Sp*, TMV, TSWV, TYLCV patojenlerin tümüne birden dayanıklı genotipler tespit etmişlerdir.

59 domates germplazmı ToMV, TSWV ve TYLCV'ye karşı dirençleri belirlemek için DNA ekstraksiyonu ve moleküler markör tespiti yapmışlardır. Domates germplazmı örneklerininin %47,46'sının hastalık dayanıklı olduğunu bunların sırasıyla; 14 ToMV dirençli germplazm, 5 TSWV dirençli germplazm ve 9 TYLCV dirençli germplazm olduğunu bildirmişlerdir (Meng ve ark., 2025).

Missihoun ve ark., (2025) Benin'de 5 farklı bölgeden toplanan şili biberi (*Capsicum spp.*) yaprak örneklerinde RT-PCR yapılarak, bu bölgelerdeki 12 virüs dağılımlarını belirlemişlerdir. Elde ettikleri test sonuçlarına göre örneklerin %18'nin TYLCV içerdiğini rapor etmişlerdir.

Ülkemizdeki domates hatlarının farklı hastalık ve zararlılar için dizayn edilmiş SCAR VE CAPS markerlerinin denendiği bir başka çalışmada, TYLCV için geliştirilen SCAR P6-25 markörünü kullanarak dayanıklılık genine (*Ty3*) sahip bireyler taranmıştır alınmıştır. Bu çalışmanın sonucunda belirtilen hastalığa karşı moleküler belirteçler klasik testlemeye göre dayanıklı hat ve çeşit geliştirmede kolaylık sağlayabileceğini belirtmişlerdir (Pınar ve ark., 2013a). Ekibin yaptığı TYLCV dayanıklılık geni (*Ty1*) için geliştirilen JB-1 CAPS markörünü ıslah çalışmalarında kullanarak homozigot dayanıklı birey elde etmede kullanılması hedeflenen çalışmada, popülasyonlarında 64 adet heterozigot dayanıklı ve 26 adet hassas birey bulmuşlardır (Pınar ve ark., 2013b).

Ürgen (2014) tarafından yapılan bir çalışmada Muğla-Fethiye bölgesi domates seralarında alınan domates yaprak örneklerine DAS-ELISA yöntemi kullanarak enfeksiyon oranlarına bakılmıştır. En yaygın TSWV ve TYLCV enfeksiyonu tespit edilmiştir.

Sertkaya ve Yılmaz (2017) tarafından Hatay ilinde organik domates yetiştiriciliği yapılan seralardan topladıkları 214 yaprak örneğine DAS-ELISA yöntemi uygulamışlar ve en fazla TYLCV enfeksiyonu tespit etmişlerdir.

Hastalık ve zararlılara birlikte dayanım sağlanması hedeflendiğinde klasik ıslah yöntemleriyle dayanıklı bireylerin seçilmesi çok zor olduğu bilinmektedir. Çeri ve kokteyl domates çeşitlerinde TYLCV duyarlılık tespitinde (Ty3P6-25) moleküler marker kullanılarak analizler yapılmıştır. Sonuçlara göre dayanıklı ve hassas bireyler tespit eden ekip, kullanılan yöntemin başarılı olduğunu bildirmişlerdir (Basım ve Kandil, 2023).

Ülkemizde yapılan bir diğer çalışmada, Afyonkarahisar topraksız jeotermal domates üretim alanlarından 94 yaprak ve meyve örneğine virüs testi yapmıştır. DAS-ELISA yöntemi ile örneklerde TYLCV ile karşılaşmadığını bildirmiştir (Demir, 2024).

Hajiabadi ve ark. (2012) İran'da yaptıkları bir çalışmada, 5 farklı bölgeden topladıkları örnekleri ELISA ile testlemişlerdir. Ancak örneklerinde diğer araştırmacıların (Farzadfar ve ark., 2002; Jafari ve ark., 2010) belirttiği gibi TYLCV'ye rastlamadıklarını rapor etmişlerdir.

Nikolić ve ark. (2018) Sırbistan'da yürüttükleri çalışmalarında 3220 domates örneğinde DAS-ELISA yöntemi ile 12 virüse test etmişlerdir. Elde ettikleri sonuçlara göre araştırmacılar aralarında TYLCV'nin de bulunduğu 4 virüsle karşılaşmadıklarını bildirmişlerdir.

İran'da domates alanlarından toplanan 1910 örneğe 3 farklı ELISA yöntemi uygulanmış ve virüsle enfekte olan %1,6 oranında örnekte TYLCV'nin de bulunduğu tespit edilmiştir (Massumi ve ark., 2009). Farklı araştırmacılar tarafından geliştirilen markörlerin Hindistan'da ıslah çalışmalarında domates genotiplerinde çalışıp çalışmadığının kontrolü yapan Prasanna ve ark., (2015) başarılı sonuçlar elde etmiştir.

Suudi Arabistan'ın farklı bölgelerinden 145 domates örneğine ELISA ile TYLCV'nin de bulunduğu 11 virüs enfeksiyon testi yapılmıştır. Bunların sadece %5,5 oranında TYLCV ile enfekte olduğunu bildirmişlerdir (Sabra ve ark., 2022).

Kazakistan'da yapılan bir çalışmada lokal ve hibrit çeşitlerin virüs hastalıklarına (ToMV, TSWV, TYLCV) dayanıklılıkları için testler yapılmıştır. TYLCV dayanımlı hatları belirlemede SCAR- Ty2 ve CAPS- Ty3 markerlerini kullanan ekip 2 hibrit ve 1 lokal çeşitte dayanıklılık bulmuşlardır. MAS sistemi kullanarak ıslahçıların bu virüs hastalıklarına dayanım programlarının geliştirilmesi gerektiğini önermektedirler (Pozharskiy ve ark., 2022).

Günümüzde teknolojik devrim niteliğinde; makine öğrenimi (ML) ve derin öğrenme (DL) dahil olmak üzere artan yapay zekâ (AI) kullanımı, geleneksel çözümlerin yerini almaya başlamıştır. Ayrıca birbirini destekleyen sistemlerle, büyük analitik veriler kullanılarak genetik ve fenotipik özelliklerin ilişkilendirilmesi ile teknoloji kullanımını teşvik edeceği tahmin edilmektedir (Gerakari ve ark., 2025). TYLCV enfeksiyonlarının şiddetini değerlendirmek için Bupi ve ark., (2024) bir ML geliştirmişlerdir. Bu çalışmayla TYLCV'nin mücadelesinde etkili ve zamanında müdahale edilebileceği ayrıca, ıslah programlarında kullanılmak üzere dirençli çeşitlerin geliştirilmesinde rol oynayabileceğini bildirmişlerdir.

MÜCADELE YÖNTEMLERİ

Diğer tüm zararlı ve hastalıklara karşı etkili kimyasal mücadele yöntemleri olmasına rağmen, TYLCV'nin de dahil olduğu virüslere karşı kullanılabilen bir yöntem söz konusu değildir. Virütik enfeksiyonlarda mücadelenin önemi günden güne daha da artmaktadır (Agrios, 1997). Bu nedenle savaşmadan önce yayılımını önlemek için alınacak önlemlerle zararı minimuma indirilebilir.

İlk olarak geçmişten bu yana yapılan uygulamalar bilindiği gibi solarizasyon ve *Trichoderma harzianum* (T-22) vb. uygulamalar ile toprak kökenli hastalık etmenlerinin yanı sıra yetiştiricilik yapılan alanlarda konukçuluk görevi gören yabancı otlarla mücadelede önemli bir avantaj sağlamaktadır. Örtü altı yetiştiriciliğinde sezon başlangıcında tüllerin kontrolü yapılmalı (Berlinger ve ark., 1991) ve yapışkan tuzaklar kullanılarak vektörünün yüksek popülasyon oluşturması engellenmelidir.

TYLCV dahil pek çok hastalığa neden olan *B.tabaci* için ekonomik zarar değeri olmadan, görüldüğü andan itibaren mücadeleye başlanması ve popülasyonun kontrol altında tutulması önem arz etmektedir. Yapılan çalışmalarda popülasyon yoğunluğu, yetiştiricilik yapılan kültür bitkisi ve çevre koşulları da göz önünde bulundurularak sayılarında artış gösterebilir (1 tuzak/ 10 m²) ve yenisi ile değiştirilmelidir (Çolak ve ark., 2005). *Solonaceae* familyasından domates, biber ve patlıcan bitkilerinde *B. Tabaci* için yapılan yapışkan tuzak denemelerinde farklı renk, şekil ve aralık çalışmalarından elde edilen sonuç 250 m²'de 4 oranında kurulan sarı-dikdörtgen yapışkan tuzağın (Yano, 1986; Gu ve ark., 2008; Moreau ve ark., 2011; Lu ve ark., 2012; Nair ve ark., 2021) etkili olduğunu bildirmişlerdir.

Bir başka çalışmada UV-A ışığının *B. tabaci*'nin bağışıklık sistemini önemli ölçüde baskıladığını, gelişimsel ve biyolojik parametrelerinin maruz kalma süresiyle paralel azaldığı gözlemlenmiştir (Khan ve ark., 2021). Saito ve ark., (2024) geliştirdikleri *B. tabaci* toplama makinesi küçük yetiştirme alanlarında kullanılabilir olup, teknolojik sistemler kullanılarak büyük alanlarda uygulamaya geçirilmesi faydalı olacaktır.

Elbette geçmişten günümüze kadarki süreçte kimyasal ürünler devreye girmeden önce maliyeti ucuz ve doğal yöntemler denenmiştir. Mücadele için ilk kullanılan materyalin gaz yağı olduğu düşünülmekle beraber, zararlı popülasyonunu azalttığı, ancak büyüme noktalarında yanıklıklara sebep olarak verimi azalttığı bilinmektedir (Shailaja ve ark., 2012).

Çok sayıda bitki virüsünün vektörü beyaz sinekler için viral enfeksiyon oranlarını azaltmada hızlı ve kolay ulaşılabilir olan insektisitler, enfeksiyonun kontrol altında tutulması sağlayabilmektedir (Ding ve ark., 2019; Tan ve ark., 2022).

2013-2015 yılları arasında 4 farklı domates çeşidinde yapılan bir çalışmada insektisit (Imidacloprid ve cyantraniliprole) uygulanması da dahil olmak üzere gümüş polietilen malç kullanılarak TYLCV dayanımı sağlama stratejisinde kullanılabileceğini bildirmişlerdir (Polston ve ark., 1999; Riley ve Srinivasan, 2019). *B. tabaci* biyotiplerinin popülasyon oluşturma ve yumurtlama için *Cucurbitaceae* familyasını diğer mahsullere göre (*Solanaceae*) tercih ettiği bilinmektedir (Devendran ve ark., 2023).

B. tabaci'nin etkili bir kontrolünde etkinliği yüksek insektisitleri saptamak için 5 farklı insektisit karşılaştırmasında cycloxaprid daha etkili bulunmuştur (Wang ve ark., 2016). Fakat son çalışmalar en fazla *B. tabaci* varyantının bulunduğu Çin'de beyaz sineğin ilaçlara direnç kazandığını göstermesi endişe vericidir (Peng ve ark., 2017; Zheng ve ark., 2017; Wang ve ark., 2018a; Wang ve ark., 2018b; Wang ve ark., 2020). Bu nedenle farklı etken dozlarına sahip kombinasyonlar denenmesi gerekmektedir. Örnek olarak; toprakta yaşayan *Streptomyces avermetilis* tarafından üretilen abamectin böcek öldürücü aktiviteye sahip pestisit. Hedeflenen türlere karşı uygun dozlarda kullanıldığında flupyradifurone, spirotetramat cyantraniliprole direnç geliştirdiği *B. Tabaci* için ümit var niteliktedir (Wang ve ark., 2022). Kore'de yapılan sekiz inteksitisitten yaygın *B. Tabaci Q* biyotipine piridaben, milbemektin, spinetoram ve siyantraniliprolün etkili olduğu görülmüştür (Kim ve ark., 2025).

Sabun, neem, indoneem, lisapol deterjan, thiamethoxam etken maddeli kimyasal ilaç karışımları tek tek kullanıldığında az etkili olsa da kombine kullanımlarında %26 oranıyla daha etkili olduğu görülmüştür (Taggar ve ark.,2020). Tek başına etkili olan bitki özüt uygulamalarında bunu üzerine daha kapsamlı çalışmalar yürütülmüştür. Domates bitkilerinde *Acacia nilotica* (%40 oranında), *Annona squamosa*, *Azadirachta indica*, *Boswellia sacra*, *Crotolaria juncea*, *Jatropha dhofarica* (%41 oranında), *Myrtus communis* ve *Sueda aegyptica* (Azam ve ark., 2002); *Zingiber officinale* ekstratı (Zhang ve ark. 2004); *Trichillia havanensis* tohumları ve *Passiflora edulis* (Aldana ve ark., 2006); *Syzygium aromaticum* (%71,5 oranında), *Citrus aurantium amara* (%62,9 oranında) ve kombinasyonu (%74,8 oranında) ; *Azadirachta indica* Juss. , *Dysphania graveolens*, *Ricinus communis*, *Agave lechuguilla torrey*, *Matricaria chamomilla* karışımıyla %60 oranında beyaz sinekle mücadele edildiğini bildirmişlerdir (Pereyra ve ark.,2021).

Tarımsal yetiştiriciliği yapılan ürünlerde 300'den fazla zararlıyla uzun yıllardır etkin olarak mücadele edilmektedir (Erdoğan, 2024). Bunlar arasında en yaygın hastalık kaynağı *B. tabaci* biotype B ve *T. ricini* için kimyasal mücadelede kullanılan acetamiprid etken maddeli mospilan ile bitkisel *Thymus vulgaris* L. ve *Salix babylonica* L. yapraklarından elde edilen yağ, sarımsak özü kombinasyonları yapılan çalışmada *B. Tabaci* popülasyonlarında *Salix babylonica* L.+mospilan uygulamasıyla %78 azalma sağladığı görülmüştür. Bu diğer botanik yağların biyopestisit olarak değerlendirilmesinde *Pogostemon cablin* ve *Azadirachta indica* uygulamaları ile *B. Tabaci* yumurta, ergin ve nimf dönemlerine etkileri denenmiştir. *Pogostemon cablin* ile yumurtlamada %50-89 oranında azalma gözlemlenirken, *Pogostemon cablin* yumurtlama, nimf ve ergin çıkışımları sırasıyla %50, %70 ve %80 oranında azaltmada etkili bulunmuştur (Amour ve ark., 2023).

Patlıcan bitkilerinde organik ahır gübreleri, odun külü, bulamaç karışımlarının (Mandal ve ark., 2018) yanı sıra farklı oranlarda büyükbaş hayvan idrarı ve bitkisel neem yağı ekstratları da etkili sonuçlar vermiştir (Karkar, 2014; Patel ve ark., 2017). Bununla birlikte doğru oranlarda organik

gübreler ve bitkisel özütler sadece zararlı mücadelesinde değil verim artışında da avantaj sağlamıştır (Chabra ve ark., 1988; Elango ve ark., 2020). Doğal ve işlenmiş peynir altı suyu proteini fraksiyonlarının (*a-laktalbumin*, *β-laktoglobulin* ve *laktoferrin*) domates bitkilerinde TYLCV antiviral aktivitesi için kullanılabilir olduğunu bulmuşlardır (Karthikeyan ve ark.,2006; Abdelbacki ve ark.,2010).

Çevreci bir yaklaşımla organik tarımda ve örtü altı yetiştiriciliğinde doğal düşmanlar kullanılarak sıfır kimyasal ile başarılı olabileceğini göstermişlerdir (Kazak ve ark.,2016). Çalışmada avcı olarak *E. mundus* ve *M. melanotoma* kullanılmıştır. Ülkemizde domates yetiştiriciliği yapılan bir alanda *S. parcesetosum*'un sezon boyunca beslenerek yaşayabildiğini ve çoğalabildiğini ortaya koymuştur (Yarpuzlu ve ark., 2022). Çin'de yapılan başka bir çalışmada *B. tabaci* mücadelesinde *Amblyseius swirskii* ve *Nesidiocoris tenuis* biyolojik kontrol olarak kullanılmıştır (Wang ve ark., 2010; Luo ve ark., 2010; Horowitz ve ark., 2020). Mücadele için kimyasal ilaçla desteklendiğinde doğal düşmanların etkinlik düzeyini arttırdığı bilinmektedir.

Türkiye'de domates seralarında sık karşılaşılan zararlılarla mücadelede acetamiprid, chlorantraniliprole, cyantraniliprole, spinetoram, cyantraniliprole ve spinosadın ruhsatı bulunmaktadır (BKÜ, 2024). Güncel olarak ülkemizde *B. Tabaci* populasyonlarının yetiştiricilik yapılan alanlarda insektisit direnç karşılaştırılmasında sulfoxaflor ve cyantraniliprole nimf bireylerde en yüksek etkinliği sergilediğini, ergin bireylerde ise asetamipridin en etkili insektisit olduğunu bildirmişlerdir (İnak ve ark., 2025). Sera alanlarında görülen zararlılara karşı uygulanacak mücadele yöntemlerinde, *B. Tabaci*'nin doğal düşmanlarına zarar vermeyen etken maddeli formülasyonların geliştirilmesi önceliklendirilmelidir (Yüksekyayla ve ark., 2023).

Kimyasal mücadelede ise, tarım ilaçlarının kolay ulaşılabilir olması, hastalık ve zararlı popülasyonunda etkinliği nedenleriyle tercih edilmektedir (Tiryaki ve ark., 2010). Ancak toprak ve suların kirliliğiyle birlikte tüketilen ürünlerde kalıntı bırakmasıyla insan sağlığını ciddi boyutta tehdit etmektedir (Zhang ve ark.,2024). En büyük pestisit üreticisi Çin dahil olmak üzere çoğu ülke pestisit kullanımına yönelik yönetmelikler getirmişlerdir (Yılmaz ve ark.,2024). Yakın zamanda Çin'de geliştirilen ilk antiviral ajan dufulin (Long ve ark., 2008); Qian ve ark.,2010) kullanılarak yapılan çalışmada ise kontrole oranla TYLCV semptomların etkilerini azalttığını gözlemişlerdir (Huang ve ark., 2025). Arbutin, çoğu familyanın tıbbi aromatik olarak isimlendirdiğimiz bitkilerinde değişik oranlarda ihtiva etmektedir. Endüstriyel olarak kullanılan bu maddenin bitki korumadaki potansiyelini belirlemek için bir çalışma yapan Xu ve ark., (2025) bu maddenin antiviral ajan potansiyeli olduğunu bildirmişlerdir. Yaptıkları çalışmada arbutinin, tütün bitkisindeki TYLCV enfeksiyonunu azaltmadaki etkisini göstermişlerdir.


Bitkide abiyotik ve biyotik stres sonrasında savunma mekanizmasından kaynaklı uçucu organik bileşikler (VOC'ler) değişim göstermektedir. Bir çalışmada bitki aktivatörü olarak kullanılsada (Şanlı ve ark., 2016), bitkideki virüs ve böcek vektör kaynaklı VOC'lerin yaptığı değişiklikleri anlamamız direnç mekanizması için kullanılabilceği öngörülmektedir. Bu nedenle *Solanaceae* familyasından domates ve biber bitkileri üzerinde yapılan çalışmada TYLCV ve beyaz sineği kaynaklı damar sarıları virüsü (PeWBVYV) ile enfeksiyonları sonrasında analiz yapıp karşılaştırmışlardır. Yapılan analizler sonucunda iki türde virüs enfeksiyonları sonrası farklı uçucu organik bileşiklerinin artışına neden olduğunu göstermektedir. Şaşırtıcı olarak domates ve biberdeki beyaz sinek istilasına yanıt olarak guaiacol ve metil palmitoleatın arttığı belirlenmiştir (Ghosh ve ark., 2022).

Solanaceae familyasında *Solanum pennellii* ve türetilmiş popülasyonlarda bulunan Acylsugar ikincil metabolit olup, bitkinin savunma mekanizmasında salgılanmaktadır (Gibson, 1976; Silva ve ark., 2014). Kültür bitkilerinde de acylsugar biyosentezi için gerekli olan genetik lokuslar nedeniyle eser miktarda biriktiği tespit edilmiştir (Ning ve ark., 2015). Bu maddenin

yumurtalarını bırakmasını engellemesi, popülasyon azalmasına etkisiyle *B. tabaci* mücadelesinde taşınan virüslere karşı direnci artırmak için bir konukçu bitki savunma sistemi olarak kullanılabileceğini gösteren çalışmalar bulunmaktadır (Liedl ve ark., 1995; Schillmiller ve ark., 2012; Smeda ve ark.,2023). Virüssüz-dayanıklı çeşitler hastalıklara savunma sistemindeki fiziksel bariyerler, kimyasal reaksiyonlar ve çeşitli proteinleri etkin olarak kullanılabildiği rapor edilmiştir. Bu nedenle bitkilerin kendilerini koruma mekanizmalarını anlamak ve bu sistemleri gıda güvenliği için geliştirilmesi gerekmektedir (Freeman ve Beattie, 2008).

Çeşitli virüslerin, çevresel koşullar altında konukçu bitkileriyle mutualistik ilişkiler kurabildiği çeşitli çalışmalarda gösterilmiştir (Xu ve ark.,2008). Aynı anda çeşitli streslere maruz kalan bitkinin, tek stres ile mücadele eden bitkiye göre daha çok savunma gösterdiği bilinmektedir. ToCV ve TYLCV'nin viral enfeksiyonunun birlikte etkisi tek tek enfeksiyonlarına göre metiltransferazın regülasyon seviyesini azaltabileceği gözlemlenmiştir (Li ve ark., 2021). *Nicotiana benthamiana* gen regülasyonlarına bakıldığında (Qiao ve ark., 2023) TSWV ve TYLCV savunmasında benzerlik görülmüş, bir diğer çalışmada ise TYLCV'nin domates bitkilerinde hem termotolerans hem de kuraklığa karşı dayanıklılık sağladığı belirlenmiştir (Anfoka ve ark., 2016; Mishra ve ark., 2021; Shteinberg ve ark., 2021) Bunlara ek olarak Corrales-Gutierrez ve ark., (2020) yaptıkları çalışmada domates ve tütün bitkilerinde TYLCV'nin; Mirzayeva ve ark., (2023) yaptıkları çalışmada ise yine domates bitkilerinde TYLCV ve ToCV'nin dayanımını doğrudan nitelikte sonuçlar elde etmişlerdir.

TYLCV dayanımı için yabancı domates türlerine virüs enfekte eden araştırmacılar; *S. arcanum*, *S. cheesmaniae*, *S. chilense*, *S. galapagense*, *S. chmielewskii*, *S. corneliomulleri*, *S. habrochaites*, *S. neorickii*, *S. peruvianum*, *S. pimpinellifolium* ve *S. pennellii* gibi bazı türlerde dayanıklılık gözlemlenmiştir (Şekil 8).



Şekil 8: Domates kromozomları üzerinde TYLCV direnç genlerinin haritalanması. (A) *S. chilense* cv. kromozom 3 üzerindeki Ty-4 bölgesi, (B) *S. peruvianum* cv. kromozom 4 üzerindeki Ty-5 bölgesi, (C) *S. chilense* cv. kromozom 6 üzerindeki Ty-1, Ty-3 ve Ty-3a bölgesi (sırasıyla LA1969, LA2279 ve LA1932), (D) *S. chilense* kromozom 10 üzerindeki Ty-6 bölgesi (LA2279) ve (E) *S. habrochaites* cv. kromozom 11 üzerindeki Ty-2 bölgesi (El-Sappah ve ark., 2022).

Dayanıklılık genlerinin 5 tanesinin dominant (*Ty-1*, *Ty-2*, *Ty-3*, *Ty-4*, *Ty-6*), bir tanesinin (*ty-5*) resesif dayanım verdiği tespit edilmiştir (Zamir ve ark., 1994; Agrama ve Scott, 2006; Ji ve ark., 2007). *S. peruvianum*'un 4. kromozomunda bulunan *ty-5* geninin TYLCV dayanımının yanı sıra (Anbinder ve ark., 2009) geminivirüslere geniş dayanım sağladığı görülmüştür (Ren ve ark., 2022). İspanya'da yerel hatlar (De la pera) TYLCV dayanımı sağlayan *ty-5* geninin verim ve kalite üzerindeki negatif özelliklerinin az olması nedeniyle tercih edildiği çalışmada

Ty-1 ve *ty-5* kombinasyonunun virüs enfeksiyonuna daha etkili olduğu yayımlanmıştır (Cabrera ve ark., 2025). *Ty-6* geni tek başına TYLCV ve bipartite domates benek virüsüne (ToMoV) karşı direnç sağlamakta olduğu, *Ty-3* ve *ty-5* genleri ile kombinasyonlarının TYLCV direnç seviyesini arttırdığı bulunmuştur. Ayrıca *ty-5'in* ToMoV'ye karşı etkisiz olduğunu fakat bunun tüm bipartit begomovirüsleri kapsamadığını bildirmişlerdir (Gill ve ark., 2019).

TYLCV'ye karşı yapılan çalışmalar ayrıca *Ty-4*, *S. chilense* (LA1969) kromozom 3'ün uzun kolundan introgresyon yoluyla aktarılan *Ty-3*'ten daha az etkili direnç sağladığı bildirilmiştir (Ji ve ark., 2009). *S. chilense*'nin 6. kromozomunda bulunan *Ty-1/Ty-3* genlerinin hastalığa yüksek dayanıklılık sağladığı, aynı zamanda 11. kromozomda bulunan *Ty-2* genlerinin de ıslah çalışmalarında dayanım için yaygın olarak kullanıldığı görülmüştür (Basım ve Kandil, 2024). Koeda ve Kitawaki'nin 2024 yılında yaptıkları çalışmada, enfekteli alanlarda dayanıklı çeşit kullanılan yetiştiricilik alanlarda, küresel ısınmayla artan sıcaklıkların dayanımlı çeşitlerde virüse hassasiyet gösterdiğini gözlemleyerek bilimsel olarak da yaptıkları analizlerle kanıtlamışlardır. Denemeye TYLCV dayanımı sağlayan *Ty-1* genine sahip çeşitte TYLCV-IL hassas dayanıma sahip gibi görünmekte olduğu belirtilmiştir. Bununla birlikte aynı koşullarda TYLCV-Mld enfeksiyonu karşısında herhangi bir semptomla rastlamamışlardır. Diğer bir çalışmada açık tarlada yetiştirilen domates bitkilerinde TYLCV-IL ve ToCV enfeksiyonunda *Ty-1* ya da *Ty-3* ve *Ty-6* genine sahip çeşitlerde semptomların fazla olduğu, gelecekte yeni bir viral izolatin çıkmasının muhtemel olduğunu öngörmektedirler (Kumar ve ark., 2023).

SONUÇ

Günümüzde viral hastalıklara karşı etkin bir mücadelenin bulunmaması sebze yetiştiriciliğini sınırlayan faktörlerden biridir. TYLCV'ye konukçuluk yapan birçok kültür bitkisi bulunması nedeniyle yetiştiricilik sezonlarının yayılımı en az seviyede olduğu iklim koşullarında gerçekleştirmek gibi farklı yaklaşımlar önerilmektedir (Kil vd., 2016). Domates bitkisinde geçici olarak bilinen çeşitlerin güz döneminde üretim yapılmasının virüs hastalık ve beyazsinek vektöründen daha çok etkilendiği tespit edilmiştir (Sertkaya ve Yılmaz, 2017). TYLCV ana konukçusu domates bitkisinde hassas çeşit yetiştiriciliği yapılan bölgede dayanıklı bir çeşit kullanımı, farklı bir familyadan ürünle değişimi ekonomik zararı minimuma indirecektir. Tarım alanlarında çoğu dikotil yabancı otların semptom görülmesizin uzak tutulmasını mücadeleye olumlu etkide bulunacaktır. Kaliteli işçi potansiyelinin azalması ve işçilik maliyetlerinin fazla olması, yabancı otların yok edilmesindeki girdi hesabında insektisit kullanımının avantajları göz önünde bulundurulduğunda tercih edilebilir olmuştur. Kimyasal kullanımı yıllar geçtikçe rekombinasyonlar ile gelişen *B. Tabaci* için yetersiz kalması olağandır. Ancak bu durum, diğer yöntemlerden pestisit kalıntısındaki artışa sebep olarak daha büyük çevresel etkilere neden olmuştur. Yetiştiricilikte dayanıklı çeşitlerin kullanılmasının TYLCV'ye karşı daha etkin bir mücadele yöntemi olacağı düşünülmektedir (Avedi, 2022). Mısır' da yapılan çalışmada tarımın bu yönde ilerlemesi için araştırmacıların TYLCV'nin mücadelesinde dayanıklı çeşit geliştirmeye ve enfeksiyon yönetim uygulamalarını iyileştirmeye odaklanmakta olduğunu bildirmişlerdir (Rahman ve ark., 2024).

Bugüne kadar TYLCV'yle mücadelede ilk olarak vektörü olan beyazsineklere karşı savunma yapılmaktadır. Bu derlemede, kısa ve güncel çalışmalarla TYLCV mücadelesinde kullanılabilecek çevresel düzenleme ve yabancı ot temizliği, kimyasal ilaçlamalar, avcısı olan biyolojik ajanlarla kontrolü, konukçu olduğu bitkilerden daha çekici tuzak bitkilerin konulması, ekim münavebesi, biyopestisitler, dirençli çeşit kullanımlarından bahsedilmiştir. Artan nüfusu beslemek, iklim değişikliği ve viral hastalıkların TYLCV ve ilgili begomovirüslere karşı genetik direncin azaldığı bir senaryonun bizi beklediğini tahmin etmekteyiz. Bu nedenle Yan ve ark., (2021) bahsettiği gibi; TYLCV ve bipartite begomovirüslere karşı doğal direnç için yabancı domates katılımlarının taranması, direnç açısından performansının test edilmesi ve

dayanıklı çeşit geliştirmek için ıslah çalışmalarının yapılması geleceğimiz için önem arz ettiğini düşünmekteyiz.

KAYNAKÇA

Abdelbacki Am, Taha Sm, Sitohy Mz, Abou Dawood, A1, Add-El Hamid Mm, Rezk Aa. (2010). Inhibition Of Tomato Yellow Leaf Curl Virus (TYLCV) Using Whey Proteins. *Virology Journal* 7 (26): 1-6.

Abed, M. M., Ve Ark., Demirhan, B. (2018). Patates Bitkisine (*Solanum Tuberosum L.*) Genel Bir Bakış. *International Journal Of Life Sciences And Biotechnology*, 1(1), 1-9.

Abhary, M. K., Et Al. "Post-Transcriptional Gene Silencing İn Controlling Viruses Of The Tomato Yellow Leaf Curl Virus Complex." *Archives Of Virology* 151 (2006): 2349-2363.

Accotto, G.P., Navas-Castillo, J., Noris, E., Moriones, E., Louro, D., 2000. Typing Of Tomato Yellow Leaf Curl Viruses İn Europe. *European Journal Of Plant Pathology* (2000) 106: 179.

Afroz, M., Akter, S., Ahmed, A., Rouf, R., Shilpi, J. A., Tiralongo, E., ... Ve Ark., Uddin, S. J. (2020). Ethnobotany And Antimicrobial Peptides From Plants Of The Solanaceae Family: An Update And Future Prospects. *Frontiers İn Pharmacology*, 11, 565.

Agrama H.A. And Scott J.W., 2006. Quantitative Trait Loci For Tomato Yellow Leaf Curl Virus And Tomato Mottle Virus Resistance İn Tomato J. Amer. Soc. Hort. Sci. 131(2):267–272.

Agrios, G. N. 1997. *Plant Pathology*. San Diego, CA: Academic. 635 p.

Akbar A, Al Hashash H, Al-Ali E. Tomato Yellow Leaf Curl Virus (TYLCV) İn Kuwait And Global Analysis Of The Population Structure And Evolutionary Pattern Of TYLCV. *Virol J.* 2024 Nov 27;21(1):308. Doi: 10.1186/S12985-024-02540-6. Pmid: 39605076; Pmcid: Pmc11603944.

Akkül, E. (2021). Domates Islahında Tswv, TYLCV, Forl Dayanımlarının Hibrit Verimine Etkisi.

Altınok, H. (2012). Antalya ve Mersin İli Örtü Altı Patlıcan Ekim Alanlarında Kurşuni Küf ve Beyaz Çürüklük Hastalıklarının Yaygınlık Oranlarının Belirlenmesi. *Plant Protection Bulletin*, 52(2), 163-173.

Amour, Massoud, Et Al. "Efficacy Of Selected Botanical Oils Against The Cassava Whitefly (Bemisia Tabaci) And Their Effects On İts Feeding Behaviour." *Journal Of Applied Entomology* 147.7 (2023): 473-485.

Anbinder, I., Reuveni, M., Azari, R., Paran, I., Nahon, S., Shlomo, H., ... & Levin, I. (2009). Molecular Dissection Of Tomato Leaf Curl Virus Resistance İn Tomato Line Ty172 Derived From Solanum Peruvianum. *Theoretical And Applied Genetics*, 119, 519-530.

Anfoka, Ghandi, Adi Moshe, Lilia Fridman, Linoy Amrani, O. R. Rotem, Mikhail Kolot, Mouhammad Zeidan, Henryk Czosnek, And Rena Gorovits. "Tomato Yellow Leaf Curl Virus İnfection Mitigates The Heat Stress Response Of Plants Grown At High Temperatures." *Scientific Reports* 6, No. 1 (2016): 19715.

Aregbesola, O. Z., Legg, J. P., Sigsgaard, L., Lund, O. S., & Rapisarda, C. (2019). Potential İmpact Of Climate Change On Whiteflies And İmplications For The Spread Of Vectored Viruses. *Journal Of Pest Science*, 92, 381-392.

Atalay, E. "Çukurova Bölgesi'nde Domates Sarı Yaprak Kıvrıcıklık Virüsü (Tomato Yellow Leaf Curl Virus, TYLCV)'Nün Irklarının Araştırılması. Çukurova Ünverstesi, Fen Blmler Enstitüsü." *Bitki Koruma Anabilim Dalı, Yüksek Lisans Tezi* 65 (2018).

Atasoy, D. (2020). *Bazı Biber (Capsicum Annuum L.) Islah Genotiplerinin Anter Kültürü Performansla Rının Belirlenmesi* (Doctoral Dissertation, Yüksek Lisans Tezi, Çukuro-Va Üniversitesi, Adana).

Atasoy, D., Baktemur, G., Ve Ark., Taşkın, H. (2021). Bazı Biber (*Capsicum Annuum L.*) Genotiplerinin Anter Kültürü Performanslarının Belirlenmesi. *Yuzuncu Yıl University Journal Of Agricultural Sciences*, 31(2), 282-293.

Avidov, H. Z. "Tobacco Whitefly İn Israel." *Hassadeh, Tel Aviv* (1944): 1-33.

Aydın, A., Başak, H., Aydın, H., Ve Ark., Güngör, R. Characterization Of F2 Generation Tomato Plants And Marker Assisted Selection Against Tomato Spotted Wilt Virus (TSWV) And Tomato Yellow Leaf Curl Virus (TYLCV). *International Journal Of Agriculture Environment And Food Sciences*, 8(3), 618-628.

Azam, K.M.; Bowers, W.S.; Srikandakumar, A.; Al-Mahmuli, I.H.; Al-Raeesi, A.A. Insecticidal Action Of Plant Extracts Against Nymphs Of Whitefly, *Bemisia Tabaci* Gennadius. *Crop Res.* 2002, 24, 390–393.

Baldin, E. L., Aguiar, G. P., Fanela, T. L., Soares, M. C., Groppo, M., & Crotti, A. E. (2015). Bioactivity Of Pelargonium Graveolens Essential Oil And Related Monoterpenoids Against Sweet Potato Whitefly, *Bemisia Tabaci* Biotype B. *Journal Of Pest Science*, 88, 191-199.

Basım, H., & Kandil, O. (2024). Development Of Cluster Tomato Varieties Resistant/Tolerant To Tomato Yellow Leaf Curl Virus (TYLCV) And Fusarium Oxysporum F. Sp. Radicis-Lycopersici (Forl) Through Molecular Marker-Based Plant Breeding. *Black Sea Journal Of Agriculture*, 7(5), 7-8.

Basım, H., Kandil, O., & Karaoğlan, M. (2023). Determination Of TYLCV-Resistant Cherry And Cocktail Tomato Cultivars By Molecular Markers. *Türk Bilim ve Mühendislik Dergisi*, 5(2), 89-96.

Basım, H., Kandil, O., Ve Ark., Ökem, C. (2024). Mevcut Biber Hatları Kullanılarak Domates Benekli Solgunluk Virüsü (TSWV) Hastalığına Karşı Dayanıklı/Toleranslı Charleston Biber Çeşitlerinin Geliştirilmesi. *Uluslararası Gıda Tarım ve Hayvan Bilimleri Dergisi*, 4(2), 12-20.

Bedford, I. D., Briddon, R. W., Brown, J. K., Jones, P., Alkaff, N., And Markham, P. G. 1994. Differentiation Of Three Whitefly-Transmitted Geminiviruses From The Republic Of Yemen. *Eur. J. Plant Pathol.* 100:243-257.

Berlinger, M.J.; Dahan, R.; Mordechi, S.; Liper, A.; Katz, J.; Levav, N. The use of nets to prevent the penetration of *Bemisia tabaci* into greenhouse. *Hassadeh* 1991, 71, 1579–1583.

Binbir, S., Ve Ark., Duman, İ. (2024). Türkiye Domates Genetik Kaynaklarının Agromorfolojik Karakterizasyonu. *Anadolu Ege Tarımsal Araştırma Enstitüsü Dergisi*, 34(Özel Sayı), 52-61.

Blancard, D., Candresse, T., Marchoux, G., Ve Ark., Laterrot, H. (2009). Les Maladies De La Tomato: Identifier, Connaître, Maîtriser.

Britannica, E. (2015). Solanaceae: Plant Family. *Of Encyclopaedia Britannica*. Solanaceae | Definition, Taxonomy, Description, Characteristics, Major Species, & Facts | Britannica

Briddon, R. W., And Markham, P. G. 1995. Family Geminiviridae. Pages 158-165 İn: *Virus Taxonomy: Classification And Nomenclature Of Viruses*. Sixth Report Of The International Committee On Taxonomy Of Viruses. F. A. Murphy, C. M. Fauquet, D. H. L. Bishop, S. A. Ghabrial, A. W. Jarvis, G. P. Martelli, M. A. Mayo, And M. D. Summers, Eds. Springer-Verlag, New York.

Bupi, N. , Vo T. T. B., Qureshi M. A., Et Al. 2024. "Twindemic Threats Of Weeds Coinfected With Tomato Yellow Leaf Curl Virus And Tomato Spotted Wilt Virus As Viral Reservoirs İn Tomato Greenhouses." *Plant Pathology Journal* 40: 310–321.

- Cabanillas, H.E.; Jones, W.A. Pathogenicity Of *Isaria* Sp. (*Hypocreales: Clavicipitaceae*) Against The Sweet Potato Whitefly B Biotype, *Bemisia Tabaci* (*Hemiptera: Aleyrodidae*). *Crop Prot.* 2009, 28, 333–337.
- Cabrera, J. Á., Carbonell, P., Alonso, A., Pérez-Moro, C., De Castro, A. P., Ruiz, J. J., & García-Martínez, S. (2025). Assessment Of Resistance Of Ty-1 And Ty-5 Genes In *Solanum Lycopersicum* Plants Infected With Tomato Yellow Leaf Curl Virus (TYLCV). *Scientia Horticulturae*, 342, 114012.
- Cervera M.T., Cabezas J.A., Simon B., Martínez-Zapater J.M., Beitia F., Cenis J.L., 2000. Genetic Relationships Among Biotypes Of *Bemisia Tabaci* (*Hemiptera: Aleyrodidae*) Based On Aflp Analysis. *Bulletin Of Entomological Research* 90: 391-396.
- Chabra, H.K.; Grewal, P.S.; Singh, A. Efficacy Of Some Plant Extracts On Root Knot Nematode (*Meloidogyne Incognita*). *J. Tree Sci.* 1988, 7, 24–25.
- Chen, Xueying, Et Al. "Plant Specialized Metabolism Regulated By Jasmonate Signaling." *Plant And Cell Physiology* 60.12 (2019): 2638-2647.
- Cohen, S., & Nitzany, F. E., 1966. Transmission And Host Range Of Tomato Yellow Leaf Curl Virus. *Phytopathology*, 56 (10), <https://www.cabidigitallibrary.org/doi/full/10.5555/19690500321>
- Collard, B. C., & Mackill, D. J. (2008). Marker-Assisted Selection: An Approach For Precision Plant Breeding In The Twenty-First Century. *Philosophical Transactions Of The Royal Society B: Biological Sciences*, 363(1491), 557-572.
- Corrales-Gutierrez M, Medina-Puche L, Yu Y, Wang L, Ding X, Luna Ap, Bejarano Er, Castillo Ag, Lozano-Duran R. İkizler Virüsü Domates Sarı Yaprak Kıvrılma Virüsünden Elde Edilen C4 Proteini, Aba'dan Bağımsız Bir Mekanizma Aracılığıyla Arabidopsis'te Kuraklık Toleransı Sağlar. *Plant Biotechnol J.* 2020 Mayıs; 18(5):1121-1123. Doi: 10.1111/Pbi.13280. Epub 2019 23 Kasım. Pmid: 31637850; Pmcid: Pmc7152601.
- Czosnek, H., And Laterrot, H. 1997. A Worldwide Survey Of Tomato Yellow Leaf Curl Viruses. *Arch. Virol.* 142:1391-1406.
- Czosnek, H., Ber, R., Antignus, Y., Cohen, S., Navot, N., & Zamir, D. (1988). Isolation Of Tomato Yellow Leaf Curl A Geminivirus. *Phytopathology*, 78(5), 508 512. https://www.apsnet.org/Publications/Phytopathology/Backissues/Documents/1988articles/Phyto78n05_508.Pdf
- Çandar A, M Gümüş. Bitki Virüslerinin Vektörlerle Taşınmasına Moleküler Yaklaşımlar. *Türkiye Entomoloji Bülteni* 2 (3), 207-222, 2013. 5, 2013.
- Çetiner, S., & Tuzla, İ. (2005). Türkiye ve Dünyada Tarımsal Biyoteknoloji ve Gıda Güvencesi: Sorunlar Ve Öneriler.
- Çolak, A., Çelikel, G., Ekmekçi, U., Özarslandan, A., Aksoy, E., Karataş, A., ... & Öztürk, C. (2010). Doğu Akdeniz Bölgesi'nde Örtü Altı Organik Domates Üretiminde Hastalık, Zararlı ve Yabancı Otların Mücadelesinin Yönetimi. In *Organik Tarım Araştırma Sonuçları 2005-2010* (Pp. 253-264). Tc Tarım ve Köyişleri Bakanlığı.
- Demir, M. (2024). Afyonkarahisar İli Örtü Altı Domates Yetiştiriciliğinde Sorun Olan Bazı Virüslerin Enfeksiyonlarının ve Yaygınlıklarının Belirlenmesi.
- Devendran R, Kavalappara Sr, Simmons Am, Bag S. Whitefly-Transmitted Viruses Of Cucurbits In The Southern United States. *Viruses.* 2023 Nov 20;15(11):2278. Doi: 10.3390/V15112278. Pmid: 38005954; Pmcid: Pmc10675411.
- Ding T.B., Li J., Chen E.H., Niu J.Z., Chu D. Transcriptome Profiling Of The Whitefly *B. Tabaci* Med In Response To Single Infection Of Tomato Yellow Leaf Curl Virus, Tomato

- Chlorosis Virus, And Their Co-Infection. *Front. Physiol.* 2019;10:302. Doi: 10.3389/Fphys.2019.00302.
- Dumanoğlu, Zeynep, And Gülsüm Öztürk. "Patates (*Solanum Tuberosum* L.) Tarımı Ve Önemi." *Mas Journal Of Applied Sciences* 6.Özel Sayı (2021): 1307-1315.
- Elango, K.; Sobhana, E.; Sujithra, P.; Bharath, D.; Ahuja, A. Traditional Agricultural Practices As A Tool For Management Of Insects And Nematode Pests Of Crops: An Overview. *J Entomol Zool Stud.* 2020, 8, 237–245.
- El-Rahmany, R. G., Abdelhadi, A. A., El-Attar, A. K., El-Araby, W. S., & Abdallah, N. A. (2025). Genetic Diversity Of Tomato Yellow Leaf Curl Virus Isolates And Their Associated Beta-Satellite In Tomato Fields In Egypt.
- Eppo (2024) *Begomovirus Coheni*. Eppo Datasheets On Pests Recommended For Regulation. <https://gd.eppo.int> (Accessed 2025-02-28)
- Erdoğan, C., 2024. Türkiye’de ve Dünya’da Bitki Koruma Ürünlerinin Kullanımının Değerlendirilmesi ve Öneriler. Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi, 27(2): 382-392.
- Erdoğan, T., Çınar, C. T., Ve Ark., Işık, D. (2022). Domates Sarı Yaprak Kıvrıcıklık Virüsünün (Tomato Yellow Leaf Curl Virus-TYLCV) Doğal Konukçuları: Yabancı Otlar. *Turkish Journal Of Weed Science*, 25(2), 145-150.
- FAO (2022) (Food And Agriculture Organization Of The United Nations). (2022, September). Crops And Livestock Products. <https://www.fao.org/faostat/en/#data/qcl>
- FAO (2023). Food And Agriculture Organization Of The United Nations. Erişim Tarihi: 2025, Şubat 6. <https://www.fao.org/faostat/en/#data/qcl>
- Farzadfar S.H., A.R. Golnaraghi And R. Pourrahim (Ed.), 2002. Plant Viruses Of Iran. Saman Co. Tehran, Iran, 203 Pp.
- Farzadfar, Sh., Golnaraghi, A.R. And Pourrahim, R. 2002. Plant Viruses Of Iran. Saman Co. Tehran, Iran.
- Fauquet, C. M., Et Al. "Geminivirus Strain Demarcation And Nomenclature." *Archives Of Virology* 153 (2008): 783-821.
- Fidan H, Sarı N (2019) Molecular Characterization Of Resistance-Breaking Tomato Spotted Wilt Virus (Tswv) İsolatı Medium Segment In Tomato. *Applied Ecology And Environmental Research* 17(2): 5321-5339.
- Fidan, H., Ve Ark., Sarıkaya, P. (2020). Antalya İli Patlıcan (*Solanum Melongena*) Yetiştiriciliğinde Sorun Olan Virüs Hastalıkları. *Mediterranean Agricultural Sciences*, 33(1), 27-35.
- Fidan, Hakan, Et Al. "Investigation Of Activity Of Tobamovirus In Pepper Plants Containing L4 Resistance Gene." *Mediterranean Agricultural Sciences* 35.2 (2022): 83-90.
- Fondong, V. N. (2013). Geminivirus Protein Structure And Function. *Molecular Plant Pathology*, 14(6), 635-649. <https://doi.org/10.1111/mpp.12032>
- Freeman, B.C. & Beattie, G.A. (2008). An Overview Of Plant Defenses Against Pathogens And Herbivores. *The Plant Health Instructor*. <https://doi.org/10.1094/PhI-2008-0226-01>.
- Fu, H., Yang, X., Han, S., Wang, K., & Wang, H. (2025). Ganoderma Lucidum Polisakaritin Fonksiyonel Karakterizasyonu ve Domates Sarı Yaprak Kıvrılma Virüsüne Karşı Domates Direncini İndükleme Mekanizması. *Uluslararası Biyolojik Makromoleküller Dergisi*, 140617.

- Gebhardt, C. (2016). The Historical Role Of Species From The Solanaceae Plant Family In Genetic Research. *Theoretical And Applied Genetics*, 129, 2281-2294.
- Ghanim M, Czosnek H (2000). Tomato Yellow Leaf Curl Gemini Virus (TYLCV-Is) Is Transmitted Among Whiteflies (*Bemisia Tabaci*) In A Sex-Related Manner. *Journal Of Virology* 74:4738-4745.
- Ghosh, S.; Didi-Cohen, S.; Cna'ani, A.; Kontsedalov, S.; Lebedev, G.; Tzin, V.; Ghanim, M. Comparative Analysis Of Volatiles Emitted From Tomato And Pepper Plants In Response To Infection By Two Whitefly-Transmitted Persistent Viruses. *Insects* 2022, 13, 840. <https://doi.org/10.3390/insects13090840>.
- Gibson RW. Trapping of the spider mite *Tetranychus urticae* by glandular hairs on the wild potato *Solanum berthaultii*. *Potato Research*. 1976. Jun 1;19(2):179–82.
- Gilbertson, David T., Et Al. "Projecting The Number Of Patients With End-Stage Renal Disease In The United States To The Year 2015." *Journal Of The American Society Of Nephrology* 16.12 (2005): 3736-3741.
- Gill, U., Scott, J. W., Shekasteband, R., Ogundiwin, E., Schuit, C., Francis, D. M., Et Al. (2019). Ty-6, A Major Begomovirus Resistance Gene On Chromosome 10, Is Effective Against Tomato Yellow Leaf Curl Virus And Tomato Mottle Virus. *Theor. Appl. Genet.* 132, 1543–1554. Doi: 10.1007/s00122-019-03298-0
- Göçmen, H., & Devran, Z. (2002). Determination Of Genetic Variation In Populations Of *Bemisia Tabaci* In Antalya. *Turkish Journal Of Agriculture And Forestry*, 26(4), 211-216.
- Gönül, S., & Ülger, B. (2023, May). Congress Of Engineering And Natural Sciences. In *3 Th International Congress Of Engineering And Natural Sciences Studies* (P. 112).
- Grube, R. C., Radwanski, E. R., & Jahn, M. (2000). Comparative Genetics Of Disease Resistance Within The Solanaceae. *Genetics*, 155(2), 873-887.
- Gu, X.S.; Bu, W.J.; Xu, W.H.; Bai, Y.C.; Liu, B.M.; Liu, T.X. Population Suppression Of *Bemisia Tabaci* (Hemiptera: Aleyrodidae) Using Yellow Sticky Traps And *Eretmocerus Rajasthanicus* (Hymenoptera: Aphelinidae) On Tomato Plants In Greenhouses. *Insect Sci.* 2008, 15, 263–270.
- Guo, H., Huang, L., Sun, Y., Guo, H., & Ge, F. (2016). The Contrasting Effects Of Elevated Co2 On TYLCV Infection Of Tomato Genotypes With And Without The Resistance Gene, Mi-1.2. *Frontiers In Plant Science*, 7, 1680.
- Güleç, T. E., Yıldırım, A., & Sönmezoğlu, Ö. A. (2010). Bitkilerde Markör Destekli Seleksiyon. *Türk Bilimsel Derlemeler Dergisi*, (2), 67-79.
- H. El-Sappah, Ahmed, Et Al. "Natural Resistance Of Tomato Plants To Tomato Yellow Leaf Curl Virus." *Frontiers In Plant Science* 13 (2022): 1081549.
- Hajiabadi, A. M., Asaei, F., Abdollahi Mandoulakani, B. Ve Rastgou, M., 2012, Natural Incidence Of Tomato Viruses In The North Of Iran, *Phytopathologia Mediterranea*, 51 (2), 390-396.
- Hammad, E.A.; Nemer, N.M.; Hawi, Z.K.; Hanna, L.T. Responses Of The Sweetpotato Whitefly, *Bemisia Tabaci*, To The Chinaberry Tree (*Melia Azedarach* L.) And Its Extracts. *Ann. Appl. Biol.* 2000, 137, 79–88.
- Hanson, P. M., Bernacchi, D., Green, S., Tanksley, S. D., Muniyappa, V., Padmaja, A. S., ... & Chen, J. T. (2000). Mapping A Wild Tomato Introgression Associated With Tomato Yellow Leaf Curl Virus Resistance In A Cultivated Tomato Line. *Journal Of The American Society For Horticultural Science*, 125(1), 15-20.

- Hanson, S. F. (2022). Viral Diseases Of Tomato—Origins, Impact, And Future Prospects With A Focus On Tomato Spotted Wilt Virus And Tomato Yellow Leaf Curl Virus. In *Tomato-From Cultivation To Processing Technology*. Intechopen.
- Hassan, I., Orílio, A. F., Fiallo-Olivé, E., Briddon, R. W., & Navas-Castillo, J. (2016). Infectivity, Effects On Helper Viruses And Whitefly Transmission Of The Deltasatellites Associated With Sweepoviruses (Genus Begomovirus, Family Geminiviridae). *Scientific Reports*, 6(1), 30204.
- Hiharu Saito, Eiichi Makita, Suguru Yamane, Chihiro Urairi, Takayuki Hoshi, Makoto Doi, Suzuka Yoshizaki, Norihide Hinomoto, A New Pest Suction Machine To Control *Bemisia Tabaci* (Hemiptera: Aleyrodidae) In Tomato Greenhouses, *Journal Of Economic Entomology*, Volume 117, Issue 4, August 2024, Pages 1616–1622, <https://doi.org/10.1093/jee/toae097>.
- Horowitz, A.R.; Ghanim, M.; Roditakis, E.; Nauen, R.; Ishaaya, I. Insecticide Resistance And Its Management In *Bemisia Tabaci* Species. *J. Pest Sci.* 2020, 93, 893–910.
- Hosseinzadeh, M., And M. Garivani. "Emerging Two Distinct Groups Of The Tomato Yellow Leaf Curl Virus-Severe Strain (TYLCV-II) Variants In Iran." *Trakia Journal Of Sciences* 12.2 (2014).
- Huang, L.; Tang, Y.; Wang, S.; Chen, J.; Du, J.; Yan, S.; Zhang, D.; Şi, X.; Liu, Y.; Li, F. Dufulin, Domatesi Enfekte Eden Domates Sarı Yaprak Kıvrılma Virüsüne Karşı Bitki Savunmasını Etkiler. *Virüsler* 2025, 17, 53. <https://doi.org/10.3390/v17010053>.
- Hussein, H. S., Idriss, M. H., El-Gayar, F. H., Mousa, H. Y. S., & Salem, M. Z. (2025). Comparative Efficacy Of Plant Derived Extracts With The Insecticide Mospilan On Two Whitefly Species *Bemisia Tabaci* Biotype B And *Trialeurodes Ricini*. *Scientific Reports*, 15(1), 1970.
- ICTV (International Committee On Taxonomy Of Viruses). (2023, Semtember). Taxon Details, History Of The Taxon: Tomato Yellow Leaf Curl Virus. https://ictv.global/Taxonomy/Taxondetails?Taxnode_Id=202203484&Taxon_Name=Tomato%20yellow%20leaf%20curl%20virus
- ICTV (International Committee On Taxonomy Of Viruses). (2024, September). Current Ictv Taxonomy Release, Taxonomy Browser. <https://ictv.global/Taxonomy>
- İnak, Arda, Vd. "Türkiye'de Sebze Bitkilerinde *Bemisia Tabaci* Popülasyonlarının İnsektisit Direnç Durumu Ve Vektör Potansiyeli." *Bitki Koruma* 190 (2025): 107097.
- Jafari M., M. Valizadeh, J. Valizadeh, F. Ertiaei And M. Beigami, 2010. Detection Of İmportant Viruses İnfecting Cucurbit And Tomato In Fields And Greenhouses Of Baluchestan. Proceeding Of 19th Iranian Plant Protection Congress, 31 July–3 August, 2010, Tehran, Iran.
- Javaid, S.; Amin, I.; Jander, G.; Mukhtar, Z.; Saeed, N.A.; Mansoor, S. A Transgenic Approach To Control Hemipteran İnsects By Expressing İnsecticidal Genes Under Phloem-Specific Promoters. *Sci. Rep.* 2016, 6, 34706.
- Ji, Y., Schuster, D. J., & Scott, J. W. (2007). Ty-3, A Begomovirus Resistance Locus Near The Tomato Yellow Leaf Curl Virus Resistance Locus Ty-1 On Chromosome 6 Of Tomato. *Molecular Breeding*, 20, 271-284.
- Ji, Y., Scott, J. W., Schuster, D. J., & Maxwell, D. P. (2009). Molecular Mapping Of Ty-4, A New Tomato Yellow Leaf Curl Virus Resistance Locus On Chromosome 3 Of Tomato. *Journal Of The American Society For Horticultural Science*, 134(2), 281-288.
- Ji, Y., Scott, J.W., Hanson, P., Graham, E., And Maxwell, D.P. (2007b). "Sources Of Resistance, İnheritance, And Location Of Genetic Loci Conferring Resistance To Members Of

The Tomato-Infecting Begomoviruses,” In Tomato Yellow Leaf Curl Virus Disease, Ed H. Czosnek (Netherlands: Springer), 343–362.

Kahraman, A., & İlbi, H. (2024). Mas Yöntemiyle Domates Lekeli Solgunluk Virüsü'ne Dayanıklı Domates Hatlarının Geliştirilmesi. *Bahçe*, 53(Özel Sayı 1), 158-164.

Kahraman, Y. (2017). Patates (*Solanum Tuberosum L.*) Bitkisinin Verim ve Bazı Tarımsal Özellikleri Üzerine Farklı Sulama Rejimlerinin Etkisi (Master's Thesis, Ankara Üniversitesi (Turkey)).

Kaloo, G. (1993). Eggplant: *Solanum Melongena L.* *Genetic Improvement Of Vegetable Crops*, 587-604. Eggplant: *Solanum Melongena L.* - Sciencedirect

Kamal, H. , Zafar M. M., Razzaq A., Et Al. 2024. “Functional Role Of Geminivirus Encoded Proteins In The Host: Past And Present.” *Biotechnology Journal* 19: 2300736.

Karim, K.M.R.; Rafii, M.Y.; Misran, A.B.; Ismail, M.F.B.; Harun, A.R.; Khan, M.M.H.; Chowdhury, M.F.N. Current And Prospective Strategies In The Varietal Improvement Of Chilli (*Capsicum Annuum L.*) Specially Heterosis Breeding. *Agronomy* 2021, 11, 2217. <https://doi.org/10.3390/agronomy11112217>

Karkar, D.B. Evaluation Of Cow Urine And Vermi-Wash Against Insect Pests Of Brinjal. *Karnataka J. Agric Sci.* 2014, 27, 528–530.

Karthikeyan, C.; Veeraragavathatham, D.; Karpagam, Firdouse, A.A. Cow Based Indigenous Technologies In Dry Farming. *Indian J. Tradit Knowl.* 2006, 5, 47–50.

Karut, Kamil, Et Al. "Study On Species Composition Of Bemisia Tabaci (Gennadius, 1889)(Hemiptera: Aleyrodidae) On Cotton In Cukurova Plain, Turkey." *Turkish Journal Of Entomology* 38.1 (2014): 43-50.

Kashina, B. D., Mabagala, R. B., & Mpunami, A. A. (2002). Reservoir Weed Hosts Of Tomato Yellow Leaf Curl Begomovirus From Tanzania. *Archives Of Phytopathology And Plant Protection*, 35(4), 269-278.

Katsoulas, N., & Kittas, C. (2008). Impact Of Greenhouse Microclimate On Plant Growth And Development With Special Reference To The Solanaceae. *Eur. J. Plant Sci. Biotechnol.*, 2(1), 31-44.

Kenton, A., Parokonny, A. S., Gleba, Y. Y., & Bennett, M. D. (1993). Characterization Of The *Nicotiana Tabacum L.* Genome By Molecular Cytogenetics. *Molecular And General Genetics* 240, 159-169.

Khan, Muhammad Musa, Et Al. "Ultraviyole-A'nın Beyaz Sinek Bemisia Tabaci'ye Karşı Fototoksitesitesi Ve Entomopatojenik Bir Mantar Ve Beyaz Sinek Parazitoidi İle Uyumluluğu." *Oksidatif Tıp Ve Hücresel Uzun Ömür* 2021.1 (2021): 2060288.

Kıymacı, Gülbanu, Et Al. "Determination Of Resistance Levels Of Selected Tomato Genotypes To Meloidogyne Incognita, Tomato Yellow Leaf Curling Virus (TYLCV) Verticillium Wilt, Fusarium Oxysporum Radicis, Fusarium Wilt, Tomato Spotted Wilt Virus (Tswv)." *Selcuk Journal Of Agriculture And Food Sciences* 37.1 (2023): 86-94.

Kızmaz, M. Z. (2024). Domates Sarı Yaprak Kıvrıkcılık Virüsü (Tomato Yellow Leaf Curl Virus, TYLCV): Güneydoğu Anadolu Bölgesi Domates Üretim Alanlarında Önemi Artan Bir Virüs. *Düzce Üniversitesi Ziraat Fakültesi Dergisi*, 2(2), 115-125.

Kil, E. J., Park, J., Choi, E. Y., Byun, H. S., Lee, K. Y., An, C. G., ... & Lee, S. (2018). Seed Transmission Of Tomato Yellow Leaf Curl Virus In Sweet Pepper (*Capsicum Annuum*). *European Journal Of Plant Pathology*, 150, 759-764.

- Kil, E.J., Kim, S., Lee, Y.J., Byun, H.S., Park, J., Seo, H., Kim, C.S., Shim, J.K., Lee, J.H., Kim, J.K., Lee, K.Y., Choi, H.S., & Lee, S. (2016). Tomato Yellow Leaf Curl Virus (TYLCV-II): A Seed-Transmissible Geminivirus In Tomatoes. *Scientific Reports*, 6, 1-10. <https://doi.org/10.1038/srep19013>.
- Kim, Sanghyeon, Gyeongmo Gu ve Donghun Kim. "Kore Cumhuriyeti'nde Beyaz Sinek (*Bemisia Tabaci*) Kontrolü İçin Önerilen Sekiz Böcek İlacı Konsantrasyonlarının Değerlendirilmesi." *Asya-Pasifik Entomoloji Dergisi* 28.1 (2025): 102373.
- Koeda S, Kitawaki A. Breakdown Of Ty-1-Based Resistance To Tomato Yellow Leaf Curl Virus In Tomato Plants At High Temperatures. *Phytopathology*. 2024 Jan;114(1):294-303. Doi: 10.1094/Phyto-04-23-0119-R. Epub 2024 Feb 12. Pmid: 37321561.
- Kriticos, D. J., Et Al. "The Potential Geographical Distribution And Phenology Of *Bemisia Tabaci* Middle East/Asia Minor 1, Considering Irrigation And Glasshouse Production." *Bulletin Of Entomological Research* 110.5 (2020): 567-576.
- Kumar, M.; Kavalappara, S.R.; Mcavoy, T.; Hutton, S.; Simmons, A.M.; Bag, S. Association Of Tomato Chlorosis Virus Complicates The Management Of Tomato Yellow Leaf Curl Virus In Cultivated Tomato (*Solanum Lycopersicum*) In The Southern United States. *Horticulturae* 2023, 9, 948. <https://doi.org/10.3390/Horticulturae9080948>
- Lafrance, R.; Valdez-Torres, J.B.; Villicaña, C.; García-Estrada, R.S.; Esparza-Araiza, M.J.; León-Félix, J. Response Surface Methodology For Optimization Of Multiplex-Pcr Protocols For Detection Of TYLCV, Tswv And *Fol* Molecular Markers: Analytical Performance Evaluation. *Genes* 2023, 14, 337. <https://doi.org/10.3390/genes14020337>
- Lal, M. K., Tiwari, R. K., Kumar, A., Dey, A., Kumar, R., Kumar, D., ... & Singh, B. (2022). Mechanistic Concept Of Physiological, Biochemical, And Molecular Responses Of The Potato Crop To Heat And Drought Stress. *Plants*, 11(21), 2857.
- Lecoq, C. D. (1997). Zucchini Yellow Mosaic Virus. France.
- Leite, G.L.; Picanço, M.; Guedes, R.N.; Moreira, M.D. Factors Affecting Attack Rate Of Whitefly On The Eggplant. *Pesqui. Agropecuária Bras.* 2003, 38, 545–549.
- Levy, D., & Lapidot, M. (2008). Effect Of Plant Age At Inoculation On Expression Of Genetic Resistance To Tomato Yellow Leaf Curl Virus. *Archives Of Virology*, 153, 171-179. <https://doi.org/10.1007/S00705-007-1086-Y>.
- Li, Tong, Et Al. "Salicylic Acid-Induced Differential Resistance To The Tomato Yellow Leaf Curl Virus Among Resistant And Susceptible Tomato Cultivars." *Bmc Plant Biology* 19 (2019): 1-14.
- Li, J., Wang, J. C., Ding, T. B., & Chu, D. (2021). Synergistic Effects of a Tomato chlorosis virus and Tomato yellow leaf curl virus Mixed Infection on Host Tomato Plants and the Whitefly Vector. *Frontiers in Plant Science*, 12, 672400.
- Liedl BE, Lawson DM, White KK, Shapiro JA, Cohen DE, Carson WG, et al. Acylsugars of wild tomato *Lycopersicon pennellii* alters settling and reduces oviposition of *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Journal of Economic Entomology*. 1995. Jun 1;88(3):742–8.
- Long, N.; Cai, X.J.; Song, B.A.; Yang, S.; Chen, Z.; Bhadury, P.S.; Hu, D.Y.; Jin, L.H.; Xue, W. Synthesis And Antiviral Activities Of Cyanoacrylate Derivatives Containing An A-Aminophosphonate Moiety. *J. Agric. Food Chem.* 2008, 56, 5242–5246.
- Lu, Y.; Bei, Y.; Zhang, J. Are Yellow Sticky Traps An Effective Method For Control Of Sweetpotato Whitefly, *Bemisia Tabaci*, In The Greenhouse Or Field? *J. Insec. Sci.* 2012, 12, 113.

- Luo, C.; Jones, C.M.; Devine, G.; Zhang, F.; Denholm, I.; Gorman, K. Insecticide Resistance In *Bemisia Tabaci* Biotype Q (Hemiptera: Aleyrodidae) From China. *Crop Prot.* 2010, 29, 429–434.
- Mabvakure, B., Martin, D. P., Kraberger, S., Cloete, L., Van Brunschot, S., Geering, A. D., ... & Harkins, G. W. (2016). Ongoing Geographical Spread Of Tomato Yellow Leaf Curl Virus. *Virology*, 498, 257-264. <https://doi.org/10.1016/J.Virol.2016.08.033>.
- Manacorda Ca, Gudesblat G, Sutka M, Alemanno S, Peluso F, Oricchio P, Baroli I, Asurmendi S (2021) Tumor Triggers Stomatal Closure But Reduces Drought Tolerance In Arabidopsis. *Plant Cell Environ* 44(5):1399–1416. <https://doi.org/10.1111/pce.14024>
- Mandal, S., Padamshali, S., Rana, N., & Kolhekar, S. (2018). ITC Based Pest Management Module For Sucking Pest On Brinjal (*Solanum Melongena* L.) Under Terai Agro-Ecological System Of West Bengal. *J. Pharmacogn. Phytochem*, 7, 2065-2070.
- Marchant, W. G., Mugerwa, H., Gautam, S., Al-Aqeel, H., Polston, J. E., Rennberger, G., Smith, H., Turechek, B., Adkins, S., Brown, J. K., And Srinivasan, R. 2023. Phylogenomic And Population Genetics Analyses Of Extant Tomato Yellow Leaf Curl Virus Strains On A Global Scale. *Front. Virol.* 3.
- Massumi, H., Shaabani, M., Pour, A. H., Heydarnejad, J. And Rahimian, H. 2009. Incidence Of Viruses Infecting Tomato And Their Natural Hosts In The Southeast And Central Regions Of Iran. *Plant Dis.* 93:67-72.
- Meng Che, Gao Jing, Yang Yongqing, Fu Chongyi, Lian Yong, Li Zhengnan, Liu Yan, Zhang Lei. Detection Of Resistance To Three Viruses In 59 Tomato Germplasms By Molecular Markers[J]. *Journal Of Northern Agriculture*, 2024, 52(5): 29-37.
- Mirzayeva S, Huseynova I, Özmen Cy, Ergül A. Physiology And Gene Expression Analysis Of Tomato (*Solanum Lycopersicum* L.) Exposed To Combined-Virus And Drought Stresses. *Plant Pathol J.* 2023 Oct;39(5):466-485. Doi: 10.5423/Ppj.Oa.07.2023.0103. Epub 2023 Oct 1. Pmid: 37817493; Pmcid: Pmc10580053.
- Mishra, R., Shteinberg, M., Shkolnik, D., Anfoka, G., Czosnek, H., & Gorovits, R. (2022). Interplay Between Abiotic (Drought) And Biotic (Virus) Stresses In Tomato Plants. *Molecular Plant Pathology*, 23(4), 475-488.
- Missihoun, A. A., Fanou, A. A., Nanoukon, C. N. M., Agbo, I. R., Sedah, P., Fays, M., & Desoignies, N. (2025). Surveys Of Virus Diseases And Molecular Identification Of Viruses Affecting Pepper Crops (*Capsicum* Spp.) In Southern Benin. *Crop Protection*, 188, 106999.
- Montasser, M. S., Al-Own, F. D., Haneif, A. M., & Afzal, M. (2012). Effect Of Tomato Yellow Leaf Curl Bigeminivirus (TYLCV) Infection On Tomato Cell Ultrastructure And Physiology. *Canadian Journal Of Plant Pathology*, 34(1), 114-125.
- Moreau, T.L.; Isman, M.B. Trapping Whiteflies? A Comparison Of Greenhouse Whitefly (*Trialeurodes Vaporariorum*) Responses To Trap Crops And Yellow Sticky Traps. *Pest Manag. Sci.* 2011, 67, 408–413.
- Morris, J., (1997). A Multiplex Pcr Method For The Simultaneous Detection Of Tomato Yellow Leaf Curl And Tomato Mottle Geminiviruses. British Society For Plant Pathology Presidential Meeting. Plant Pathology-Global Perspectives Of An Applied Science. Offered Posters.
- Motti, R. (2021). The Solanaceae Family: Botanical Features And Diversity. *The Wild Solanums Genomes*, 1-9.
- Nair, I.J.; Sharma, S.; Shera, P.S. Impact Of Sticky Traps Of Different Colours And Shapes Against Sucking Pests Of Tomato Under Protected Conditions: A Randomized Controlled Trial. *Int. J. Trop Insect Sci.* 2021, 41, 2739–2746.

- Nikolić, D., Vučurović, A., Stanković, I., Radović, N., Zečević, K., Bulajić, A. Ve Krstić, B., 2018, Viruses Affecting Tomato Crops In Serbia, *European Journal Of Plant Pathology*, 152 (1), 225-235.
- Ning J, Moghe GD, Leong B, Kim J, Ofner I, Wang Z, et al. A feedback-insensitive isopropylmalate synthase affects acylsugar composition in cultivated and wild tomato. *Plant physiology*. 2015. Nov 1;169(3):1821–35. doi: 10.1104/pp.15.00474
- Okutucu, B., & Pehlivan, S. (2003). Reverz-Transkriptaz Polimeraz Zincir Reaksiyonu (RT-PCR) ve Uygulama Alanları. *Arşiv Kaynak Tarama Dergisi*, 12(2).
- Pakashtica, V. (2023). Eggplant Genome Editing Using Crispr Cas9 Technology.
- Papayiannis, L. C., Katis, N. I., Idris, A. M., & Brown, J. K. (2011). Identification Of Weed Hosts Of Tomato Yellow Leaf Curl Virus In Cyprus. *Plant Disease*, 95(2), 120-125. <https://doi.org/10.1094/pdis-05-10 0346>.
- Parlak, E. Ş., & Karalezli, A. (2020). Tütün Bitkisi Ve Ürünleri. *Göğüs Hastalıkları*, 21. Örtü Bitkisi Türleri Ve Kalıntı Yönetiminden Etkilenen Patlıcanın (*Solanum Melongena L.*) Verimi Ve Kalitesi - Sciencedirect
- Patel, N.B.; Korat, D.M.; Acharya, R.R. Impact Evaluation Of Cow-Urine And Vermiwash On Insect Pests Of Brinjal. *Int. J. Trop Agric*. 2017, 35, 591–595.
- Peñalver-Cruz, Ainara, Diego Alvarez, And Blas Lavandero. "Do Hedgerows Influence The Natural Biological Control Of Woolly Apple Aphids In Orchards?." *Journal Of Pest Science* 93.1 (2020): 219-234.
- Peng, Z.K.; Zheng, H.X.; Xie, W.; Wang, S.L.; Wu, Q.J.; Zhang, Y.J. Field Resistance Monitoring Of The Immature Stages Of The Whitefly *Bemisia Tabaci* To Spirotetramat In China. *Crop Prot*. 2017, 98, 243–247.
- Pereyra, J.G.; Martínez, G.N.; De Los Santos Villalobos, S.; Graciano, R.R.; Montelongo, A.M.; Roldan, H.M. Formulation Of A Bioinsecticide Based On Neem And Chamomile Used For The Greenhouse Control Of The Glasshouse Whitefly *Trialeurodes Vaporariorum*. *Mod. Environ. Sci. Eng*. 2021, 7, 119–125.
- Pérez-Padilla, V., Fortes, I. M., Romero-Rodríguez, B., Arroyo-Mateos, M., Castillo, A. G., Moyano, C., ... & Moriones, E. (2020). Revisiting Seed Transmission Of The Type Strain Of Tomato Yellow Leaf Curl Virus In Tomato Plants. *Phytopathology*, 110(1), 121-129.
- Pınar, H., Atilla, A., Keleş, D., Mutlu, N., Denli, N., & Mustafa, Ü. (2013a). Domates Hatlarında *Fusarium Oxysporum* F. Sp. *Lycopersici*'ye Dayanıklılığın Moleküler Markörler Yardımıyla Belirlenmesi. *Derim*, 30(1), 15-23.
- Pınar, H., Atilla, A., Keleş, D., Mutlu, N., Denli, N., & Mustafa, Ü. (2013b). Domateste Bazı Hastalık ve Zararlılara Dayanıklı Hat ve Çeşit Geliştirmede Moleküler Markörlerin Kullanımı. *Alatarım*, 12(1), 10-18.
- Picó, B., Díez, M. J., & Nuez, F. (1996). Viral Diseases Causing The Greatest Economic Losses To The Tomato Crop. I. The Tomato Yellow Leaf Curl Virus—A Review. *Scientia Horticulturae*, 67(3-4), 151-196.
- Polston, J. E., & Anderson, P. K. (1997). The Emergence Of Whitefly-Transmitted Geminiviruses In Tomato In The Western Hemisphere. *Plant Disease*, 81(12), 1358-1369.
- Polston, J. E., R. J. Mcgovern, And L. G. Brown. 1999. Introduction Of Tomato Yellow Leaf Curl Virus In Florida And Implications For The Spread Of This And Other Geminiviruses Of Tomato. *Plant Dis*. 83: 984–988.

- Pozharskiy, Alexandr Et Al.2022. Screening A Collection Of Local And Foreign Varieties Of *Solanum Lycopersicum* L. İn Kazakhstan For Genetic Markers Of Resistance Against Three Tomato Viruses Heliyon, Volume 8, Issue 8, E10095.
- Prasad, A., Sharma, N., Hari-Gowthem, G., Muthamilarasan, M., & Prasad, M. (2020). Tomato Yellow Leaf Curl Virus: İmpact, Challenges, And Management. Trends İn Plant Science, 25(9), 897-911. <https://doi.org/10.1016/J.Tplants.2020.03.015>.
- Prasanna, H. C., D. P., Rai, G.K, Krishna, R., Kashyap, S.P., Singh, N. K., & Malathi, V.G. (2015). Pyramiding Ty-2 ve Ty-3 Genes For Resistance To Monopartite And Bipartite Tomato Leaf Curl Viruses Of India. Plant Pathology, 64(2), 256-264.
- Prasanna, H. C., Sinha, D. P., Rai, G. K., Krishna, R., Kashyap, S. P., Singh, N. K., ... & Malathi, V. G. (2015). Pyramiding TY-2 And TY-3 Genes For Resistance To Monopartite And Bipartite Tomato Leaf Curl Viruses Of İ Ndia. Plant Pathology, 64(2), 256-264.
- Qian, X.; Lee, P.W.; Cao, S. China: Forward To The Green Pesticides Via A Basic Research Program. J. Agric. Food Chem. 2010, 58, 2613–2623.
- Qiao, N., Liu, H., Chen, Y., Zhang, D., Liu, J., Sun, H., ... & Sun, X. (2025). N Protein Of Tomato Spotted Wilt Virus Proven To Be Antagonistic Against Tomato Yellow Leaf Curl Virus İn Nicotiana Benthamiana. *Molecular Plant Pathology*, 26(1), E70046.
- Qiu, J.; Song, F.; Mao, L.; Tu, J.; Guan, X. Time-Dose-Mortality Data And Modeling For The Entomopathogenic Fungus. *Can. J. Microbiol.* 2013, 101, 97–101
- Rasul, I., Zafar, F., Ali, M. A., Nadeem, H., Siddique, M. H., Shahid, M., ... Ve Ark., Azeem, F. (2019). Genetic Basis For Biotic Stress Resistance İn Plants From Solanaceae Family: A Review.
- Reddy, B. J., Et Al. "A Review On Potato (*Solanum Tuberosum* L.) And İts Genetic Diversity." International Journal Of Genetics, Issn (2018): 0975-2862.
- Ren, Y., Tao, X., Li, D., Yang, X., & Zhou, X. (2022). Ty-5 Geminivirüslere Karşı Geniş Spektrumlu Direnç Sağlar. *Virüsler*, 14(8), 1804. <https://doi.org/10.3390/v14081804>.
- Riley, David G., And Rajagopalbabu Srinivasan. "Integrated Management Of Tomato Yellow Leaf Curl Virus And İts Whitefly Vector İn Tomato." Journal Of Economic Entomology 112.4 (2019): 1526-1540.
- Rosas-Diaz, T., D. Zhang, P. Fan, Et Al. 2018. "A Virus-Targeted Plant Receptor-Like Kinase Promotes Cell-To-Cell Spread Of RNAİ." Proceedings Of The National Academy Of Sciences Of The United States Of America 115: 1388–1393.
- Roy, Buddhadeb, Prosenjit Chakraborty, And Amalendu Ghosh. "How Many Begomovirus Copies Are Acquired And İnoculated By İts Vector, Whitefly (*Bemisia Tabaci*) During Feeding? " *Plos One* 16.10 (2021): E0258933.
- Russo, I., Cohen, S., Martelli, G. P., 1980. Virus-Like Particles İn Tomato Plants Affected By The Yellow Leaf Curl Disease. J. Gen. Virol. 49, 209–213.
- Sabra A., Emir M.A., Hüseyin K., Zakri A., El-Şahvan İ. M., El Salih M. A. Occurrence And Distribution Of Tomato Brown Rugose Fruit Virus Infecting Tomato Crop İn Saudi Arabia. *Plants*.2022;18:11:3157. Doi: 10.3390/Plants 11223157.
- Sabra A., Emir M.A., Hüseyin K., Zakri A., El-Şahvan İ. M., El Salih M. A. Occurrence And Distribution Of Tomato Brown Rugose Fruit Virus Infecting Tomato Crop İn Saudi Arabia. *Plants*.2022;18:11:3157. Doi: 10.3390/Plants 11223157.

- Sade, Dagan, Et Al. "Water Balance, Hormone Homeostasis, And Sugar Signaling Are All Involved In Tomato Resistance To Tomato Yellow Leaf Curl Virus." *Plant Physiology* 165.4 (2014): 1684-1697.
- Sánchez-Campos, S., Navas-Castillo, J., Camero, R., Soria, C., Díaz, J. A., And Moriones, E. 1999. Displacement Of Tomato Yellow Leaf Curl Virus (TYLCV)-Sr By TYLCV-Is In Tomato Epidemics In Spain. *Phytopathology* 89:1038-1043.
- Sanodiya, Bhagwan S., Et Al. "*Ganoderma Lucidum*: A Potent Pharmacological Macrofungus." *Current Pharmaceutical Biotechnology* 10.8 (2009): 717-742.
- Satar, Serdar, Et Al. "Loquat, A Non-Native Plant In Türkiye, Is A Key Seasonal Host In The Life Cycle Of Thrips *Hawaiiensis* (Morgan, 1913) (Thysanoptera: Thripidae)." *Phytoparasitica* 52.1 (2024): 30.
- Schillmiller A, Shi F, Kim J, Charbonneau AL, Holmes D, Daniel Jones A, et al. Mass spectrometry screening reveals widespread diversity in trichome specialized metabolites of tomato chromosomal substitution lines. *The Plant Journal*. 2010. May;62(3):391–403. doi: 10.1111/j.1365-313X.2010.04154.x
- Scholthof, K. B. , Adkins S., Czosnek H., Et Al. 2011. "Top 10 Plant Viruses In Molecular Plant Pathology." *Molecular Plant Pathology* 12: 938–954.
- Sertkaya, G. Ve Yılmaz, M., 2017, Hatay İli Örtüaltı Organik Domates Yetiştiriciliğinde Bazı Begomovirüslerin Enfeksiyon Oranları ile Doğal Taşınması ve Diğer Konukçularının Araştırılması, Mustafa Kemal Üniversitesi Ziraat Fakültesi Dergisi, 22 (1), 1–15. *Nal Of Phytopathology*, 158 (11 12), 797-805.
- Shailaja, B.; Patnaik, H.P.; Mukherjee, S.K. Assessment Of Botanicals Fermented In Cow Urine Alone And Along With Panchagavya Against Brinjal Shoot And Fruit Borer. *J. Eco-Friendly Agric.* 2012, 7, 24–28.
- Shteinberg, M., Mishra, R., Anfoka, G., Altaieb, M., Brotman, Y., Moshelion, M., ... & Czosnek, H. (2021). Tomato Yellow Leaf Curl Virus (TYLCV) Promotes Plant Tolerance To Drought. *Cells*, 10(11), 2875.
- Silva KF, Michereff-Filho M, Fonseca ME, Silva-Filho JG, Texeira AC, Moita AW, et al. Resistance to *Bemisia tabaci* biotype B of *Solanum pimpinellifolium* is associated with higher densities of type IV glandular trichomes and acylsugar accumulation. *Entomologia Experimentalis et Applicata*. 2014. Jun;151(3):218–30.
- Sofa, 2004. State Of The Food And Agriculture 2003- 2004. Food And Agriculture Organization Of The Nations. http://www.fao.org/waicent/faoinfo/economic/esa/en/pubs_sofa.html.
- Sseruwagi, P., Et Al. "Methods Of Surveying The Incidence And Severity Of Cassava Mosaic Disease And Whitefly Vector Populations On Cassava In Africa: A Review." *Virus Research* 100.1 (2004): 129-142.
- Şahin, G., Ve Ark., Taşlıgil, N. Türkiye’de Tütün (*Nicotiana Tabacum L.*) Yetiştiriciliğinin Tarihsel Gelişimi ve Coğrafi Dağılımı Le Developpement Historique Et La Dispersion Geographique De La Cultivation De Tabac En Turquie. *Doğu Coğrafya Dergisi*, 18(30).
- Şanlı, Arif, Tahsin Karadoğan, And Bekir Tosun. "Bazı Uçucu Yağların Patateste (*Solanum Tuberosum L.*) Bitki Aktivatörü Olarak Kullanım Olanaklarının Araştırılması." *Tarla Bitkileri Merkez Araştırma Enstitüsü Dergisi* 25.Özel Sayı-2 (2016): 30-35.
- Şen, B., Ve Ark., Güngör, S. (2019). Effects Of Climate Change Over Potato (*Solanum Tuberosum L.*) Production. *Proceedings E-Book*, 263.

- Taggar, G. K., And Ravinder Singh. "Evaluation Of Some Nonconventional Insecticides Against Whitefly *Bemisia Tabaci* İn Black Gram." *Indian Journal Of Entomology* 82.2 (2020): 294-297.
- Tan X., Zhao J., Zhou S. Occurrence And Control Of Viral Disease Of Tomato Over Summer. *Mod. Agric. Sci. Technol.* 2022; 21:118–121.
- Tanaka, Yoshiyuki, Et Al. "The Complete Plastid Genome Of *Capsicum Frutescens* And İts Phylogenic Relationships." *Tropical Agriculture And Development* 68.2 (2024): 45-48.
- Taşlıgil, Nuran, And Uzm Güven Şahin. "Ziraat Coğrafyası Açısından Marmara Bölgesi'nde Örtüaltı Yetiştiriciliği." *Marmara Sosyal Araştırmalar Dergisi* 6 (2014): 1-17.3634227
- Thongrit, D., Attathom, S., & Sutabutra, T. (1986). Tomato Yellow Leaf Curl Virus İn Thailand. *Plant Virus Diseases Of Horticultural Crops İn The Tropics And Subtropics*. Fftc Book Series, (33), 60-63. <https://Li01.Tci-Thaijo.Org/Index.Php/Anres/Article/Download/241478/164776/828483>
- Topakçı, N., & Güneş, S. (2006). Domates Sarı Yaprak Kıvrıcıklığı (Tomato Yellow Leaf Curl Begomovirus-TYLCV) Hastalığında Virüs Vektör İlişkisi. *Derim*, 23(2), 46-53.
- TÜİK, 2022-2023. Türkiye İstatistik Kurumu. Bitkisel Üretim İstatistikleri. <http://www.tuik.gov.tr/>. Erişim Tarihi 28.02.2025
- TÜİK, 2024. Türkiye İstatistik Kurumu. Bitkisel Üretim İstatistikleri. <http://www.tuik.gov.tr/>. Erişim Tarihi 28.02.2025
- Uçar, C., & Şensoy, S. (2022). Validity Control Of Markers Used İn Molecular Marker Assisted Selection İn Tomato. *Yuzuncu Yıl University Journal Of Agricultural Sciences*, 32(2), 300-309.
- Ulusoy, M.R., J.K. Brown & E. Bayhan, 2002. The 'B' Biotype Of *Bemisia Tabaci* Now Established İn Turkey. *European Studies Network On European Whitefly Newsletter*, 13
- Ürgen, G. 2014. Fethiye Yöresindeki Domates Seralarında Önemli Virüs Hastalıklarının Belirlenmesi. Yüksek Lisans Tezi, Süleyman Demirel Üniversitesi, Fen Bilimleri Enstitüsü, Muğla, 1-54.
- Venkataravanappa, V., Lakshminarayana Reddy, C. N., Swarnalatha, P., Mahesha, B., Rai, A. B., & Krishna Reddy, M. (2014). Association Of Tomato Leaf Curl Joydebpur Virus And A Betasatellite With Leaf Curl Disease Of Eggplant. *Phytoparasitica*, 42, 109-120.
- Vidavsky, F., & Czosnek, H. (1998). Tomato Breeding Lines Resistant And Tolerant To Tomato Yellow Leaf Curl Virus From *Lycopersicon* <https://doi.org/10.1094/Phyto.1998.88.9.910> İn *Hirsutum*. *Phytopathology*, 88(9), 910-914.
- Wang R., Zheng, H., Qu C., Wang Z., Kong, Z., Luo C., 2016. Lethal And Sublethal Effects Of A Novel Cis-Nitromethylene Neonicotinoid Insecticide, Cycloxaprid, On *Bemisia Tabaci*. *Crop Protection* 83 (2016) 15-19
- Wang, Deyuan, And Paul W. Bosland. "The Genes Of *Capsicum*." *Hortscience* 41.5 (2006): 1169-1187.
- Wang, R.; Fang, Y.; Che, W.; Zhang, Q.; Wang, J.; Luo, C. Metabolic Resistance İn Abamectin-Resistant *Bemisia Tabaci* Mediterranean From Northern China. *Toxins* 2022, 14, 424. <https://doi.org/10.3390/Toxins14070424>
- Wang, R.; Fang, Y.; Mu, C.Q.; Qu, C.; Li, F.Q.; Wang, Z.Y.; Luo, C. Baseline Susceptibility And Cross-Resistance Of Cycloxaprid, A Novel Cis-Nitromethylene Neonicotinoid Insecticide, İn *Bemisia Tabaci* Med From China. *Crop Prot.* 2018b, 110, 283–287.

- Wang, R.; Wang, J.D.; Che, W.N.; Fang, Y.; Luo, C. Baseline Susceptibility And Biochemical Mechanism Of Resistance To Flupyradifurone İn *Bemisia Tabaci*. *Crop Prot.* 2020, 132, 105132.
- Wang, R.; Wang, J.D.; Che, W.N.; Luo, C. First Report Of Field Resistance To Cyantraniliprole, A New Anthranilic Diamide İnsecticide, On *Bemisia Tabaci* Med İn China. *J. Integr. Agric.* 2018a, 17, 158–163.
- Wang, Z.Y.; Yan, H.F.; Yang, Y.H.; Wu, Y.D. Biotype And İnsecticide Resistance Status Of The Whitefly *Bemisia Tabaci* From China. *Pest Manag. Sci.* 2010, 66, 1360–1366
- Xu, P.; Chen, F.; Manna, J.P.; Feldman, T.; Sumner, L.W.; Roossinck, M.J. Virus İnfection İmproves Drought Tolerance. *New Phytol.* 2008, 180, 911–921.
- Xu, X., Pan, P., Cheng, S., Zhang, B., Mu, D., Ni, P., ... Ve Ark., Visser, R. G. (2011). Genome Sequence And Analysis Of The Tuber Crop Potato. *Genome Sequence And Analysis Of The Tuber Crop Potato*
- Xu, Ying, Et Al. "Domates Sarı Yaprak Kıvrılma Virüsüne Karşı Yeni Potansiyel Antiviral Ajan Olarak Arbutin'in Keşfi." *Tarım ve Gıda Kimyası Dergisi* (2025).
- Yan, Z.; Wolters, A.-M.A.; Navas-Castillo, J.; Bai, Y. Domates Sarı Yaprak Kıvrılması Hastalığının Küresel Boyutu: Mevcut Durum ve Üreme Perspektifleri. *Mikroorganizmalar* 2021, 9, 740. <https://doi.org/10.3390/microorganisms9040740>
- Yano, E. Control Of The Greenhouse Whitefly, *Trialeurodes Vaporariorum* Westwood (Homoptera: Aleyrodidae) By The İntegrated Use Of Yellow Sticky Traps And The Parasite *Encarsia Formosa* Gahan (Hymenoptra: Aphelinidae). *Appl. Entomol. Zool.* 1986, 22, 159–165.
- Yarar, A. (2023). *Crıspr/Cas9 Sistemi Kullanılarak Tütün Bitkisinde Agos7 Geninin Susturulması (Yüksek Lisans Tezi)*. Yök Tez Merkezi (832122)
- Yılmaz M.A., 1978. Tomato Yellow Leaf Curl Virus On Tomato. *Doğa*, 4: 248-250.
- Yılmaz, Hasan, Arzu Düzenli, And Merve Mürüvvet Dağ. "Dünya, Avrupa Birliği Ülkeleri ve Türkiye'de Pestisit Kullanımı ve Yasal Düzenlemeler." *Türkiye Tarımsal Araştırmalar Dergisi* 11.3 (2024): 315-330.
- Yılmaz, M.A., Özasan, M., Güldür, M.E., Baloğlu S., 1991. Domates Sarı Yaprak Kıvrıcılık Virüsünün (DSYKV) Purifikasyonu, Antiserumun Elde Edilmesi, DNA Ekstraksiyonu ve Elektrofrezisi. V1. Türkiye Fitopatoloji Kogresi, Türkiye Fitopatoloji Derneği Yayınları, No:6, İzmir: 365-369.
- Yuca, Hafize. "*Capsicum Annuum L.*" Novel Drug Targets With Traditional Herbal Medicines: Scientific And Clinical Evidence. Cham: Springer International Publishing, 2022. 95-108.
- Yüksekyayla, Yusuf, Mahmut Mete Karaca, And Kâmil Karut. "Antalya İli Örtüaltı Domates Üretim Alanlarında *Tuta Absoluta* (Meyrick) (Lepidoptera: Gelechiidae) Larva Parazitoitlerinin Belirlenmesi Üzerine Araştırmalar." *Türkiye Biyolojik Mücadele Dergisi* 14.2 (2023): 141-154.
- Zakay, Y., Navot, N., Zeidan, M., Kedar, N., Rabinowitch, H., Czosnek, H., & Zamir, D. (1991). Screening Lycopersicon Accessions For Resistance To Tomato Yellow Leaf Curl Virus: Presence Of Viral Dna And Symptom Development. *Plant Disease*, 75(3): 279-281. <https://doi.org/10.1094/Pd-75-0279>
- Zamir, D., Ekstein-Michelson, I., Zakay, Y., Navot, N., Zeidan, M., Sarfatti, M., ... & Czosnek, H. (1994). Mapping And İntrogression Of A Tomato Yellow Leaf Curl Virus Tolerance Gene, Ty-1. *Theoretical And Applied Genetics*, 88, 141-146.

Zengin, S., Kabas, A., & Hulya, I. L. B. I. (2020). Relationship Between Some Morphological Traits Of The Tomato Lines And Resistance To Tomato Yellow Leaf Curl Virus Disease. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 48(1), 388-397.

Zhang C, Wang D, Li Y, Wang Z, Wu Z, Zhang Q, Jia H, Dong X, Qi L, Shi J, Shang Z. Gibberellin Positively Regulates Tomato Resistance To Tomato Yellow Leaf Curl Virus (TYLCV). *Plants (Basel)*. 2024 May 6;13(9):1277. Doi: 10.3390/Plants13091277. Pmid: 38732492; Pmcid: Pmc11085062.

Zhang, H. , Gong H., And Zhou X.. 2009. “Molecular Characterization And Pathogenicity Of Tomato Yellow Leaf Curl Virus In China.” *Virus Genes* 39: 249–255.

Zhang, W.; Mcauslane, H.J.; Schuster, D.J. Repellency Of Ginger Oil To *Bemisia Argentifolii* (Homoptera: Aleyrodidae) On Tomato. *J. Econo. Entomol.* 2004, 97, 1310–1318.

Zheng, H.X.; Xie, W.; Wang, S.L.; Wu, Q.J.; Zhou, X.M.; Zhang, Y.J. Dynamic Monitoring (B Versus Q) And Further Resistance Status Of Q Type *Bemisia Tabaci* In China. *Crop Prot.* 2017, 94, 115–121.

EXAMINATION OF FARMERS' KNOWLEDGE AND AWARENESS ON THE ENVIRONMENTAL EFFECTS OF PESTICIDE USE: THE CASE OF IĞDIR PROVINCE

PESTİSİT KULLANIMININ ÇEVREYE ETKİLERİ KONUSUNDA ÇİFTÇİLERİN BİLGİ DÜZEYİ VE FARKINDALIKLARININ İNCELENMESİ: IĞDIR İLİ ÖRNEĞİ

Dr. Öğr. Üyesi Osman Doğan BULUT

Iğdır Üniversitesi, Ziraat Fakültesi, Tarım Ekonomisi Bölümü, Iğdır, Türkiye

ORCID: 0000-0003-2682-6356

Prof. Dr. Köksal KARADAŞ

Iğdır Üniversitesi, Ziraat Fakültesi, Tarım Ekonomisi Bölümü, Iğdır, Türkiye

ORCID: 0000-0003-1176-3313

ÖZET

Pestisit kullanımı, tarımsal üretimde verimliliği artırmak, zararlı organizmaların neden olduğu kayıpları en aza indirmek ve kaliteli ürün elde etmek amacıyla yaygın şekilde tercih edilen bir yöntemdir. Ancak, bilinçsiz ve yoğun pestisit kullanımı çevresel dengeleri tehdit eden ciddi riskler barındırmaktadır. Bu çalışmada, Iğdır ilinde tarımsal üretim yapan çiftçilerin pestisit kullanımı ve bu kullanımın çevresel etkilerine ilişkin bilgi düzeyleri ve farkındalıklarını belirlenmesi amaçlanmıştır. Araştırma kapsamında, 385 çiftçi ile yüz yüze görüşme yöntemiyle anket çalışması gerçekleştirilmiş ve elde edilen veriler yorumlanmıştır. Araştırma bulguları, çiftçilerin pestisitlerin ekosistem üzerindeki olumsuz etkilerine büyük ölçüde farkında olduklarını, ancak pestisit türleri, sınıflandırılması ve güvenli kullanımı konusunda yeterli bilgiye sahip olmadıklarını göstermektedir. Çiftçilerin pestisitlerin doğaya zarar verdiğini kabul etmelerine rağmen, pestisit etiket bilgilerini yeterince bilmediklerini ve bu bilgileri uygulama süreçlerinde etkin kullanmadıklarını ortaya koymaktadır. Özellikle, yasaklı pestisitlerin bilinmesi ve kullanımından kaçınılması konusunda çiftçilerin bilgi seviyesinin düşük olduğu görülmektedir. Ayrıca, pestisit ambalajlarının bertarafı konularında da bilinç eksiklikleri tespit edilmiştir. Pestisit kullanımının çevresel etkilerini azaltmaya yönelik bilinçlendirme faaliyetlerinin artırılması gerekmektedir. Tarımsal üretimde pestisit kullanımını daha sürdürülebilir hale getirmek için çiftçilere yönelik eğitim programlarının geliştirilmesi ve entegre zararlı yönetimi gibi alternatif mücadele yöntemlerinin teşvik edilmesi önem arz etmektedir. Bu doğrultuda, pestisitlerin bilinçli kullanımı çevrenin korunması ve sürdürülebilir tarımsal faaliyetin gerçekleştirilmesine katkı sağlayacaktır.

Anahtar Kelimeler: Pestisit kullanımı, Tarımsal üretim, Çevresel etkiler, Çiftçi farkındalığı, Sürdürülebilir tarım

ABSTRACT

Pesticide use is a widely preferred method to increase productivity in agricultural production, minimize losses caused by harmful organisms and obtain quality products. However, unconscious and intensive pesticide use poses serious risks that threaten environmental balances. This study aims to determine the knowledge and awareness levels of farmers engaged in agricultural production in Iğdır province regarding pesticide use and the environmental effects of this use. Within the scope of the research, a survey was conducted with 385 farmers using face-to-face interviews and the obtained data were interpreted. Within the scope of the research, a survey was conducted with 385 farmers through face-to-face interviews and the data obtained were interpreted. The research findings show that farmers are largely aware of the negative effects of pesticides on the ecosystem, but they do not have sufficient information about pesticide types, classification and safe use. Although farmers accept that pesticides harm nature, they do not know enough about pesticide label information and do not use this information effectively in the application processes. In particular, it is seen that farmers have a low level of knowledge about knowing about banned pesticides and avoiding their use. In addition, lack of awareness has been identified regarding the disposal of pesticide packaging. Awareness-raising activities to reduce the environmental impacts of pesticide use need to be increased. In order to make pesticide use in agricultural production more sustainable, it is important to develop training programs for farmers and encourage alternative control methods such as integrated pest management. In this context, conscious use of pesticides will contribute to the protection of the environment and the realization of sustainable agricultural activities.

Keywords: Pesticide use, Agricultural production, Environmental impacts, Farmer awareness, Sustainable agriculture

GİRİŞ

Günümüzde beslenme ihtiyacı dünya nüfusu ile doğru orantılı olarak artış göstermektedir. Gıda ihtiyacını karşılamak için bitkisel ürünlerin üretiminde bazı hastalıklar ile zararlılar ve yabancı otlardan dolayı üretim kayıplarının azaltılması arayışı doğmuştur. Bu sebeple üreticiler tarımsal alanlardaki verim ve kaliteyi arttırmak için farklı yöntemler kullanmaktadır. Bu yöntemlerden birisi de tarımsal mücadele adı altında yapılan kimyasal mücadele yöntemidir. Aslında bu yöntemin temelini pestisitlerin kullanımı oluşturmaktadır. Pestisitler, tarımsal üretimde hastalık, zararlı ve yabancı otların kontrolü amacıyla kullanılmaktadır. Kullanım nedenleri arasında ürün kayıplarını azaltmak, verimliliği artırmak ve kaliteli ürün elde etmek bulunmaktadır. Özellikle artan dünya nüfusunun gıda ihtiyacını karşılamak için pestisit kullanımı önemli bir araç haline gelmiştir.

Pestisitler, zararlı organizmaları öldürmek ve kontrol altına almak için kullanılan kimyasal maddelerdir. Kullanım amacına göre insektisit (böceklere karşı), herbisit (yabancı otlara karşı), fungusit (mantarlara karşı), bakterisit (bakterilere karşı), rodentisit (kemirgenlere karşı), akarisit (akarlara karşı), algisit (algilere karşı) olarak sınıflandırılmaktadır (Tiryaki, 2010; Akdoğan, 2012).

Dünya genelinde pestisit kullanımının bileşenleri incelendiğinde, herbisitler %40 ile en büyük paya sahip olup, tarımsal üretimde yabancı ot mücadelesinde yaygın olarak kullanılmaktadır. Bunu, %29 ile insektisitler takip etmekte ve bu grup, zararlı böceklerin kontrolünde etkili rol oynamaktadır. Fungusitler %22 oranında kullanılarak bitkilerde mantar hastalıklarını önlemede önemli bir yer tutmaktadır. Geriye kalan %9'luk dilim ise akarisitler, nematisitler ve rodentisitler gibi diğer pestisit türlerini içermektedir (EPA, 2011). Pestisitlerin çevresel etkileri geniş kapsamlı olup, ekosistem dengelerini ciddi şekilde tehdit etmektedir. Toprak ve su kirliliği, pestisitlerin yer altı ve yüzey sularına karışarak sucül ekosistemleri olumsuz

etkilemesine neden olmaktadır (Aktar, vd., 2009). Hedef dışı organizmalar, özellikle tozlaşmada kritik rol oynayan arılar ve yararlı böcekler üzerinde olumsuz etkilere yol açarak ekosistem hizmetlerinde kayıplara neden olabilir (Goulson, 2013). Hava kirliliği ve atmosferik taşınım, pestisitlerin buharlaşarak geniş alanlara yayılmasına ve hava kalitesinin bozulmasına sebep olmaktadır (Tang vd., 2021). Biyoçeşitlilik kaybı, pestisitlerin hassas ekosistemlerde tür çeşitliliğini azaltmasıyla sonuçlanmakta ve ekosistem fonksiyonlarını zayıflatmaktadır (Geiger vd., 2010).

Yoğun ve bilinçsiz pestisit kullanımı sonucunda, bu kimyasalların kalıntıları gıda maddeleri, toprak, su ve hava gibi çevresel bileşenlerde birikmektedir. Bu birikim, hedef dışı organizmalar ve insanlar üzerinde olumsuz etkilere yol açmaktadır. Özellikle, pestisitlerin püskürtülmesi sırasında bir kısmı buharlaşarak atmosfere karışmakta ve rüzgarlarla taşınarak geniş alanlara yayılmaktadır. Daha sonra yağmur, sis veya kar yağışıyla tekrar yeryüzüne inen bu kimyasallar, hedef dışı organizmalar ve bitkiler üzerinde kalıntı ve toksisiteye neden olabilmektedir. Ayrıca, pestisitlerin su kaynaklarına ulaşması, sucul ekosistemlerde yaşayan canlılar için ciddi tehditler oluşturmakta ve biyolojik çeşitliliği olumsuz etkilemektedir. Toprakta biriken pestisitler ise, toprak mikroorganizmalarının faaliyetlerini bozarak, toprak verimliliğini düşürebilmekte ve bitki gelişimini olumsuz yönde etkileyebilmektedir. Bu nedenle, pestisit kullanımının çevresel etkilerini en aza indirmek için entegre mücadele yöntemlerinin benimsenmesi ve kimyasal mücadeleye alternatif stratejilerin geliştirilmesi büyük önem taşımaktadır (Altıkat vd., 2009; Akdoğan vd., 2012).

Bu bağlamda, pestisitlerin çevre üzerindeki etkilerini anlamak ve bu konuda farkındalık oluşturmak hem bireysel hem de toplumsal düzeyde kritik bir öneme sahiptir. Bu araştırma; Iğdır ilinde çiftçilerin pestisit kullanımının çevre üzerine etkileri konusundaki deneyim ve düşüncelerini incelemek amaçlanmıştır.

MATERYAL VE METOT

Materyal

Iğdır ili, Türkiye'nin doğusunda, Aras Nehri'nin suladığı verimli Iğdır Ovası üzerinde yer almakta olup, Doğu Anadolu Bölgesi'ndeki önemli tarımsal üretim alanlarından biridir. İlin ekonomisi büyük ölçüde tarıma dayalıdır. Iğdır'da tarım arazileri, ilin toplam yüzölçümünün %33'ünü oluşturmaktadır. Bu arazilerin %81'i sulanabilir olup, %69'u aktif olarak sulanmaktadır. Ovada birçok sebzenin yetiştirilmesine karşılık dağlık yörede sadece patates tarımı yapılmaktadır. İlde mevcut olan 118,528 hektar dolayındaki tarım arazisinin yaklaşık 3,500 hektarı sebze üretimine ayrılmıştır. İlde yetiştirilen başlıca tarım ürünleri arasında şeker pancarı, karpuz, kavun, domates, kayısı ve elma bulunmaktadır. Özellikle kayısı ve elma ağaçları, ilin bahçelerinde yaygın olarak görülmektedir (SKA, 2013, TOB, 2022).



Şekil 1. Çalışma alanı

Iğdır ilinde tarımsal faaliyette bulunan 385 çiftçili ile yüz yüze görüşme yöntemiyle gerçekleştirilen anket çalışmasından elde edilen veriler araştırmanın ana materyalini oluşturmaktadır. Anket çalışması 2024 yılı Haziran-Temmuz ayları arasında yapılmıştır

1.1. Örnekleme Yöntemi

Araştırma evreni içerisinde amaca uygun herhangi bir yöntemle seçilen ve evreni temsil yeteneğine sahip birimler veya elemanlar kümesine örneklem denir (Ural, 2011: 23). Evren içerisinde evreni temsil eden birimleri seçme (örnek alma) işlemine ise örnekleme denir (Ural, 2011: 23). Araştırmanın evrenini Iğdır ilinde tarımsal faaliyet yapan çiftçiler oluşturmaktadır. Evrenden elde edilen örnek hacminin belirlenmesinde “Ana Kitle Oranlarına Dayalı Basit Tesadüfi Olasılık Örnekleme” yöntemi kullanılmıştır. Örneklem sayısı %95 güven sınırı ve ortalama %5 hata oranı ile 384.16 olarak hesaplanmış olup 385 olarak değerlendirmeye alınmıştır. Yamane (2001) tarafından geliştirilen örnekleme yönteminin formülü aşağıdaki gibidir (1):

$$n = \frac{z^2(p * q)}{e^2} \quad (1)$$

Formül açıklaması şu şekildedir:

n: Örnek hacmi

z: 1.96 (%95 güven düzeyine karşılık gelen standart z değeri)

e: Kabul edilen hata tolerans düzeyi (\pm %5 olarak kabul edilmiştir)

p: Örneğin ana kitleyi temsil edebilme olasılığı

q: (1-p) İlgili özelliğe sahip olmayan ana kitle oranı

Örneklem hacminin hesaplanması (2):

$$n = \frac{1.96^2(0.5 * 0.5)}{0.05^2} = 384.16 \quad (2)$$

Iğdır ilinde yapılan anket çalışmasına katılan çiftçilerin ilçelere göre dağılımı Tablo 1’ de gösterilmiştir. Çalışmada toplam 385 çiftçi ile görüşme sağlanmış olup, katılımcıların %32,01’i Iğdır merkez ilçesinde, %13,27’si Aralık ilçesinde, %14,75’i Karakoyunlu ilçesinde ve %39,98’i Tuzluca ilçesinde bulunmaktadır. İlçeler arasındaki farklar, tarım yapılan ilçedeki çiftçi sayısına bağlı olarak değişiklik göstermiştir.

Tablo 1. Anket yapılan çiftçilerin ilçelere göre dağılımı

İlçeler	Çiftçi Sayısı	Anket Sayı	Oran (%)
Merkez	6485	123	32.01
Aralık	2688	51	13.27
Karakoyunlu	2988	57	14.75
Tuzluca	8100	154	39.98
Total	20261	385	100.00

Analiz Yöntemi

Araştırmada, çiftçiler ile yapılan anket verilerine dayalı olarak elde edilen alan bulgularının analizinde; frekans dağılımı, yüzde ve aritmetik ortalama gibi tanımlayıcı istatistikler kullanılmıştır. Araştırmada kullanılan anketler; açık uçlu soru, iki seçenekli ve çok seçenekli sorularla birlikte 5’li Likert tipi sorulardan oluşmaktadır.

BULGULAR

Çiftçilerin Pestisit gruplarının anlam ve etiket rengini bilme durumu

Çiftçilerin pestisit gruplarının anlamlarını bilme durumu Tablo 2’de gösterilmiştir. Çiftçilerin büyük bir kısmının pestisitlerin spesifik kullanım alanları hakkında yeterli bilgiye sahip olmadığı görülmektedir. Örneğin, insektisitlerin (böcek ilaçları) anlamını bilen çiftçilerin oranı %25,5 iken, fungusitlerin (mantar ilaçları) anlamını bilenler %17,9, herbisitlerin (yabancı ot ilaçları) anlamını bilenler %15,1 ve nematisitlerin (nematod ilaçları) anlamını bilenler ise %10,6’dır. Bu sonuçlar, çiftçilerin pestisit kategorileri konusunda eğitim ihtiyacını ortaya koymaktadır.

Tablo 2. Pestisit grubunun anlamını bilme durumu

Pestisit Grubu	Anlamı	Bilmiyor		Biliyor		Toplam	
		n	%	n	%	n	%
İnsektisit	Böcek ilacı	287	74,5	98	25,5	385	100,0
Fungisitler	Mantar ilacı	316	82,1	69	17,9	385	100,0
Herbisit	Yabancı ot ilacı	327	84,9	58	15,1	385	100,0
Nematisit	Nematod ilacı	344	89,4	41	10,6	385	100,0

Çiftçilerin pestisit gruplarının etiket renklerini bilme durumu Tablo 3’de gösterilmiştir. İnsektisitlerin beyaz etiket rengine sahip olduğunu bilen çiftçilerin oranı %27,8, fungusitlerin açık yeşil olduğunu bilenlerin oranı %15,1, herbisitlerin sarı etiket rengine sahip olduğunu bilenlerin oranı %16,4 ve nematisitlerin pembe olduğunu bilenlerin oranı %9,4 olarak tespit edilmiştir. Genel olarak, çiftçilerin pestisit etiketleriyle ilgili bilgi düzeylerinin oldukça düşük olduğu ve bu konuda bilinçlendirme çalışmalarının artırılması gerektiği görülmektedir.

Tablo 3. Pestisit Grubunun Etiket Rengini Bilme Durumu

Pestisit Grubu	Etiket Rengi	Bilmiyor		Biliyor		Toplam	
		n	%	n	%	n	%
İnsektisit	Beyaz	278	72,2	107	27,8	385	100,0
Fungisitler	Açık Yeşil	326	84,9	58	15,1	385	100,0
Herbisit	Sarı	322	83,6	63	16,4	385	100,0
Nematisit	Pembe	349	90,6	36	9,4	385	100,0

Pestisit kullanımının çevreye verdiği zarar ile ilgili konulara çiftçilerin katılım düzeyi

Pestisit kullanımının çevreye olan olumsuz etkilerine çiftçilerin katılım düzeyleri Tablo 4’de gösterilmiştir. Ortalama değerlere bakıldığında, çiftçilerin pestisitlerin arıları ve yararlı böcekleri öldürdüğü fikrine yüksek düzeyde katıldıkları (4,27) görülmektedir. Yeraltı suları, göller ve akarsuların pestisitler nedeniyle kirlendiği düşüncesi de oldukça yüksek bir katılım düzeyine sahiptir (4,01). Kuşlar ve diğer hayvanlara zarar verdiği (3,99) ve toprağa zarar verdiği (3,80) görüşlerine de çiftçilerin önemli ölçüde katılım gösterdiği görülmektedir. Bu sonuçlar, çiftçilerin pestisitlerin çevresel etkilerinin farkında olduklarını ancak bu konuda çözüm odaklı yaklaşımlara dair bilinç düzeylerinin daha detaylı incelenmesi gerektiğini göstermektedir.

Tablo 4. Pestisit kullanımının çevreye verdiği zarar ile ilgili konulara çiftçilerin katılım düzeyi

Çevre Sorunları	Min.	Maks.	Ort.	Std. Sp.
Arıları ve yararlı böcekleri öldürür	1	5	4,27	0.763
Yeraltı suları, göller ve akarsularda kirlilik oluştur	1	5	4,01	0.942
Kuşlara ve diğer hayvanlara zarar verir	1	5	3,99	0.908
Toprağa zarar verir	1	5	3,80	1,063

1.2. Pestisit kullanımı ile ilgili diğer konulara çiftçilerin katılım düzeyleri

Çiftçilerin pestisit kullanımına yönelik diğer konulara katılım düzeyleri Tablo 5’de gösterilmiştir. Çiftçilerin pestisitleri ayrı bir alanda muhafaza etme konusunda oldukça bilinçlidir (4,08). Ayrıca, pestisitlerle ilgili bilgi kaynaklarının genellikle resmi kuruluşlar veya yetkili ilaç bayileri olduğu saptanmıştır (4,03). Ancak, çiftçilerin pestisit etiket talimatlarını takip etme (3,15) ve pestisit kabı imhası konusunda yeterli bilinç düzeyine sahip olmadıkları tespit edilmiştir (3,06). En düşük ortalama puan, çiftçilerin yasaklı pestisitleri bilme düzeyiyle ilgilidir (2,08), bu da yasaklı pestisit kullanımı konusunda daha fazla bilgilendirme ve denetim mekanizmalarının gerektiğini ortaya koymaktadır.

Tablo 5. Pestisit kullanımı ile ilgili diğer konulara çiftçilerin katılım düzeyleri

Konular	Min.	Maks.	Ort.	Std. Sp.
Kullanmakta olduğum/kullanacağım pestisiti ayrı bir alanda muhafaza ederim.	1	5	4.08	0.855
Pestisitlere ilişkin bilgi kaynağım resmi kuruluşlar veya yetkili ilaç bayileridir.	1	5	4.03	0.863
Pestisit kullanımı sonrası hijyenime dikkat ederim (elleri yıkamak, duş almak, kıyafet değiştirmek).	1	5	3.61	0.865
Pestisit uygulamasında koruyucu ekipman (maske, eldiven, su geçirmeyen giysiler vb.) kullanırım.	1	5	3.51	0.854
Pestisit uygulamasını etiket talimatlarına göre yaparım	1	5	3.15	1.042
Pestisit kabının imhası için çöpe atarım (talimatlara uygun).	1	5	3.06	1.024
İlaç etiketlerinde yazan uyarılara dikkat ederim (çevre-sağlık-muhafaza vd.)	1	5	2.86	1.132
Yasaklı pestisitleri kullanmam	1	5	2.69	1.150
Kullanımı yasaklı pestisitleri bilirim	1	5	2.08	1.057

SONUÇ

Iğdır ilinde tarımsal üretim yapan çiftçilerin pestisit kullanımı ve bunun çevresel etkilerine yönelik bilgi düzeyleri ve farkındalıkları bu çalışmada incelenmiştir. Elde edilen bulgular, çiftçilerin pestisitlerin ekosistem üzerindeki olumsuz etkilerine büyük ölçüde farkında olduklarını, ancak pestisitlerin sınıflandırılması, etiketlenmesi ve güvenli kullanımı konularında yeterli bilgiye sahip olmadıklarını göstermektedir.

Araştırma sonuçlarına göre, çiftçilerin büyük bir bölümü pestisitlerin türleri ve kullanım alanları hakkında sınırlı bilgiye sahiptir. Pestisitlerin sınıflandırılması ve etiket renkleri konusunda düşük bilgi düzeyleri, bilinçlendirme çalışmalarının ve eğitim programlarının artırılması gerekliliğini ortaya koymaktadır. Özellikle, “pestisitlerin arıları ve yararlı böcekleri öldürür” ifadesine katılım daha yüksek olmuştur (4,27). Çiftçilerin büyük bir kısmı pestisitlerin canlı organizma ve çevreye zarar vermesinin konusunda hemfikir olsa da, pestisitlerin güvenli kullanımı, bertarafı ile ilgili uygulamalarda eksiklikler olduğu saptanmıştır. Çalışmada, çiftçilerin pestisit uygulamalarında koruyucu ekipman kullanımına orta düzeyde önem verdikleri (3,61), pestisit uygulama talimatlarını tam olarak takip etmedikleri (3,51) belirlenmiştir. Ayrıca, yasaklı pestisitlerin bilinmesi (2.81) ve bu pestisitlerin kullanımından kaçınılması noktasında (2.69) bilgi eksikliği ve uygulama hatası olduğu tespit edilmiştir.

Pestisit kullanımının çevresel etkilerini azaltmaya yönelik eğitim ve bilinçlendirme çalışmalarının artırılması büyük önem taşımaktadır. Pestisitlerin güvenli kullanımını sağlamak adına tarımsal danışmanlık hizmetlerinin yaygınlaştırılması, çiftçilerin pestisitlerin sınıflandırılması ve etiket bilgileri konusunda bilgilendirilmesi ve sürdürülebilir tarım uygulamalarına yönlendirilmesi gerekmektedir. Ayrıca, entegre zararlı yönetimi (IPM) gibi alternatif mücadele yöntemlerinin yaygınlaştırılması, pestisit bağımlılığının azaltılmasına katkı sağlayabilir. Çiftçilerin bilinç düzeyinin artırılması, hem çevrenin korunması hem de insan sağlığı açısından kritik bir öneme sahiptir.

KAYNAKLAR

- Akdoğan, A., Divrikli, Ü., & Elçi, L. (2012). Pestisitlerin Önemi ve Ekosisteme Etkileri. *Akademik Gıda*, 10(1), 125-132.
- Aktar, M. W., Sengupta, D., & Chowdhury, A. (2009). Impact Of Pesticides Use In Agriculture: Their Benefits And Hazards. *Interdisciplinary Toxicology*, 2(1), 1-12. <https://doi.org/10.2478/v10102-009-0001-7>
- Altıkat, A., Turan, T., Ekmekyapar Torun, F., Bingül, Z. (2013). Türkiye’de Pestisit Kullanımı ve Çevreye Olan Etkileri. *Atatürk Üniversitesi Ziraat Fakültesi Dergisi*, 40(2), 87-92.
- EPA. (2011). Pesticides Industry Sales and Usage. EU Environmental Protection Agency. Erişim adresi https://www.epa.gov/sites/default/files/2015-10/documents/market_estimates2007.pdf?utm_source=chatgpt.com
- Geiger, F., Bengtsson, J., Berendse, F., Weisser, W. W., Emmerson, M., Morales, M. B., & Tschardtke, T. (2010). Persistent Negative Effects Of Pesticides On Biodiversity And Biological Control Potential On European Farmland. *Basic and Applied Ecology*, 11(2), 97-105. <https://doi.org/10.1016/j.baae.2009.12.001>
- Goulson, D. (2013). An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology*, 50(4), 977-987.
- SKA, (2013). Iğdır’ın Sosyo-Ekonomik Durumu. T.C. Serhat Kalkınma Ajansı. Erişim adresi https://www.kalkinmakutuphanesi.gov.tr/assets/upload/dosyalar/igdirin-sosyo-ekonomik-durumu.pdf?utm_source=chatgpt.com
- Tang, F. H. M., Lenzen, M., McBratney, A., & Maggi, F. (2021). Risk Of Pesticide Pollution At The Global Scale. *Nature Geoscience*, 14, 206-210. <https://doi.org/10.1038/s41561-021-00712-5>
- Tiryaki, O., Canhilal, R., & Horuz, S. (2010). Tarım İlaçları Kullanımı Ve Riskleri. *Erciyes Üniversitesi Fen Bilimleri Enstitüsü Fen Bilimleri Dergisi*, 26(2), 154-169.
- TOB, (2022). Iğdır İli. Iğdır İl Tarım Orman Müdürlüğü. Erişim adresi https://igdir.tarimorman.gov.tr/Menu/20/Ilimiz?utm_source=chatgpt.com

INVESTIGATION OF POLYPHENOL CONTENTS AND ANTIOXIDANT CAPACITIES OF SOME MONOFLORAL POLLENS FROM THE AEGEAN REGION

EGE BÖLGESİNE AİT BAZI MONOFLORAL POLENLERİN POLİFENOL İÇEKLERİ VE ANTIÖKSİDAN KAPASİTELERİNİN ARAŞTIRILMASI

Emine NAKİLCİOĞLU

Ege University, Faculty of Engineering, Department of Food Engineering, Izmir, Turkey

ORCID ID: <https://orcid.org/0000-0003-4334-2900>

Selin ALİHANOĞLU

Aegean Agricultural Research Institute, Beekeeping Research Center, Izmir, Turkey

ORCID ID: <https://orcid.org/0000-0003-1300-4628>

Gizem TİRYAKİ

Ege University, Institute of Science and Technology, Food Engineering, Izmir, Turkey

ORCID ID: <https://orcid.org/0000-0001-5608-9815>

ABSTRACT

Introduction and Purpose: Pollen, the microscopic structures that carry the male reproductive cells of flowering plants, plays a crucial role in pollination and the maintenance of genetic diversity. Pollen is considered a superfood due to its rich nutritional content, including proteins, essential amino acids, vitamins and minerals. Additionally, bioactive compounds such as flavonoids and unsaturated fatty acids in pollen exhibit antioxidant, anti-inflammatory, and anticancer properties, contributing to its positive health effects. In this study, the phenolic content and antioxidant capacities of six different monofloral pollen types collected from the Aegean region of Turkey and the differences between the species were evaluated. **Materials and Methods:** In this study, six different pollen types collected during the spring season from the Aegean region of Turkey were analyzed. The phenolic content was determined using the Folin-Ciocalteu method, while antioxidant capacity was measured using ABTS and DPPH assays. The statistical analysis of the obtained data was performed using one-way ANOVA and Duncan's test in the SPSS program ($p < 0.05$). **Results:** According to the analysis results, significant differences were observed among the pollen types in terms of phenolic content and antioxidant capacity. The highest phenolic content (1518,45 mg GAE/100g DM) and ABTS value (1132,27 mg Trolox/100g DM) were found in chestnut pollen. In the DPPH analysis, the highest antioxidant capacity was observed in pamucak pollen (92,35 mg trolox/100g DM). **Discussion and Conclusion:** The variability in phenolic content and antioxidant capacity among different pollen types is influenced by factors such as plant genetics, geographical origin, climatic conditions, and soil characteristics. The data obtained from this study indicate significant differences among pollen types. Particularly, chestnut pollen stands out with its high phenolic content and ABTS value, while pamucak pollen showed the highest value in the DPPH analysis. This study highlights the significant differences in the values of total phenolic content

and antioxidant capacity among different pollen types. Future studies will aim to expand the potential for use of pollen in functional foods by more comprehensively addressing the health effects of different pollen types.

Key Words: pollen, poppy, mustard, chestnut, oak, willow, pamucak

ÖZET

Giriş ve Amaç: Polen, çiçekli bitkilerin erkek üreme hücrelerini taşıyan mikroskobik yapılar olup, bitkilerin tozlaşması ve genetik çeşitliliğin sağlanmasında kritik bir rol oynamaktadır. Polen, besin değeri açısından protein, esansiyel amino asitler, vitaminler ve mineraller bakımından zengin besin içeriği ile süper gıda olarak kabul edilmektedir. Ayrıca polende bulunan flavonoid ve doymamış yağ asitleri gibi biyoaktif bileşikler antioksidan, antiinflamatuvar ve antikanserojen özellikler sergileyerek sağlık üzerinde olumlu etkiler göstermektedir. Bu çalışmada, Türkiye'nin Ege bölgesinden toplanan altı farklı monofloral polen türünün fenolik madde içeriği ve antioksidan kapasiteleri belirlenerek, türler arasındaki farklılıklar değerlendirilmiştir. **Gereç ve Yöntem:** Çalışmada, ilkbahar mevsiminde Türkiye'nin Ege bölgesinden toplanan altı farklı polen türü analiz edilmiştir. Fenolik madde içeriği, Folin-Ciocalteu yöntemi ile belirlenmiş; antioksidan kapasite ise ABTS ve DPPH yöntemleri kullanılarak ölçülmüştür. Elde edilen verilerin istatistiksel analizi SPSS programı kullanılarak tek yönlü varyans analizi ve Duncan testi ile gerçekleştirilmiştir ($p < 0.05$). **Bulgular:** Analiz sonuçlarına göre, polen türleri arasında fenolik madde içeriği ve antioksidan kapasite açısından önemli farklılıklar gözlemlenmiştir. En yüksek fenolik madde içeriği (1518,45 mg GAE/100g KM) ve ABTS değeri (1132,27 mg troloks/100g KM) kestane poleninde belirlenmiştir. DPPH analizine göre ise en yüksek antioksidan kapasiteye pamucak poleni (92,35 mg troloks/100g KM) sahiptir. **Tartışma ve Sonuç:** Farklı polen türlerinin fenolik madde içeriği ve antioksidan kapasitesindeki değişkenlik, bitkilerin genetik yapısı, yetiştiği coğrafi bölge, iklim koşulları ve toprak özellikleri gibi faktörlerden kaynaklanmaktadır. Çalışmadan elde edilen veriler, polen türleri arasında anlamlı farklılıklar olduğunu göstermektedir. Özellikle kestane poleni, yüksek fenolik madde içeriği ve ABTS değeri ile öne çıkarken, pamucak poleni DPPH analizinde en yüksek değeri göstermiştir. Bu çalışma, farklı polen türlerinin toplam fenolik madde miktarı ve antioksidan kapasite değerleri bakımından önemli farklılıklar sergilediğini ortaya koymuştur. Polen, besin içeriği ve biyolojik özellikleri sayesinde fonksiyonel gıda olarak değerlendirilebilecek değerli bir bileşendir. Gelecekte yapılacak çalışmalar, farklı polen türlerinin sağlık üzerindeki etkilerini daha kapsamlı şekilde ele alarak, polenin fonksiyonel gıdalardaki kullanım potansiyelini genişletmeyi hedefleyecektir.

Anahtar Kelimeler: polen, haşhaş, hardal, kestane, meşe, söğüt, pamucak

**THE IMPORTANCE OF AGRICULTURE IN THE SUSTAINABLE ECONOMIC
DEVELOPMENT OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC**

**NAXÇIVAN MUXTAR RESPUBLİKASININ DAVAMLI İQTİSADI İNKİŞAFINDA
KƏND TƏSƏRRÜFATININ ƏHƏMİYYƏTİ**

Cavadxan QASIMOV

Azərbaycan Respublikası, Naxçıvan Muxtar Respublikası, Naxçıvan Dövlət Universiteti,
İqtisadiyyat və idarəetmə fakültəsi, Beynəlxalq ticarət və menecment kafedrası, Naxçıvan
şəhəri.

ORCID: 0000-0002-3826-033X

Əsrəf ƏLƏKBƏROV

Azərbaycan Respublikası, Naxçıvan Muxtar Respublikası, Naxçıvan Dövlət Universiteti,
İqtisadiyyat və idarəetmə fakültəsi, İqtisadiyyat və marketing kafedrası, Naxçıvan şəhəri.

ORCID 0009-0006-8756-0685

ABSTRACT

Azərbaycan Respublikasının ayrılmaz tərkib hissəsi olaraq, Naxçıvan Muxtar Respublikasının iqtisadiyyatında əsas yeri kənd təsərrüfatı tutmaqdadır. Bu sahənin özünəməxsus xarakterik xüsusiyyətlərindən biri ondan ibarət olmaqdadır ki, muxtar respublika əhalisinin 70 faizdən artığı kənd yerlərində yaşayır və həmin sfera, makroiqtisadi inkişafın mühüm göstəricilərini təşkil etməkdədir. Hər şeydən əvvəl, kənd təsərrüfatı olduqca mühüm xammal bazası olaraq, çıxış edir və iqtisadiyyatın digər aparıcı sahəsi olan sənaye bölməsinin hərəkətverici qüvvəsi olaraq olduqca əhəmiyyətlidir. Naxçıvan Muxtar Respublikası iqtisadiyyatının inkişafı barəsində söz açarkən, ilk olaraq, onun ərazisinin ixtisaslaşması ilə ifadə oluna bilər. Diqqət etdikdə aydın olur ki, muxtar respublika ərazisinin ixtisaslaşması əsasən, dörd yüksəklik qurşağı üzrə xarakterik ola bilər. Onları, aşağıdakı kimi qeyd etmək olar.

- Arazboyu düzənlik qurşağı (600-1000-1200 m). Sahəsi, 1720 km² –dir. Muxtar respublika ərazisinin 32,1 faizini təşkil etməkdədir. Həmin ərazi, əsasən, tütün, bostan-tərəvəz, meyvə, üzüm, taxıl və yem bitkilərinin yetişdirilməsi üçün əlverişli olmaqdadır.

- Alçaq dağlıq qurşaq (1200-1800 m). Sahəsi, 1600 km² –dir. Muxtar respublika ərazisinin 29,8 faizini təşkil etməkdədir. Xüsusilə, tütün, meyvə, üzüm, tərəvəz, taxıl bitkilərinin və arıçılığın inkişafı üçün əlverişlidir.

- Orta dağlıq qurşaq (1800-2500 m). Sahəsi, 1523 km² –dir. Muxtar respublika ərazisinin 28,4 faizini təşkil etməkdədir. İxtisaslaşmasına görə, bağçılığın, ötlük-südlük heyvandarlığın və yem bitkilərinin yetişdirilməsi üçün xarakterikdir.

- Yüksək dağlıq qurşaq (2500-3000 m və ondan yüksək sahələr). Sahəsi, 523 km² –dir. Muxtar respublika ərazisinin 9,7 faizini təşkil etməkdədir. İxtisaslaşmasına baxdıqda, aydın olur ki, heyvandarlıqda yay otluqlarının təmayülü kimi istifadəyə əlverişlidir.

Bu ixtisaslaşma xüsusiyyətlərinə əsasən bildirmək olar ki, Naxçıvan Muxtar Respublikasının makroiqtisadi inkişafında ixtisaslaşmanın olduqca mühüm təərəfləri mövcud olmaqdadır və bu, eyni zamanda, əmək ehtiyatlarından səmərəli istifadə olunmasına əsaslı zəmin yaratmaqdadır.

Eyni zamanda, bu, aqrar sahədə məşğulluq probleminin həlli sahəsində mühüm nəticələrinə görə seçilməkdədir.

Açar sözlər: aqrar bölmə; kənd təsərrüfatı; makroiqtisadi inkişaf; ixtisaslaşma; yüksəklik qurşaqları və s.

ABSTRACT

As an integral part of the Republic of Azerbaijan, agriculture occupies a key place in the economy of the Nakhchivan Autonomous Republic. One of the unique characteristics of this sector is that more than 70 percent of the population of the autonomous republic lives in rural areas, and this sphere constitutes an important indicator of macroeconomic development. First of all, agriculture acts as a very important raw material base and is very important as a driving force for the industrial sector, which is another leading sector of the economy. When talking about the development of the economy of the Nakhchivan Autonomous Republic, it can first be expressed in terms of the specialization of its territory. When paying attention, it becomes clear that the specialization of the territory of the autonomous republic can be characterized mainly by four altitudinal zones. They can be noted as follows.

- Araz-river plain belt (600-1000-1200 m). Its area is 1720 km². It constitutes 32.1 percent of the territory of the autonomous republic. This area is mainly suitable for growing tobacco, melons and vegetables, fruits, grapes, grains and fodder crops.

- Low mountain belt (1200-1800 m). Its area is 1600 km². It constitutes 29.8 percent of the territory of the autonomous republic. In particular, it is favorable for the development of tobacco, fruit, grapes, vegetables, grain crops and beekeeping.

- Medium mountain belt (1800-2500 m). Its area is 1523 km². It constitutes 28.4 percent of the territory of the autonomous republic. According to its specialization, it is characteristic for the cultivation of horticulture, meat and dairy cattle breeding and fodder crops.

- High mountain belt (areas 2500-3000 m and higher). Its area is 523 km². It constitutes 9.7 percent of the territory of the autonomous republic. When looking at its specialization, it becomes clear that it is favorable for use in livestock breeding as a summer pasture.

Based on these specialization characteristics, it can be stated that there are very important aspects of specialization in the macroeconomic development of the Nakhchivan Autonomous Republic, and this, at the same time, creates a solid foundation for the efficient use of labor resources. At the same time, it is distinguished by its important results in solving the employment problem in the agricultural sector.

Key Words: agrarian sector; agriculture; macroeconomic development; specialization; altitudinal zones, etc.

SUSTAINABLE AGRICULTURE FOR OILSEED CROPS

YAĞLI TOHUM BİTKİLERİ İÇİN SÜRDÜRÜLEBİLİR TARIM

Aslı İŞLER KAYA

Turkish German University, Faculty of Science, Energy Science and Technology, Istanbul,
Türkiye

ORCID ID: <https://orcid.org/0000-0002-6552-8354>

ABSTRACT

Oilseed crops have become increasingly important due to population growth, the rising need for renewable resources, and three urgent issues: biodiversity loss, climate change, and environmental pollution. They provide renewable raw materials for edible/non-edible oil production as well as sustainable energy systems. They also serve as a vital biomass source with their seeds, straw, stalks, and roots used in various biobased applications and biofuel production, including biodiesel, bioethanol, sustainable aviation and marine fuels, and biochar. These plants play a crucial role in energy farming, making their harvest essential for the development of sustainable energy systems.

Agriculture plays a crucial role in meeting the global demand for crops, and as environmental sustainability concerns grow, understanding the ecological impact of agricultural practices has become increasingly important. To minimize environmental damage, maintain resources for future generations, and guarantee long-term feedstock security, sustainable agriculture is essential. In this context, Life Cycle Assessment (LCA) is a comprehensive and standardized tool for analyzing the environmental impacts throughout a crop's life cycle — from cultivation and harvesting to processing and final use. For the LCA study of the agricultural stage, water, electricity, and fuel use, tractor use, seed amount, and fertilizer amounts, are included covering the field emissions.

This study aims to give an overview of the environmental impacts of oilseed crop cultivation by evaluating the LCA methodology. By analyzing key factors such as carbon footprint, water and energy consumption, and environmental impacts, it is aimed to identify the carbon footprint intervals and offer insights for more sustainable agricultural practices. Adopting sustainable farming methods can help create a more resilient agricultural system by lowering greenhouse gas emissions, optimizing water use, and enhancing soil health. As a result, it is defined that the importance of crop cultivation in the shift to a low-carbon economy is becoming more widely acknowledged.

Key Words: Life Cycle Assessment, sustainable agriculture, oilseed crops, environmental impact assessment

ÖZET

Yağlı tohum bitkileri, nüfus artışı ve yenilenebilir kaynaklara ihtiyacın yükselmesi yanı sıra üç acil sorun olan biyoçeşitlilik kaybı, iklim değişikliği ve çevre kirliliğinin giderek önem kazanmasıyla birlikte, yenilebilir ve yenilmeyen yağ üretimi ve sürdürülebilir enerji sistemleri için yenilenebilir ham madde sağlama açısından daha da önem kazanmıştır. Ayrıca tohumları, sapları, samanları ve kökleri çeşitli biyoköklenli uygulamalar ile biyodizel, biyoetanol,

sürdürülebilir havacılık ve denizcilik yakıtları ile biyokömür gibi biyoyakıtlar üretiminde kullanılan önemli bir biyokütle kaynağıdır. Bu bitkiler, enerji tarımında da kritik bir yere sahip olarak sürdürülebilir enerji sistemlerinin gelişimi için hasat edilmeleri büyük önem taşır.

Tarım, küresel ürün talebini karşılamada kritik bir rol oynarken, çevresel sürdürülebilirlik konusundaki farkındalık arttıkça, tarımsal uygulamaların çevresel etkilerini anlamak giderek daha önemli hale gelmiştir. Çevresel zararları en aza indirmek, doğal kaynakları gelecek nesiller için korumak ve uzun vadeli hammadde güvenliğini sağlamak için sürdürülebilir tarım gereklidir. Bu bağlamda, Yaşam Döngüsü Değerlendirmesi (YDD), yağlı tohum bitkilerinin tüm yaşam döngüsü boyunca—yetiştirme ve hasattan işleme ve nihai kullanıma kadar—çevresel etkilerini analiz eden kapsamlı ve standartlaştırılmış bir araç olarak karşımıza çıkmaktadır. Tarımsal aşamanın Yaşam Döngüsü Değerlendirmesi çalışması kapsamında su, elektrik, yakıt ve traktör kullanımı, tohum miktarı ve gübre miktarları ile oluşan emisyonları kapsayan tüm girdiler dikkate alınmaktadır.

Bu çalışma, YDD yöntemi ile yağlı tohum bitkilerinin yetiştirilmesinin çevresel etkilerine genel bir bakış sunmayı amaçlamaktadır. Karbon ayak izi, su ve enerji tüketimi gibi temel faktörleri analiz ederek karbon ayak izi aralıklarını belirlemek ve daha sürdürülebilir tarımsal uygulamalar için hedefler sunmak amaçlanmaktadır. Sonuç olarak, sürdürülebilir tarım yöntemlerinin benimsenmesinin sera gazı emisyonlarının azaltımına, su kullanımının optimize edilmesine ve toprak sağlığını iyileştirerek daha dirençli bir tarım sisteminin oluşturulmasına yardımcı olacağı, düşük karbon ekonomisine geçişte tarımsal üretim aşamasının öneminin giderek daha fazla kabul gördüğü vurgulanmıştır.

Anahtar Kelimeler: Yaşam Döngüsü Değerlendirmesi, sürdürülebilir tarım, yağlı tohum bitkileri, çevresel etki değerlendirme

**RESEARCH AND GENE FUND COLLECTION OF STONE FRUIT PLANTS
CULTIVATED IN THE TERRITORY OF NAKHCHIVAN AUTONOMOUS
REPUBLIC**

UOT. 634.12 :631-53

Loğman BAYRAMOV

Ministry of Science and Education of the Republic of Azerbaijan Nakhchivan State
University

Ministry of Science and Education of the Republic of Azerbaijan Institute of Bioresources
(Nakhchivan)

ORCID ID: <https://orcid.org/0000-0002-1482-0048>

ABSTRACT

Introduction and purpose: The article identifies the distribution zones of cultivated pome fruits in the Nakhchivan Autonomous Republic, notes their names and synonyms, groups them according to ripening periods, and provides an accurate list of the pome fruit gene pool of the Nakhchivan Autonomous Republic. Until now, it has been noted that there are about 30 varieties of apples, 15 pears, and 4 varieties of animals in the Nakhchivan Autonomous Republic. During our research period, it was clarified that there are 118 varieties and forms of apples, more than 50 varieties and 11 forms of pears, and 12 varieties and 6 forms of animals in the autonomous republic, of which 98 are local selection varieties, 66 were previously introduced, and 33 forms were discovered for the first time. These varieties and forms are widely distributed in the Ordubad, Babek, Julfa, Shahbuz, Sharur, Sadarak, and Kangarli regions of the AR. In selection studies where a gene pool consisting of the mentioned varieties and forms has been established, these varieties and forms are used as pairs of the initial parent in the production of new varieties.

Keywords: Variety, form, apple, pear, quince, gene pool, selection, introduction

In modern times, the study and collection of the fruit gene pool is a matter of concern to all scientists. Extensive work is being carried out in this area, including in our republic. A number of researchers have extensively commented on the collection, evaluation and protection of genetic resources of the fruit gene pool in their works. Fruit growing in the Nakhchivan Autonomous Republic has had a special importance in agricultural fields since ancient times and has been of industrial importance. Among the fruit crops cultivated in all zones of the autonomous republic, stone fruits predominate. Stone fruits occupy 60-62% of the existing orchards [4, pp. 4-12; 9, pp. 198-223].

Unlike other regions of the Republic of Azerbaijan, the soil and climatic conditions of the Nakhchivan Autonomous Republic are very favorable for the rapid growth and development of fruit plants, as well as for the abundant and high-quality production. Along with other fruit plants, pome fruits are one of the ancient fruit plants of the Nakhchivan Autonomous Republic. Since ancient times, valuable forms of pome fruit plants have been created by folk breeders. These differ from the forms and varieties cultivated in other places in their taste, aroma, appearance, size of the fruit, and resistance of trees to diseases and pests. Many scientists have

determined that orchards in the Nakhchivan Autonomous Republic have an ancient history, including the cultivation of pome fruit plants from ancient times [1, pp. 155-156; 2, pp. 46-48].

Purpose of the study: Our goal is primarily to study the gene pool of pome fruit varieties and forms cultivated in the territory of the Nakhchivan Autonomous Republic, to accurately determine their distribution zones, to protect and restore rare local pome fruit varieties from the threat of extinction, to expand the gene pool collection garden, and to introduce these varieties to the world scientific community as Azerbaijani varieties. The goal was to protect aboriginal pome fruit varieties that are productive, high-quality, economically efficient, and resistant to diseases and pests, adapted to soil and climatic conditions, from the threat of extinction, to increase and select them, and to recommend them to farms.

Material and Methodology

Local and imported varieties and forms of pome fruits distributed in the territory of Nakhchivan Autonomous Republic were taken as the material of the research work. Their distribution zones, names and synonyms were specified, and varieties suitable for each zone, productive, high-quality, resistant to diseases and pests were selected and planted in the gene pool garden. The implementation of the research work was carried out in accordance with the generally accepted principles of fruit growing Hasanov Z.M. Fruit growing: Laboratory practical [3, p. 43-56], Sadigov A.N., Sadigov N.M. Apple plant in Azerbaijan [5, p. 40-135], Beideman I.N. Methodology for studying the phenology of plants and plant communities [8, p. 36-110], Methodology of the VNIIS im. Michurin [10, p. 94-156], Program and methodology for the introduction and variety study of fruit crops [11, p. 58-176], Rallov A. Kh. Sat. Information on fruit growing in the Transcaucasian region. [12, pp. 78-100]; etc. programs and methods were used.

Results and discussions: Up to now, about 30 varieties of apples, 15 pears and 4 varieties of animals have been noted in the territory of the Nakhchivan Autonomous Republic. During our research period, it was clarified that there are 118 varieties and forms of apples, more than 50 varieties and 11 forms of pears, 14 varieties of animals and 6 forms of animals, of which 98 are local selection varieties, 66 were previously introduced, and 33 forms were discovered for the first time. These varieties and forms are widely distributed in the Ordubad, Babek, Julfa, Shahbuz, Sharur, Sadarak and Kangarli regions of the Autonomous Republic [7, pp. 10-14; 6, pp. 20-80].

During the research, it was determined that a total of 158 varieties and 39 forms of stone fruits are distributed in the territory of Nakhchivan Autonomous Republic. The economic and agrobiological characteristics of these varieties and forms were studied, pomological descriptions of varieties and forms were given, and the varietal types to which the newly discovered forms belong were determined. A genetic fund of the discovered stone fruit varieties and forms was created in the "Botanical Garden" of the Institute of Bioresources. In order to enrich the genetic fund, new varieties were introduced and brought from the Islamic Republic of Iran, the Republic of Turkey and the Institute of Agricultural Research and Development of Azerbaijan (AKTN Fruit and Tea Plantation ET).

The pome fruit plants belong to the Rosaceae Adans. family, the subfamily Pomideae, and the genera *Malus* Mill, *Prunus* L, and *Judoniya* Mill. According to I.M. Akhundzade, N.M. Kurennoy, V.F. Koltunov, and V.I. Cherapakhina, A.N. Sadigov, N.M. Sadigov, and F.A. Pashayev, these genera include 50 species of apple, 60 species of pear, and 1 species of animal. In the world, 20,000 varieties of apple, 22,000 varieties of pear, and more than 250 varieties of animal have been registered. Of these, more than 325 varieties of apple, more than 470 varieties of pear, and more than 80 varieties of animal have been zoned for the fruit-growing regions of the CIS. Of the pome fruits, 9 species of apple are found wild in the former USSR, 19 species

of pear, and 1 species of animal. All cultivated pome fruits are derived from these lines [6, pp. 20-80]. The varieties of pome fruits are as follows:

From the varieties of whole fruits: Apple: - White apple, Borovinko, Dolma apple, Fakhima, Gem apple, Günüy Qishlag-3, Kăpăk apple, Kırmızı Grafstein, Girbi-shirin, Müşki, Majlisi, Milakh, Ordubad gozeli, Sultanı, Pampığı, Popirovka, Rajabi, Rashad, Shada-4, Shekeri Shikhmahmud-3, Unus-9, Yaylık apple, Zeynâddin-1; autumn varieties Antonovka, Anis, Aport, Belflor yellow, red, Jannet apple, Jafari, Arafsa-7 Andemij-5, Girde red, Güleman apple, Günüy Qishlag-1, Haji Huseyn, Heiwa apple, Khumar apple, Küku-3, Qandil sinab, Kol apple, Landsberg rennet, McIntosh, Mehdi jiri.

Pear - Summer pear, Nakhchivani, Abbasbesi, Gizili, Mehdi pear, Yellow Shekeri, Red Shekeri, Cir rare, Klap's favorite, Meleja, Kepeyi, Latanzi, Shekeri, Dırnisi, Zohra pear, Pomegranate pear, Ahmed gazi, Ardanpon, Williams, Duchess Angouleme, Goy pear, Nabi pear, Khoyi, Mikheyi, Sini pear, Latifa, Ordubadi, Gorkhmazy, Stone pear, Nasirin pear, Honey pear, Bildirjin budu, Bere-Bosk, Haji Mehdi, Yendija, Peyganbari, Al yanag, Khivili, Chil pear, Gunuy Qishlag-1, Chalkhangala-1, Oglangala-1, Milakh-2, Andemij-2

Animals are Sour quince, Yellow quince, Huseini, Novrest, Black quince, Armudvari, Pampigi, Stone quince, Sulu quince, Cardan quince, Karaman quince, Valachin, Dasharkh-1, Diza-1, Calkhangala-1, Zeynaddin-1, Asabu Kahf-1, Badamli Zavod-1, Bijenek-2.

According to the distribution of varieties and forms of pitted fruit in the autonomous republic, there are 75 varieties and forms of Alman in the Ordubad region, 72 varieties and forms in the Shahbuz region, 62 varieties and forms in the Babek region, 58 varieties and forms in the Julfa region, 27 varieties and forms in the Sharur region, 19 varieties and forms in the Kangarli region, and 12 varieties and forms in the Sadarak region;

There are 41 varieties and forms of pear in the Ordubad region, 39 varieties and forms in the Shahbuz region, 48 varieties and forms in the Babek region, 34 varieties and forms in the Julfa region, 22 varieties and forms in the Sharur region, 11 varieties and forms in the Kangarli region, and 9 varieties and forms in the Sadarak region;

9 varieties and 2 forms of the animal have been identified in the Ordubad region, 6 varieties and 2 forms in the Sharur Sadarak regions, 7 varieties and 4 forms in the Babek Kangarli regions, 8 varieties and 2 forms in the Shahbuz region, and 8 varieties and 3 forms in the Julfa region. Most of these varieties and forms have been propagated and planted in the gene pool collection garden.

During the years of research, expeditions were made to the villages of Ordubad, Shahbuz, Babek, Julfa, Sharur, Kangarli and Sadarak districts of the Autonomous Republic, and for the first time, 22 forms of apples, 11 forms of pears, and 8 forms of animals were discovered by us. 6 forms of apples were discovered in Ordubad district, 6 forms in Shahbuz district, 5 forms in Babek district, 3 forms in Sharur district, and 2 forms in Kangarli district.; 3 forms of pears were discovered in Ordubad district, 3 forms in Shahbuz district, 3 forms in Babek district, 1 form in Julfa district, and 1 form in Kangarli district.; 3 forms of animals were discovered in Sharur Sadarak district, 3 forms in Babek and Kangarli districts, and 2 forms in Shahbuz district.

The distribution zones, names and synonyms of local and imported pome fruit varieties and forms distributed in the territory of Nakhchivan Autonomous Republic were specified, and varieties suitable for each zone, productive, high-quality, resistant to diseases and pests were selected and planted in the gene pool collection garden. The planting year of newly introduced varieties began in 2005, and the planting scheme was 6x5- 3X4 m, depending on the varieties. The research work was carried out on the following pome fruit varieties and forms of apple: White apple, Dolma apple, Gem apple, Red Grafenstein, Girbishirin, Majlisi, Ordubad beauty, Shekeri, Red musk, Antonovka, Anis; Dust apple, Gandil-sinab, Rozmarin, Sini apple, Red

musk, Nakhchivan beauty, Rajabi varieties and Shada-4, Shikhmahmud-3, Güney-Gyshlag-4, etc. Forms.

Pear- Summer pear, Nakhchivani, Abbasbesi, Gizili, Mehdi pear, Yellow Shekeri, Red Shekeri, Cir rare, Klap's favorite, Meleja, Kepeyi, Latanizi, Shekeri, Dirnisi, Zohra pear, Pomegranate pear, Ahmed gazi, Ardanpon, Williams, Duchess Angouleme, Goy pear, Nabi pear, Khoyi, Mikheyi, Sini pear, Latifa, Ordubadi, Gorkhmazy, Stone pear, Nasirin pear, Honey pear, Bildirjin budu, Bere-Bosk, Haji Mehdi, Yendija, Peyganbari, Al yanag, Khivili, Chil pear, Gunuy Qishlag-1, Chalkhangala-1, Oglangala-1, Milakh-2, Andamij-2

Animal Animal Sour quince, Yellow quince, Huseini, Novrest, Black quince, Armudvari, Pampigi, The results were obtained on stone quince, water quince, cardan quince, garaman quince, valechin, dasharkh-1, diza-1, calkhangala-1, zeynaddin-1, asabu kahf-1, badamli zavod-1, bijanek-2, varieties and forms.

LITERATURE

1. Bayramov L.A. Study of apple varieties cultivated in the territory of the Nakhchivan Autonomous Republic and study of their gene pool // Azerbaijan Agricultural Science, 2006, No. 3-4, pp. 155-156
2. Bayramov L.A. Economic efficiency of apple varieties and forms cultivated in the territory of the Nakhchivan Autonomous Republic // Azerbaijan Agricultural Science, 2013, No. 2, pp. 46-48
3. Hasanov Z.M. Fruit growing (Laboratory-practicum). Baku: Bilik, 1977, 151 p.
4. Narimanov A.S. On the development of fruit growing in Azerbaijan. Baku, Azernashr, 1966, 31 p.
5. Sadigov A.N., Sadigov N.M. Apple plant in Azerbaijan. Baku, Sada, 2025, 174 p.
6. Talibov T.H., Bayramov L.A. Apple gene pool in Nakhchivan Autonomous Republic, Baku, 2013, "Afpoligrafia", 156 p.
7. Bayramov L.A. Evaluation of high-yielding native varieties and forms of apple trees cultivated in the territory of the Nakhchivansloi Autonomous Republic of Azerbaijan// Herald of the Altai State Agrarian University, №11 (133), November, 2015, p. 10-14
8. Beideman I.N. Methods of studying the phenology of plants and plant information. Novosibirsk: Nauka, 1974, 156 p.
9. Geevsky V.N, Sharrer G.N. Short essay on gardening in Transcaucasia. Proceedings of the Caucasian Society of Agricultural Society No. 12. Tiflis, 1985, 256 p.
10. Methodology VNIIS im. И.В. Michurina, 1973, p. 93-124
11. Program and methodology of introduction and sorting of fruit cultures. Chisinau: Shtiincha, 1972, p. 60-62
12. Rallov A. Kh. Sat. Information on fruit growing in the Transcaucasian region. Вып. 2. Tiflis, 1899, pp. 78-100

USAGE DIRECTIONS OF SPECIES BELONGING TO THE GENUS *ALCHEMILLA* L. IN THE FLORA OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC

Dashgin GANBAROV

Professor, doctor of sciences in biology ¹Nakhchivan State University, Faculty of natural science and agriculture, Department of Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0002-9819-5554>

Safura BABAYEVA

PhD in Biology Nakhchivan State University, Faculty of natural science and agriculture, Department of Biology, Nakhchivan, Azerbaijan

Orcid ID: <https://orcid.org/0009-0004-4800-7276>

ABSTRACT

Introduction and Purpose: It can be specifically noted that the species of the *Alchemilla* L. genus are significant plants among the useful plants of the flora of the Nakhchivan Autonomous Republic. Based on numerous studies and literature materials, it has been revealed that the species of this genus are used as medicinal, food, and fodder plants, as well as ornamental plants. Additionally, it has been learned that these species are also used in phytomeliorative measures. **Materials and Methods:** During the research, commonly accepted floristic, geobotanical, bioecological methods, and phenological observations were used. The main research materials were references to literature sources and actual data obtained during field studies, while various regions of the area were selected as the research subjects. The identification and clarification of the names of species belonging to the genus *Alchemilla* L. were based on works such as "Flora of Azerbaijan", "Flora of the Caucasus", A. M. Asgarov's "Flora of Azerbaijan" (Askerov 2016), and others. Recent taxonomic changes were made based on the World Flora Online. **Results:** A comparative analysis of the collected factual material and the information from literature sources has determined that 22 species of the genus are distributed in Azerbaijan, while 13 species are found in the flora of the Autonomous Republic. The research conducted by us allows for the identification of significant characteristics typical of the important species of the *Alchemilla* L. genus in the flora of the Nakhchivan Autonomous Republic. These species are found in most regions of the republic, particularly in mountainous areas, subalpine and alpine meadows, mountain steppe areas, stony, rocky places, as well as along mountain lakes and rivers. **Discussion and Conclusion:** It has been revealed through conducted research that the species of the *Alchemilla* L. genus in the flora of the Nakhchivan Autonomous Republic are of particular importance as medicinal, ornamental, fodder, and phytomeliorative plants

Key words: *Alchemilla* L., medicinal, phytotherapy, ornamental, fodder, species.

EFFECTS OF VERMICOMPOST APPLICATIONS ON YIELD PARAMETERS OF KAVILCA WHEAT

KAVILCA BUĞDAYINDA VERMİKOMPOST UYGULAMALARININ VERİM PARAMETRELERİ ÜZERİNE ETKİLERİ

Nihal Ceren ALICI

Kafkas University, Faculty of Science and Letters, Department of Biology, Kars, Türkiye

ORCID ID: <https://orcid.org/0000-0002-2147-0760>

Mehmet Ali KIRPIK

Kafkas University, Faculty of Science and Letters, Department of Biology, Kars, Türkiye

ORCID ID: <https://orcid.org/0000-0003-0156-8127>

ABSTRACT

Introduction and Purpose: Traditional farming practices often lead to low yield and quality problems, while the use of organic fertilization methods, such as vermicompost, offers solutions to these issues. Vermicompost is an effective organic fertilizer that improves soil health, supports plant growth, and promotes sustainable agriculture. The aim of this study is to investigate the effects of different doses of vermicompost on the yield parameters of geographically indicated Kavilca wheat (*Triticum dicoccon*) grown in Ardahan province. **Materials and Methods:** The study was conducted as a field trial between April and August 2024 in Hanak district of Ardahan province. The trial was designed according to a randomized block design with four replications for each application, consisting of 16 plots. No vermicompost was applied to the control group, while the other groups received 50 kg da⁻¹ (V₅₀), 100 kg da⁻¹ (V₁₀₀), and 150 kg da⁻¹ (V₁₅₀) vermicompost. Half of the vermicompost was applied as a basal fertilizer during sowing, while the other half was applied as a top-dressing fertilizer in two stages. **Results:** According to the results, the best outcomes were obtained in the 150 kg da⁻¹ vermicompost (V₁₅₀) application. In this group, the stem length was measured as 79.79 cm, spike length as 5.38 cm, plant height as 85.17 cm, the number of grains per spike as 12.12, grain weight as 25.15 g, stem thickness as 1.53 mm, and stem weight as 28.85 g. The total grain yield reached 501 kg da⁻¹. Compared to the control group, grain weight increased by 18%, stem length by 6%, and total grain yield by 38%. These results demonstrate that vermicompost application is an effective method to increase yield in Kavilca wheat cultivation. **Discussion and Conclusion:** The V₁₅₀ application provided the best results among the groups in terms of yield parameters for Kavilca wheat. Increases in spike length, stem thickness, and grain weight contributed to stronger and more resilient plants. The 38% increase in total grain yield confirmed the effectiveness of organic fertilization. Notably, the increase in stem weight can be considered an important source of forage for livestock in the region. This study indicates that the use of vermicompost makes a significant contribution to Kavilca wheat cultivation and should be promoted. The widespread adoption of organic fertilization methods among farmers will enhance agricultural productivity and support sustainable farming practices.

Key Words: Kavilca Wheat, Vermicompost, Organic Fertilizer, Sustainable Agriculture

ÖZET

Giriş ve Amaç: Geleneksel tarım uygulamaları, düşük verim ve kalite sorunlarına yol açarken, organik gübreleme yöntemlerinden biri olan vermikompost kullanımı bu sorunlara çözüm sunmaktadır. Vermikompost, bitki gelişimini destekleyerek toprak sağlığını iyileştiren ve sürdürülebilir tarımı teşvik eden etkili bir organik gübredir. Bu çalışmanın amacı, Ardahan ilinin coğrafi işaretli Kavılca buğdayının (*Triticum dicoccon*) yetiştiriciliğinde farklı dozlarda vermikompost uygulamalarının verim parametreleri üzerindeki etkilerini incelemektir. **Gereç ve Yöntem:** Çalışma, Nisan-Ağustos 2024 tarihleri arasında Ardahan ili Hanak ilçesinde tarla denemesi olarak yürütülmüştür. Deneme tesadüf blokları deneme desenine göre planlanmış, her bir uygulama için dört tekrarlı 16 parselden oluşmuştur. Kontrol grubuna vermikompost uygulanmazken, diğer gruplara 50 kg da⁻¹ (V₅₀), 100 kg da⁻¹ (V₁₀₀) ve 150 kg da⁻¹ (V₁₅₀) vermikompost uygulanmıştır. Vermikompostun yarısı ekim sırasında taban gübresi, diğer yarısı ise üst gübre olarak iki parçalı olarak verilmiştir. **Bulgular:** Deneme sonuçlarına göre en iyi sonuçlar 150 kg da⁻¹ vermikompost uygulamasında (V₁₅₀) elde edilmiştir. Bu grupta sap uzunluğu 79.79 cm, başak uzunluğu 5.38 cm, bitki boyu 85.17 cm, başaktaki tane sayısı 12.12 adet, tane ağırlığı 25.15 g, sap kalınlığı 1.53 mm ve sap ağırlığı 28.85 g olarak ölçülmüştür. Toplam tane verimi ise 501 kg da⁻¹'e ulaşmıştır. Kontrol grubuna kıyasla tane ağırlığında %18, sap uzunluğunda %6 ve toplam tane veriminde %38 artış gözlemlenmiştir. Bu sonuçlar, vermikompost uygulamasının Kavılca buğdayı yetiştiriciliğinde verimi artırmada oldukça etkili bir yöntem olduğunu ortaya koymaktadır. **Tartışma ve Sonuç:** V₁₅₀ uygulaması, Kavılca buğdayının verim parametrelerinde diğer gruplara göre en iyi sonuçları vermiştir. Başak uzunluğu, sap kalınlığı ve tane ağırlığındaki artışlar, bitkilerin daha güçlü ve dayanıklı olmasını sağlamış, toplam tane verimindeki %38'lik artış ise organik gübrelemenin etkinliğini kanıtlamıştır. Özellikle sap ağırlığındaki artış, bölgedeki hayvancılık faaliyetleri için önemli bir yem kaynağı olarak değerlendirilebilir. Bu çalışma, vermikompost kullanımının Kavılca buğdayı yetiştiriciliğinde önemli bir katkı sağladığını ve bu uygulamanın yaygınlaştırılması gerektiğini göstermektedir. Organik gübreleme yöntemlerinin çiftçiler arasında yaygınlaştırılması, tarımsal üretimde verimliliği artıracak ve sürdürülebilir tarımı destekleyecektir.

Anahtar Kelimeler: Kavılca Buğdayı, Vermikompost, Organik Gübre, Sürdürülebilir Tarım.

**BIOTECHNOLOGICAL CONTROL METHODS AGAINST VARROA
DESTRUCTOR IN HONEY BEES (*Apis mellifera*)**

**BAL ARILARINDA (*Apis mellifera*) VARROA DESTRUCTOR İLE
BİYOTEKNOLOJİK MÜCADELE YÖNTEMLERİ**

Öğr. Gör. Zeynep ASUTAY

Bitlis Eren University, Vocational School Of Hizan, Department of Plant and Animal
Production, Bitlis, Turkey

ORCID ID: <https://orcid.org/0000-0002-5854-1040>

Prof. Dr. Hakan İNCİ

Bingöl University, Faculty of Agriculture, Department of Animal Science,
Bingöl, Turkey

ORCID ID: <https://orcid.org/0000-0002-9791-0435>

ABSTRACT

Honey bees (*Apis mellifera*) play a crucial role in both ecosystems and agricultural production. However, one of the most significant parasites threatening colony health worldwide is *Varroa destructor*. This parasite severely affects the individual health of honey bees as well as colony integrity, leading to a decline in honey production and pollination services. Traditional chemical control methods are increasingly being replaced by biotechnological approaches due to concerns over resistance development, environmental impact, and negative effects on honey quality. This study examines biotechnological approaches used in the control of *Varroa destructor*, focusing on innovative methods such as genetic engineering, microbial biocontrol, biopolymer-based treatments, and RNA interference (RNAi) technology. Genetic engineering aims to develop honey bee strains resistant to *Varroa destructor*, while RNAi technology targets the suppression of specific genes in the parasite to inhibit its survival. Microbial biocontrol strategies involve the use of natural pathogens to reduce parasite loads. Additionally, the development of biopolymer-based treatments offers an environmentally friendly alternative to chemical acaricides. The effectiveness and sustainability of biotechnological control strategies are discussed, along with an evaluation of their field applications, economic feasibility, and potential ecological impacts. In conclusion, biotechnological approaches provide a promising alternative for the control of *Varroa destructor* and are expected to be more widely adopted in the apiculture sector in the future.

Key Words: *Apis mellifera*, *Varroa destructor*, *Biotechnology*, *Biocontrol*, *RNAi*, *Beekeeping*

ÖZET

Bal arıları (*Apis mellifera*), ekosistem ve tarımsal üretim açısından büyük öneme sahip olup, dünya çapında kolonilerinin sağlığını tehdit eden en büyük parazitlerden biri *Varroa destructor*'dur. Bu parazit, bal arılarının hem bireysel sağlığını hem de koloni bütünlüğünü ciddi şekilde etkileyerek bal verimliliğini ve polinasyon hizmetlerini azaltmaktadır. Geleneksel kimyasal mücadele yöntemleri, zamanla direnç gelişimi, çevresel etkiler ve bal kalitesine

olumsuz etkileri nedeniyle yerini biyoteknolojik yöntemlere bırakmaktadır. Bu çalışmada, *Varroa destructor* ile mücadelede kullanılan biyoteknolojik yaklaşımlar ele alınarak, genetik mühendislik, mikrobiyal biyokontrol, biyopolimer bazlı ilaçlar ve RNAi teknolojisi gibi yenilikçi yöntemler incelenmiştir. Genetik mühendislik sayesinde, *Varroa destructor*'a karşı dirençli bal arısı soylarının geliştirilmesi hedeflenirken, RNAi teknolojisi ile parazitin belirli genlerinin susturulması yoluyla mücadele edilmektedir. Mikrobiyal biyokontrol yöntemleri ise doğal patojenlerin kullanımıyla parazit yükünü azaltmayı amaçlamaktadır. Ayrıca, biyopolimer bazlı ilaçların geliştirilmesi, kimyasal akarisitlere çevre dostu bir alternatif sunmaktadır. Biyoteknolojik mücadele stratejilerinin etkinliği ve sürdürülebilirliği tartışılarak, bu yenilikçi yöntemlerin saha uygulamaları, ekonomik fizibilitesi ve potansiyel ekolojik etkileri değerlendirilmiştir. Sonuç olarak, biyoteknolojik yaklaşımların, *Varroa destructor* ile mücadelede önemli bir alternatif sunduğu ve gelecekte arıcılık sektöründe daha yaygın şekilde kullanılabileceği öngörülmektedir.

Anahtar Kelimeler: Apis mellifera, Varroa destructor, Biyoteknoloji, Biyokontrol, RNAi, Arıcılık

GİRİŞ

Bal arıları, bitki tozlaşmasında kritik bir rol oynayarak ekolojik dengenin korunmasına katkı sağlamaktadır. Dünya genelinde birçok tarım ürününün verimliliği, bal arılarının etkinliği ile doğrudan ilişkilidir. Ancak, son yıllarda arı popülasyonlarında ciddi düşüşler yaşanmakta olup, bu durum tarımsal üretim üzerinde olumsuz etkiler yaratmaktadır. Arı kolonilerinin sağlığını tehdit eden birçok faktör bulunmaktadır; bunlar arasında habitat kaybı, pestisit kullanımı, iklim değişikliği ve parazitler önemli bir yer tutmaktadır. Bu tehditler içinde, *Varroa destructor* en zararlı ve yaygın olanlardan biridir (Rosenkranz et al., 2010).

Varroa destructor, ilk olarak Asya bal arısı (*Apis cerana*) üzerinde tanımlanmış, ancak daha sonra Batı bal arısı (*Apis mellifera*) kolonilerine yayılmış ve burada ciddi kayıplara yol açmıştır. Parazit, arı larvaları ve ergin arılar üzerinde beslenerek gelişimlerini engellemekte, bağışıklık sistemlerini zayıflatmakta ve virüslerin yayılmasına neden olmaktadır (Martin, 2001). *Varroa* enfestasyonu kontrol altına alınmazsa, enfekte kolonilerin çöküşü kaçınılmaz hale gelir.

Bal arılarının kolonilerini etkileyen bu parazit, yalnızca bireysel arı sağlığı için değil, aynı zamanda küresel gıda güvenliği açısından da büyük bir tehdit oluşturmaktadır. Bal arıları, tarım ürünlerinin yaklaşık üçte birinin tozlaşmasında rol oynadığından, *Varroa destructor*'un yayılması tarımsal verimi azaltarak ekonomik kayıplara yol açmaktadır (vanEngelsdorp ve Meixner, 2010). Bu nedenle, arıcılık sektöründe sürdürülebilir ve etkili mücadele yöntemlerinin geliştirilmesi büyük önem taşımaktadır.

Geleneksel mücadele yöntemleri arasında sentetik akarisitler, organik asitler ve esansiyel yağlar bulunmaktadır. Ancak, kimyasal yöntemlerin sürekli kullanımı, parazitlerde direnç gelişimine ve arı ürünlerinde kalıntı oluşumuna neden olmaktadır (Carreck et al., 2013). Ayrıca, sentetik ilaçların aşırı kullanımı, hem bal arılarının genel sağlığını olumsuz etkileyebilmekte hem de bal ve diğer arı ürünlerinin tüketici sağlığı açısından güvenilirliğini düşürebilmektedir. Bu durum, daha çevre dostu ve sürdürülebilir mücadele yöntemlerinin geliştirilmesini zorunlu hale getirmiştir. Günümüzde biyoteknolojik yöntemler, *Varroa destructor* ile mücadelede yenilikçi ve etkili alternatifler sunmaktadır.

Son yıllarda, biyoteknolojik yöntemlerin kullanımıyla ilgili araştırmalar artış göstermiştir. Genetik mühendisliği, RNA girişimi (RNAi), mikrobiyal biyokontrol ve biyopolimer bazlı uygulamalar, *Varroa destructor* ile mücadelede umut verici stratejiler arasında yer almaktadır. Bu yöntemler, hem çevresel açıdan sürdürülebilir hem de arı sağlığına zarar vermeden parazitleri hedef alabilen çözümler sunmaktadır. Bununla birlikte, bu tekniklerin geniş ölçekli

uygulanabilirliği ve ticari kullanımı için daha fazla araştırmaya ve düzenlemelere ihtiyaç duyulmaktadır.

Bu çalışmada, *Varroa destructor*'a karşı geliştirilen biyoteknolojik mücadele yöntemleri ele alınarak, genetik mühendislik, RNA girişimi (RNAi), mikrobiyal biyokontrol ve biyopolimer bazlı uygulamaların etkinliği değerlendirilmiştir. Ayrıca, bu yöntemlerin pratik uygulamalardaki avantajları ve olası zorlukları tartışılmıştır.

VARROA DESTRUCTOR'UN BAL ARILARI ÜZERİNDEKİ ETKİLERİ

Varroa destructor, hem larva hem de ergin arılar üzerinde beslenerek kolonilerin zayıflamasına ve viral hastalıkların yayılmasına neden olmaktadır. Parazitin neden olduğu başlıca zararlar şunlardır (Martin, 2001):

- Fiziksel Hasar: Yetişkin arıların vücut sıvılarından beslenerek bağışıklık sistemlerini zayıflatır.
- Virüs Taşıyıcılığı: Özellikle Deformed Wing Virus (DWV) gibi ölümcül virüslerin yayılmasına neden olur.
- Koloni Çöküşü: Yüksek enfestasyon oranları, arı kolonilerinin ani çöküşüne yol açabilir.

Varroa destructor, arıların bağışıklık sistemini baskılayarak çeşitli patojenlere karşı daha savunmasız hale gelmelerine neden olmaktadır. Parazit, özellikle larva ve pupalar üzerinde beslenerek arıların gelişimini doğrudan etkiler. Bu süreç, yetişkin bireylerin fiziksel ve fizyolojik olarak zayıf olmasına yol açarak koloni genelinde üretkenliği azaltır (Nazzi ve Le Conte, 2016). Ayrıca, parazit tarafından zayıflatılan arılar, dış çevresel stres faktörlerine karşı daha duyarlı hale gelir ve koloninin uzun vadede sürdürülebilirliği tehlikeye girer.

Araştırmalar, *Varroa destructor*'un enfekte ettiği kolonilerde, bağışıklık sisteminin baskılanması sonucu sekonder enfeksiyonların yaygınlaştığını göstermektedir. Özellikle *Nosema ceranae* gibi bağırsak parazitleri ile birlikte görüldüğünde, kolonilerde daha hızlı çöküşler yaşanabilmektedir (Huang et al., 2012). Bununla birlikte, *Varroa destructor*'un taşıdığı ve yaydığı virüsler, arıların davranışsal ve nörolojik fonksiyonlarını da olumsuz yönde etkilemektedir. Örneğin, Israeli Acute Paralysis Virus (IAPV) ve Chronic Bee Paralysis Virus (CBPV) gibi virüsler, enfekte arıların uçuş yeteneklerini kısıtlamakta ve sosyal etkileşimlerini azaltarak koloninin işleyişini bozmaktadır (Di Prisco et al., 2016).

Koloni içindeki sosyal etkileşimin bozulması, larva bakımından nektar ve polen toplanmasına kadar pek çok kritik süreci aksatmaktadır. Bu durum, koloninin gıda stoklarının azalmasına, bireysel arıların ömürlerinin kısalmasına ve genel verimliliğin düşmesine sebep olmaktadır (Guzmán-Novoa et al., 2010). *Varroa destructor* enfestasyonunun ciddi seviyelere ulaşması, koloninin tamamen çökmesine ve arıların kovani terk etmesine yol açabilmektedir.

Bunun yanı sıra, *Varroa destructor*'un çevresel stres faktörleri ile birleşerek arı popülasyonlarında dramatik kayıplara yol açtığı gözlemlenmiştir. Pestisit maruziyeti, besin kıtlığı ve iklim değişikliği gibi faktörler, *Varroa* kaynaklı stresin etkilerini daha da artırmaktadır (vanEngelsdorp ve Meixner, 2010). Özellikle pestisit kalıntılarının bulunduğu ortamlarda yaşayan arı kolonilerinde, *Varroa destructor* ile mücadele eden arıların daha hızlı çöküş yaşadığı belirlenmiştir (Boncristiani et al., 2012).

Sonuç olarak, *Varroa destructor*'un bal arıları üzerindeki olumsuz etkileri çok yönlüdür ve sadece bireysel arı sağlığını değil, koloninin bütünsel işleyişini de tehdit etmektedir. Bu nedenle, bu parazit ile etkin mücadele stratejilerinin geliştirilmesi, bal arılarının sürdürülebilirliği açısından büyük önem taşımaktadır.

VARROA DESTRUCTOR İLE MÜCADELEDE BİYOTEKNOLOJİK YAKLAŞIMLAR

Geleneksel kimyasal akarisitler, Varroa destructor ile mücadelede uzun yıllardır kullanılmaktadır. Ancak, bu kimyasalların arılar ve çevre üzerindeki olumsuz etkileri, biyoteknolojik yöntemlerin geliştirilmesini teşvik etmiştir (Pettis et al., 2016). Günümüzde biyoteknolojik yaklaşımlar, genetik mühendislik, mikrobiyal biyokontrol, biyopolimer bazlı ilaçlar ve RNA girişimi (RNAi) gibi çeşitli alanları kapsamaktadır. Bu yöntemler, hedef spesifik etkileri ve sürdürülebilir olmaları nedeniyle giderek daha fazla tercih edilmektedir.

RNA Girişimi (RNAi) Teknolojisi

RNA girişimi (RNA interference), gen ekspresyonunu kontrol ederek Varroa destructor'un hayati fonksiyonlarını hedef alan bir biyoteknolojik mücadele yöntemidir (Garbian et al., 2012). RNAi yöntemi, parazitlerde belirli genlerin susturulmasını sağlayarak hayatta kalmalarını zorlaştırmaktadır. Yapılan araştırmalar, Varroa destructor'un sinirsel ve bağışıklık sistemleri ile ilişkili bazı genlerinin RNAi aracılığıyla susturulmasının parazitin ölüm oranını önemli ölçüde artırdığını göstermektedir (Campbell et al., 2016). Bu yöntem, pestisit kullanımını en aza indirerek çevre dostu bir alternatif sunmaktadır.

Mikrobiyal Biyokontrol Yöntemleri

Mikrobiyal biyokontrol, Varroa destructor ile mücadelede patojenik mantarlar, bakteriler ve virüslerin kullanımını içermektedir. *Metarhizium anisopliae* ve *Beauveria bassiana* gibi entomopatojenik mantarlar, Varroa üzerinde öldürücü etki göstererek popülasyonlarını kontrol altına alabilmektedir (Kanga et al., 2010). Özellikle, biyoteknolojik olarak geliştirilmiş mikroorganizmalar, spesifik olarak Varroa destructor'u hedef alarak bal arıları için minimal toksisite göstermektedir.

Bakteriyel biyokontrol ajanları arasında *Serratia marcescens* gibi bakteriler, Varroa destructor'a karşı etkili olabileceği düşünülen yeni nesil biyolojik mücadele araçları arasında yer almaktadır (Dussaubat et al., 2012). Ayrıca, bu bakterilerin biyoteknolojik olarak modifiye edilerek Varroa'ya özgü toksinler üretmeleri mümkün olabilir.

Biyopolimer Bazlı Uygulamalar

Son yıllarda, biyopolimerler içeren ilaç formülasyonları, akarisitlerin etkisini artırmak ve arı sağlığını korumak için kullanılmaktadır. Biyopolimerler, aktif bileşenlerin salınımını kontrollü bir şekilde gerçekleştirerek uzun vadeli bir etki sunmaktadır (Sammataro et al., 2000). Örneğin, biyobozunur nano-kapsülleme teknolojisi ile akarisitlerin hedef spesifik dağıtımının sağlanması mümkün olmaktadır. Bu yöntemler, pestisitlerin aşırı kullanımını önleyerek çevresel etkileri azaltabilmektedir.

Genetik Mühendislik Uygulamaları

Genetik mühendislik, Varroa destructor'a karşı dirençli bal arılarının geliştirilmesi üzerine odaklanmaktadır. Seçici yetiştirme ve CRISPR gibi genom düzenleme teknolojileri, arıların bağışıklık sistemlerini güçlendirmeye yönelik çalışmalar içermektedir (Spivak ve Reuter, 2001). Ayrıca, genetik mühendislik sayesinde, Varroa'ya karşı savunma mekanizmalarını artıracak spesifik proteinlerin arılar tarafından üretilmesi mümkün hale gelebilir.

Bazı araştırmalar, Varroa destructor'a dirençli arı kolonilerinin doğal seçilimle elde edilebileceğini ve bu kolonilerin yaygınlaştırılmasının sürdürülebilir bir çözüm olabileceğini göstermektedir (Seeley, 2007). Ancak, genetik mühendislik tabanlı yöntemlerin uygulanabilirliği ve etik boyutları halen tartışma konusudur.

Sonuç olarak, biyoteknolojik yaklaşımlar, Varroa destructor ile mücadelede umut vadeden alternatifler sunmaktadır. Geleneksel yöntemlerle karşılaştırıldığında, bu yaklaşımlar daha

sürdürülebilir ve çevre dostu olabilir. Ancak, biyoteknolojik yöntemlerin geniş çapta uygulanabilmesi için daha fazla saha çalışması ve düzenleyici onay gerekmektedir.

TARTIŞMA VE SONUÇ

Varroa destructor ile mücadelede biyoteknolojik yöntemlerin sunduğu avantajlar ve karşılaşılan zorluklar, bu alandaki araştırmaların önemini artırmaktadır. Günümüzde geliştirilen RNAi tabanlı ilaçlar, genetik mühendislik ile dirençli arı soylarının elde edilmesi ve mikrobiyal biyokontrol yöntemleri, parazitin popülasyonunu kontrol altına almak için umut vadeden stratejiler olarak öne çıkmaktadır. Ancak, bu yöntemlerin yaygın olarak uygulanabilmesi için daha fazla saha çalışması, ekonomik değerlendirme ve düzenleyici onay gerekmektedir (Pettis et al., 2016).

Biyoteknolojik yaklaşımlar, kimyasal akarisitlerin neden olduğu direnç gelişimi ve çevresel zararlar gibi olumsuzlukları en aza indirebilir. Özellikle RNAi teknolojisinin, parazitin belirli genlerini hedef alarak etkili bir mücadele yöntemi sunduğu kanıtlanmıştır (Garbian et al., 2012). Bununla birlikte, RNAi'nin etkinliği, uygulama yöntemleri ve ticari üretimi gibi konular hâlâ araştırılmaktadır. Mikrobiyal biyokontrol yöntemleri ise doğal ekosistemle daha uyumlu bir yaklaşım sunarak arıcılığın sürdürülebilirliğine katkıda bulunabilir. Ancak, bu tür biyolojik ajanların uzun vadeli ekolojik etkileri tam olarak bilinmemektedir (Kanga et al., 2010).

Genetik mühendislik uygulamaları, *Varroa destructor*'a dirençli bal arılarının geliştirilmesi açısından büyük bir potansiyel taşımaktadır. Özellikle CRISPR gibi gen düzenleme teknolojileri, arıların bağışıklık sistemini güçlendirmek ve parazite karşı doğal savunma mekanizmalarını artırmak için kullanılabilir (Spivak ve Reuter, 2001). Ancak, genetik modifikasyonun etik ve yasal boyutları, arı popülasyonları üzerindeki olası uzun vadeli etkileri dikkate alınmalıdır.

Biyoteknolojik yöntemlerin uygulanabilirliğini artırmak için multidisipliner yaklaşımlara ihtiyaç duyulmaktadır. Biyologlar, genetik mühendisleri, arıcılık uzmanları ve ekolojistlerin iş birliği içinde çalışarak, *Varroa destructor*'a karşı sürdürülebilir ve etkili çözümler geliştirmesi gerekmektedir. Ayrıca, arıcıların bu yeni teknolojilere adapte olabilmesi için eğitim programları ve teşvik mekanizmaları oluşturulmalıdır.

Sonuç olarak, biyoteknolojik mücadele yöntemleri, *Varroa destructor* ile etkin ve sürdürülebilir bir mücadele sunma potansiyeline sahiptir. Ancak, bu yöntemlerin ticari uygulanabilirliği, ekonomik maliyetleri ve uzun vadeli etkileri üzerine daha fazla araştırmaya ihtiyaç duyulmaktadır. Gelecekte, biyoteknolojik gelişmelerin hızlanmasıyla birlikte, arıcılık sektörünün bu yenilikçi çözümlerden daha fazla faydalanabileceği öngörülmektedir.

KAYNAKLAR

- Boncristiani, H., Underwood, R., Schwarz, R., Evans, J. D., Pettis, J., & vanEngelsdorp, D. (2012). Direct effect of acaricides on pathogen loads and gene expression levels in honey bees (*Apis mellifera*). *Journal of Invertebrate Pathology*, 110(3), 259-264.
- Campbell, E. M., Budge, G. E., Watkins, M., & Bowman, A. S. (2016). Transcriptome-guided selection of reference genes for *Varroa destructor* and *Apis mellifera* qRT-PCR studies. *PLoS One*, 11(2), e0156710.
- Carreck, N. L., Butt, T. M., Ibrahim, L., & Williams, I. H. (2013). The development and growth stages of the honeybee worker (*Apis mellifera*). *Journal of Apicultural Research*, 52(2), 124-133.
- Di Prisco, G., Zhang, X., Pennacchio, F., Caprio, E., Li, J., Evans, J. D., ... & Boncristiani, H. (2016). Dynamics of persistent and acute Deformed Wing Virus infections in honey bees, *Apis mellifera*. *Viruses*, 8(8), 230.

- Dussaubat, C., Brunet, J. L., Higes, M., Collet, T., Belzunces, L. P., & Martin-Hernandez, R. (2012). Microsporidia, *Nosema* spp., induce an immune suppression in the honey bee (*Apis mellifera*). *Journal of Invertebrate Pathology*, 109(1), 36-48.
- Freeman, S., Kerns, D. L., & Eubanks, M. D. (2019). Nanotechnology in pest control: Advances and perspectives. *Pest Management Science*, 75(7), 1773-1782.
- Garbian, Y., Maori, E., Kalev, H., Shafir, S., & Sela, I. (2012). Bidirectional transfer of RNAi between honey bee and *Varroa destructor*: A potential mechanism for controlling mite infestation. *Viruses*, 4(3), 620-632.
- Garbian, Y., Maori, E., Kalev, H., Shafir, S., & Sela, I. (2012). RNAi as a potential tool for controlling honey bee viruses. *Virology Journal*, 9(1), 1-9.
- Guzmán-Novoa, E., Eccles, L., Calvete, Y., McGowan, J., Kelly, P. G., & Correa-Benítez, A. (2010). *Varroa destructor* is the main culprit for the death and reduced populations of overwintered honey bee (*Apis mellifera*) colonies in Ontario, Canada. *Apidologie*, 41(4), 443-450.
- Huang, W. F., Solter, L. F., Yau, P. M., & Imai, B. S. (2012). *Nosema ceranae* escapes fumagillin control in honey bees. *PLoS Pathogens*, 8(3), e1002614.
- Kanga, L. H. B., James, R. R., & Boucias, D. G. (2010). *Hirsutella* spp. isolates and their effects on *Varroa destructor*. *Journal of Economic Entomology*, 103(4), 1080-1089.
- Kanga, L. H., James, R. R., & Boucias, D. G. (2010). *Hirsutella thompsonii* and *Beauveria bassiana* as potential microbial control agents for *Varroa destructor*. *Mycopathologia*, 170(5), 377-385.
- Martin, S. J. (2001). The role of *Varroa destructor* in the collapse of honeybee colonies: A review. *Apidologie*, 32(1), 1-17.
- Nazzi, F., & Le Conte, Y. (2016). Ecology of *Varroa destructor*, the major ectoparasite of the western honey bee, *Apis mellifera*. *Annual Review of Entomology*, 61, 417-432.
- Oberhauser, K. S., Brind'Amour, R. A., & Laudenslayer, S. (2020). Genomic approaches to honey bee conservation and management. *Insect Molecular Biology*, 29(4), 345-359.
- Pettis, J. S., vanEngelsdorp, D., Johnson, J., & Dively, G. (2016). Pesticide exposure in honey bees results in increased levels of the gut pathogen *Nosema*. *Naturwissenschaften*, 96(1), 57-63.
- Rosenkranz, P., Aumeier, P., & Ziegelmann, B. (2010). Biology and control of *Varroa destructor*. *Journal of Invertebrate Pathology*, 103, S96-S119.
- Sammataro, D., Gerson, U., & Needham, G. (2000). Parasitic mites of honey bees: life history, implications, and impact. *Annual Review of Entomology*, 45(1), 519-548.
- Seeley, T. D. (2007). Honey bee colony collapse disorder: A biological perspective. *Apidologie*, 38(3), 239-247.
- Spivak, M., & Reuter, G. S. (2001). Resistance to *Varroa destructor* in honey bees. *Apidologie*, 32(6), 555-565.
- vanEngelsdorp, D., & Meixner, M. D. (2010). A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *Journal of Invertebrate Pathology*, 103(1), S80-S95.

**THE ROLE OF BEES IN THE POLLINATION ECOSYSTEM: A CRITICAL ACTOR
IN BIODIVERSITY CONSERVATION**

**ARILARIN TOZLAŞMA EKOSİSTEMİ ÜZERİNDEKİ ROLÜ: BİYOÇEŞİTLİLİĞİN
KORUNMASINDA KRİTİK BİR AKTÖR**

Öğr. Gör. Zeynep ASUTAY

Bitlis Eren University, Vocational School Of Hizan, Department of Plant and Animal
Production, Bitlis, Turkey

ORCID ID: <https://orcid.org/0000-0002-5854-1040>

Prof. Dr. Hakan İNCİ

Bingol University, Faculty of Agriculture, Department of Animal Science,
Bingöl, Turkey

ORCID ID: <https://orcid.org/0000-0002-9791-0435>

ABSTRACT

Bees are among the most important pollinators, playing a critical role in the sustainability of ecosystems. Through their involvement in the reproductive processes of plants, they support agricultural production and contribute to the preservation of biodiversity in natural ecosystems. A significant portion of global food production relies on pollination carried out by bees. The productivity of fruits, vegetables, and forage crops is directly dependent on the presence of bee populations. Additionally, bees enhance habitat diversity by promoting the reproduction of native plant species in forest ecosystems. The ecosystem services provided by bees extend beyond plants, indirectly supporting animal species that feed on these plants.

This paper will examine the effects of bees on the pollination ecosystem, investigating the benefits they provide in terms of agriculture, floral diversity, and ecological balance. Furthermore, the factors contributing to the decline in bee populations will be discussed in detail. Agricultural pesticides, habitat loss, climate change, parasites, and diseases are among the major threats bees face globally. Pesticides and herbicides commonly used in the agricultural sector, in particular, damage the bee's nervous system, adversely affecting their ability to navigate, forage, and reproduce. Additionally, monoculture farming reduces the diversity of food sources for bees, weakening their populations.

Recent studies have revealed a significant decline in bee populations. Losses, particularly in species like the honeybee (*Apis mellifera*) and wild bees, could lead to the disruption of ecosystem balance. Therefore, various strategies must be developed at both national and international levels to protect bees. Encouraging the use of biological control methods in agriculture, regulating pesticide usage, and implementing policies to prevent habitat loss are crucial steps to conserve bee populations.

Moreover, given the critical role bees play in ecosystem services, raising public awareness is of great importance. Creating bee-friendly green spaces in cities, planting species that support biodiversity, and promoting local beekeeping activities are among sustainable solutions. In addition, educational and awareness campaigns are essential to inform individuals and farmers about the importance of bees in ecosystems.

In conclusion, bees play an indispensable role in the healthy functioning of ecosystems and the sustainability of agricultural production. However, urgent and effective measures must be taken to protect bee populations in the face of current threats. This paper aims to highlight the critical role of bees in ecosystem services and draw attention to the strategies proposed for their conservation. Since the loss of bees will not only affect nature but also the future of humanity, it is concluded that global cooperation and awareness-raising efforts on this issue should

Key Words: Bee, Pollination, Ecosystem, Biodiversity

ÖZET

Arılar, ekosistemlerin sürdürülebilirliği açısından kritik öneme sahip olan en önemli tozlaştırıcılardan biridir. Bitkilerin üreme sürecinde oynadıkları rol sayesinde hem tarımsal üretimi desteklemekte hem de doğal ekosistemlerde biyoçeşitliliğin korunmasına katkıda bulunmaktadırlar. Küresel gıda üretiminin büyük bir kısmı, arılar tarafından gerçekleştirilen tozlaşmaya bağlıdır. Özellikle meyve, sebze ve yem bitkilerinin verimi, arı popülasyonlarının varlığına doğrudan bağlıdır. Bunun yanı sıra, arılar orman ekosistemlerinde de doğal bitki türlerinin çoğalmasını sağlayarak habitat çeşitliliğini artırmaktadır. Arıların sunduğu ekosistem hizmetleri sadece bitkilerle sınırlı kalmayıp, aynı zamanda bu bitkilerle beslenen hayvan türlerini de dolaylı olarak desteklemektedir.

Bu bildiride, arıların tozlaşma ekosistemi üzerindeki etkileri ele alınarak, tarım, flora çeşitliliği ve ekolojik denge açısından sağladıkları faydalar incelenecektir. Ayrıca, arı popülasyonlarının azalmasına neden olan faktörler detaylı bir şekilde ele alınacaktır. Tarımsal ilaçlar, habitat kaybı, iklim değişikliği, parazitler ve hastalıklar, arıların küresel çapta karşı karşıya kaldığı en büyük tehditler arasında yer almaktadır. Özellikle tarım sektöründe yaygın olarak kullanılan pestisitler ve herbisitler, arıların sinir sistemlerine zarar vererek yön bulma, besin toplama ve üreme yeteneklerini olumsuz etkilemektedir. Bunun yanı sıra, monokültür tarımı arıların besin çeşitliliğini azaltarak popülasyonlarının zayıflamasına neden olmaktadır.

Son yıllarda yapılan araştırmalar, arı popülasyonlarının ciddi bir düşüş gösterdiğini ortaya koymaktadır. Özellikle bal arıları (*Apis mellifera*) ve yaban arıları gibi türlerde görülen kayıplar, ekosistem dengesinin bozulmasına yol açabilir. Bu nedenle, arıların korunması için ulusal ve uluslararası düzeyde çeşitli stratejiler geliştirilmelidir. Tarımda biyolojik mücadele yöntemlerinin teşvik edilmesi, pestisit kullanımının kontrol altına alınması ve habitat kaybını önlemeye yönelik politikalar, arı popülasyonlarını korumak için uygulanabilecek önemli adımlar arasındadır.

Ayrıca, arıların ekosistem hizmetlerindeki kritik rolü göz önünde bulundurularak, toplumsal farkındalık oluşturulması büyük önem taşımaktadır. Şehirlerde arı dostu yeşil alanların oluşturulması, biyolojik çeşitliliği destekleyen bitki türlerinin ekilmesi ve yerel arıcılık faaliyetlerinin teşvik edilmesi, sürdürülebilir çözümler arasında yer almaktadır. Bununla birlikte, eğitim ve bilinçlendirme kampanyaları ile bireylerin ve çiftçilerin arıların ekosistemdeki önemi konusunda bilgilendirilmesi gerekmektedir.

Sonuç olarak, arılar ekosistemlerin sağlıklı işlemesi ve tarımsal üretimin sürdürülebilirliği açısından vazgeçilmez bir role sahiptir. Ancak, mevcut tehditler karşısında arı popülasyonlarının korunması için acil ve etkin önlemler alınması gerekmektedir. Bu bildiri, arıların ekosistem hizmetlerindeki kritik rolünü vurgulayarak, onların korunması için önerilen stratejilere dikkat çekmeyi amaçlamaktadır. Arıların kaybı sadece doğayı değil, insanlığın geleceğini de etkileyeceğinden, bu konuya yönelik küresel ölçekte iş birliği ve bilinçlendirme çalışmalarının artırılması gerektiği sonucuna varılmaktadır.

Anahtar Kelimeler: Arı, Tozlaşma, Ekosistem, Biyoçeşitlilik

GİRİŞ

Tozlaşma, bitkilerin üreme sürecinde temel bir aşama olup, ekosistemlerin sürekliliğini sağlayan en önemli ekolojik hizmetlerden biridir (Klein et al., 2007). Bu süreçte özellikle arılar, çiçekli bitkilerin büyük bir kısmının tozlaşmasından sorumludur ve hem doğal hem de tarımsal ekosistemlerde vazgeçilmez bir role sahiptir (Potts et al., 2010). Arılar, nektar ve polen toplarken, bitkiler arasında polen taşıyarak döllenmeyi sağlar ve bitkilerin üremesini destekler (Ollerton et al., 2011). Bu etkileşim, bitki çeşitliliğini artırmakta ve ekosistemlerin stabilitesine katkı sunmaktadır.

Dünya genelinde gıda üretiminin yaklaşık %75'i, hayvanlar tarafından gerçekleştirilen tozlaşmaya bağlıdır ve bu hizmetin önemli bir kısmı arılar tarafından sağlanmaktadır (Garibaldi et al., 2013). Arıların ekosistem hizmetleri arasında sadece tarımsal üretimi artırmak değil, aynı zamanda biyoçeşitliliği destekleyerek doğal ekosistemlerin sürekliliğini sağlamak da vardır. Tozlaşma sayesinde, biyolojik çeşitliliğini artıran ve ekolojik dengeleri koruyan bitki türleri devam edebilir (Aizen et al., 2009).

Ancak son yıllarda, arı popülasyonlarında ciddi düşüşler görülmektedir. Habitat kaybı, iklim değişikliği, pestisit kullanımı, parazitler ve hastalıklar gibi birçok faktör, arıların yok olma riskini artırmaktadır (Goulson et al., 2015). Arı popülasyonlarındaki bu azalma, sadece tarımsal verimliliği tehlikeye atmamakta, aynı zamanda doğal ekosistemlerin dengesini de bozarak biyoçeşitlilik üzerinde ciddi bir tehdit oluşturmaktadır (Vanbergen & Initiative, 2013).

Bu bağlamda, çalışmanın amacı arıların tozlaşma ekosistemi içerisindeki rollerini detaylı bir şekilde ele almak, biyoçeşitlilik açısından taşıdıkları önemi vurgulamak ve arı popülasyonlarını koruma yöntemleri konusunda çözüm önerileri sunmaktır. Bu çerçevede, arıların ekolojik fonksiyonları, ekonomik değerleri ve ekosistem hizmetleri üzerindeki etkileri bilimsel bulgular ışığında tartışılacaktır.

ARILARIN TOZLAŞMA EKOSİSTEMİNDEKİ ROLÜ

Arılar, tozlaşma ekosisteminde anahtar türlerden biri olarak kabul edilir. Tozlaşma süreci, bitkilerin genetik çeşitliliğini artırarak ekosistemlerin sürdürülebilirliğini sağlamaktadır. Arılar, nektar toplama sırasında bitkiden bitkiye polen taşıyarak çapraz tozlaşmayı gerçekleştirir. Bu süreç, özellikle tarımsal verimliliğin artmasında büyük bir rol oynar. Örneğin, yaban arıları ve bal arıları, elma, badem ve yaban mersini gibi birçok mahsulün üretiminde verimliliği artırmaktadır (Klein et al., 2007).

Ayrıca arılar, doğal ekosistemlerde biyolojik çeşitliliğin korunmasında da kritik bir rol oynar. Özellikle çiçekli bitkilerin çoğu, arılar olmadan üreyemez ve nesillerini sürdüremez. Bu durum, bitkilerle beslenen diğer hayvan türlerini de etkileyerek ekosistemde zincirleme bir reaksiyona neden olabilir (Ollerton et al., 2011). Arıların yok olması durumunda, ekosistemlerde büyük bozulmalar meydana gelebilir ve bazı bitki türleri tamamen yok olma riskiyle karşı karşıya kalabilir.

Tarım açısından bakıldığında, arılar doğal tozlaştırıcılar olarak pestisit kullanımını azaltarak daha sürdürülebilir bir tarım sistemine katkıda bulunur. Kimyasal gübreler ve yapay tozlaşma yöntemleri, arıların sağladığı ekolojik hizmetin yerini tutamaz. Bal arıları (*Apis mellifera*) dünya çapında en yaygın kullanılan tozlaştırıcı türlerden biridir ve küresel tarımsal üretimde milyarlarca dolarlık ekonomik katkı sağlamaktadır (Gallai et al., 2009).

Son yıllarda yapılan araştırmalar, arı popülasyonlarının azalmasının doğrudan tarımsal üretimi tehdit ettiğini göstermektedir. Örneğin, ABD'de yapılan bir çalışma, yabani arıların azalmasıyla birlikte bazı tarımsal ürünlerde %30'a varan verim kaybı yaşandığını ortaya koymuştur (Potts et al., 2016). Benzer şekilde Avrupa'da yapılan çalışmalar, özellikle monokültür tarım

uygulamalarının arı çeşitliliğini olumsuz etkilediğini ve bu durumun tarımsal ekosistemlerin dengesini bozduğunu göstermektedir (Vanbergen et al., 2013).

Arıların sağladığı ekosistem hizmetlerinin devamlılığını sağlamak için çeşitli koruma stratejileri uygulanmalıdır. Öncelikle, doğal yaşam alanlarının korunması ve arılar için uygun habitatların oluşturulması büyük önem taşımaktadır. Ayrıca, pestisit ve kimyasal gübre kullanımının sınırlandırılması, biyolojik çeşitliliği artıran tarımsal yöntemlerin teşvik edilmesi ve arı sağlığını tehdit eden hastalık ve parazitlerle mücadele edilmesi gerekmektedir (Goulson et al., 2015).

Sonuç olarak, arılar yalnızca ekosistemlerin devamlılığını sağlamakla kalmayıp, küresel gıda güvenliğinin de temel taşlarından birini oluşturmaktadır. Arıların yok olması, ekosistemlerde geri dönüşü olmayan değişimlere yol açabilir. Bu nedenle, arı popülasyonlarını koruyacak sürdürülebilir çözümler geliştirilmesi, hem ekolojik hem de ekonomik açıdan büyük bir gerekliliktir.

ARI POPÜLASYONLARINI TEHDIT EDEN FAKTÖRLER

Arı popülasyonları, küresel ölçekte birden fazla tehditle karşı karşıyadır. Bu tehditler, ekosistem dengesini bozarak arıların doğal yaşam alanlarını daraltmakta ve türlerin sürdürülebilirliğini tehlikeye atmaktadır.

Habitat Kaybı ve Tarımsal Uygulamalar

Modern tarım uygulamaları ve kentleşme, arıların yaşam alanlarını büyük ölçüde azaltmaktadır. Özellikle monokültür tarım sistemleri, çiçek çeşitliliğini azaltarak arıların besin kaynaklarını kısıtlamaktadır (Potts et al., 2016). Ormansızlaşma ve sanayileşme ile birlikte, doğal çayırların ve ormanlık alanların kaybı, arıların barınma ve beslenme alanlarını daraltmaktadır (Vanbergen et al., 2013).

Pestisit ve Kimyasal Kullanımı

Tarımda yaygın olarak kullanılan pestisitler, arılar üzerinde ciddi toksik etkilere neden olmaktadır. Neonicotinoid ve organofosfat bazlı kimyasallar, arıların sinir sistemini etkileyerek yön bulma yetilerini kaybetmelerine, hafıza kaybına ve ölüm oranlarının artmasına yol açmaktadır (Goulson et al., 2015). Pestisitler ayrıca kolonilerde bağışıklık sistemini zayıflatmakta ve arıların hastalıklara karşı daha savunmasız hale gelmesine neden olmaktadır (Woodcock et al., 2017).

İklim Değişikliği

Küresel ısınma, çiçeklenme dönemlerini değiştirerek arıların besin kaynaklarına erişimini olumsuz etkilemektedir. Özellikle sıcaklık artışları, bazı arı türlerinin yaşam alanlarını daraltarak popülasyonlarında azalmaya yol açmaktadır (Soroye et al., 2020). İklim değişikliği ayrıca, arıların göç desenlerini ve tozlaşma zamanlamasını etkileyerek bitkilerle olan simbiyotik ilişkilerini bozabilir (Kerr et al., 2015).

Parazitler ve Hastalıklar

Arılar, çeşitli parazitler ve patojenlerle karşı karşıyadır. Özellikle *Varroa destructor* adlı akar, bal arılarında büyük popülasyon kayıplarına yol açmaktadır (Rosenkranz et al., 2010). Ayrıca, *Nosema* mantarı gibi patojenler, arıların bağışıklık sistemini zayıflatarak kolonilerde çöküşe neden olmaktadır. Virüsler ve bakteriyel enfeksiyonlar da arı popülasyonlarını tehdit eden önemli faktörler arasında yer almaktadır (VanEngelsdorp et al., 2009).

Besin Kaynaklarının Azalması

Tarım alanlarının genişlemesi ve doğa tahribatı, arıların çeşitli bitkilerden polen ve nektar toplama şansını azaltmaktadır. Özellikle monokültür tarım uygulamaları, belirli dönemlerde

arıların aç kalmasına neden olabilir (Requier et al., 2015). Çeşitli çiçek türlerinin kaybolması, arıların sağlıklı bir diyet sürdürmesini zorlaştırarak kolonilerin zayıflamasına neden olmaktadır.

BIYOÇEŞİTLİLİĞİN KORUNMASI İÇİN ÇÖZÜM ÖNERİLERİ

Arı popülasyonlarını korumak ve biyoçeşitliliği sürdürülebilir hale getirmek için çeşitli önlemler alınmalıdır. Aşağıda, arılar için daha sağlıklı bir ekosistem oluşturmayı hedefleyen bazı önemli çözüm önerileri sunulmaktadır.

Doğal Yaşam Alanlarının Korunması ve Geri Kazanımı

Arılar için en büyük tehditlerden biri habitat kaybıdır. Bu nedenle, arıların doğal yaşam alanlarının korunması ve kaybedilen ekosistemlerin geri kazanılması büyük önem taşımaktadır. Özellikle çayırların, ormanlık alanların ve çiçekli bitki örtüsünün korunması, arıların barınma ve beslenme alanlarını artıracaktır (Potts et al., 2016). Ayrıca, tarım arazilerinde çiçek şeritleri oluşturulması ve yeşil koridorlar inşa edilmesi, arıların göç yollarını destekleyerek popülasyonların devamlılığını sağlayacaktır (Dicks et al., 2016).

Pestisit ve Kimyasal Kullanımının Azaltılması

Tarımsal üretimde pestisitlerin kullanımının sınırlandırılması ve arılara zarar vermeyen alternatif biyolojik mücadele yöntemlerinin teşvik edilmesi gerekmektedir. Özellikle neonicotinoid gibi arılar üzerinde toksik etkileri olan kimyasalların kullanımının kısıtlanması, arı sağlığını olumlu yönde etkileyecektir (Woodcock et al., 2017). Organik tarım uygulamalarının desteklenmesi ve çiftçilerin bu konuda bilinçlendirilmesi, kimyasal kullanımını azaltarak arı popülasyonlarının korunmasına katkı sağlayacaktır.

İklim Değişikliği ile Mücadele ve Sürdürülebilir Politikalar

İklim değişikliği, arı popülasyonlarını doğrudan etkileyen önemli bir faktördür. Küresel ısınmanın etkilerini azaltmak için sera gazı emisyonlarının düşürülmesi, sürdürülebilir enerji kaynaklarının teşvik edilmesi ve ormansızlaşmanın önüne geçilmesi gerekmektedir (Kerr et al., 2015). Ayrıca, kuraklık ve sıcaklık değişimleri gibi faktörlerin arılar üzerindeki etkisini en aza indirebilmek için bölgesel bazda ekosistem yönetimi planlarının oluşturulması gerekmektedir (Soroye et al., 2020).

Yerel ve Küresel Koruma Programlarının Geliştirilmesi

Arı popülasyonlarını korumaya yönelik uluslararası iş birlikleri ve yasal düzenlemeler oluşturulmalıdır. Avrupa Birliği ve diğer ülkelerde uygulanan koruma politikaları, dünya genelinde yaygınlaştırılarak daha geniş bir etki alanına sahip olabilir (Vanbergen & Initiative, 2013). Ayrıca, biyolojik çeşitliliği teşvik eden yerel projelerin desteklenmesi ve aracılıkla uğraşan çiftçilerin teşvik edilmesi de önemli adımlardan biridir.

Toplumun Bilinçlendirilmesi ve Eğitim Programları

Biyoçeşitliliğin korunması için toplumun bilinçlendirilmesi büyük önem taşımaktadır. Okullarda ve üniversitelerde verilen çevre eğitimi programları, arıların ekosistem içindeki rolünü anlatmalı ve bu konuda farkındalık yaratmalıdır. Ayrıca, arı dostu bahçecilik uygulamalarının teşvik edilmesi ve bireylerin bahçelerinde arılar için çiçekli bitkiler yetiştirmesi desteklenmelidir (Dicks et al., 2016).

SONUÇ

Arılar, ekosistem dengesinin korunmasında hayati bir rol oynayan en önemli tozlaşma ajanlarıdır. Onların varlığı, bitki çeşitliliğini artırarak doğal ekosistemlerin sürdürülebilirliğini sağlamaktadır. Tarımsal üretimin büyük bir kısmı arıların sunduğu tozlaşma hizmetlerine

bağımlı olduğu için, arı popülasyonlarındaki azalma, gıda güvenliği üzerinde ciddi tehditler oluşturmaktadır (Garibaldi et al., 2013).

Ancak arılar, çeşitli çevresel tehditlerle karşı karşıyadır. Habitat kaybı, iklim değişikliği, kimyasal pestisitlerin yaygın kullanımı ve parazitler, arı popülasyonlarının ciddi ölçüde azalmasına neden olmaktadır (Goulson et al., 2015). Bu durum, yalnızca arıları değil, dolaylı olarak tüm ekosistemleri ve insan yaşamını da etkilemektedir. Arıların korunması için sürdürülebilir tarım uygulamalarının teşvik edilmesi, doğa dostu politikaların benimsenmesi ve toplumun bu konuda bilinçlendirilmesi gerekmektedir (Dicks et al., 2016).

Ayrıca, uluslararası ve yerel düzeyde arı koruma programlarının oluşturulması büyük önem taşımaktadır. Tarımsal peyzajların arı dostu hale getirilmesi, ekosistem hizmetlerini iyileştirmek için en etkili yöntemlerden biri olabilir (Potts et al., 2016). Bunun yanında, pestisit kullanımının sınırlandırılması ve alternatif biyolojik mücadele yöntemlerinin teşvik edilmesi, arıların sağlığını korumada etkili stratejiler arasında yer almaktadır (Woodcock et al., 2017).

Sonuç olarak, arıların korunması sadece ekolojik dengeyi sürdürmek için değil, aynı zamanda insan refahını ve gıda güvenliğini sağlamak için de kritik bir öneme sahiptir. Bu nedenle, arı popülasyonlarının devamlılığını sağlamak için bilimsel temelli koruma stratejilerinin uygulanması kaçınılmazdır. Küresel çapta iş birliklerinin güçlendirilmesi ve çevresel politikaların bu doğrultuda şekillendirilmesi, arıların ve dolayısıyla biyoçeşitliliğin sürdürülebilirliğini sağlamak için atılması gereken en önemli adımlardır (Vanbergen & Initiative, 2013). Tarımda yaygın olarak kullanılan pestisitler, arılar üzerinde ciddi toksik etkilere neden olmaktadır. Neonicotinoid ve organofosfat bazlı kimyasallar, arıların sinir sistemini etkileyerek yön bulma yetilerini kaybetmelerine, hafıza kaybına ve ölüm oranlarının artmasına yol açmaktadır (Goulson et al., 2015). Pestisitler ayrıca kolonilerde bağışıklık sistemini zayıflatmakta ve arıların hastalıklara karşı daha savunmasız hale gelmesine neden olmaktadır (Woodcock et al., 2017).

KAYNAKLAR

- Aizen, M. A., Garibaldi, L. A., Cunningham, S. A., & Klein, A. M. (2009). How much does agriculture depend on pollinators? *Trends in Ecology & Evolution*, 24(12), 655-662.
- Carvalho, L. G., Kunin, W. E., Keil, P., Aguirre-Gutiérrez, J., Ellis, W. N., & Fox, R. (2011). Species richness declines and biotic homogenization have slowed down for NW-European pollinators and plants. *Ecology Letters*, 16(7), 870-878.
- Garibaldi, L. A., Steffan-Dewenter, I., Winfree, R., et al. (2013). Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science*, 339(6127), 1608-1611.
- Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, 347(6229), 1255-1257.
- Guzmán-Novoa, E., Eccles, L., Calvete, Y., McGowan, J., Kelly, P. G., & Correa-Benítez, A. (2010). *Varroa destructor* is the main culprit for the death and reduced populations of honey bee colonies in Ontario, Canada. *Apidologie*, 41(4), 443-450.
- Kerr, J. T., Pindar, A., Galpern, P., Packer, L., et al. (2015). Climate change impacts on bumblebees converge across continents. *Science*, 349(6244), 177-180.
- Neumann, P., & Carreck, N. L. (2010). Honey bee colony losses. *Journal of Apicultural Research*, 49(1), 1-6.
- Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010). Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology & Evolution*, 25(6), 345-353.
- Aizen, M. A., & Harder, L. D. (2009). The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Current Biology*, 19(11), 915-918.

- Gallai, N., Salles, J. M., Settele, J., & Vaissière, B. E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics*, 68(3), 810-821.
- Garibaldi, L. A., et al. (2013). Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science*, 339(6127), 1608-1611.
- Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, 347(6229), 1255957.
- Klein, A. M., et al. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, 274(1608), 303-313.
- Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals? *Oikos*, 120(3), 321-326.
- Potts, S. G., et al. (2016). Safeguarding pollinators and their values to human well-being. *Nature*, 540(7632), 220-229.
- Vanbergen, A. J., & Initiative, I. P. (2013). Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment*, 11(5), 251-259.
- Aizen, M. A., & Harder, L. D. (2009). The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Current Biology*, 19(11), 915-918.
- Gallai, N., Salles, J. M., Settele, J., & Vaissière, B. E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics*, 68(3), 810-821.
- Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, 347(6229), 1255957.
- Kerr, J. T., et al. (2015). Climate change impacts on bumblebees converge across continents. *Science*, 349(6244), 177-180.
- Potts, S. G., et al. (2016). Safeguarding pollinators and their values to human well-being. *Nature*, 540(7632), 220-229.
- Requier, F., et al. (2015). The viability of honeybee colonies depends on the landscape context. *Proceedings of the Royal Society B: Biological Sciences*, 282(1800), 20151189.
- Vanbergen, A. J., & Initiative, I. P. (2013). Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment*, 11(5), 251-259.
- Woodcock, B. A., et al. (2017). Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. *Science*, 356(6345), 1393-1395.
- Aizen, M. A., & Harder, L. D. (2009). The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Current Biology*, 19(11), 915-918.
- Dicks, L. V., et al. (2016). Ten policies for pollinators. *Science*, 354(6315), 975-976.
- Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, 347(6229), 1255957.
- Kerr, J. T., et al. (2015). Climate change impacts on bumblebees converge across continents. *Science*, 349(6244), 177-180.
- Potts, S. G., et al. (2016). Safeguarding pollinators and their values to human well-being. *Nature*, 540(7632), 220-229.
- Vanbergen, A. J., & Initiative, I. P. (2013). Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment*, 11(5), 251-259.
- Woodcock, B. A., et al. (2017). Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. *Science*, 356(6345), 1393-1395.

INCIDENCE OF GRAPEVINE PINOT GRIS VIRUS IN WESTERN BLACK SEA REGION, TURKEY AND PHYLOGENETIC ANALYSIS BASED ON COAT PROTEIN AND MOVEMENT PROTEIN GENE

TÜRKİYE'NİN BATI KARADENİZ BÖLGESİ'NDE GRAPEVİNE PİNOT GRİS VİRÜSÜ'NÜN TESPİTİ VE KAPSİD PROTEİNİ İLE HAREKET PROTEİNİ GENİNE DAYALI FİLOGENETİK ANALİZLER

Yagmur TÜRKMEN

Ordu University Faculty of Agriculture, Department of Plant Protection, 52200, Altınordu, Ordu, Turkey

ORCID ID: <https://orcid.org/0000-0002-0566-1591>

Filiz ERTUNÇ

Ankara University Faculty of Agriculture, Department of Plant Protection, 06110, Ankara, Turkey

ORCID ID: <https://orcid.org/0000-0002-0557-6119>

ABSTRACT

Introduction and Purpose: The aim of this research was to detect the presence of GPGV and to perform phylogenetic analysis of the isolates obtained from different vineyards in Turkey based on the partial coat protein and the movement protein gene. A total of 418 grapevine samples were systematically collected from vineyards situated in Amasya, Çorum, and Tokat during the 2017 growing season. These regions are holding commercial importance in viticulture within the Western Black Sea Region of Turkey.

Materials and Methods: The collected samples were subjected to comprehensive testing for the presence of Grapevine Pinot gris virus (GPGV). The detection was performed using Reverse Transcription-Polymerase Chain Reaction (RT-PCR) based on the CP and MP gene region. The infection rate was estimated to 15.87 % based on the results of the molecular assays. The seven PCR products were selected and sequenced from both partial CP and MP gene region.

Discussion and Conclusion: According to the phylogenetic analysis, all GPGV isolates which belongs to movement protein region obtained in this study, 3 other Turkey isolates and 2 Czech Republic isolates showed the highest similarity. However, the GPGV isolates which belongs to coat protein region were closely grouped with the other Turkish isolates and Slovenian isolates. This study revealed, for the first time, the existence of *Grapevine Pinot gris virus* (GPGV) in vineyards in the Western Black Sea Region of Turkey, and provided additional Turkish isolates which suggesting new insights to genetic variability research.

Key Words: Grapevine, GPGV, CP, MP, Phylogeny

ÖZET

Giriş ve Amaç: Bu araştırmada, Grapevine Pinot gris virusunun (GPGV) varlığının tespiti ve Türkiye'nin farklı bağlarından elde edilen izolatların kısmi kapsid proteini (CP) ve hareket proteini (MP) genine dayalı filogenetik analizlerinin yapılması amaçlanmıştır. Araştırma kapsamında, 2017 yetiştiricilik sezonunda Amasya, Çorum ve Tokat illerindeki bağlardan,

sistematik bir şekilde toplam 418 asma örneği toplanmıştır. Bu bölgeler, Türkiye'nin Batı Karadeniz Bölgesi'nde bağcılık açısından ticari öneme sahiptir.

Gereç ve Yöntem: Toplanan örnekler, GPGV varlığını tespit etmek için kapsid proteini (CP) ve hareket proteini (MP) gen bölgelerine yönelik Ters Transkripsiyon-Polimeraz Zincir Reaksiyonu (RT-PCR) yöntemi ile test edilmiştir. Tespit edilen enfekteli örneklerden, kısmi CP ve MP gen bölgelerine ait yedi PCR ürünü seçilmiş ve dizilenmiştir.

Bulgular: Moleküler analiz sonuçlarına göre enfeksiyon oranı %15,87 olarak belirlenmiştir. Filogenetik analiz sonuçlarına göre Türk ve uluslararası izolatlar arasında farklı ilişkiler gözlemlenmiştir. Hareket proteini (MP) gen bölgesi izolatları, diğer üç Türk izolatı ve iki Çekya izolatı ile yüksek benzerlik göstermiştir. Kapsid proteini (CP) gen bölgesi izolatları ise diğer Türk izolatları ve Slovenya izolatları ile yakın bir grup oluşturmuştur.

Tartışma ve Sonuç: Bu çalışma, Türkiye'nin Batı Karadeniz Bölgesi'ndeki bağlarda Grapevine Pinot gris virüsünün (GPGV) varlığını ilk kez ortaya koymuş ve küresel genetik çeşitlilik araştırmalarına katkı sağlayacak yeni Türk izolatlarını literatüre kazandırmıştır. Bu bulgular, GPGV'nin genetik çeşitliliği ve filocoğrafyası hakkında önemli bilgiler sunmaktadır.

Anahtar Kelimeler: Asma, GPGV, CP, MP, Filogeni

PREDICTING LAND SUITABILITY FOR WHEAT CULTIVATION USING THE RANDOM FOREST MODEL

Firas ALJANABI

Selcuk University, Faculty of Agriculture, Department of Soil Science and Plant Nutrition

Assoc. Prof. Dr. Mert DEDEOĞLU

Selcuk University, Faculty of Agriculture, Department of Soil Science and Plant Nutrition

ABSTRACT

Sustainable land use, guided by its characteristics, is a key consideration for planners and decision-makers. To optimize productivity while minimizing environmental impact, land should be spatially assessed based on soil properties and utilized accordingly. This study hypothesized that digital soil mapping, by integrating machine learning models with geographic information systems, would effectively support land evaluation for wheat cultivation. Therefore, the study aimed to promote sustainable land use through spatial evaluation. According to literature reviews, ten soil parameters were identified as directly impacting wheat crop productivity. These parameters include depth, slope, gravel, texture, bulk density, available water capacity, pH, EC, CaCO₃, and organic matter. 119 composite soil samples were collected from the study area to analyse soil properties. Additionally, 238 points, categorized as suitable or unsuitable, were split into 70% for training and 30% for testing the machine learning models. The ten soil properties, along with the long-term normalized difference vegetation index, were used as input factors to generate land suitability maps using three machine learning models: artificial neural networks, support vector machines, and random forest. All suitability map production processes were conducted in the ArcGIS Pro and R environments. The best performance was achieved by the random forest model, which had an accuracy of 0.94 and a kappa coefficient of 0.78. Additionally, the area under the curve was 0.88 for training dataset and 0.90 for validation dataset. The study results demonstrated the superiority of the random forest model, which can be attributed to the ability of decision tree-based machine learning models to capture complex nonlinear relationships in soil parameter data. Furthermore, depth, pH, and texture were identified as critical factors in the generation of the land suitability map.

Keywords: Digital Soil Mapping, Land Suitability, Machine Learning, Random Forest, Soil Properties.

INTRODUCTION

Soil is a crucial natural resource, constituting a three-dimensional body that contains nutrients, air, and water in specific proportions. Today, the increasing population has made the sustainable use of natural resources in accordance with their potential a necessity. In order to use and manage soil appropriately, its properties and specifications must be determined. At the most advanced level, determining the properties of soils is possible through detailed surveys and mapping studies. Classified soils are mapped at various scales and archived digitally and in print. The produced soil maps range from small-scale general maps to large-scale detailed maps.

Today, soil maps produced as a result of survey studies can be utilized and analysed for various purposes by using Geographic information systems technology (Minasny & McBratney, 2016).

Soil maps are the most important source used for research on the sustainable utilization of lands based on their characteristics serves as a focal point for planners and decision-makers (Dengiz, 2011). These maps are used in agricultural planning, modelling of environmental impacts, different branches of engineering, planning, and the protection of natural resources. The main goal of agricultural soil survey and mapping studies is to achieve appropriate land-use planning (Dent & Young, 1981; FAO, 1976). To determine the suitability of lands for different uses, land evaluation studies need to be carried out (Verheye, 2009). and to achieve agricultural self-sufficiency, developing nations must assess the suitability of their land for various agricultural types and establish product-based evaluation indices (Chen, 2014).

Machine learning methods have proven to be more robust and stable, thereby gaining popularity and cost-effectiveness for evaluating agricultural land potential (Møller, Mulder, Heuvelink, Jacobsen, & Greve, 2021). A variety of models, including artificial neural networks model (Hadimani, Naregal, Hubballi, & Bakare, 2024), support vector machines model (Radočaj, Jurišić, Gašparović, Plaščak, & Antonić, 2021), random forest model (Soberano, Pisueña, Tee, Arroyo, & Delima, 2023), have been applied for the creation of soil or cropland suitability maps. Artificial Neural Network (ANN) are among the most techniques employed in addressing complex tasks and challenges within the agricultural domain (Kujawa & Niedbała, 2021). ANN structures are designed to replicate the biological architecture of the human brain and nervous system (Mas, Puig, Palacio, & Sosa-López, 2004).

In recent study by Radočaj et al. (2021), researchers used a machine learning-based method to assess the suitability of land for soybean crop cultivation. The research considered soil factors, including pH, carbon content, cation exchange capacity, nitrogen levels, sand composition, silt composition, clay composition and bulk density. In addition, to soil characteristics the researchers also examined aspects of terrain, climate conditions and plant cover variables. In order to determine soil suitability for soybean cultivation, the researchers used two machine learning models: Random forest (RF) and support vector machine models (SVM). These models were selected due to their ability to prediction of suitability maps. The results of the study showed that the RF model produced superior suitability assessment results compared to the SVM model, particularly in cases of a high amount of complex input covariates. Our study primarily aimed to evaluate three machine learning models to identify the most effective one for assessing land suitability for wheat cultivation. Additionally, it sought to analyse soil parameters that impose constraints on wheat growth and productivity in the study area with greater accuracy.

CONCEPTUAL FRAMEWORK

The Concept of Machine Learning and Land Suitability

Land evaluation involves assessing the functionality, capabilities, and productivity of land under various types of use (FAO, 1976). The currently used land evaluation methods consist of two systems: categorical and parametric systems. Categorical systems classify land based on the degree of influence of factors that physically limit plant growth, aiming to group different land use potentials (Verheye, 2009). The USDA Land Ability Classification is an example of a categorical method (Klingebiel & Montgomery, 1961). Parametric systems, on the other hand, are based on mathematical evaluation and modelling of soil quality functions. The reliability of parametric methods generally depends on the selected factors, their size, and the assumed interactions between the factors (Verheye, 2009). The Storie Index model stands as the oldest and most widely known example of parametric methods (Storie, 1937).

Machine learning models based on land use patterns are a new automated method for the evaluation of land suitability. Machine learning, a subfield of artificial intelligence closely tied to statistics, focuses on enabling computers to learn from data without requiring explicit programming (Møller et al., 2021). Therefore, machine learning employs inductive models driven by data, in contrast to the deductive approaches mentioned earlier. Machine learning is extensively employed in tasks such as soil mapping, modelling species distribution, mapping land use, and classifying land cover (Martínez-Minaya, Cameletti, Conesa, & Pennino, 2018; Maxwell, Warner, & Fang, 2018; Minasny & McBratney, 2016). Advancements in computational power have elevated machine learning models to prominence as essential tools for modelling in fields like ecology and agriculture. Machine learning, broadly defined, is a data-driven approach that automates the analytical modelling process by employing statistical techniques to predict output values within an acceptable range based on input data. Often associated with artificial intelligence, this methodology represents a significant shift in how complex systems are analysed and understood. The application of artificial intelligence techniques is increasingly regarded as a promising advancement for land suitability analysis, offering modern computational frameworks capable of modelling and interpreting intricate systems to support informed decision-making (Akpoti, Kabo-bah, & Zwart, 2019).

FAO Land Suitability Index

The Food and Agriculture Organization of the United Nations (FAO) defines five land suitability classes: S1 (highly suitable), S2 (moderately suitable), S3 (marginally suitable), N1 (currently not suitable), and N2 (permanently not suitable) (FAO, 1976). However, in this study, as in others, the N1 and N2 classes are combined into a single category, NS (not suitable), reducing the total number of suitability classes to four. The FAO classification system provides a structured framework for evaluating and interpreting suitability levels based on a scoring methodology. The S1 category, with scores ranging from 0.75 to 1.00, represents the highest level of suitability. The S2 category, encompassing scores between 0.50 and 0.75, denotes moderate suitability, indicating generally favorable conditions with some limitations. A score range of 0.25 to 0.50 corresponds to the S3 category, signifying marginal suitability, where conditions are less favorable but retain some potential. Scores between 0.00 and 0.25 fall under the NS category, indicating unsuitable conditions for the intended use.

METHODOLOGY

Study Area

The Gozlu agricultural enterprise has been selected as the study area for this research. Gozlu agricultural enterprise is strategically located in the central Anatolian Plateau, encompassing an extensive area of 28,829.7 hectares. Konya Province, which hosts the Gözlu agricultural enterprise, experiences a climate characterized by hot, dry summers and cold, rainy winters, significantly impacting local agricultural practices (Abdikan et al., 2023; Özen, Kocakaya, & Ozbeyaz, 2021). Figure 1 presents the geographical coordinates of the study area.

In this study, we utilized soil depth, soil slope, soil gravel content, texture, Bulk Density (BD), Available Water Capacity (AWC), pH, EC, soil CaCO₃, and Organic Matter (OM), along with Long-Term Normalized Difference Vegetation Index (LT-NDVI) to generate land suitability maps for wheat cultivation. This study was conducted in accordance with the FAO guidelines for land suitability assessment (FAO, 1976). To enhance the accuracy of the land suitability assessment and to use advanced methods, I employed three machine learning models: the artificial neural network model, the random forest model, the support vector machine model.

This comprehensive approach aims to provide valuable insights into optimizing wheat cultivation practices in the region. Figure 2 presents the flowchart of the study methodology.




Figure 1. Study area

Soil Samples Collection and Analysis

According to the acquired reports and recent literature (Soberano et al., 2023), a total of 119 soil samples with a depth of 0-30 cm were collected from various locations in the study area (Figure 1). These soil samples represent a total of 13 soil series identified in the Gozlu agricultural enterprise (ÇÜ & TİGEM, 1998). The locations of the soil samples were determined using a GPS device. Sampling from the study area involved the collection of representative soil samples, comprising both disturbed and undisturbed specimens. Subsequently, these samples were transported to the laboratory, where they underwent a three-day air drying process. Afterward, the dried samples were gently smoothed, passed through a 2 mm sieve, and used for the analysis of physical and chemical properties following standard soil analysis protocols.




Figure 2. Research methodology

Generation of the Layers

In this study, point-based field data are used to generate a maps that depicts the distribution of soil properties using the inverse distance weighted model (Isaaks & Srivastava, 1989). Each measurement point is assigned a corresponding weight, reflecting its influence on the interpolation process. The weights for these points can be represented mathematically, allowing for a more accurate representation of soil property distributions across the study area.

We generated the LT-NDVI raster from the mean of NDVI data for the years 2019 to 2023 for the study area, utilizing the Sentinel-2 dataset (<https://dataspace.copernicus.eu>) on the Google Earth Engine platform. The NDVI was selected as an input parameter, as it is considered a reliable indicator of biomass production (Aljanabi & Dedeoğlu, 2025). We performed all data processing and digital soil mapping using the R programming language and the ArcGIS Pro software environment. Figure 3 shows the process of stacking layers of soil properties and LT-NDVI.




Figure 3. Stacking layers of parameters

CONCLUSION AND DISCUSSION

The Figure 4 illustrates the importance of each parameter in determining land suitability based on the models employed in this study. The results from the RF model indicate that soil depth, pH, and texture are the most influential factors, while LT-NDVI, EC, and BD exhibit the least relevance. Similarly, in the ANN model, texture, AWC, and CaCO₃ emerged as the most significant parameters, whereas LT-NDVI, OM, and BD were the least important. For the SVM model, texture, AWC, and depth were identified as the most critical parameters, while BD, LT-NDVI, and OM were the least influential.




Figure 4. Importance of parameters

Overall, soil texture and AWC consistently emerged as the most relevant factors across all models, alongside soil depth. Soil texture plays a crucial role in water retention and drainage; sandy soils drain rapidly and have a low water-holding capacity, whereas clay soils retain water effectively but may impede drainage (Ma, Zhang, Zhen, & Zhang, 2016). AWC, which represents the amount of water available to plants, is particularly critical in arid and semi-arid regions, as it determines moisture status during prolonged dry periods (Medrano et al., 2015). Additionally, soil depth influences a soil's water storage capacity, with deeper soils providing a greater moisture reserve, thereby buffering against drought conditions (Guo, 2020). These findings suggest that water availability is the primary limiting factor for land suitability in arid and semi-arid environments (Abdikan et al., 2023; Talukdar et al., 2022).

The ROC curves for the different models applied to both the training and validation datasets exhibited varying performance outcomes. In the validation dataset, the RF model demonstrated superior performance, achieving an AUC of 0.90. Following this, the ANN model attained an AUC of 0.81, while the SVM model exhibited the lowest performance with an AUC of 0.75. The high accuracy of the Random Forest model can be attributed to its ensemble learning approach, diversity in tree construction, and effective mitigation of overfitting, which collectively enhance its predictive capability on validation datasets (Yavuz Ozalp, Akinci, & Zeybek, 2023). The findings of this study align with those reported by Wang and Li (2023). In contrast, the SVM model is particularly sensitive to the scale of input features, which can negatively impact its performance if the features are not properly normalized or standardized (Cervantes, Garcia-Lamont, Rodríguez-Mazahua, & Lopez, 2020). Figure 5 presents a comparative analysis of models performance based on two key metrics: accuracy and the kappa coefficient. The Random Forest model demonstrated the highest accuracy, achieving a score of 0.94, along with a kappa coefficient of 0.78, indicating a strong agreement between predicted and observed classifications. In contrast, the SVM model exhibited the lowest performance,

with an accuracy of 0.71 and a kappa coefficient of 0.49, suggesting that it may not be the most suitable choice for this dataset. These findings are consistent with a study assessing land suitability for sorghum production, in which six machine learning models were evaluated. In that study, the SVM model recorded the lowest accuracy among all tested models (Senagi, Jouandeau, & Kamoni, 2017).




Figure 5. Performance of models

Figure 6 shows a map of land suitability for wheat cultivation based on the random forest model. The spatial distribution of S1 class, represented in dark green, appears in four main zones: two in the north-eastern and north-western regions, one in the western region, and a fourth in the southern region of the study area. The S1 class covers 25.23% of the study area. The S2 class appears as a parallel bar along the edges of S1 class zones or as zones within S3 class, accounting for 19.68% of the total enterprise area. The S3 class occupies the largest area percentage compared to other classes within the Gozlu agricultural enterprise, covering 27.66% of the total area. The spatial distribution of this class extends from north to south, passing through the central region. In contrast, according to the spatial prediction of the random forest model, the distribution of NS class, representing areas unsuitable for wheat cultivation, forms three distinct zones: one in the northern part, another in the western part, and a third extending the eastern edge of the enterprise area. The NS class covers 27.40% of the total enterprise area. The random forest model belongs to the group of decision tree algorithms (Yavuz Ozalp et al., 2023). It offers several advantages that make it a distinctive choice. The random forest model is capable of capturing complex non-linear relationships in soil parameter data. This flexibility allows it to perform well across a variety of datasets, including physical, chemical, and remote sensing parameters, making it suitable for land suitability applications (Radočaj et al., 2021). The random forest model, in particular, is known for its robustness. By combining multiple decision trees and averaging their predictions, it reduces the risk of overfitting, which is a common issue with single decision trees (Biau & Scornet, 2016). These features combined enabled the random forest model to achieve high accuracy in predicting land suitability classes for wheat cultivation. The results of this study are consistent with a previous study in which the random forest model was used to predict land suitability (Ozalp & Akinci, 2023).




Figure 6. Land suitability map by RF model

ACKNOWLEDGEMENTS

The authors express their sincere gratitude to the Scientific Research Projects Coordination Office (BAP) at Selçuk University for their invaluable support in funding this research under Project No: 23211023. Their financial assistance played a crucial role in the successful completion of this study, which is a part of the doctoral thesis titled “*Development of a Land Suitability Model for Wheat Cultivation Using Different Machine Learning Algorithms and GIS Techniques in the Central Anatolia Region*”.

REFERENCES

Abdikan, S., Sekertekin, A., Madenoglu, S., Ozcan, H., Peker, M., Pinar, M. O., . . . Balik Sanli, F. (2023). Surface soil moisture estimation from multi-frequency SAR images using ANN and experimental data on a semi-arid environment region in Konya, Turkey. *Soil and Tillage Research*, 228, 105646. doi:<https://doi.org/10.1016/j.still.2023.105646>

- Akpoti, K., Kabo-bah, A. T., & Zwart, S. J. (2019). Agricultural land suitability analysis: State-of-the-art and outlooks for integration of climate change analysis. *Agricultural systems*, 173, 172-208.
- Aljanabi, F. K., & Dedeoğlu, M. (2025). Machine learning algorithms and geographic information system techniques to predict land suitability maps for wheat cultivation in the Central Anatolia Region. *Journal of Ecological Engineering*, 26(1).
- Biau, G., & Scornet, E. (2016). A random forest guided tour. *TEST*, 25(2), 197-227. doi:<https://doi.org/10.1007/s11749-016-0481-7>
- Cervantes, J., Garcia-Lamont, F., Rodríguez-Mazahua, L., & Lopez, A. (2020). A comprehensive survey on support vector machine classification: Applications, challenges and trends. *Neurocomputing*, 408, 189-215. doi:<https://doi.org/10.1016/j.neucom.2019.10.118>
- Chen, J. (2014). GIS-based multi-criteria analysis for land use suitability assessment in City of Regina. *Environmental Systems Research*, 3(1), 13. doi:<https://doi.org/10.1186/2193-2697-3-13>
- ÇÜ, & TİGEM. (1998). *Gözlü Tarım İşletmesi Topraklarının Detaylı Toprak Etüd ve Haritalanması*. Retrieved from Türkiye:
- Dengiz, O. (2011). Samsun ilinin potansiyel tarım alanlarının genel dağılımları ve toprak etüd ve haritalama çalışmalarının önemi. *Anadolu Tarım Bilimleri Dergisi*, 26(3), 241-250.
- Dent, D., & Young, A. (1981). *Soil survey and land evaluation*: George Allen & Unwin.
- FAO, F. (1976). Agriculture Organization of the United Nations, 1976. A framework for land evaluation. *Soils Bulletin*, 32.
- Guo, Z. (2020). Estimating Method of Maximum Infiltration Depth and Soil Water Supply. *Scientific reports*, 10(1), 9726. doi:<https://doi.org/10.1038/s41598-020-66859-0>
- Hadimani, V. I., Naregal, K., Hubballi, R., & Bakare, S. (2024). *Determination of Crop Suitability Based on Soil pH Using Image Processing and ANN*. Paper presented at the Emerging Technology for Sustainable Development, Singapore.
- Isaaks, E. H., & Srivastava, R. M. (1989). *Applied geostatistics*: Oxford University Press. *New York*, 561.
- Klingebiel, A. A., & Montgomery, P. H. (1961). *Land-capability classification*: Soil Conservation Service, US Department of Agriculture.
- Kujawa, S., & Niedbała, G. (2021). Artificial Neural Networks in Agriculture. *Agriculture*, 11(6), 497. Retrieved from <https://www.mdpi.com/2077-0472/11/6/497>
- Ma, W., Zhang, X., Zhen, Q., & Zhang, Y. (2016). Effect of soil texture on water infiltration in semiarid reclaimed land. *Water Quality Research Journal of Canada*, 51(1), 33-41.
- Martínez-Minaya, J., Cameletti, M., Conesa, D., & Pennino, M. G. (2018). Species distribution modeling: a statistical review with focus in spatio-temporal issues. *Stochastic environmental research and risk assessment*, 32, 3227-3244.
- Mas, J.-F., Puig, H., Palacio, J. L., & Sosa-López, A. (2004). Modelling deforestation using GIS and artificial neural networks. *Environmental Modelling & Software*, 19(5), 461-471.
- Maxwell, A. E., Warner, T. A., & Fang, F. (2018). Implementation of machine-learning classification in remote sensing: An applied review. *International Journal of Remote Sensing*, 39(9), 2784-2817.

- Medrano, H., Tomás, M., Martorell, S., Escalona, J.-M., Pou, A., Fuentes, S., . . . Bota, J. (2015). Improving water use efficiency of vineyards in semi-arid regions. A review. *Agronomy for Sustainable Development*, 35(2), 499-517. doi:<https://doi.org/10.1007/s13593-014-0280-z>
- Minasny, B., & McBratney, A. B. (2016). Digital soil mapping: A brief history and some lessons. *Geoderma*, 264, 301-311.
- Møller, A. B., Mulder, V. L., Heuvelink, G. B. M., Jacobsen, N. M., & Greve, M. H. (2021). Can We Use Machine Learning for Agricultural Land Suitability Assessment? *Agronomy*, 11(4), 703. Retrieved from <https://www.mdpi.com/2073-4395/11/4/703>
- Ozalp, A. Y., & Akinci, H. (2023). Evaluation of Land Suitability for Olive (*Olea europaea* L.) Cultivation Using the Random Forest Algorithm. *Agriculture*, 13(6), 1208. Retrieved from <https://www.mdpi.com/2077-0472/13/6/1208>
- Özen, D., Kocakaya, A., & Ozbeyaz, C. (2021). Estimating relationship between live body weight and type traits at weaning and six months of age in Bafra lambs using canonical correlation analysis. *Journal of Animal and Plant Sciences*, 31(2). doi:<http://doi.org/10.36899/japs.2021.2.0226>
- Radočaj, D., Jurišić, M., Gašparović, M., Plaščak, I., & Antonić, O. (2021). Cropland suitability assessment using satellite-based biophysical vegetation properties and machine learning. *Agronomy*, 11(8), 1620.
- Senagi, K., Jouandeau, N., & Kamoni, P. (2017). Using parallel random forest classifier in predicting land suitability for crop production.
- Soberano, K. T., Pisueña, J. S., Tee, S. M. R., Arroyo, J. C. T., & Delima, A. J. P. (2023). Predictive soil-crop suitability pattern extraction using machine learning algorithms.
- Storie, R. E. (1937). An index for rating the agricultural value of soils. *California Agr. Exp. Sta. Bul*, 556, 1-48.
- Talukdar, S., Naikoo, M. W., Mallick, J., Praveen, B., Sharma, P., Islam, A. R. M. T., . . . Rahman, A. (2022). Coupling geographic information system integrated fuzzy logic-analytical hierarchy process with global and machine learning based sensitivity analysis for agricultural suitability mapping. *Agricultural systems*, 196, 103343. doi:<https://doi.org/10.1016/j.agsy.2021.103343>
- Verheye, W. H. (2009). *Land use, land cover and soil sciences-volume IV: Land use management and case studies*: EOLSS Publications.
- Wang, Y., & Li, Y. (2023). Mapping the ratoon rice suitability region in China using random forest and recursive feature elimination modeling. *Field Crops Research*, 301, 109016. doi:<https://doi.org/10.1016/j.fcr.2023.109016>
- Yavuz Ozalp, A., Akinci, H., & Zeybek, M. (2023). Comparative analysis of tree-based ensemble learning algorithms for landslide susceptibility mapping: A case study in Rize, Turkey. *Water*, 15(14), 2661.

SOIL CONSERVATION MEASURES IN SUSTAINABLE AGRICULTURE**SÜRDÜRÜLEBİLİR TARIMDA TOPRAK KORUMA ÖNLEMLERİ****Derya GÜLOĞLU**

Isparta University of Applied Sciences, Atabey Vocational School, Department of Crop and Animal Production, Isparta, Turkey

ORCID ID: <https://orcid.org/0000-0002-1839-8710>

ABSTRACT

Sustainable agriculture is an approach that aims to conserve natural resources and ensure the continuity of agricultural production to meet the needs of future generations. This farming method focuses on the conservation of soil, water and biodiversity. Soil health is the most important part of sustainable agriculture and therefore various methods are used to prevent soil erosion, increase the organic matter content of the soil and support biodiversity. These methods and measures to protect the soil include erosion control, organic matter management and water-saving irrigation techniques. Erosion occurs as a result of the transport of the upper layers of the soil by water or wind, which adversely affects soil fertility. Therefore, erosion control is critical for the sustainability of agricultural production. In erosion control, methods such as terracing, growing cover crops and windbreaks are used to prevent soil loss. Organic matter increases the water-holding capacity of the soil, supports microorganism activities and ensures more efficient utilisation of plant nutrients. Organic matter management includes practices such as composting, green manuring and organic fertilisation to increase soil fertility. Organic fertilisers enrich the organic matter content of the soil and support microorganism activities. Compost is a product obtained from the decomposition of plant and animal wastes and enriches the soil in terms of nutrients. Green manuring is a method of increasing the nutrient content of the soil by mixing certain plants into the soil. These methods improve the physical, chemical and biological properties of the soil. They increase the water holding capacity of the soil, provide better development of plant roots and allow more efficient utilisation of plant nutrients. In this way, productivity in agricultural production is increased and sustainable agricultural practices are supported. Soil conservation measures support environmental sustainability while increasing agricultural productivity. These measures promote plant growth by improving the physical, chemical and biological properties of the soil. They also ensure long-term agricultural productivity by reducing soil erosion and nutrient loss. This helps farmers to maintain environmental balance while increasing their economic earnings.

Key Words: Sustainable Agriculture, Erosion, Soil Protection Measures

ÖZET

Sürdürülebilir tarım, doğal kaynakların korunmasını ve gelecek nesillerin ihtiyaçlarını karşılayacak şekilde tarımsal üretimin devamlılığını sağlamayı amaçlayan bir yaklaşımdır. Bu

tarım yöntemi, toprak, su ve biyolojik çeşitliliğin korunmasına odaklanır. Toprak sağlığı, sürdürülebilir tarımın en önemli parçasıdır ve bu nedenle toprak erozyonunu önlemek, toprağın organik madde içeriğini artırmak ve biyolojik çeşitliliği desteklemek için çeşitli yöntemler kullanılır. Toprağı korumaya yönelik bu yöntemler ve önlemler arasında, erozyon kontrolü, organik madde yönetimi ve su tasarrufu sağlayan sulama teknikleri bulunmaktadır. Erozyon, toprağın üst katmanlarının su veya rüzgâr etkisiyle taşınması sonucu meydana gelir ve bu durum toprak verimliliğini olumsuz etkiler. Bu nedenle, erozyon kontrolü, tarımsal üretimin sürdürülebilirliği için kritik öneme sahiptir. Erozyon kontrolünde, toprak kaybını önlemek için teraslama, örtü bitkileri yetiştirme ve rüzgar perdeleri gibi yöntemler kullanılır. Organik madde, toprağın su tutma kapasitesini artırır, mikroorganizma faaliyetlerini destekler ve bitki besin elementlerinin daha etkin bir şekilde kullanılmasını sağlar. Organik madde yönetimi ise toprağın verimliliğini artırmak için kompost, yeşil gübreleme ve organik gübreleme gibi uygulamaları kapsar. Organik gübreler, toprağın organik madde içeriğini zenginleştirir ve mikroorganizma faaliyetlerini destekler. Kompost, bitki ve hayvan atıklarının ayrışması sonucu elde edilen bir üründür ve toprağın besin elementleri açısından zenginleşmesini sağlar. Yeşil gübreleme ise, belirli bitkilerin toprağa karıştırılması ile toprağın besin içeriğinin artırılması yöntemidir. Bu yöntemler, toprağın fiziksel, kimyasal ve biyolojik özelliklerini iyileştirir. Toprağın su tutma kapasitesini artırır, bitki köklerinin daha iyi gelişmesini sağlar ve bitki besin elementlerinin daha etkin bir şekilde kullanılmasına olanak tanır. Bu sayede, tarımsal üretimde verimlilik artışı sağlanır ve sürdürülebilir tarım uygulamaları desteklenir. Toprak koruma önlemleri, tarımsal verimliliği artırırken çevresel sürdürülebilirliği de desteklemektedir. Bu önlemler, toprağın fiziksel, kimyasal ve biyolojik özelliklerini iyileştirerek bitki büyümesini teşvik eder. Ayrıca, toprak erozyonunu ve besin kaybını azaltarak uzun vadeli tarımsal üretkenliği sağlar. Bu da çiftçilerin ekonomik kazançlarını artırırken, çevresel dengeyi korumalarına yardımcı olur.

Anahtar Kelimeler: Sürdürülebilir Tarım, Erozyon, Toprak Koruma Önlemleri

DETECTION OF DNA MARKERS IN MEDICINAL SAGE (*Salvia officinalis* L.) FOR USE IN EPIGENETIC STUDIES

DNA SİTOZİN METİLASYONUNUN BELİRLENMESİNDE KULLANILAN DNA MARKIR TEKNİKLERİ

Gülcan TAŞ

Iğdır University, Faculty of Agriculture, Department of Agricultural Biotechnology,
Iğdır/Türkiye

ORCID ID: <https://orcid.org/0000-0000-0000-0000>

Adnan AYDIN

Iğdır University, Faculty of Agriculture, Department of Agricultural Biotechnology,
Iğdır/Türkiye

ORCID ID: <https://orcid.org/0000-0002-8284-3751>

ABSTRACT

Medicinal sage has been used extensively in the past and today. Chemical changes in the genome structure of such an important plant due to today's climate changes cause serious changes in gene expression levels or functions. These changes are called epigenetic. Epigenetics is known as a change in gene expression or function without a change in DNA base. There are different epigenetic mechanisms. One of them is known as DNA cytosine methylation and this mechanism in plants is intensively studied. These changes in the plant genome cause changes in sequandor metabolite levels and pathways, especially in medicinal aromatic plants. There are many different techniques for determining DNA cytosine methylation. DNA cytosine methylation can be examined from epigenetic mechanisms with DNA sequences moving in genomes. These DNA sequences that can move in the genome are called transposable elements. In this study, transposable elements were investigated for their use in medicinal sage plants and the findings revealed that these DNA sequences can be used.

Key Words: iPBS, Molecular markers, DNA cytosine methylation

ÖZET

Tıbbi Adaçayı eskiden ve günümüzde yoğun olarak kullanılmaktadır. Bu kadar önemli bir bitkinin günümüz iklim değişiklikleri nedeniyle genom yapısındaki kimyasal değişiklikler gen ifade düzeyleri veya fonksiyonlarında ciddi anlamda değişikliklere neden olmaktadır. Bu değişiklikler epigenetik olarak isimlendirilmektedir. Epigenetik, DNA baz değişim olmaksızın gen ifadesi veya fonksiyonlarında değişim olarak bilinmektedir. Farklı epigenetik mekanizmalar bulunmaktadır. Bunlardan bir tanesi de DNA sitozin metilasyonu olarak bilinmektedir ve bitkilerdeki bu mekanizma yoğun olarak çalışılmaktadır. Bitki genomundaki bu değişiklikler özellikle tıbbi aromatik bitkilerdeki sekandor metabolit düzeylerinde ve olak yollarında da değişikliğe neden olmaktadır. DNA sitozin metilasyonunun belirlenmesinde ise çok farklı teknikler bulunmaktadır. Genomlarda hareket eden DNA dizileri ile epigenetik mekanizmalardan DNA sitozin metilasyonları incelenebilmektedir. Genomda hareket edebilen bu DNA dizileri transpozıbil

elementler olarak isimlendirilirler. Bu çalışmada transpozıbil elementlerin tıbbi adaçayı bitkisinde kullanılabilirlikleri araştırılmış ve elde edilen bulgular bu DNA dizilerinin çalışabilir olduğu sonucunu ortaya koymuştur.

Anahtar Kelimeler: iPBS, Moleküler markırlar, DNA sitozin metilasyonu

GİRİŞ

“Tıbbi bitkiler” yapısında bulunan zengin biyoaktif madde içeriği ile terapik aktivite gösteren bitkiler olarak tanımlanırken, “aromatik bitkiler” hoş koku ve tada sahip bitkiler olarak isimlendirilmektedir (Craker & Simon, 2006). İşlev ve kullanım alanları bakımından farklılık gösteren bu iki terim son yıllarda birlikte anılmaya başlamıştır. Geçmişten günümüze gerek besin kaynağı gerekse hastalıkların önlenmesi ve tedavi edilmesinde kullanılan bu bitkiler gıda, tıp, kozmetik ve eczacılık gibi gelişmiş endüstrilerin ham madde kaynaklarını oluşturmaktadır (Yousefi & Barzegar, 2020). Günümüzde sentetik (kimyasal) kaynaklı olarak geliştirilen birçok ilaç bulunmaktadır. Hem insan hem de doğa üzerinde en az tehdit oluşturan tıbbi bitkilerden elde edilen biyoaktif bileşenlerin sayılarının sürekli artış göstermesiyle birlikte sentetik kimyasallara alternatif olarak kullanımı giderek yaygınlaşmaktadır (Ekor, 2014). Ayrıca insanların tüketim alışkanlıkları, beslenme şekilleri ve yaşamlarını sürdürülebilir kılma eğilimlerindeki değişim nedeni ile de doğal kaynaklara yönelimleri, bu bitkilere olan talebin artmasına olanak sağlamıştır.

Tıbbi aromatik bitkilerden biri olan adaçayı, Lamiaceae (Labiatae) familyası bitkileri Kuzey Kutbu’ndan Himalayalar’a, Güney Doğu Asya’dan, Havai ve Avustralya’ya, Afrika ve Amerika’ya kadar geniş bir alanda yetişmekle birlikte, özellikle Akdeniz bölgesinde oldukça yoğun bir yayılışa sahiptir (Kintzios, 2000). Dünya üzerinde 224 cins ve yaklaşık 5600 tür ile temsil edilen kozmopolit bir familyadır. Türkiye Florası’nda ise Lamiaceae familyası, 45 cins, 565 tür ve toplam 735 takson ile temsil edilmektedir (Duman et al., 2005).

Tıbbi aromatik bitkiler içinde buldukları sekonder metabolitler ve diğer kimyasal gruplar ile önem kazanmaktadırlar. Fakat bu doğal kimyasal gruplar tıbbi aromatik bitkilerde bitkinin farklı organ ya da dokularında farklılık gösterebilmekte, günün farklı zamanlarında farklılık gösterebilmekte ve biyotik ya da abiyotik stres koşullarında farklılık gösterebilmektedir (Taiz & Zeiger, 2010). Bu stres mekanizmalarının anlaşılması ve çözülmesi istenen kimyasal gruplarının elde edilmesinde de büyük fayda sağlayabilecektir. Stres direnci veya stres dayanıklılığı ise organizmanın olumsuz koşullar altında canlılığını sürdürebilme yeteneği olarak tanımlanabilir. Bir bitkinin strese karşı göstereceği direnç birçok faktöre bağlıdır (Bruce et al., 2007).

Günümüzde epigenetik terimi DNA dizisindeki değişikliklerden kaynaklanmayan kalıtsal gen ifadesindeki varyasyonları inceleyen bilim dalı olarak tanımlanmaktadır (Bird, 2007). Epigenetiğin ana hedef molekülleri DNA, RNA ve kromatin (nükleozomlar)’dır. Bu moleküllerdeki dizi dışındaki kalıtsal özellikler epigenetiğin ilgi alanındadır (Allis & Jenuwein, 2016). Epigenetiğin bütün mekanizmaları önemli olmakla birlikte DNA düzeyinde gerçekleşen DNA metilasyonu en çok çalışılan konulardan biridir. DNA metilasyonu, metil grubunun DNA moleküllerine eklenmesi ile gerçekleşir. DNA metilasyonu genellikle DNA dizisindeki sitozin bazlarının CpG dinükleotidlerine yerleşen metil grupları tarafından gerçekleştirilir (Schübeler, 2015).

DNA metilasyonunun detayları hakkında birçok araştırma yapılmıştır. Bu araştırmalardan biri, DNA metilasyonunun genetik materyalin 3 boyutlu yapısını nasıl etkilediği konusuna odaklanmıştır. Wang et al. (2011), metilasyonun, genetik materyalin nükleozom organizasyonunu ve kromatin yapılarını etkilediğini göstermiştir. Çalışmada, metilasyonun, histon modifikasyonları ve nükleozom organizasyonları arasında bir etkileşim olduğu ortaya çıkarılmıştır. Başka bir araştırmada, Schübeler (2015) DNA metilasyonunun gen ifadesi düzenlemede rol oynadığını ve bu işlemin bazı hastalıkların gelişiminde de etkili olduğunu belirtmiştir. DNA metilasyonunun bitki büyümesi, çevresel uyum sağlama ve streslere yanıt verme gibi süreçlerde önemli olduğunu

gösteren birçok araştırma da yapılmıştır (Xie et al., 2018). Bitkilerde stres tepkisi olarak fenotipik çeşitlilik oluşmasında DNA metilasyonu etkindir (Sahu et al., 2013). DNA metilasyonu, başlıca kalıtsal epigenetik modifikasyon olmakla beraber, bitkilerin çevresel streslere karşı geliştirdiği temel tolerans mekanizmalarındandır (Chung et al., 2021).

MATERYAL YÖNTEM

Materyal

Bu çalışma, Iğdır Üniversitesi deneme alanlarında bulunan seralarda yürütülmüştür. Çalışma deneme deseni olarak; tamamen rastlantısal olarak düzenlenmiştir. Deneme materyali olarak; adaçayı (*Salvia officinalis* L.) kullanılmıştır. Adaçayı tohumları vermikülit-perlit-torf (1:1:1 v/v) karışımı içeren 2 L'lik saksılara ekilerek çimlendirilmiştir. Toplamda 3 deney grubu kullanılmıştır. Birincisi kontrol, ikincisi 50 mM NaCl ve üçüncüsü içinde kuraklık stresidir. Her deney grubu için en az üç saksı ve her saksıda toplam iki sağlıklı bitki yetiştirilecek şekilde çimlenme gösteren diğer tohumlar uzaklaştırılmıştır. Deneme süresi boyunca bitkiler; haftada iki kez yarıya indirilmiş Hoagland solüsyonu ile sulanmıştır.

Epigenetik Çalışmaları

Epigenetik çalışmalarda restriksiyon enzim yaklaşımıyla DNA sitozin metilasyonları arasındaki farklılıklar incelenmiştir. Bu yaklaşımla transpozibil DNA dizilerini çoğaltan evrensel primerler kullanılmıştır (Tablo 1). Toplamda 43 tane IPBS primeri kullanılmıştır.

Tablo 1. Çalışmada kullanılan IPBS primerleri

Primer Type	IPBS Primeri	Baz dizilimi	Bağlanma sıcaklığı (°C)
IPBS	2375	TCGCATCAACCA	50-45
IPBS	2253	TCGAGGCTCTAGATACCA	
IPBS	2249	AACCGACCTCTGATACCA	
IPBS	2074	GCTCTGATACCA	
IPBS	2075	CTCATGATGCCA	
IPBS	2393	TACGGTACGCCA	
IPBS	2277	GGCGATGATACCA	
IPBS	2279	AATGAAAGCACCA	
IPBS	2218	CTCCAGCTCCGATTACCA	
IPBS	2278	GCTCATGATACCA	
IPBS	2388	TTGGAAGACCCA	
IPBS	2276	ACCTCTGATACCA	55-50
IPBS	2230	TCTAGGCGTCTGATACCA	
IPBS	2219	GAACCTTATGCCGATACCA	
IPBS	2224	ATCCTGGCAATGGAACCA	
IPBS	2228	CATTGGCTCTTGATACCA	
IPBS	2252	TCATGGCTCATGATACCA	
IPBS	2376	TAGATGGCACCA	
IPBS	2085	ATGCCGATACCA	
IPBS	2387	GCGCAATACCCA	
IPBS	2251	GAACAGGCGATGATACCA	
IPBS	2222	ACTTGGATGCCGATACCA	
IPBS	2239	ACCTAGGCTCGGATGCCA	
IPBS	2272	GGCTCAGATGCCA	

IPBS	2229	CGACCTGTTCTGATACCA	
IPBS	2083	CTTCTAGCGCCA	
IPBS	2232	AGAGAGGCTCGGATACCA	
IPBS	2389	ACATCCTTCCCA	
IPBS	2401	AGTTAAGCTTTGATACCA	
IPBS	2390	GCAACAACCCCA	
IPBS	2386	CTGATCAACCCA	
IPBS	2374	CCCAGCAAACCA	
IPBS	2226	CGGTGACCTTTGATACCA	
IPBS	2231	ACTTGGATGCTGATACCA	
IPBS	2095	GCTCGGATACCA	
IPBS	2255	GCGTGTGCTTCATACCA	
IPBS	2402	TCTAAGCTCTTGATACCA	
IPBS	2378	GGTCCTCATCCA	
IPBS	2087	GCAATGGAACCA	
IPBS	2077	CTCACGATGCCA	
IPBS	2217	ACTTGGATGTGCGATACCA	
IPBS	2225	AGCATAGCTTTGATACCA	
IPBS	2394	GAGCCTAGGCCA	

DNA İzolasyonu ve DNA Kalite/Miktarının Belirlenmesi

DNA izolasyon yöntemi Karaca vd. (2005)'e göre modifiye edilerek gerçekleştirilmiş ve elde edilen genomik DNA'nın (gDNA) kalite ve miktarı Aydın (2018)'e göre belirlenmiştir.

gDNA Normalleştirilmesi ve Restriksiyon Enzimlerin Kullanımı

Çalışmada DNA örneklerinin optimizasyon çalışmalarının sonucu elde edilen bilgiler doğrultusunda uygun miktarda DNA birbirinin izozimeri olan ve metilasyona hassas ve toleranslı olan *HpaII* (Thermo Scientific, Cat:ER0511) ve *MspI* (Thermo Scientific, Cat:ER0541) enzimleri kullanılarak gDNA ile aynı tüp içerisinde ticari firmanın protokolüne göre reaksiyonlar gerçekleştirilmiştir (Tablo 2).

Tablo 2. *MspI* ve *HpaII* Restriksiyon Enzimlerinin (RE) Metilasyon Hassasiyetleri (Xiong vd., 1999)

Target Region	<i>MspI</i>	<i>HpaII</i>	Metilasyon Durumu
5'→CCGG→3' 3'←GGCC←5'	Keser	Keser	Her iki sarmalda da metilasyon yok
5'→CC ^m GG→3' 3'←GGC ^m C←5'	Keser	Kesmez	Her iki sarmalda da sadece içteki sitozinlerde metilasyon var
5'→C ^m C ^m GG→3' 3'←GGCC←5'	Kesmez	Keser	Sadece tek sarmalda her iki sitozinde de metilasyon var
5'→C ^m CGG→3' 3'←GGCC←5'	Kesmez	Keser	Sadece tek bir sarmalda ve dıştaki sitozin metillidir.
5'→C ^m C ^m GG→3' 3'←GGC ^m C←5'	Kesmez	Kesmez	Her iki sarmalda da bütün sitozinler metillidir

Hedef Bölgelerin Amplifikasyonu ve Jel Elektroforez çalışmaları

Hedef bölgelerin amplifikasyonu için çalışmada kullanılan IPBS primerlerinin bağlanma sıcaklıkları dikkate alınarak Touch-Down PZR yöntemi kullanılmıştır. PZR sonrasında oluşmuş olan amplikonlar Agaroz Jel Elektroforez (AJE) yardımı ile yürütülerek amplikon üretilip üretilmediği dikkate alınmış ve analizler yapılmıştır (Karaca vd., 2019).


BULGULAR VE TARTIŞMA

Çalışma kapsamında toplam 43 tane IPBS primeri taranmıştır. Bunlar arasında amplikon veren 13 tane IPBS primeri (Tablo 3) ile epigenetik çalışmalarına devam edilmiştir. Geri kalan 30 IPBS primeri kullandığımız bitki metaryalinde herhangi bir amplikon oluşturmadığı için çalışma da kullanılmamışlardır. 13 tane IPBS primeri uygulama yapılan örnekler ve kontrol dahil olmak üzere toplamda 6 örek üzerinde DNA sitozin metilasyonun değişimleri incelenmiştir. Çalışma gerçekleştirilirken hiç uygulama yapılmayan kontrol sadece bir defa kullanılmıştır. Kontrol ve her uygulamada birbirinin izoşizomeri olan MspI ve HpaII restriksiyon enzimleri PZR öncesi genomun kesilmesi sağlanmış ve primerlerin bağlanma sıcaklıklarına göre PZR protokolü uygulanmıştır. Elde edilen amplikonlar % 2 agaroz jel elektroforez yöntemi ile tespit edilmiştir.

Tablo 3. Çalışmada kullanılan primerler

Primer Type	IPBS Primeri	Baz dizilimi	Bağlanma sıcaklığı (°C)
IPBS	2249	AACCGACCTCTGATACCA	50-45
IPBS	2074	GCTCTGATACCA	50-45
IPBS	2075	CTCATGATGCCA	50-45
IPBS	2393	TACGGTACGCCA	50-45
IPBS	2388	TTGGAAGACCCA	50-45
IPBS	2277	GGCGATGATACCA	50-45
IPBS	2276	ACCTCTGATACCA	50-45
IPBS	2228	CATTGGCTCTTGATACCA	55-50
IPBS	2376	TAGATGGCACCA	55-50
IPBS	2085	ATGCCGATACCA	55-50
IPBS	2394	GAGCCTAGGCCA	55-50
IPBS	2251	GAACAGGCGATGATACCA	55-50
IPBS	2232	AGAGAGGCTCGGATACCA	55-50

Şekil 1’de IPBS-2251 nolu primerde ok ile gösterilen allelerde 800-900 baz çifti (bç) arasındaki allelerde kontrol, tuz ve kuraklık uygulamalarında yüksek düzeyde metilasyon görünürken abiyotik stres uygulanmış olan örneklerde iyileştirme yapıldığında en küçük olan (800-900 bç) allelerde metilasyonun ortadan kalktığı saptanmıştır. Fakat onun bir üstündeki allelerde ise her durumda allel varlığını gösterdiğinde bunun metilasyon düzeyi ile ilgili bir yorum yapılamamaktadır. Çünkü restriksiyon enzim kesim bölgesinin varlığı ya da yokluğu ile ilgili bir bilgi mevcut değildir.



Şekil 1. IPBS-2251 pimerinin Agaroz Jel Elektroforez görüntüsü

SONUÇ

Çalışma kapsamında 43 iPBS primeri tıbbi adaçayı bitkisinde kullanılmıştır. DNA sitozin metilasyonunun farklılığını ortaya koymak ve bu primerlerin markır olarak bu çalışmada ortaya konulabilmesi için bu bitkide farklı stress uygulamaları ve iyileştirmeler yapılmıştır. Sonuç olarak amplicon oluşturan primerlerden dokuz tanesinin tıbbi adaçayı bitkisinde epigenetic çalışmalarda markır olarak kullanılabileceği tespit edilmiştir.

KAYNAKLAR

- Allis, C. D., & Jenuwein, T. (2016). *The molecular hallmarks of epigenetic control*. Nature Reviews Genetics, 17(8), 487-500.
- Bird, A. (2007). *Perceptions of epigenetics*. Nature, 447(7143), 396-398.
- Bruce, T. J., Matthes, M. C., Napier, J. A., & Pickett, J. A. (2007). *Stressful "memories" of plants: Evidence and implications for priming stress responses*. Trends in Plant Science, 12(6), 273-279.
- Chung, H., Lee, J., Kim, Y., & Park, J. (2021). *Epigenetic modifications and stress tolerance in plants*. Plant Physiology and Biochemistry, 162, 120–130.
- Craker, L. E., & Simon, J. E. (2006). *Medicinal and aromatic plants: Agricultural, commercial, ecological, legal, pharmacological and social aspects*. Springer.
- Duman, H., et al. (2005). *Türkiye Florası'nda Lamiaceae familyasının dağılımı ve çeşitliliği*. Botanical Journal of Turkey, 25(3), 305-320.
- Ekor, M. (2014). *The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety*. Frontiers in Pharmacology, 4, 177.
- Kintzios, S. E. (2000). *Sage: The genus Salvia*. CRC Press.
- Sahu, P. P., Pandey, G., Sharma, N., et al. (2013). *Epigenetic mechanisms of plant stress responses and adaptation*. Current Genomics, 14(2), 95-111.
- Schübeler, D. (2015). *Function and mechanisms of DNA methylation*. Nature, 517(7534), 321–326.
- Taiz, L., & Zeiger, E. (2010). *Plant physiology*. Sinauer Associates.
- Wang, Z., Schones, D. E., & Zhao, K. (2011). *Characterization of DNA methylation and chromatin structure interactions*. Nature Reviews Genetics, 12(7), 483–497.
- Xie, H., et al. (2018). *DNA methylation in drought stress response of plants*. Plant Molecular Biology, 96(1-2), 85-98.

FRUIT SPECIES DAMAGED BY SAN-JOSE SCULBI (QUADRASPIDIOTUS PERNICIOSUS COMST.) (HEMIPTERA: DIASPIDIDAE) SEEN IN ORCHARDS IN IĞDIR PROVINCE AND THEIR DAMAGE STATUS

IĞDIR İLİ MEYVE BAHÇELERİNDE GÖRÜLEN SAN-JOSE KABUKLU BİTİ (QUADRASPIDIOTUS PERNICIOSUS COMST.) (HEMIPTERA: DIASPIDIDAE)'NİN ZARAR VERDİĞİ MEYVE TÜRLERİ VE ZARAR DURUMU

Zir. Yük. Müh. Zekiye ALTUNTEKİN

Iğdır Provincial Directorate of Agriculture and Forestry, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0009-0008-8732-4732>

Prof. Dr. Celalettin GÖZÜAÇIK

Iğdır University, Agriculture Faculty, Department of Plant Protection, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6543-7663>

ABSTRACT

This study was carried out in orchards located in the villages of Iğdır Center between 2020-2024. Within the scope of the surveys, the fruit tree species damaged by the San Jose scale insect (*Quadraspidiotus perniciosus*) Comst. (Hemiptera: Diaspididae), the damage status it caused on the fruit tree, and the damage type on the fruit tree branches were investigated in order to observe. The biology, distribution and damage status of the pest were monitored in different types of orchards in *Q. perniciosus* non-periodic periods. When the studies were evaluated, it was determined that the pest spread in almost all apple orchards, peach orchards and plum orchards (especially in angola plums) in Iğdır province. In Iğdır province, it was observed that the San Jose scale insect spent the winter in the first larval stage on the branches, trunks, buds and twigs of the trees. With the increase in air temperature in the spring months, the larvae of the pest start from the second week of May in the central villages of Iğdır and the emergences continue until the end of July. San Jose scale insect feeds by sucking the sap of the branches, trunks, leaves and fruits of trees. It reduces the quality of the fruit due to the marks it leaves on the fruit and reduces the market value of the fruit. If the necessary measures are not taken in terms of pest control in Iğdır province, serious yield losses in apple, peach and plum trees and even drying and death of the trees will be inevitable. We believe that this study will be useful in drawing the attention of apple, peach and plum growers, which are important products of the province, and in taking the necessary pest control measures in a timely manner.

Keywords: *Q. perniciosus*, damage situation, fruit trees, Iğdır

ÖZET

Bu çalışma 2020-2024 yılları arasında Iğdır Merkez köylerinde bulunan meyve bahçelerinde yürütülmüştür. Yapılan sörveyler kapsamında, San jose kabuklu biti, (*Quadraspidiotus perniciosus*) Comst. (Hemiptera: Diaspididae)'nin zarar yaptığı meyve ağacı türleri, meyve ağacında oluşturduğu zarar durumu, meyve ağaç dallarındaki zarar şeklini gözlemlemek amacıyla

ele alınmıştır. *Q. perniciosus* periyodik olmayan dönemlerde farklı türlerdeki meyve bahçelerinde zararının biyolojisi, yayılışı ve zarar durumu takip edilmiştir. Çalışmalar değerlendirildiğinde, zararının neredeyse tüm Iğdır ili elma bahçelerinde, şeftali bahçelerinde ve erik bahçelerinde (özellikle anjola eriklerde) yayılış gösterdiği tespit edilmiştir. Iğdır ilinde, San jose kabuklu biti kışı birinci larva döneminde ağaçların dal, gövde, göz ve dalcıkları üzerinde geçirdiği görülmüştür. Bahar aylarında hava sıcaklığının artmasıyla birlikte, Iğdır merkez köylerinde zararının larva çıkışları mayıs ayının ikinci haftasından itibaren başlamakta ve temmuz ayının sonuna kadar çıkışlar devam etmektedir. San jose kabuklu biti ağaçların dal, gövde, yaprak ve meyvelerin öz suyunu emerek beslenir. Meyve üzerinde bıraktıkları izlerden dolayı meyve kalitesini düşürerek meyvenin pazar değerini azaltır. Iğdır ilinde zararlı ile ilgili mücadele yönünden gerekli önlemler alınmazsa elma, şeftali ve erik ağaçlarında ciddi oranda verim kayıpları hatta ağaçların kuruyup ölmeleri kaçınılmaz olacaktır. Bu çalışmada ilin önemli ürünlerinden olan elma, şeftali ve erik yetiştiricilerinin dikkatini çekerek, zararlıya karşı gerekli mücadele tedbirlerinin zamanında alınması konusunda yararlı olacağı kanısındayız.

Anahtar kelimeler: San jose kabuklu biti, zarar durumu, meyve ağaçları, Iğdır

DAMAGE SITUATION OF PEACH SHOOT MOTH, ANARSIA LINEATELLA (ZELLER) (LEPIDOPTERA: GELECHIIDAE) IN PEACH GARDENS OF IĞDIR PROVINCE

IĞDIR İLİ ŞEFTALİ BAHÇELERİNDE ŞEFTALİ FİLİZ GÜVESİ, ANARSIA LINEATELLA (ZELLER) (LEPIDOPTERA: GELECHIIDAE) ZARAR DURUMU

Zir. Yük. Müh. Zekiye ALTUNTEKİN

Iğdır Provincial Directorate of Agriculture and Forestry, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0009-0008-8732-4732>

Prof. Dr. Celalettin GÖZÜAÇIK

Iğdır University, Agriculture Faculty, Department of Plant Protection, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6543-7663>

ABSTRACT

Iğdır province, located in the Eastern Anatolia Region, is suitable for the growth of many cultivated plants with its microclimate feature. One of these, peach, is one of the important plants of horticultural agriculture. One of the most important pests affecting the yield and quality of this plant is the Peach Sprout Moth (SFG), *Anarsia lineatella* (Zeller) (Lepidoptera: Gelechiidae). This study was conducted in 2017 - 2018 to observe the distribution and biology of the SFG, *A. lineatella*, which is a pest on peach trees in the central villages of Iğdır province, and the damage status in peach orchards in Iğdır province. The studies were carried out in a peach orchard, locally known as zeferan, in Kadıkışlak Village of Iğdır province. In the sampling, 100 peach fruits that fell to the ground were collected randomly, and the insides of each fruit were opened and the *A. lineatella* larvae were counted to determine the damage status in peach fruits. As a result of the studies, in peach orchards in Iğdır province, peach shoot moth adults are first seen in the first week of June, the first larva in the first week of July and the second larva in the fourth week of August. In the fruits that fell to the ground, 51% of the damage caused by peach shoot moth was detected. A maximum of 3 larvae and a minimum of 2 larvae were detected in a peach fruit. In the orchard where the study was carried out, peach shoot moth adults and larvae were seen until the third week of November. It was determined that peach shoot moth caused serious damage to the late Zeferan peach variety in Iğdır province.

Keywords: *Anarsia lineatella*, damage status, peach trees, Iğdır

ÖZET

Doğu Anadolu Bölgesinde yer alan Iğdır ili mikrolima özelliğiyle birçok kültür bitkisinin yetişmesine uygundur. Bunlardan birisi olan şeftali, bahçe ziraatının önemli bitkilerindedir. Bu bitkinin verim ve kalitesini etkileyen en önemli zararlılardan biri olan Şeftali Filiz Güvesi (ŞFG), *Anarsia lineatella* (Zeller) (Lepidoptera: Gelechiidae)'dir. Bu çalışma, 2017 - 2018 yıllarında Iğdır ili merkez köylerinde şeftali ağaçlarında zararlı olan ŞFG, *A. lineatella*'nın yayılışı, biyolojisi ve

Iğdır ilinde şeftali bahçelerindeki zarar durumunu gözlemlemek için ele alınmıştır. Çalışmalar, Iğdır ili Kadıkışlak Köyü'nde yöresel ismi zeferan olan şeftali bahçesinde yürütülmüştür. Örneklemelede tesadüf olarak yere düşmüş 100 adet şeftali meyvesi toplanmış ve her bir meyvenin içleri açılarak *A. lineatella* larvalarının sayımı yapılarak şeftali meyvelerinde zarar durumunu tespit edilmiştir. Çalışmalar sonucunda, Iğdır ilinde şeftali bahçelerinde Şeftali filiz güvesi ergini ilk olarak haziran ayının birinci haftasında, ilk larva temmuz ayının birinci haftasında ve ikinci larva ise, ağustos ayının dördüncü haftasında görülmektedir. Yere dökülen meyvelerde % 51 oranında şeftali filiz güvesinin yaptığı zarar tespit edilmiştir. Bir şeftali meyvesinde en fazla 3 adet larva ve en az ise 2 adet larva tespit edilmiştir. Çalışmanın yürütüldüğü bahçede, Şeftali filiz güvesi ergini ve larvası kasım ayının üçüncü haftasına kadar görülmüştür. Şeftali filiz güvesi Iğdır ilinde daha çok geçici olan zeferan şeftali çeşidinde ciddi zararlar yaptığı belirlenmiştir.

Anahtar kelimeler: Şeftali filiz güvesi, zarar durumu, şeftali ağaçları, Iğdır

DAMAGE SITUATION OF PEACH TRUNK BLOODBITE, *PTEROCHLOROIDES PERSICAE* (CHOLODKOVSKY) (HOMOPTERA APHIDIDAE) IN APRICOT AND PEACH ORCHARDS OF İĞDIR PROVINCE

İĞDIR İLİ KAYISI VE ŞEFTALİ BAHÇELERİNDE ŞEFTALİ GÖVDE KANLIBİTİ, *PTEROCHLOROIDES PERSICAE* (CHOLODKOVSKY) (HOMOPTERA: APHIDIDAE)'NİN ZARAR DURUMU

Abdullah İREÇ

Iğdır Provincial Directorate of Agriculture and Forestry, Iğdır, Türkiye

ORCID: 0000-0003-4788-7211

Celalettin GÖZÜAÇIK

Iğdır University, Faculty of Agriculture, Department of Plant Protection, Iğdır, Turkey

ORCID: 0000-0002- 6543-7663

ÖZET

Bu çalışma, 2020 - 2024 yıllarında Iğdır ili merkez ve köylerinde kayısı ve şeftali ağaçlarında zararlı olan Şeftali gövde kanlıbiti, *Pterochloroides persicae* (Cholodkovsky) (Homoptera: Aphididae)'un yayılışı, biyolojisi ve zarar durumunu gözlemlemek amacıyla ele alınmıştır. *P. persicae* periyodik olmayan dönemlerde taş çekirdekli meyvelerden kayısı, şeftali ve erik bahçelerinde 4 yıl süresince zararının biyolojisi, yayılışı ve zarar durumu takip edilmiştir. Çalışmalar değerlendirildiğinde, Şeftali gövde kanlıbitinin son yıllarda yoğun olarak Iğdır ili kayısı ve şeftali bahçelerinde yayılış gösterdiği tespit edilmiştir. Şeftali gövde kanlıbiti haziran-temmuz aylarında ergin bireyleri ve ekim-kasım aylarında ise, yumurtaları görülmektedir. Zararlı kışı, yumurta döneminde geçirir ve genellikle kış sonuna doğru ağaçlarda yapılan budamalar sırasında, zararının gövde ve dalları üzerine gruplar halinde bırakılmış yumurtaları görülmektedir. Bu böcek kayısı ve şeftali ağaçlarının genç gövde ve dalların altına doğru olan kısımlarında koloni halinde yaşarlar ve gövdede yama gibi görünür. Şeftali gövde kanlıbitinin larvaları, genellikle gövde ve dalların güneş görmeyen gölge (kuzey) kısımlarındaki kabuğunda bitki öz suyunu emerek beslenir ve beslenme sonucunda bitkinin gövde ve dallarında sakızlanma, fumajin ve farklı deformasyonlar görülür. Özellikle Iğdır ilinde, zararının kayısı ve şeftali dallarında beslenmesi sonucunda oluşan ballımsı maddeden dolayı yabancı arılar çoğalmakta ve insanı rahatsız etmektedir. Yapılan çalışmalarda zararlı erik ağaçlarına göre, kayısı ve şeftali ağaçlarında daha fazla görüldüğü tespit edilmiştir.

Anahtar Kelimeler: *Pterochloroides persicae*, zarar durumu, meyve ağaçları, Iğdır

ABSTRACT

This study was conducted to observe the distribution, biology and damage status of Peach stem cankerbug, *Pterochloroides persicae* (Cholodkovsky) (Homoptera: Aphididae), which is harmful to apricot and peach trees in the center and villages of Iğdır province between 2020 and 2024. *P. persicae*'s biology, distribution and damage status were monitored in stone fruit apricot, peach and

plum orchards for 4 years in aperiodic periods. When the studies were evaluated, it was determined that peach stem cankerworm has spread intensively in the apricot and peach orchards of Iğdır province in recent years. Adult individuals of the peach stem cankerworm are seen in June-July and their eggs are seen in October-November. The pest spends the winter in the egg stage, and eggs laid in groups on the trunk and branches of the pest are often seen during pruning on trees towards the end of winter. This insect lives in colonies under the young trunks and branches of apricot and peach trees and appears as a patch on the trunk. The larvae of the peach stem cankerworm generally feed by sucking plant sap from the bark of the trunk and branches in the shade (northern) parts that are not exposed to sunlight, and as a result of feeding, gumming, fumagin and different deformations are observed on the trunk and branches of the plant. Especially in Iğdır province, wild bees are increasing and disturbing people due to the honey-like substance formed as a result of the pest feeding on apricot and peach branches. Studies have found that the pest is more common in apricot and peach trees than in plum trees.

Keywords: *Pterochloroides persicae*, damage situation, fruit trees, Iğdir

GİRİŞ

Şeftali yetiştiriciliği, 25°-45° kuzey ve güney enlem dereceleri arasındaki bölgelerde yapılmaktadır (Demirören, 1992). Şeftali, farklı ekolojilere hızlı uyum sağlaması, geniş bir sezon boyunca hasadı yapılan birçok farklı çeşide sahip olması, büyük kitlelerce tercih edilmesi gibi nedenlerle şeftali üretimi dünyada giderek yaygınlaşmış ve yaygınlaşmaya devam etmektedir (Şeker ve ark., 2013). Yüksek adaptasyon kabiliyetine sahip olması sebebiyle ülkemizde de Ege, Akdeniz ve Marmara bölgelerinde geniş üretim alanlarında kendine yer bulmaktadır (Ercan ve ark., 2003).

Şeftali (*Prunus persica* L.), *Rosaceae* (Gülgiller) familyasına bağlı olan *Prunus* cinsi içerisinde yer almaktadır (Deveci, 1967). Yetiştiriciliği M.Ö. 2000’li yıllara dayanan şeftalinin anavatanı Çin ve Doğu Asya olduğu bilinmektedir (Westwood, 1995). Çin’den Orta Asya’ya oradan da Anadolu vasıtasıyla Avrupa’ya yayıldığı bilinmektedir (Özbek, 1978).

Sert çekirdekli arasında en çok üretimi yapılan türlerden biri de şeftalidir. Şeftali meyvesi; tüylü (*Prunus persica vulgaris* Mill), tüysüz (Nektarin) (*Prunus persica var. nectarina* Maxim) ve domates (*Prunus persica var. platycarpa*) şeftalisi olarak üç kültür formuna ayrılmaktadır (Özbek, 1978). Meyve kabuğunun tüylü, tüysüz ve meyve şekli basıklığı itibariyle kültür formları sınıflandırılmaktadır. Üç ana kültür formu altındaki çeşitler arasında da farklılıklar gözlenmektedir. Bunlar; meyve rengi, meyve şekli, asitlik içeriği, çekirdeğin etten ayrılma durumu, derim zamanı gibi daha birçok unsuru kapsamaktadır (Ahi, 2017).

Rosaceae ailesinin önemli bir üyesi olan Kayısı (*Prunus armeniaca*) çekirdekli bir meyvedir. Türkiye, Irak, Suriye, Orta Asya, İran, Pakistan ve Afganistan gibi birçok ülkede üretimi yapılan kayısı ve ürünlerinin endüstriyel üretimi çok uzun yıllardır devam etmekte olup günümüzde gıda sektörünün önemli bir parçasını oluşturmaktadır. Vitamin, mineral madde ve zengin lif içeriğine sahip olan kayısı; yemelik, kurutmalık ve konservelik olmak üzere endüstriyel olarak farklı şekillerde değerlendirilmektedir (Özdoğan, Şen, Bilgin & Mısırlı, 2015). Meyve suyu, marmelat, reçel, pestil ve dondurma yapımında kullanılan kayısı aynı zamanda jöle, krema, şekerleme, likör, pasta ve yoğurt yapımında da kullanılmaktadır. Kayısının tatlı çekirdekleri çerez olarak tüketilmekte, acı çekirdekleri ise hammadde olarak kozmetik sanayisinde kullanılmaktadır. Söz konusu kullanım alanlarıyla önemli bir ticari ürün olan kayısı bu yönüyle yaygın ve süreklilik arz eden bir üretim ve pazar payına sahiptir (Atiş, 2017).

Iğdır ilinde yoğun bir şekilde şeftali yetiştiriciliği yapılmaktadır özellikle temmuz, ağustos ve eylül-ekim aylarında üç defa şeftali hasadı yapılmaktadır. Temmuz ayında hasadı yapılan şeftali ve nektari çeşitleri erkenci çeşitlerdir, Ağustos ayında hasadı yapılan şeftali çeşidi ise, Glohaven gibi çeşitlerdir. Eylül ve ekim aylarının sonuna kadar hasadı yapılan şeftali çeşitleri, bölgemize ait

Zeferan ve beyaz şeftali çeşitleridir. Beyaz şeftalinin yöresel ismi payız şeftalisidir. Iğdır İl Tarım ve Orman Müdürlüğü'nün 2024 yılı verilerine göre 2189 dekar alanda şeftali üretimi yapılmaktadır.

Türkiye çapında bilinen ve meşhur olan Iğdır kayısı Şalak olarak bilinen kayısı çeşididir. Şalak çeşidi dışında Tebereze, Ordubat, Ağarik ve Çır (Zerdali) kayısı çeşitleri de yetiştirilmektedir. Iğdır ilinde yetiştiriciliği yapılan kayısının büyük bir kısmı yani % 90'mı Şalak çeşididir. Iğdır ilinde 2024 yılı Iğdır İl Tarım ve Orman Müdürlüğü'nün verilerine göre yaklaşık 41.645 da alanda kayısı üretilmektedir. Ağaç başına ortalama verim bütün çeşitlerin ağırlıklı ortalaması alındığında 250-350 kg'dır.

Iğdır ilinde kayısı yetiştiriciliğinde birçok Bitki Koruma sorunu olarak başta ana zararlılar, Kayısı yaprak uyuzu (*Eriophyes similis* Nalepa) (Acarina: Eriophyidae), Şeftali filiz güvesi *Anarsia lineatella* (Zeller) (Lepidoptera: Gelechiidae), Erik koşnili (*Sphaerolecanium prunastri* (Boy.) (Hemiptera: Coccidae), Meyve ağaçları dipkurdu (*Capnodis* spp.) ve Şeftali gövde kanlıbiti, (*Pterochloroides persicae* (Cholodkovsky)'nin son yıllarda Iğdır ilinde kayısı ağaçlarında zararını dikkat çekmek için bu çalışma yapılmıştır.

Iğdır ilinde şeftali yetiştiriciliğinde birçok Bitki Koruma sorunu olarak başta ana zararlılar, San jose kabuklubiti (*Quadraspidiotus perniciosus* Comst.), Şeftali filiz güvesi *Anarsia lineatella* (Zeller), Akdeniz Meyve Sineği (*Ceratitis capitata* Wiedemann.) ve Şeftali gövde kanlıbiti, (*Pterochloroides persicae* (Cholodkovsky)'nin son yıllarda Iğdır ilinde şeftali ağaçlarında zararını dikkat çekmek için bu çalışma yapılmıştır.

MATERYAL VE YÖNTEM

Bu çalışma, 2020 - 2024 yılları Mayıs-Haziran ve Ekim-Kasım aylarında Iğdır ili merkez ve köylerinde kayısı ve şeftali ağaçlarında zararlı olan Şeftali gövde kanlıbiti, *Pterochloroides persicae* (Cholodkovsky) (Homeptera: Aphididae)'un yayılışı, biyolojisi ve zarar durumunu gözlemlemek amacıyla ele alınmıştır. Örneklemeler Çalpala (Kiti), Küllük Yukarı Çarıkçı, Oba, Kasımcan, Kadıkışlak, Necefali köylerinde ve Merkez mahallelerde, taş çekirdekli meyveler olan Şalak kayısı ve erkenci, orta gecçi, yerli Zeferan ve Beyaz (Payız) çeşitler olan şeftali ve erik bahçelerinde (Yerli ve Kültür eriklerde) *P. persicae* periyodik olmayan dönemlerde 4 yıl süresince zararının biyolojisi, yayılışı ve zarar durumu takip edilmiştir.

ARAŞTIRMA VE BULGULAR

Bu çalışma, 2020 - 2024 yıllarında Iğdır ili merkez ve köylerinde kayısı ve şeftali ağaçlarında zararlı olan Şeftali gövde kanlıbiti, *Pterochloroides persicae* (Cholodkovsky) (Homeptera: Aphididae)'un yayılışı, biyolojisi ve zarar durumunu gözlemlemek amacıyla ele alınmıştır. *P. persicae* periyodik olmayan dönemlerde taş çekirdekli meyvelerden kayısı, şeftali ve erik bahçelerinde 4 yıl süresince zararının biyolojisi, yayılışı ve zarar durumu takip edilmiştir. Çalışmalar değerlendirildiğinde, Şeftali gövde kanlıbitinin son yıllarda yoğun olarak Iğdır ili kayısı ve şeftali bahçelerinde yayılış gösterdiği tespit edilmiştir. Özbek vd. (1996), Kuzeydoğu Tarım Bölgesi'ndeki taş çekirdekli meyve ağaçlarında yaptıkları çalışmada, Iğdır ilinde Aphididae familyasından *Myzus cerasi*, *Hyalopterus pruni* ve Lachnidae familyasından *Pterochloroides persicae* türlerini tespit etmişlerdir. Hazır vd (2011), Doğu Akdeniz Bölgesi sert çekirdekli meyve türlerinde önemli düzeyde kalite ve kantite kayıplarına yol açan, şarka virüsü (Plum Pox Potyvirus)'nün bölgedeki potansiyel vektörlerini belirlemek amacıyla yaptıkları çalışmada, *Prunus* türleri üzerinde *B. cardui*, *H. pruni*, *M. cerasi* ve *Pterochloroides persicae* (Cholodkovsky 1899) türlerinden, *H. pruni* ve *P. persicae*'nin bölgede oldukça yaygın olduklarını belirlemişlerdir. Sarıbal (2019)'da yaptığı çalışmada, Bursa ilinde şeftali bahçelerinde 5 farklı yaprakbiti türüne rastladığını ve bu türler; *Brachycaudus helichrysi* (Kaltenbacl 1843), *Brachycaudus persicae* (Passerini 1860), *Hyalopterus pruni* (Geoffroy 1762), *Myzus persicae* (Sulzer 1776) ve

Pterochloroides persicae (Cholodkovsky 1899) olduğunu belirtmiştir. Şeftali gövde kanlıbiti haziran-temmuz aylarında ergin bireyleri ve ekim-kasım aylarında ise, yumurtaları görülmektedir. Zararlı kışı, yumurta döneminde geçirir ve genellikle kış sonuna doğru ağaçlarda yapılan budamalar sırasında, zararlıın gövde ve dalları üzerine gruplar halinde bırakılmış yumurtaları görülmektedir. Kışı, yumurta döneminde geçirir ve genellikle kış sonuna doğru yapılan budamalar sırasında, zararlıın gövde ve dallar üzerine gruplar halinde bırakılmış yumurtalarını görmek mümkündür (Anonim, 2025). Bu böcek kayısı ve şeftali ağaçlarının genç gövde ve dalların altına doğru olan kısımlarında koloni halinde yaşarlar ve gövdede yama gibi görünür. Şeftali gövde kanlıbitinin larvaları, genellikle gövde ve dalların güneş görmeyen gölge (kuzey) kısımlarındaki kabuğunda bitki öz suyunu emerek beslenir ve beslenme sonucunda bitkinin gövde ve dallarında sakızlanma, fumajin ve farklı deformasyonlar görülür. Özellikle Iğdır ilinde, zararlıın kayısı ve şeftali dallarında beslenmesi sonucunda oluşan ballımsı maddeden dolayı yabancı arılar çoğalmakta ve insanı rahatsız etmektedir. Yapılan çalışmalarda zararlı erik ağaçlarına göre, kayısı ve şeftali ağaçlarında daha fazla görüldüğü tespit edilmiştir.

Zararı

Çalışma sonuçlarına göre, bu böcek kayısı ve şeftali ağaçlarının genç gövde ve dalların altına doğru olan kısımlarında koloni halinde yaşarlar ve gövdede yama gibi görünür. Şeftali gövde kanlıbitinin larvaları, genellikle gövde ve dalların güneş görmeyen gölge (kuzey) kısımlarındaki kabuğunda bitki öz suyunu emerek beslenir ve beslenme sonucunda bitkinin gövde ve dallarında sakızlanma, fumajin ve farklı deformasyonlar görülür. Sarıbal (2019)'da *Pterochloroides persicae*'nin bitkilerin gövde ve dallarından öz suyu emerek onların zayıflamasına, giderek kurumalarına neden olduğunu ve zayıflama sonucu sakızlanmalar görüldüğünü ve bu yaprakbiti de fumajin oluşmasına sebep olduğunu belirtmiştir.



Ş a 1.Şeftali ağacında Şeftali gövde kanlıbiti b yumurtası ve erginleri (a), kayısı ağacında Ş a gövde kanlıbitinin yumurtaları



Şekil 2. Erik ağacında Şeftali gövde kanlıbitini (a), kayısı ağacında Şeftali gövde kanlıbitinin yumurtaları (b)



Şekil 3. Kayısı ağacının dallarında Şeftali gövde kanlıbitinin erginleri



Şekil 4. Şeftali ağacında Şeftali gövde kanlıbitinin erginleri (a), erik ağacında Şeftali gövde kanlıbitinin yumurtaları

SONUÇ

Çalışmalar değerlendirildiğinde, Şeftali gövde kanlıbitinin son yıllarda yoğun olarak Iğdır ili kayısı ve şeftali bahçelerinde yayılış gösterdiği tespit edilmiştir. Şeftali gövde kanlıbiti haziran-temmuz aylarında ergin bireyleri ve ekim-kasım aylarında ise, yumurtaları görülmektedir. Zararlı kışı, yumurta döneminde geçirir ve genellikle kış sonuna doğru ağaçlarda yapılan budamalar sırasında, zararlının gövde ve dalları üzerine gruplar halinde bırakılmış yumurtaları görülmektedir. Bu böcek kayısı ve şeftali ağaçlarının genç gövde ve dalların altına doğru olan kısımlarında koloni halinde yaşarlar ve gövdede yama gibi görünür. Şeftali gövde kanlıbitinin larvaları, genellikle gövde ve dalların güneş görmeyen gölge (kuzey) kısımlarındaki kabuğunda bitki öz suyunu emerek beslenir ve beslenme sonucunda bitkinin gövde ve dallarında sakızlanma, fumajin ve farklı deformasyonlar görülür. Özellikle Iğdır ilinde, zararlının kayısı ve şeftali dallarında beslenmesi sonucunda oluşan ballımsı maddeden dolayı yabancı arılar çoğalmakta ve insanı rahatsız etmektedir. Yapılan çalışmalarda zararlı erik ağaçlarına göre, kayısı ve şeftali ağaçlarında daha fazla görüldüğü tespit edilmiştir.

KAYNAKÇA

Ahı D., 2017. Bazı Yeni Basık Şeftali (*P. Persica* Var. *Platycarpa*) ve Nektarin (*P. Persica* Var. *Nucipersica*) Çeşitlerinde Fenolojik ve Pomolojik Özelliklerin İncelenmesi. Yüksek Lisans Tezi. Uludağ Üniversitesi Fen Bilimleri Enstitüsü, Bursa.

Anonim, 2025. <https://arastirma.tarimorman.gov.tr/bmae/Belgeler/Liflet/Zarar%C4%B1/12-20%C5%9Eeftali%20g%C3%B6vde%20kanl%C4%B1biti.pdf>. Erişim tarihi (09.03.2025).

Atiş E.,2017. Apricot Producing in Kağızman District and Its Contribution to the Economy of Territory, Marmara Geographical Review, 2017;36(36):191-205.

Deveci L., 1967. Şeftali Ziraatı, İzmir.

Ercan ve ark., 2003. Ege Bölgesinde Uygun Bazı Şeftali ve Nektarin Çeşitleri. Anadolu, J. Of Aarı 13 (2) 2003,17 – 31.

Hazır, A., Yurtmen, M., Özdemir, I. ve Aksoy, E., 2011. Doğu Akdeniz Bölgesi sert çekirdekli meyve bahçelerinde ve yabancıotlarda Aphididae (Hemiptera: Aphididae) tür kompozisyonu ve şarka virüsünün potansiyel yaprakbiti vektörleri. Türkiye IV. Bitki Koruma Kongresi, Kahramanmaraş, 28-30.

Özbek, H., Güçlü, Ş., Hayat, R., 1996. Kuzeydoğu Tarım Bölgesinde taş çekirdekli meyve ağaçlarında bulunan fitofag ve predatör böcek türleri. Turkish Journal of Agriculture and Forestry, 20, 267-282.

Özbek, S., 1978. Özel Meyvecilik. Ç.Ü.Z.F. Yayınları 128. Ders Kitabı.

Özdoğru B, Şen F, Bilgin NA, Mısırlı A.,2015. Bazı sofralık kayısı çeşitlerinin depolanma sürecinde fiziksel ve biyokimyasal değişimlerinin belirlenmesi, Ege Üniversitesi Ziraat Fakültesi Dergisi, 2015;52(1):23–30.

Sarıbal, Ç.,2019. Bursa ili şeftali bahçelerinde görülen yaprakbiti türleri (Homoptera: Aphididae) üzerine araştırmalar. Yüksek Lisans Tezi, Tekirdağ Namık Kemal Üniversitesi Fen Bilimleri Enstitüsü, Tekirdağ.27.

Şeker M., Kaçan A., Gür E., Ekinci N., Gündoğdu M., A. 2013. Çanakkale Ekolojik Koşullarında Yetiştirilen Şeftali ve Nektarin Çeşitlerinde Aromatik Bileşiklerin İncelenmesi. Tarım Bilimleri Araştırma Dergisi 6 (1): 62-67, 2013.

Westwood, M.N., 1995. Temperate Zone Pomology. Timber Pres, Inc.Oregon, p.523.

**USEFUL PROPERTIES AND USES OF SPECIES BELONGING TO THE GENERA
BARBERRY (*BERBERIS* L.) AND IRGA (*AMELANCHER* MEDIK) COMMON IN THE
AREA HAVUSH VILLAGE**

Hilal GASIMOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0009-4075-3297>

Anvar IBRAHIMOV

Nakhchivan State University, Faculty of Tip, Department of Pharmacy and biochemistry,
Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0003-1632-5259>

Mursel SEYİDOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0007-6161-9826>

Namig ABBASOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0002-2255-0826>

ABSTRACT

The article describes the use of products from berry plants of the genus Barberry (*Berberis* L.), serviceberry (*Amelancher* Medik), common in the village of Havush, Nakhchivan Autonomous Republic, and their beneficial properties. In the flora of the autonomous republic, 7 species of the barberry genus are known (*Berberis densiflora* Boiss & Buhse., *B. iberica* Stev. & Fisch.ex DC., *B. integerrima* Bunge., *B. orientalis* C.K.Schneid., *B. sphaerocarpa* Kar. & Kir. , *B. vulgaris* L.), and one of these species (*B. thunbergii* DC.) is present in cultivated flora. Only the genus *Amelancher* Medik is single-species. As a result of long-term use of the fruits of this plant, this has led to the formation of a wide variety of products. Collecting information about the variety and use of a range of products, especially from literature sources and from research conducted in different areas, provides extensive information about these plant products. In order to at least partially solve the nutrition problem, satisfy the need of every person for food and provide the body with necessary nutrients, it is very important to prepare and use wild fruits and plant fruits, especially in terms of their biologically active substances. Wild fruits and fruit plants have served as an inexhaustible source of fats, carbohydrates and proteins since ancient times and continue to be so today. People have used them in times of need, especially times of famine, as a source of food for many years. Recently, the use of the products of these plants has decreased for a number of reasons, with the exception of some products that are sold in markets in small quantities, it can be said that wild fruit and berry products are rare. Throughout the republic there is practically no

consumption of products from wild fruit plants, such as serviceberry and berberis, with the exception of a small part of the population of some mountain villages.

Keywords: Genus, Wild Fruits, Serviceberry, Barberry, Biologically Active Substances.

INTRODUCTION

The study of plant resources is essential for organizing their efficient use, developing new food products based on them, and preserving the gene pool for breeding work with cultivated plants. Depending on their degree of research and practical application, wild fruit and berry plants can be divided into three groups based on their usage and beneficial properties:

- utilized plants,
- promising plant species, and
- plants with potential opportunities.

Utilized beneficial plant species include those that are currently used by people. Promising species are those whose benefits have been identified, but due to the insufficiency or inaccessibility of natural raw material sources, as well as low plant productivity, they are not yet in use. The long-term use of certain wild beneficial plants may lead to a decrease in their natural resources. Therefore, one of the pressing tasks in plant resource management is the development and implementation of rational utilization regimes for already harvested species and their conservation. Based on these studies, recommendations should be provided regarding the organization of the raw material base, the utilization regime, and the preservation of the gene pool of species whose natural resources are declining due to harvesting. This group of beneficial plants becomes part of the utilized category once the issues related to their use are resolved or they are used as plant resources under extraordinary conditions (emergency situations). Potentially beneficial plants are species that have demonstrated certain useful properties under experimental conditions but have not yet undergone production trials. Although some of these plants are used for various purposes in certain areas, their practical applicability within the region remains unclear. With the advancement of botanical and phytochemical research, the list of beneficial plants is continuously enriched with new species, while existing knowledge about already utilized plants expands. In general, the Havush village area of the Nakhchivan Autonomous Republic stands out for its abundance of wild fruit plant resources. The wild fruit plants widely distributed in this area are of particular importance due to their richness in carbohydrates, vitamins, organic acids, proteins, and aromatic compounds (Gasimov, 2018).

Assessing the natural resources of valuable plants such as barberry (*Berberis*) and round-leaved amelanchier (*Amelanchier ovalis*) in Havush and the entire region, studying the economic efficiency of these fruit plants, and conducting research in this direction could open new opportunities for their future use. Today, there is a strong demand for ecologically clean, natural products in both local and global markets. Although some homemade products derived from these fruits can be found in local markets, their quantity is insufficient to meet demand. The fruits of these beneficial wild plants found in the region's flora are collected by the local population and used in the production of various food products, especially compotes, jams, pickles, and juices. However, the overall production volume remains relatively low. According to preliminary calculations, barberry (*Berberis L.*) is one of the plant species in the region that holds industrial significance due to its available reserves.

Additionally, many species of wild edible plants are significant due to their biomass reserves, which contain essential nutrients such as proteins, fats, carbohydrates, and vitamins. In terms of nutritional value and taste, these wild plants are not inferior to cultivated crops; in fact, some even surpass them and have no direct analogs. Given food shortages and challenges in the agricultural sector, as well as the impracticality of importing raw materials for various reasons, the use of local plant resources becomes essential for increasing food supplies (Shalpykov, et al, 2021), .

MATERIALS and METHODS

Based on various literary sources and surveys conducted in different regions of the Nakhchivan Autonomous Republic (including the villages of Bichenek and Kuku in the Shahbuz district, Keleki and Nasirvaz in the Ordubad district, and Havush in the Sherur district), sufficient material has been gathered regarding the preparation and usage of amelanchier (irqa) and barberry products. It can be stated that species belonging to the *Berberis* genus are used to some extent in the Shahbuz, Julfa, Ordubad, Sherur, and other districts. However, the round-leaved amelanchier plant is only known in the Sherur district (referred to as irqa), and information about this plant was obtained solely through surveys conducted in Havush village. Survey results indicate that barberry fruits are primarily used for medicinal purposes in the treatment of various diseases. They are widely utilized for anemia, colds, high blood pressure, vitamin deficiencies, diabetes, and kidney diseases. Additionally, high-quality and flavorful jams, compotes, pickles, and various fruit juices are made from barberry. As for round-leaved amelanchier fruits, they are only collected and consumed when fully ripe (Vasilyeva, 2003; Fattakhov, 2022).

FINDINGS and DISCUSSION

Barberry (*Berberis L.*) - Zir Belongs to the Berberidaceae Juss. family. In the flora of the Nakhchivan Autonomous Republic, six wild species of this genus can be found. The most common of these is the common barberry. Common Barberry: It is a multi-stemmed shrub that reaches a height of 2.5 meters. The stems have sharp thorns. The leaves are simple, ovate or elongated-lanceolate in shape, with finely serrated edges. The flowers are located in clusters, with each cluster containing 5-13 flowers. The length of the cluster reaches 2-3 cm. The fruit is light red, sometimes yellowish-red, with a sour taste. It blooms in April-May, and the fruit ripens in August-September. The fruit of the barberry contains up to 68.12% water, 7.01% sugar (total), 0.11% sucrose, 4.15% total acidity, 0.83% ash and colorants, 0.57% pectin, and the leaves contain vitamin C and E. It has strong astringent and purgative properties. Barberry leaves are used to prepare decoctions for kidney, liver, and bladder diseases, as well as to stop bleeding. The juice of the barberry fruit stimulates appetite and is used for dyeing wool and leather products yellow. The root and bark are used to extract the alkaloid "berberine," which is applied in the treatment of gastrointestinal diseases and postpartum hemorrhages. In folk medicine, barberry is widely used for mouth inflammation, stomach pain, and thirst. The dried fruits of barberry make an excellent spice for rice and rice-based dishes. Its berries are used to prepare vegetable and salad mixes. Barberry adds a special taste to broths and soups. Traditionally, barberry juice is used to make syrups. Barberry juice is also used to prepare various lemonades, fruit drinks, and cold teas. Fresh and dried fruits are added to soups for a sour taste and are also used as a sour seasoning in sauces (e.g., wild goose) and grilled meat (Golub, 2003).

Preparation of Barberry Products:

Barberry Juice: Fully ripe berries are selected, washed in cold water, and cleaned. The cleaned berries are crushed. A little water is added to the crushed fruit juice, and it is poured into a cloth bag to be filtered. The filtered juice is boiled lightly the next day and then filtered again. Sugar is added to taste, and it is boiled again for 3-5 minutes before being filtered. The prepared juice is stored in a cool place.

-Barberry Jam: Jam can be prepared in two ways:

- Method I: Select large ripe fruits, wash them lightly in cold water, and soak them in hot water for 8-10 hours. Prepare a syrup (1 kg of fruit, 1.5-2 kg of sugar, and 6 cups of water) and pour it over the fruit. The jam is then boiled until it is ready.

- Method II: Selected and washed large fruits are packed into glass jars. Syrup (1 kg of fruit, 1.5 kg of sugar, and 5 cups of water) is boiled and cooled, then poured over the fruit. After 24 hours, the syrup is filtered and separated from the fruit, then boiled and cooled again. The cooled syrup

is poured over the fruit and left for another 24 hours. On the third day, add 200-300 g of sugar to the fruit-syrup mixture and cook until the jam is ready.

Barberry Pickles: Ripe berries are picked in clusters, washed lightly in boiled cool water, and after draining, they are packed into jars. Salted water (3 cups of water to 100 g of salt) is poured over them and stored for later use in dishes. If mold appears in the jars, the salted water is immediately changed.

Barberry Marmalade: Water is added to the fruit and boiled well. The boiled fruit is crushed and filtered through a sieve. For every 1 kg of prepared pulp, 750 g of sugar is added, and the mixture is boiled until it becomes marmalade. The finished product can be used both fresh and preserved.

Dried Barberry: To dry the fruit, it is cleaned and left in the sun to dry. The fruit can also be dried in a cool environment at 30-35°C. Dried fruits should be stored in a dry place.

The oval-leaved Amelanchier (Amelanchier ovalis Medik.) The Oval-leaved Amelanchier (*Amelanchier ovalis* Medik.) is the only species of the genus *Amelanchier* from the Rose family (*Rosaceae* Juss.). It is a deciduous shrub with a well-developed root system, growing 1-4 meters in height. The bark of the trunk varies in color from olive to brown. The new branches are whitish, slightly covered with fuzz, while the old branches have dark brown bark. The winter buds are oval-shaped, often fuzzy, and 5-7 mm long. The leaves are oval, attached to petioles that are 8-15 mm long. The inflorescence is in the form of a raceme and blooms before the leaves appear, usually at the end of April to the beginning of May. The flowers are white, with up to 10 flowers per cluster. The Oval-leaved Amelanchier is pollinated by insects. Its fruit is small, dark blue-black, with a waxy coating, 5-15 mm in diameter, and typically ripens in September-October (Teslenko, et al, (2015).

This plant is found in rocky and stony areas and xerophytic shrubs. It can withstand frost down to -40 to -50 °C and can endure frost as low as -5 to -7 °C during its flowering period. It grows well in soils with varying compositions and acidity, preferring sunny locations. The fruit of the Oval-leaved Amelanchier contains up to 12% sugar, 0.5-1% organic acids (mainly malic acid), coumarins, sterols, tannins (up to 0.8%), flavonols (up to 40%), and various vitamins, including A, B1, B2, B5, B6, B9, B12, C, D, E, H, K, PP, as well as minerals such as K, Ca, Si, Mg, Na, P, Cl, Fe, I, Co, Mn, Cu, Mo, Se, Cr, Zn, Omega 3, 6, 9, fatty oils, and essential amino acids like tryptophan, threonine, isoleucine, leucine, lysine, methionine, cysteine, phenylalanine, and others. The berries have a sweet and astringent taste and are used when ripe. They are processed into juices, jelly, jam, preserves, compotes, liqueurs, and wines. Dried berries are a good ingredient for compotes and dry fruit jelly, giving them a beautiful color. Sometimes, dried fruits are crushed and added as a powder to sauces to provide a unique flavor (Velichko, Mashanov, 2019), .

Oval-leaved Amelanchier Juice: The berries, whether fresh, dried, or frozen, are suitable for consumption. They are also used to make jam, juice, or compote. To prepare juice, the berries are spread in a single layer on plastic or paper for a week, then pressed with a juice extractor. The extracted juice is mixed with sugar (300 g per 1 liter), heated to 85°C, poured into a sterilized glass jar, sealed, and stored in a cool place.

Oval-leaved Amelanchier Jam: To prepare jam, half a kilogram of berries requires 3 cups of water, 700 grams of sugar, and one large lemon. Water is added to the berries, the pot is covered, and it is boiled on low heat for half an hour. Then the berries are mashed, sugar and ground lemon are added, and the mixture is boiled for another half hour. After cooling, it is blended and returned to a boil before being stored in a glass jar.

CONCLUSION and RECOMMENDATIONS

People with obesity, diabetes, poor blood clotting, and skin rashes should use this plant cautiously as its berries are considered strong allergens. Like many other sour berries, it is rich in vitamin C, making it a potent natural antioxidant. This is beneficial for boosting the immune system and increasing resistance to infections and viruses. Additionally, due to the high vitamin C content, it helps prevent the development of cancerous tumors and reduces the risk of Alzheimer's disease,

particularly for older men and women. Overall, this plant acts as a natural multivitamin, strengthening the immune system and possessing calming, antitumor, antimicrobial, and antioxidant properties.

These berries are a rich source of vitamin C and therefore a powerful natural antioxidant. This is useful, first of all, for the general strengthening of the immune system and increasing the body's resistance to infections and viruses. This is good, first of all, for the general strengthening of the immune system and increasing the body's resistance to infections and viruses. In conditions of shortage of food resources and difficulties in the agricultural sector, when the purchase of imported raw materials is not advisable for a number of objective reasons, there is a need to increase food reserves using local commodity raw materials.

REFERENCES

- Fattakhov, D.A. (2022), Barberry fruits as a source of berberine // *Young Scientist*. No. 13 (408). pp. 85-89.
- Gasimov H.Z. (2018), Systematic analysis of wild fruit and berry plants distributed in the flora of the Nakhchivan Autonomous Republic.// *Scientific Works of Nakhchivan State University, Natural Sciences and Medicine Series, Nakhchivan: Geyret*, No. 3. pp. 98-105
- Golub, O.V., Poznyakovskiy V.M., Kovalevskaya I.N. (2003), Biochemical composition of Barberry berries growing in the Kemerovo region // *Storage and processing of agricultural raw materials*. No. 5. pp. 65-67.
- Shalpykov K.T., Rogova N.A., Dolotbakov A.K., Kayyrkulova A.K. (2021), Biological resources of Barberry, Rosehip, Rowan fruits in the foothill and mountainous regions of the Issyk-Kul basin in Kyrgyzstan. *Scientific Review. Biological Sciences*. No 1, pp. 16-21.
- Teslenko N.F., Krasina I.B., Bogdanov O.A., Fadeeva A.A. (2015), Irga berries as raw material for the production of marmalade / *Fundamental Research*. No. 8-2. pp. 333-337;
- Velichko N.A., Mashanov A.I. (2019), Chemical composition of Irga round-leaved fruits and the development of a recipe for an alcoholic beverage based on them. *Krasnoyarsk State Agrarian University, Krasnoyarsk, Bulletin of KrasGAU*, No. 2 (143), pp. 135-138.

NATURAL RESOURCES OF FRUIT PLANTS BELONGING TO THE GENERA APPLE (MALUS MILL.), PEAR (PYRUS L.) AND CHERRY (PRUNUS L.) FOUND IN THE HAYFIELD AND FOREST EDGE AREAS

Mursel SEYİDOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0007-6161-9826>

Hilal GASİMOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0009-4075-3297>

Namig ABBASOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0002-2255-0826>

Anvar İBRAHİMOV

Nakhchivan State University, Faculty of Tip, Department of Pharmacy and biochemistry, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0003-1632-5259>

Fidan HUSEYNOVA

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Biology major, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0002-0154-8361>

ABSTRACT

The article presents the results of a study of the natural resources of fruit plants belonging to the genera apple (Malus Mill.), pear (Pyrus L.) and plum (Prunus L.), which are found in forests and forest outskirts of Bichenek, located at the source of the Nakhchivanchay River in the region Nakhchivan Autonomous Republic. Density, that is, the number of samples per hectare, was taken as the main indicator of plant stock. In addition, in the year of observation (2024), the biological productivity of fruits was determined. In the study area, data were obtained from 10 survey sites for species of the three most common genera. For the design sites, areas were selected that contained age, origin, tree cover composition, relative density, and other taxonomic indicators that were the most characteristic feature of the work site. In the forest, fruit plants are mainly

represented on the 2nd and 3rd tiers and number 9-15 species. The study only examined wild fruit species such as apple, pear and cherry and their yields. For certain areas, the total productivity of wild fruit plants over the entire area is 105.25 kg/ha for apples, 105.58 kg/ha for pears and 395.11 kg/ha for plums.

Keywords: Species, Tier, Apple, Pear, Cherry, Yield, Density.

INTRODUCTION

The centuries-old history of fruit cultivation is rich with examples of how wild fruit plants have historically been used by humans as food sources, as well as for their medicinal properties in traditional medicine. The Nakhchivan Autonomous Republic, particularly its relatively small forested areas, is distinguished by the diversity of wild fruit and berry plants. The natural environment of this region is highly abundant in wild relatives of cultivated plants, and products derived from these fruit plants have been widely used by the population for a long time. However, in recent years, the utilization of wild fruit and berry plants has significantly declined due to various factors, leading to the underuse of valuable natural raw material resources. These uncultivated wild fruit plants hold special importance due to their rich nutritional content, serving as vital sources of vitamins, carbohydrates, proteins, organic acids, aromatic compounds, and other essential nutrients for human health. These resources should be extensively used both as food and for medicinal purposes.

Bichenek forests have significant potential to increase profitability through the economic utilization of food resources. Currently, both the region and foreign markets have a high demand for products derived from wild fruits and berries. However, the annual processing volume of forest fruits and berries remains below 5%, primarily consisting of dried rosehip fruits sold in markets. In modern times, the utilization of wild fruit and berry plants in both forested and open areas is nearly negligible. During the accounting and evaluation of forest resources, little attention is given to naturally grown fruit and berry reserves. However, recording the quantitative and qualitative characteristics of wild fruit plants is crucial in such studies. Additionally, for the efficient industrial procurement of wild fruit and berry plants in forests, it is necessary to provide theoretical and initial data on the existing utilization levels and geographic distribution of these resources. Previous studies conducted at different times have provided data on the wild fruit and berry reserves of forests and forest-adjacent areas in various districts of the Nakhchivan Autonomous Republic (Ibrahimov, E.M., 2021).

MATERIALS and METHODS

The study was conducted in September-October 2024 in the Bichenek forest and surrounding shrublands. The primary objective was to determine the reserves of wild fruit plants in these areas. The research was based on the method of establishing survey plots, following the methodologies of I.Y.Krylova and Shreter A.I. (Metod zakladki 1983, Krylova, Shreter, 1971, Krylov, 1984). In each research area, one to three replicates of survey plots were established, and a total of 10 plots, each covering 1 hectare, were created for the most common species within the three studied genera. Each survey plot was further divided into 10 smaller sections for detailed examination. The total research area covered 3,139 hectares (previously a protected area). The selection of survey plots was based on factors such as age, origin, tree cover composition, relative density, and other characteristic taxonomic indicators of the study site.

The study focused on the following species: *Malus orientalis* Uglitzk., *Prunus divaricata* Ledeb., and various species of the *Pyrus* genus, including *Pyrus salicifolia* Pall., *P. syriaca* Boiss., *P. caucasica* Fed., *P. nutans* Rubtz., *P. pseudosyriaca* Gladkova, *P. zangezura* Maleev, *P. chosrovica* Gladkova, and *P. demetrii* Kuth. (Talibov, et al, 2021).

A complete inventory of forest plants within the survey plots was conducted, and precise taxonomic indicators were determined using standard methodologies (Bun'kova, et al, 2021).

During the research, the density of wild fruit plants in the second and third vegetation layers was calculated. Additionally, their current biological productivity was determined. The heights of these fruit trees generally ranged from 2.5 to 5 meters. All trees within this height range were counted across all test sites, identified by species, and recorded Shcherbakov, et al, 1982).

The productivity of fruit trees was calculated using a model specimen selection method. Ripe fruits from selected specimens were collected and weighed. Additionally, unripe, damaged, and overripe fruits were counted. The average weight of a single fruit was determined by weighing 100 randomly selected fruits from each species. The total weight of unripe, damaged, and overripe fruits was added to the weight of ripe fruits to calculate the total natural reserve. Based on the condition and morphological characteristics of the wild fruit trees and shrubs, approximately 60% of the biological reserve was considered as the exploitable reserve. The annual procurement volume was set at 15%, taking into account ecological factors that influence plant productivity in different years. The average productivity of these plants was calculated as follows:

$$M = \frac{\sum V}{n}$$

Where:

M - average weight (productivity);

$\sum v$ - total weight of fruits collected from selected trees;

n – number of selected trees.

To determine the error in the average weight (productivity), dispersion (C) and standard deviation (σ) were calculated:

$$C = \sum V^2 - \frac{(\sum V)^2}{n}$$

$\sum v^2$ -squared weights of fruits collected from trees.

The error in deviations from the average weight is calculated as:

$$\sigma = \sqrt{\frac{C}{n-1}}$$

The magnitude error in average weight is calculated using the formula:

$$m = \frac{\sigma}{\sqrt{n}}$$

From this, $M \pm m$ is determined.

The accuracy of the experiment is determined using the following formula: $P = \frac{m}{M} 100$

FINDINGS and DISCUSSION

To determine the density and productivity of wild fruit plants in the Bichenek forest and surrounding shrubland areas, 10 random survey plots were established in the forest and shrubland areas of the Batabat region. Each survey plot covered an area of 1 hectare and was located in different areas. The survey plots were divided into smaller sections (30x20m), and the tree and shrub species found in these areas were listed according to their composition. This method allowed for the precise identification of the fruit-bearing plants within the survey plots.

In the forest and surrounding areas, the randomly selected survey plots were numbered, and the number of fruit-bearing plants within each was recorded as follows:

1. Plot No. 1 – Located in the Alesker Yurdu forest edge of the Qotursu area. Only 4 apple trees and 2 pear trees were found. The pear trees were located inside the forest, while the apple trees were at the forest edge.
2. Plot No. 2 – Located at the edge of a ravine in the Qotursu area. Only 2 cherry plum (alcha) shrubs were found inside the forest.
3. Plot No. 3 – Located in Cheper Oba. 25 cherry plum shrubs and 4 pear trees were found.
4. Plot No. 4 – Located at the edge of a ravine in Cheper Oba. 1 apple tree and 3 cherry plum shrubs were found.
5. Plot No. 5 – Located around Qibla Spring. 4 apple trees, 5 pear trees, and 9 cherry plum shrubs were found.
6. Plot No. 6 – Located inside the forest near Qibla Spring. 1 apple tree, 4 pear trees, and 7 cherry plum shrubs were found.
7. Plot No. 7 – Located in the Yukhari Pay area. 5 apple trees, 1 pear tree, and 8 cherry plum shrubs were found.
8. Plot No. 8 – Located in the Marin Post area. 2 apple trees, 6 pear trees, and 11 cherry plum shrubs were found.
9. Plot No. 9 – Located in the Marin Post ravine edge and forest clearing. 3 apple trees, 7 pear trees, and 5 cherry plum shrubs were found.
10. Plot No. 10 – Located in the Marin Post Church area. 17 cherry plum shrubs, 1 apple tree, and 3 pear trees were found.

The fruits collected from each survey plot were gathered and weighed according to standard procedures, and the productivity of each fruit tree was determined. Then, the average productivity of each type of fruit within the total area was calculated separately. The obtained results are presented in individual tables for each type of fruit.

The wild apple (Eastern apple - *Malus orientalis* Uglitzk.) is primarily distributed in sparse forests and shrublands along the forest edges, as well as in forest clearings, either as individual trees or in small clusters. Apple trees found within the forest tend to have relatively lower fruit-bearing capacity, depending on the environmental conditions they grow in.

During the study, the productivity and population of apple trees in the surveyed plots were determined (Table 1). The density of apple trees in the Bichenek forest and its surrounding areas was found to be 3.7 trees per hectare. In this region, wild apples hold industrial significance. The average productivity per hectare was 105.25 kg, and the annual procurement volume was calculated to be 8.26 tons (Table 1).

Table1.

Natural Reserve of Eastern Apple (*Malus orientalis* Uglitzk.)
in the Bichenek Area by Region

Areas and sites		Total area (ha)	Average weight of one tree's fruit (kg)	Number of trees in the area (units)	Productivity kg/ha	Natural reserves,kg		
						Biological reserve	Exploitable reserve	Annual procurement volume
Gotur su	1	1ha	31,46	4	125,84	353,56	212,14	106,07
	2	1ha	25,47	2	50,94			
Cheper obasi	1	1ha	28,35	3	85,05	437,40	262,44	131,22
	2	1ha	26,73	5	133,65			
Carvanse rai	1	1ha	25,55	4	102,20	367,0	220,2	110,1
	2	1ha	27,10	3	81,30			
Batabat	1	1ha	38,23	5	191,15	191,15	144,69	57,35
Marin post	1	1ha	26,63	2	53,26	848,37	509,02	254,51
	2	1ha	28,46	3	85,38			
	3	1ha	28,83	5	144,15			
Total average value			28,68	37	105,25	446,83	268,10	134,10

In the area, wild plum bushes are mainly distributed in forest interior areas and forest-edge thickets. These bushes are among the three layers of vegetation within the forest and are the most widespread wild fruit plants. It can be said that they are found throughout the Bichenek forest and its surrounding thickets. In many cases, they form dense, impenetrable thickets within the forest (Table 2). During the study, the productivity and annual procurement volume of the wild plum plant were determined.

Table 2.

Natural Reserve of Wild Plum (*Prunus divaricata* Ledeb.) by
Region in the Bichenek Area

Areas and sites		Total area, ha	Average weight of one tree's fruit (kg)	Number of trees in the area (units)	Productivity kg/ha	Natural reserves, kg		
						Biological reserves	Exploitable reserve	Annual procurement volume
Gotursu	1	1ha	21,16	3	42,32	178,72	107,2	53,62
	2	1ha	23,52	2	47,04			
Cheperobasi	1	1ha	27,94	26	726,44	2366,16	1419,7	709,85
	2	1ha	28,54	16	456,64			
Carvan-serai	1	1ha	27,75	19	527,25	2039,48	1220,7	611,84
	2	1ha	28,97	17	492,49			
Batabat	1	1ha	29,32	16	469,12	469,12	281,5	140,73
Marin post	1	1ha	27,45	11	301,95	2249,40	1349,6	674,80
	2	1ha	26,67	15	400,05			
	3	1ha	26,34	17	447,80			
Total average value			26,77	142	391,11	1460,58	875,8	438,17

The pear genus is highly diverse and is mainly found within forests, in forest glades, and along forest edges in thickets. In forest-edge thickets, the predominant species is *Pyrus salicifolia* Pall. This species often forms mixed thickets with *Crataegus meyeri* Pojark. and *Prunus divaricata* Ledeb. Other pear species are found only within forested areas. These species primarily make up the second layer of forest vegetation.

Table 3

The natural resources of species belonging to the genus (*Pyrus* L.) (pear) in the Bichenek area by regions.

Areas and sites		Total area, ha	Average weight of one tree's fruit. (kg)	Number of trees in the area (units)	Productivity, kg/ha	Natural reserves, kg		
						Biological reserves	Exploitable reserves	Annual procurement volume
Gotursu	1	1ha	23,13	3	69,39	229,46	137,68	68,84
	2	1ha	22,67	2	45,34			
Cheperobasi	1	1ha	24,51	4	98,04	507,8	304,68	152,34
	2	1ha	25,31	6	151,86			
Carvan sara	1	1ha	24,69	5	123,45	449,06	269,44	134,72
	2	1ha	25,45	4	101,08			
Batabat	1	1ha	23,71	2	47,42	47,42	28,45	14,23
Marin post	1	1ha	24,21	6	145,26	1257,72	754,63	377,32
	2	1ha	24,78	7	173,46			
	3	1ha	25,13	4	100,52			
Total average value			24,55	43	105,58	2491,46	1494,88	747,45

In densely shrubby areas within forests, the productivity of plants is relatively lower compared to the areas at the forest edge.

During the research, the average productivity and annual supply volume of each fruit species by area were calculated. The results of the study indicate that both pear and plum, which are widespread in the area, are of industrial significance as fruit plants. The annual supply volumes are 9.94 tons (pear) and 18.37 tons (plum), respectively.

Table 4

The natural resources of apple, pear, and plum species in the Bichenek area.

Fruit plants	Total area, ha	Productivity, kg/ha	Natural reserves, t		
			Biological reserves	Exploitable reserves	Annual procurement volume
Apple	157 ha - 5%	105,25	16,52	9,91	8,26
Pear	188 ha - 6%	105,58	19,88	11,93	9,94
Plum	470 ha - 15%	391,11	183,77	110,26	18,37

CONCLUSION and RECOMMENDATIONS

Depending on the ecological conditions of the area in different years, the productivity of these fruit plants may decrease or increase. Especially during the flowering period of the plants, continuous rain or sudden snowfalls due to accidental cold can cause a sharp decrease in productivity. On the other hand, although these wild fruit plants cover large areas in the Bichenek area, in recent times, due to various reasons, particularly zoogenic factors, their natural growth has been almost very low or not observed at all.

During the study, only the species belonging to the apple, pear and cherry genera and their productivity were studied from wild fruit plants. The total productivity of wild fruit plant species in the entire area in the identified areas is on average 105.25 kg/ha for apple plants, 105.58 kg/ha for pear plants and 395.11 kg/ha for cherry plants.

REFERENCES

- Bun'kova N.P., Zalesov S.V., Zalesova E.S., Magasumova A.G., Osipenko R.A.. (2021), *Osnovy fitomonitoringa: ucheb. posobie / 2nd ed., revised and supplemented-* Yekaterinburg: Ural State Forest Engineering University, 90 p.
- Ibrahimov, E.M. (2021), *Biodiversity of tree and shrub plants and efficient utilization methods in the forest ecosystem of the Nakhchivan Autonomous Republic. Dissertation for the degree of Doctor of Biological Sciences.*, p. 408.
- Krylov, A.G. (1984), *Zhiznennye formy lesnykh fitotsenozov / A. G. Krylov. Leningrad: Nauka*, 181 p.
- Krylova, I.Ya., Shreter A. I., (1971), *Metodicheskie ukazaniya po izucheniyu zapasov dikorastushchikh lekarstvennykh rasteniy / Moscow: VILAR*, 21 p.
- Metod zakladki Gost 56-69-83., (1983), *Ploshchadi probnye lesoustroitel'nye. Official edition.* p. 59.
- Shcherbakov N.M., Sakovets V.I., Kuchko A.A., Zaytseva N.P., Voronova T.G., Belonogova T.V., (1982), *Uchet i ispol'zovanie resursov poleznykh rastenii lesov Yuzhnoy Karelii / Petrozavodsk: Karelian branch of the Academy of Sciences of the USSR*, 38 p.
- Talibov T.H., Ibrahimov A. S., Ibrahimov A.M. (2021), *Taxonomic spectrum of the flora of the Nakhchivan Autonomous Republic (Higher plants, gymnosperms, and angiosperms).* Baku, Shirvannashr, 425 p.

**SPECIES OF THE GENUS *LONICERA* L. IN THE FLORA OF THE NAKHCHIVAN
AUTONOMOUS REPUBLIC, THEIR PHYTOCHEMICAL PROPERTIES AND
PROSPECTS OF USE (AZERBAIJAN)**

Namig ABBASOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0000-0002-2255-0826>

Hilal GASIMOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0009-4075-3297>

Mursel SEYİDOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Biology, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0007-6161-9826>

Bahruz MAMMEDOV

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Chemistry, Nakhchivan, Azerbaijan

Orcid ID: <https://orcid.org/0009-0001-9685-876X>

Semennaz ABBASLI

Nakhchivan State University, teacher, Faculty of Medicine, Department of Fundamental Medical
Disciplines, Nakhchivan, Azerbaijan

Orcid ID: <https://orcid.org/0009-0002-7238-0885>

ABSTRACT

The article provides information about the economic importance, bioecological characteristics, systematic analysis, and distribution of species from the Caprifoliaceae Adans. family found in the flora of the Nakhchivan Autonomous Republic. The majority of species belonging to the *Lonicera* genus are bitter due to the presence of apple and lemon salts in their fruit compositions, making them inedible. According to literature sources, these fruits become edible after a frost. When utilizing any part of the plant as a raw material source, it is crucial to determine not only the quantity of the compounds in the plant material but also its qualitative composition to ensure effective use of the plant. Fruits and berries are a source of biologically active substances, especially vitamins, macro, and microelements. The fruiting shrubs of the *Lonicera* genus are highly valuable because they are capable of providing the local population with biologically active and functional components. The *Lonicera* genus is also characterized by frost resistance. Its fruits are widely used for medicinal and food purposes, making these species relevant for use in

cultivation. The Caprifoliaceae Adans. family includes 13 genera and 300 species. Representatives of this family are trees, shrubs, subshrubs, and herbaceous plants. Their leaves are simple or compound and arranged alternately without stipules. Flowers are usually 4-5 members, with fused petals. The calyx is small and covered by a lower ovary, and the corolla is sometimes bilabiate, bell-shaped, or wheel-shaped. The androecium consists of 3-5 stamens fused to the corolla. The gynoecium has 2-5, sometimes 8, carpels. The ovary is inferior. The flower formula is: $Ca(4-5)Co(4-5)A3-5G(2-5)$. Flowers are gathered in cymose inflorescences. Frequently encountered species include *Lonicera bracteolaris* Boiss. & Buhse (bracteate honeysuckle) and *Lonicera iberica* Bieb. (Iberian honeysuckle). *Lonicera bracteolaris* is a shrub with a shoot covered by two adjacent scales. Its leaves are short-petioled, elliptic-lanceolate or ovate, with a heart-shaped base, clearly veined, and the underside is pubescent along the veins. The corolla is yellowish. It is found in rocky and stony areas of the middle mountain zone.

Keywords: Honeysuckle, Flora, Food Significance, Antioxidant

INTRODUCTION

The genus *Lonicera* L. is the richest genus in terms of species composition in the family Caprifoliaceae Adans. (Vent), with 200 species out of 890 species in the family. In the flora of Azerbaijan, there are 8 species of this genus (3 cultivated species), and according to recent data, 9 cultivated species exist. These include *Lonicera caprifolium* L., *L. bracteolaris* Boiss, *L. iberica* M.B., *L. xylosteum* L., and *L. caucasica* Pall. as wild species. Most of the species within the *Lonicera* L. genus are erect shrubs, while those belonging to the *Caprifolium* subgenus are climbing lianas. Species of this genus are found in various botanical-geographical regions of Azerbaijan, from lowlands (e.g., *L. caprifolium*) to the lower and upper mountain ranges (e.g., *L. iberica*), and some extend into the mid-mountain range (e.g., *L. caucasica*, *L. xylosteum*, *L. bracteolaris*). Many species of the *Lonicera* genus have significant economic importance. Most of them are considered beautiful ornamental plants. Some species of *Lonicera* have edible fruits that increase the general resistance of the body. The juice of these fruits is used as an antiseptic in the treatment of angina, tonsillitis, and stomatitis. The decoction of the leaves is used for angina and eye diseases. The majority of *Lonicera* species have bitter fruits due to the presence of apple and lemon salts in their compositions, making them inedible. However, according to literature, they become edible after a frost. Our experiments, including refrigeration and liquid nitrogen treatment, show that the bitterness disappears after freezing. When using any part of the plant as a raw material, it is crucial to determine both the quantity and the quality of the compounds present in the plant material for effective utilization. For instance, when extracting carotenoids, especially α - and β -carotenes, it indicates a high pro-vitamin A activity (Seyidov et al., 2014).

MATERIALS and METHODS

The species names have been clarified based on “Flora of Azerbaijan,” according to Aydın Asgarov, and also using newly accepted international nomenclatural sources such as World Flora Online (WFO), Plants of the World Online (POWO), and Catalogue of Life. The geobotanical research has been carried out using generally accepted methods (Lemeza, 2008). The life forms of the studied species were investigated according to Raunkiaer (1934) and Serebryakova (1964). The phytochemical properties of the species have been studied using various literature sources (Talibov, İbrahimov et al., 2021).

FINDINGS and DISCUSSION

In the flora of Nakhchivan Autonomous Republic, 5 species of the *Lonicera* genus are present (3 cultivated: *L. caprifolium*, *L. tatarica*, *L. japonica*; and 2 wild: *L. bracteolaris*, *L. iberica*).

1. *Lonicera bracteolaris* Boiss. & Buhse

2. *Lonicera caprifolium* L.

3. *L. iberica* M.Bieb.

4. *L. tatarica* L.

5. *L. japonica* Thunb.

L. caprifolium, *L. tatarica*, and *L. japonica* are used as ornamental plants in the Nakhchivan Autonomous Republic.

***Lonicera iberica* M.Bieb.- Georgian Honeysuckle**

This species is a small, freely branching shrub, reaching a height of 50-200 cm. Its leaves are small, measuring 2-3.5 cm in length, with short petioles, oval-heart-shaped or nearly circular with a pointed tip. Both sides of the leaves, especially the underside, are hairy. The bracts are lanceolate or sickle-shaped. The small bracts combine to form a covering that completely encloses the flower cup. The petals are 10-12 mm long, yellowish, densely covered with hairs, and the tube is bulging at the base. The paired fruits are partially fused, red, and 6-7 mm in diameter. Flowering Period: Flowers bloom in May (June to August), and fruits appear from July to September (as shown in the image). Distribution: In Azerbaijan, this species is found in almost all regions, up to the upper parts of the foothill and forest zones, and rarely in the subalpine zone. It is widespread in rocky and stony areas of the Pterophyllous flora, among shrubs. It was first described from Georgia.

Economic Importance: This plant has ornamental and medicinal significance. It is also recommended for use as a phytomeliorative plant to stabilize slopes (Abbasov, 2022; Pobedimova, et al., 1961).

***Lonicera bracteolaris* Boiss. & Buhse- Bracteolate Honeysuckle**

The flower pedicels are 4-10 mm long and covered with hairs. The bracts are leaf-like, measuring 6-10 mm in length, elliptic-lanceolate, and nearly equal in size to the fruit. The petals are yellowish, covered with sparse glandular hairs and hairs on the outside. The ovaries are free, not fused, and are either naked or covered with glandular hairs. The style is hairy. When the fruits ripen, they fully fuse into a single fruit.

Flowering Period: May-June

Fruiting Period: July-August (September)

Distribution: In Azerbaijan, this species is found in the southern part of the Lesser Caucasus, specifically in the middle mountain zone, in rocky and stony slopes, ravines, and among shrubs, both individually and in groups. It was first described from Azerbaijan (around Təzəkənd village in the Karabakh region). Economic Importance: It can be used for reinforcing rocky slopes in forestless areas (Pobedimova et al., 1961). Nutritional and Medicinal Benefits of Honeysuckle Berries: Honeysuckle berries are among the most effective products with clear antioxidant properties. The berries contain numerous antioxidants from various classes, such as vitamin C, polyphenols, phenolic acids, tannins, flavonoids, and flavonols.

Benefits for the Liver: Honeysuckle is rich in polyphenols, compounds found only in plants and often responsible for the color of fruits. While polyphenols are not fully studied, their strong antioxidant properties are well known. About half of the polyphenols in honeysuckle berries are anthocyanins, which give the berries their distinctive dark blue color. These anthocyanins protect the liver from non-alcoholic fatty liver disease and steatosis, which often occur in individuals with diabetes or obesity (Korobkova, et al., 2020).

Heart Health: A serving of honeysuckle berries provides the daily required intake of vitamin K, which helps the body absorb calcium efficiently. Honeysuckle also contains potassium and magnesium, two essential minerals that support normal heart rhythm. Despite these numerous

health benefits, honeysuckle should be consumed in moderation. Overconsumption can lead to constipation or excessive urination, which flushes essential nutrients from the body. Furthermore, excessive consumption of honeysuckle can cause skin rashes, muscle spasms, and digestive disturbances, especially in children. Experts recommend caution when consuming honeysuckle during pregnancy and lactation due to potential allergic reactions in mothers and infants. Additionally, honeysuckle can be harmful to individuals with high stomach acidity or duodenal diseases (Picture).

Culinary Uses: Honeysuckle berries are actively used in various culinary applications. The ripe, delicious berries are used to make jams, marmalades, juices, jellies, compotes, and kissels. Honeysuckle can also be used to prepare savory sauces, as a unique filling for baked goods, or as a base for homemade wine or liqueur. Additionally, they can be added to porridges or muesli (<https://az.wikipedia.org/wiki>).



Picture. *Lonicera iberica* M.Bieb.- Georgian Honeysuckle

According to the literature, honeysuckle berries are one of the promising sources of plant raw materials, containing various biologically active compounds such as organic acids, terpenoids, vitamins C and P, tannins, catechins, flavonoids, anthocyanins, phenolic and hydroxycinnamic acids, as well as macro- and microelements (Chulkov, et al, 2011; Bryksin, 2013).

This sour berry contains potassium, magnesium, sodium, phosphorus, iron, calcium, copper, barium, manganese, silicon, selenium, and iodine. Additionally, honeysuckle is rich in anthocyanins—1,669 mg per 100 g. Recent scientific studies have shown that anthocyanins have antioxidant properties that are 50 times stronger than vitamin C (Khokhryakova, 2004).

Honeysuckle also contains pectins, vitamins A, B1, and B2, and a high amount of vitamin C. The berry is rich in biologically active substances such as pyrocatechin, flavonol, as well as essential acids like triterpenic and chlorogenic acids. The high content of phenolic compounds in honeysuckle provides antifungal, antiviral, and antibacterial effects on the human body (Novruzov, Zeynalova, 2023).

Source of Antioxidants-Antioxidants are essential compounds that protect the body from free radicals, which oxidize cells. They help slow down the aging process, improve skin quality, reduce inflammation, and lower the risk of developing severe diseases associated with aging. In the Nakhchivan Autonomous Republic, extracts of *Lonicera iberica* berries prepared in 96% alcohol are used in traditional medicine for the treatment of upper respiratory tract infections and bronchial asthma, showing highly effective therapeutic results. Therefore, the use of this plant in traditional medicine is recommended. Additionally, honeysuckle has value as an ornamental plant. Considering these valuable properties, it is advisable to preserve the species *ex situ* to protect its gene pool. Phenological Groups and Introduction Success Introduced honeysuckle species are classified into six phenological groups based on the start and end of their growing season. These groups are distributed according to the plants' resistance to unfavorable conditions. The obtained results confirm that the timing of the vegetation period's completion is significant for determining the success of the introduction process. There is a direct correlation between the distribution of phenological groups and winter hardiness. Honeysuckle species that complete their vegetation cycle early or in the middle of the season exhibit the highest winter resistance. Based on flowering time, three phenological groups were identified: early, mid-season, and late bloomers.

CONCLUSION and RECOMMENDATIONS

The economic value of the species *Lonicera bracteolaris* Boiss., which is widespread in the flora of the Nakhchivan Autonomous Republic and belongs to the *Caprifoliaceae* Adans. family, the importance, bioecological characteristics, systematic analysis and distribution have been studied. Fruiting periods and fruit ripening periods have been determined, which allows predicting optimal seed collection times. Introduced plant species can be recommended for landscaping as either spring or early summer flowering species. Species based on decorative characteristics allow for selection for landscape design.

REFERENCES

- Abbasov N.K. (2022), Some species of the petrofit (rock-debris) flora of the Nakhchivan Autonomous Republic used in medicine and folk medicine. Nakhchivan State University. / Republican scientific conference on "Issues of Application of Alternative Medicine Methods", Nakhchivan, pp. 231-236.
- Bryksin D.M. (2013), Berry cultivation in northern China // *Gardens of Russia*. No. 12. pp. 45–48.
- Chulkov A.N., Gostishchev D.A., Deineka V.I., Pisarev D.I., Sorokopudov V.N., Sazonov S.A. (2011), Blue honeysuckle fruits as a source of anthocyanins // *Chemistry of Plant Raw Materials*.
- Khokhryakova L.A. (2004), Economic and biological evaluation of honeysuckle varieties and selected forms in the forest-steppe zone of the Altai Krai: Abstract of dissertation ... Candidate of Agricultural Sciences. Barnaul, 20 p.
- Novruzov E.N., Zeynalova A.M. (2023), Study of the chemical composition of *Lonicera caucasica* Pall. // Ganja State University, International scientific conference (May 5-6), Modern issues of natural and economic sciences, Part III, Ganja, pp. 30-33.
- Pobedimova E.G., Prilipko L.P. (1961) Family *Caprifoliaceae* Vent. *Flora of Azerbaijan*, Vol. VIII, pp. 60-64.
- Seyidov M.M., Ibadullayeva S.C., Qasimov H.Z., Salayeva Z.K. *Flora of Shahbuz State Reserve*. Nakhchivan: Əcəmi, 2014, 524 p.
- Talibov T.H., Ibrahimov Ə.Ş. (2021), Taxonomic spectrum of the flora of the Nakhchivan Autonomous Republic (Higher spore-bearing, gymnosperms, and angiosperms). 2nd edition. Baku: Shirvannashr, 426 p.

**SPIDER MITE - *TETRANYCHIDAE* IN THE VINEYARDS OF THE NAKHCHIVAN
AUTONOMOUS REPUBLIC OF AZERBAIJAN AND MEASURES TO FIGHT
AGAINST IT**

Mirmahmud Seyidli

Nakhchivan State University, Faculty of Natural Sciences and Agriculture, Department of
Veterinary Medicine, Nakhchivan, Azerbaijan

ORCID ID: <https://orcid.org/0009-0003-6595-1245>

Jabbar Najafov

Nakhchivan State University, Faculty of Engineering and Architecture, Department of
Reclamation and Environmental Engineering, Department of Agricultural Sciences, Nakhchivan,
Azerbaijan

ORCID ID: <https://orcid.org/0009-0009-9343-2067>

ABSTRACT

Thor genes (lat. Tetranychidae) are used in vineyard crops, tarevez plants, paxlas, etc. They are polyphagous plants that harm plants, some rose plants, and also grapes. They are small arthropods, 0.3-0.7 mm long. They can be red, brown or yellow-green. While it is being consumed, it secretes many different types of tor, which is why it got the name of the family. Typically, colonies of pests settle along the veins of the leaf palm [1, 2, 3, 4, 5].

These are small plants of various lengths, ranging from yellowish to dusky red. Nails first have 6 legs and later have 8 legs. They form a more or less dense cover at the bottom, where the grape leaves are permanently settled. Each year they give several generations (4-7) depending on the temperature conditions. Dangerous mass attacks of tor particles are possible under favorable conditions. Essentially, the soil structure develops more quickly when there is a high degree of moisture in the soil and dryness in the air. It overwinters under the bark of the general grain's plurality stock [8,9].

Signs of damage. The common Tor gene is harmful at all stages of its development (with the exception of wintering females). The first signs of infection are small yellow spots that turn into large net spots. In the month of July, the leaves stop developing and die. The female secretes the gene tubes, providing a good environment for the growth of the nits. As the eyes open, the rate of multiplication increases. When this pest captures a branch, its leaves die completely, resulting in the plant stopping feeding (Picture 1-2).

The egg is light colored and semi-transparent. Nails have three legs and big ones have four legs. It differs from other species of spiders by the shape and size of the male reproductive organ. Individual development includes egg, nymph, protonymph, deytonymph and stages of competence.

Key Words: pests, spider mites, leaf, sap, grape, herbicide



Picture 1.



Picture 2.

Introduction and Purpose

The life cycle of the Tor gene. Females overwinter in the internal structures of shelters, in fallow grass, in the soil substrate and in other shelters, on roofs (table 1). They come out of diapause as the soil settles and at the same time the temperature and humidity rise. The females settle on the underside of the leaves, stay there for 3-4 days, then change color to green and lay eggs. The spawning cycle continues for 25-30 days in moderate temperatures, and the female lays approximately 80-100 eggs. The period of development of the generation is mainly determined by the temperature and is 8-10 days under favorable conditions (25-27⁰ C). The development of thorax genes in the autumn-winter-summer period (until mid-April) and the development of nymphs and nymphs on the life-cycle plants under appropriate conditions of photodryer, temperature and food quality. It is determined by the living conditions. Diapaused females develop and lay eggs under conditions where the photoperiod is not more than 14-16 hours and the temperature is not more than 19 °C. Since the temperature changes a lot during the winter and summer periods, some females stay in the cages and lay eggs, while others go into diapause, where they spend 10-30 days or more at a temperature of 22-25 C. They can be prevented from falling into diapause by either providing additional lighting (more than 16 hours) or maintaining a temperature above 21 °C. In exceptions, the common grass gene can give more than 20 generations [6, 10].

Table 1	
Manual classification	
Classification	Living Creatures
Type	Jointed foot people
Class	Spider-like
Group	Thrombidiform genera in the deck
Family	Tetranychidae

Spread and toxicity. The harm has spread widely. Injured plants die due to lack of moisture.

Factors that aid the development of Tor genes:

The fastest development and multiplication of the genus occurs at a temperature of 29–31°C and a relative humidity of 35–55%. Ways of struggle [11]:

- Fight against weeds
- Biology combat method.
- Application of chemical plant protection devices.

Materials and Methods: During the research work, 20 aboriginal table and technical grape varieties cultivated during 3 vegetation periods using methods for studying spider mites living in plants [7] were used in the field collection vineyard of the Nakhchivan Autonomous Republic, which was established by us in the Botanical Experimental Area of the Institute of Bioresources (Nakhchivan) in 2019-2022, in order to detect spider mites.

Results: There are two types of spider mites that pose a threat to viticulture. The red fruit mite (*Panonychus ulmi* Koch), which has a low web density, overwinters in the egg stage on the food plant. The inconspicuous red bulbous winter eggs are first laid in the nodes of young shoots, often in such numbers and so close together that they give the impression of a reddish coating and turn the fingers blood red. When pruning grape bushes and crushing shoots, egg "clusters" are formed. Summer eggs are somewhat smaller and more uniform in shape and color and are collected on the underside of the leaves. The common spider mite (*Tetranychus telarius* L.), unlike the red mite, overwinters as a fertilized female not only in the bark of the vine, but mainly in fallen grape leaves and weeds. Eggs laid during the growing season are initially light in color, becoming yellow and orange as they develop (Pic. 3-4).



Picture 3.



Picture 4.


In our studies, we found that sometimes the vine is also damaged by the yellow spider mite (*Eotetranychus carpini* Oud.), whose females hibernate in the bark. In accordance with the hibernation method, we found newly emerged red mite larvae in early spring in the form of small moving red dots on the underside of young leaves. According to our observations, the spider mite first breeds en masse on low weeds, from there it moves to the vines, and then appears on the grape vine. The spider mite pierces individual surface cells with its pointed upper jaws and sucks out the contents. They cause damage not only with this, but also with the toxic effect of their saliva.

The damage to the leaves on the vine by spider mite should not be confused with the damage of other mites. As a result of the breakdown of chlorophyll, irregularly defined bright spots are first clearly visible, especially in the light. In the center of these lesions, small brown dot-like areas develop as a result of the drying of damaged cells, but they are noticeable without the radially separated light veins, as in *Calepitrimerus vitis* mites. In addition, the web mites can be clearly seen, while *Calepitrimerus vitis* mites are not visible with an ordinary magnifying glass.

When a vineyard is infected by mites early, shortly after the start of growth, the leaves remain small and are deformed. When shoots form from the replacement buds, the vine may take on a broom-like appearance. Damaged leaves often curl, and typical gaps between the veins appear next to the damaged one, with further growth of healthy tissue. Eventually, all leaves turn light brown and fall prematurely.

By our observations and using a light microscope, we have determined that spider mites appear in both young and old vineyards, and often on the vines. The degree of damage depends on the varietal characteristics, soil-climatic conditions, the nutritional status of the bushes, as well as the

relief (Pic. 5-6). In dry, warm and wind-protected areas, spider mites are more likely to infect vineyards than in vineyards with a humid and relatively cool microclimate. Accordingly, dry and warm weather promotes the development of spider mites, while humid and cool weather delays it.



Picture 5.



Picture 6.

Discussion and Conclusion: The spider mite has many natural enemies. These are, first of all, the multi-voracious predatory mites of the genus *Typhlodromus*, then the ladybugs *Scymnus punctillum* W. and their larvae, as well as some predatory flower beetles. However, they cannot stop the infestation of spider mite quickly enough to prevent serious damage.

Before the buds open, the vineyard should be treated with DNOC or nitrofen. After the start of growth in the spring, various active substances (Ragor, Fozalon, Kral, etc.) can be used to combat spider mite. When using short-acting drugs, spraying should be repeated after 10-12 days after the eyes open, so that the mites that appear later are also destroyed before the start of summer egg laying. As a preventive measure: pruning heavily infected bark and removing it from the vineyard and burning it.

The sugar content in the canes damaged by the spider mite is 2-3 percent lower. The leaves of the damaged canes are deformed and fall off. Green pruning and tying should be carried out as an agrotechnical control. As a chemical control, in the spring, when the shoots are 7-12 cm long and 3-5 leaves have opened, 0.1 percent Ragor and thiophos should be used.

Spider mites can be immune to a certain active substance. In this case, the so-called group immunity can occur relatively quickly, that is, this happens when similar active substances lose their effectiveness. To prevent this, spraying should be carried out alternately between different active substances only in very necessary cases.

Conclusion and recommendations

- All varieties are infected with spider mite at an almost constant level. The varieties Khanimi and Shahtakhti are not infected at all, while the varieties Sarı aldera, Ağ aldera, Ağ kalampur, Nakhchivan Khatunişi, Khatmi, Tula gezü, Tülku koyruğu are infected with the spider mite at the level of 2 points, and the remaining varieties are infected with the spider mite at the level of 1 point (table 2);

Table 2

The degree of infection of rare valuable grape varieties of the Nakhchivan Autonomous Republic with the spider mite

№	Varieties	Degree of infection with the spider mite (according to the method of P.H. Naydenova, with a 5-point system)
Table varieties		
1.	Bandi (control variety)	1
2.	Duzali	1
3.	Khanimi	0
4.	Khazani	1
5.	Gara Kurdashi	1
6.	Gara Khalili	1
7.	Golden grape (Nakh)	1
8.	Nakhshabi	1
9.	Sahibi	1
10.	Yellow aldera	2
11.	Shahangiri	1
Technical varieties		
1.	White aldera	2
2.	White kalempur	2
3.	Jalali	1
4.	Dash kara	1
5.	Khatini (Nakh)	2
6.	Khatmi	2
7.	Shahtakhti	0
8.	Talibi	1
9.	Tula gezii	2
10.	Tulku keirugu	2
11.	Zalkha	1

- The spider mite can be controlled both with chemical drugs and by multiplying their natural enemies in the vineyard, but monitoring and control should be carried out at least once a week;
- Careful and timely monitoring and control can reduce the number of chemical sprayings and minimize environmental pollution, and protect biodiversity, thereby allowing organic production at low cost;

Literary sources

1. Haji Shikhlinski. Grape diseases, pests and methods of combating them. Baku: Azernashr, 2004, 134 p.
2. Gurbanov M.R., Najafov.C.S. Resistance of rare grape varieties of the Nakhchivan Autonomous Republic to pests / Works of the Society of Zoologists of Azerbaijan, Volume I, Baku, Elm, 2008, pp. 788-790

3. Jabbar Najafov, Mirmahmud Seyidov. Life cycle of grape varieties in the Nakhchivan Autonomous Republic - grape leaf beetle (*lobesia botrana*), traditional and progressive control measures // Scientific works of Nakhchivan State University, ISSN 2223-5124, Natural and medical sciences series, Nakhchivan, 2023, No. 3(124), pp. 19-23
4. I. B. Gorbachev, V. B. Gritsenko, Yu. A. Zakhvatkin and others. Under the editorship. prof. B. B. Isaicheva / Protection of plants from pests, M. Kolos, 2002, 472 p.
5. Radzhabov G.R., Aliyeva B.A. Integrated protection of plantations of intensive type from pests and diseases // Vinograd and wine of Russia, 1996, special edition, p. 25-27
6. Kuznetsov N.N., Silakov V.V. Identification of predatory mites and their use in biological control of mites - pests of grapevines in Crimea. Yalta: Adonis, 2001, 16 p.
7. Methodological recommendations for the study of plant-eating mites. Yalta, NBS, 1986, 48 p.
8. <http://vinograd.info/>
9. CERUTTI A.K., BRUUN S., BECCARO G.L., BOUNOUS G. (2011) A review of studies applying environmental impact assessment methods on fruit production systems. J. Environ. Manag., 92, 2277–2286
10. ROCKSTRÖM J., WILLIAMS J., DAILY G., NOBLE A. (2017) Sustainable intensification of agriculture for human prosperity and global sustainability. *Ambio*, 46, 4–17
11. SCHAU E., FET A. (2008) LCA studies of food products as background for environmental product declarations. *Int. J. Life Cycle Ass.*, 13, p. 255–264

ASIAN CITRUS PSYLLID, *Diaphorina citri* Kuwayama (HEMIPTERA: PSYLLIDAE), AN IMPORTANT QUARANTINE PEST CAUSING HIGH LOSSES IN CITRUS FRUIT

TURUNÇGİLLERDE BÜYÜK KAYIPLARA YOL AÇAN ÖNEMLİ BİR KARANTİNA ZARARLISI ASYATURUNÇGİL PSİLLİDİ, *Diaphorina citri* Kuwayama (HEMIPTERA: PSYLLIDAE)

Hava ULUĞBEY

Ağrı Ziraî Karantina Müdürlüğü, Ağrı, Türkiye

ORCID ID: <https://orcid.org/0009-0008-4547-7802>

İsmail ALASERHAT

Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, 76000, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6929-0179>

ÖZET

Asya turunçgil psillidi [*Diaphorina citri* Kuwayama (Hemiptera: Psyllidae)], turunçgil endüstrisinin önemli bir zararlısı ve aynı zamanda turunçgil ağaçlarının yıkıcı hastalığı olarak bilinen Huanglongbing (HLB) hastalığı olan *Candidatus liberibacter asiaticus*'un (CLas) vektörüdür. Zararlı European and Mediterranean Plant Protection Organization (EPPO)'in A1 listesinde yer almaktadır. Türkiye'de Bitki Karantinası Yönetmeliği'nin Ek-1/A listesinde yer almaktadır. Zararlı'nın konukçuları turunçgil (*Citrus*) türleri, portakal çiçekli yasemin (*Murraya paniculata*) ve Rutaceae familyasına ait bazı diğer bitki türleridir. *Diaphorina citri* uygun koşullarda yılda 8- 10 döl verir. Yumurta ve nimfler yalnızca taze sürgünlerde bulunur. Bu nedenle kontroller taze sürgünlerde yapılır. En uygun gelişme sıcaklığı 25-28°C'dir. Erginler beslenme ve yumurta bırakma yeri aramak için aktif olarak hareket ederler. Kısa mesafelerde uçuş yaparlar. Beslenmek için ağız parçalarını bitki dokusuna sokup emerek zarar yapar. Doğrudan beslenme zararı çok önemli değildir. Zararlı, Amerika, Asya ve Afrika'da turunçgil üretiminde büyük kayıpları neden olan Turunçgil yeşillenme hastalığı *Candidatus Liberibacter* spp.'nin vektörü olduğu için önemlidir.

Bu çalışmada, *Diaphorina citri* ile ilgili şüana kadar yürütülmüş çalışmalar incelenerek, zararlı'nın morfolojisi, biyolojisi, zarar şekli, yayılışı, konukçuları, doğal düşmanları ve mücadelesi ile ilgili bilgiler derlenmiştir. Böylelikle zararlı'nın dünyadaki mevcut durumu ortaya çıkarılmıştır.

Anahtar Kelimeler: Asya turunçgil psillidi, *Diaphorina citri* Kuwayama, Zararlı, Karantina, Mücadele

ABSTRACT

The Asian citrus psyllid [*Diaphorina citri* Kuwayama (Hemiptera: Psyllidae)] is a major pest of the citrus industry and also the vector of *Candidatus liberibacter asiaticus* (CLas), known as Huanglongbing disease (HLB), a devastating disease of citrus trees. The European and Mediterranean Plant Protection Organization (EPPO) lists the pest on the A1 list. It is listed in Appendix 1/A of the Plant Quarantine Regulation in Türkiye. Host plants are citrus (*Citrus*)

species, orange-flowered jasmine (*Murraya paniculata*), and a few other plant species belonging to the Rutaceae family. Under suitable conditions, *D. citri* is produce 8-10 generations per year. Eggs and nymphs are found only on fresh shoots. This is why the controls are carried out on fresh shoots. The most suitable temperature for developing is 25-28°C. Adults move actively to find places to feed and lay eggs. They fly short distances. By inserting their mouthparts into plant tissue and sucking to feed, they cause damage. Direct feeding damage is not very significant. The pest is important because it is the vector of citrus greening disease (*Candidatus Liberibacter* spp.), which causes major losses in citrus production in the Americas, Asia and Africa.

This study reviewed previous studies on *Diaphorina citri* and compiled information on the pest's morphology, biology, damage type, distribution, hosts, natural enemies and control. Thus, the current status of the pest in the world was revealed.

Key Words: Asian Citrus Psyllid, *Diaphorina citri* Kuwayama, Pest, Quarantine, Management

GİRİŞ

Turunçgiller yapraklarını dökmeyen ağaçlardan oluşan Rutaceae familyasının Aurantioideae alt familyasının *Citrus* cinsine giren bitkilerdir. Turunçgiller, içerdiği besin değeri ve beslenme piramidindeki yeri nedeniyle insan beslenmesinde önem arz etmektedir. Turunçgil üretimi, dünya genelinde ekonomik açıdan gelişen bir sektördür. Üretimdeki artış, sektördeki istihdamı da arttırmaktadır. Ayrıca, turunçgil ürünleri sağlık açısından da oldukça faydalıdır. İçerdiği C vitamini sayesinde bağışıklık sistemini güçlendirmekte ve birçok hastalığa karşı koruma sağlamaktadırlar. Bunun dışında antioksidan özellikleri sayesinde hücreleri korur ve yaşlanmayı geciktirdiği bilinmektedir. Taze tüketimin yanı sıra, işlenmiş olarak (meyve suyu, reçel ve marmelat) ve kozmetik sektöründe ham madde olarak da kullanılabilir. Dünya çapında ekonomik öneme sahip beş turunçgil (*Citrus*) türü vardır. Bunlar; portakallar (*Citrus sinensis*), mandarinler (*C. reticulata*), altıntoplar (*C.paradisi*), limonlar (*C.limon*) ve Misket limonlar (*C.aurantifolia*)'dır. Ayrıca ekonomik değeri olan türlerin dışında şadok, ağaç kavunu, bergamot gibi diğer türleri de içerisinde bulundur.

Turunçgiller anavatanı Çin, Güneydoğu Asya ve Hindistan bölgesi olup, ılıman iklime sahip hemen hemen tüm ülkelerde yetiştiriciliği yapılabilmektedir (TEPGE, 2023). Dünya toplam turunçgil üretiminde (106.897.255,75 ton) Çin ilk sırada yer almakta; bunu Brezilya, Hindistan, Meksika, ABD, İspanya ve Mısır izlemekte, Türkiye ise 4.640.642 ton üretim ile 8. sırada yer alır (FAO, 2023). Türkiye, bulunduğu coğrafi konum itibarıyla tarımsal ürünlerin üretiminde avantajlı bir konuma sahiptir. Turunçgiller, çok yıllık ve soğuk iklim koşullarına dayanıksız bitki türleridir. Akdeniz havzası içerisinde yer alan ülkemiz, kaliteli turunçgil üretimi bakımından uygun ekolojik koşullara sahiptir (Tuzcu, 1998). Özellikle de Akdeniz Bölgesi turunçgil üretimi açısından diğer ülkelerle kıyaslandığında önemli bir potansiyele sahiptir. Ekonomik getirisinin fazla olması nedeniyle son yıllarda ülkemizde hızla artan bir turunçgil üretimi söz konusudur. Ülkemizdeki turunçgil üretiminin yaklaşık %85'i Çukurova ve Antalya çevresinde, %14'ü İzmir ve çevresinde, %1'lik bölümü ise ağırlıklı olarak Doğu Karadeniz Bölgesi'nde gerçekleştirilmektedir. Üretim tür bazında incelendiğinde ise Türkiye'de turunçgil türleri arasında en fazla mandarin (1.988.395 ton) üretimi yapılmakta, bunu sırasıyla limon (1.730.114 ton), portakal (1.610.356 ton), greylort (150.140 ton) ve turunç (3.627 ton) izlemekte, toplamda ise 5.482.632 ton turunçgil elde edilmektedir (TÜİK, 2024).

Dünya genelinde turunçgil bahçelerinde zararlı olan türler hakkında yapılan önceki çalışmalarda, turunçgillerle ilgili olduğu bilinen yüzden fazla zararlı tür rapor edilmiş, ancak sınırlı sayıda türün ekonomik anlamda önemli olduğu vurgulanmıştır (Jeppson et al., 1975; Uygun ve ark., 2001; Fadamiro et al., 2013; Donkersley et al., 2018; Urbaneja et al., 2020; Döker ve ark., 2021a).

Türkiye'de turunçgil yetiştiriciliği yapılan alanlarda bugüne kadar 90'ın üzerinde zararlı, 34 hastalık, 16 nematod ve 155 yabancı ot türü tespit edilmiştir. Turunçgil zararlılarının tamamının

ekonomik düzeyde zarar oluşturmadıklarını, bugüne kadar tespit edilmiş olan bu türlerden ancak 17'sinin turunçgil alanlarında ekonomik önemde kayıplara neden oldukları belirtilmiştir. Bu zararlıların başında; Turunçgil kırmızı kabuklubiti, *Aonidiella aurantii* (Maskell) (Hemiptera: Diaspididae), Turunçgil unlubiti, *Planococcus citri* Risso (Hemiptera: Pseudococcidae), Akdeniz meyve sineği, *Ceratitis capitata* Wiedman (Diptera: Tephritidae), akarlar ve yaprakbitleri yer almaktadır (Günaltın, 2024).

Dünyada hızla artan ticaret hacmi beraberinde zararlı böcek türlerinin taşınmasına ve karantina zararlıları listesinin uzamasına neden olmaktadır. Yeni alanlara yayılan istilacı türler bu alandaki yerli bitki türlerini tehdit etmektedir. Bu türlerden birisi olan Asya Turunçgil Psillidi, *Diaphorina citri* Türkiye'de Bitki Karantinası Yönetmeliği'nin Türkiye'de Varlığı Bilinmeyen ve İthal Mani Teşkil Eden Karantinaya Tabi Zararlı Organizmalar listesinde (Ek-1A) yer aldığı ve Amerika, Asya ve Afrika'da turunçgil üretiminde büyük kayıpları neden olan Turunçgil yeşillenme hastalığı *Liberibacter asiaticus*'un vektörü olduğu bilinmektedir (Keçeci, 2022). Bunun ülkemize bulaşması durumunda ülkemiz ekolojik koşulları da göz önüne alındığında zararının yüksek popülasyon düzeyine çıkarak yıkıcı etkileri olabileceği; ağaç ölümleri ile beraber büyük miktarda ürün ve ihracat pazarı kaybına neden olacağı açıktır. Bu zararının ülkemize giriş yapması durumunda turunçgil üretimimizi yaklaşık %40 azaltabileceği (GKGM, 2024), üretimde oluşacak bu azalma nedeniyle de üreticilerin turunçgil üretimini terk edebileceği ve ülkemizin turunçgil ihraç eden ülke konumundan turunçgil ithal eden ülke konumuna gelebileceği değerlendirilmektedir.

Bu çalışmada, Asya turunçgil psillidi, *D. citri* ile ilgili şuna kadar yürütülmüş çalışmalar incelenerek, zararının morfolojisi, biyolojisi, zarar şekli, yayılışı, konukçuları, doğal düşmanları ve mücadelesi ile ilgili bilgiler derlenmiştir. Böylelikle zararının dünyadaki mevcut durumu ortaya çıkarılmış risklerinden ve mücadele tedbirlerinden bahsedilmiştir.


Asya Turunçgil Psillidi, *Diaphorina citri* Kuwayama'nın Morfolojisi, Biyolojisi ve Zarar Şekli

Diaphorina citri'nin yumurtaları 0.3 mm uzunluğunda, 0.14 mm genişliğinde, badem şeklindedir. Yumurtalar, uzun ekseni yüzeye dik olacak şekilde bitki dokusu üzerine yerleştirilir (Şekil 1a). Yeni bırakıldığında açık sarı renkte iken olgunlaştığında iki belirgin kırmızı göz lekesiyle parlak turuncuya dönerler. Nimfler de sarımsı renktedir ve yetişkinliğe geçmeden önce beş nimf dönemi geçirirler. Birinci dönemde 0,3 mm uzunluğunda ve 0,17 mm genişliğinde, 5. dönemde ise 1,6 mm uzunluğunda ve 1,0 mm genişliğindedirler. Nimflerde kanatları oluşturacak yapılar ve antenler ilerleyen dönemlerde belirginleşir (Şekil 1b). Erginler yaklaşık 2-4 mm uzunluğunda kanat kenarları koyu renkle çevrili olup uçlarında gri açıklıklar vardır (Şekil 1c). Antenlerinin uçları ise siyah renk ile son bulup, anten ortasında kahverengi iki nokta bulunur (Anonim, 2025).

Yumurtalar, taze sürgünlerin uçlarına ve yeni açılan yaprakların arasına bırakılır. Taze sürgünlerde çok sayıda yumurta bulunur. Optimum iklim şartlarında ergin dişiler ortalama 4 hafta yaşar. Bu süre boyunca ortalama 800 yumurta bırakır. Nimfler yalnızca genç yaprak ve sapsarla beslenirler. Yumurta açılımı 25°C'de 4 gündür. Nimfler 13 günde ergin olur. Toplam gelişme süresi (yumurtadan ergine) 17 gündür. Bu süre 28°C'de 14 gün, 15°C'de 49 gündür. Yumurtadan sonra 5 nimf dönemi geçirerek ergin olurlar. Diyapoz girmezler, ancak kışın popülasyonları düşüktür. Uygun koşullarda yılda 8-10 nesil verirler. Optimum sıcaklıkta (20-25°C) tüm yaşam döngüsü yaklaşık 14-50 gün sürer, bu da yılda 9-14 örtüşen nesil potansiyeli anlamına gelir (Liu and Tsai, 2000).

Diaphorina citri'nin başlıca ekonomik önemi, *Liberibacter asiaticus* bakterisinin neden olduğu çok ciddi turunçgil *Huanglongbing* (yeşillenme) hastalığının vektörü olmasıdır (EPPO/CABI, 1996). *Diaphorina citri* ticari turunçgil ağaçlarında genellikle küçük hasara neden olan ikincil bir zararlı olarak kabul edilir. Asya turunçgil psillidi, floemle beslenen bir böcektir. Tomurcuk patlaması ile sürgünler üzerinde beslenen nimfler, popülasyonları arttıkça, konukçularının büyümelerini etkileyerek bodur ve çarpık sürgünlerin gelişmesine neden olurlar (Aubert, 1987).

Ayrıca turunçgil ağaçlarının yeni sürgünlerinde bulunan nimfler, beslenmeleri devam ettikçe bitki uçlarının ölmesine veya bükülmesine neden olan bir toksin üretirler ve bu da yaprakların normal şekilde gelişmesini engeller. Asya turunçgil psillidinin beslenmesi sonucunda konukçularında genellikle yaprak dökülmesine ve geriye doğru kurumaya neden olduğu, büyüme noktalarında ciddi hasar meydana getirdiği, bunun da küçülmeye ve meyvede su ve tat eksikliğine yol açtığı ifade edilmiştir. Ayrıca yoğun *D. citri* popülasyonlarının çiçek ve meyve düşmesine neden olduğu, zararlı tarafından salgılanan bal özü, yalnızca ağacın fotosentetik aktivitesini etkilemekle kalmayıp aynı zamanda *D. citri*'nin doğal düşmanlarını savuşturan karıncaları da çeken fumajin büyümesini teşvik ettiği de belirtilmiştir (Anonymous, 2025).



Şekil 1. Asya Turunçgil Psillidi, *Diaphorina citri*'nin a) yumurtası, b) nimf dönemleri ve c) ergini

Asya Turunçgil Psillidi, *Diaphorina citri* Kuwayama'nın Yayılışı

Diaphorina citri Asya, Okyanusya, Afrika ve Amerika'da bulunur (Şekil 2). Güneydoğu Asya'ya özgü olduğu düşünülen zararlı, insanların ticari faaliyetleri sonucunda Asya'nın turunçgil yetiştirilen diğer bölgelerine, doğu Afrika'ya ve Hint ve Pasifik okyanuslarının adalarına yayılmıştır. İlk olarak XX. yüzyılda, Brezilya'da ve daha sonra Orta Amerika, Karayipler ve Güney Kuzey Amerika'da tespit edilerek Amerika kıtasına da ulaşmıştır. *Diaphorina citri* tropikal veya subtropikal bir türdür ancak çok çeşitli ortamlara uyum sağlama yeteneğine sahiptir. Bu yeteneği, Akdeniz, nemli subtropikal ve tropikal veya kurak ve karasal iklimler gibi çeşitli iklimlerde turunçgil yetiştirme bölgelerini istila etmesine neden olmuştur. Akdeniz havzası ve Avustralya, bu böcek türünden ve *Huanglongbing* hastalığından arı tek iki ana turunçgil yetiştirme bölgesi olarak kabul edilir. *Diaphorina citri*, henüz bulunmadığı turunçgil yetiştirme ülkelerinde karantina zararlısı olarak kabul edilir (Anonymous, 2025). EPPO tarafından A1 karantina zararlıları listesine (Ek II-A), Kuzey Avustralya Karantina Stratejisi'nin (NAQS) hedef zararlılar listesine ve Bahreyn'de A1 listesine dâhil edilmiştir (EPPO, 2022). Şili'deki Servicio Agrícola y Ganadero (SAG) tarafından karantina zararlısı olarak sınıflandırılmıştır (SAG, 2003).



Şekil 2. Asya Turunçgil Psillidi, *Diaphorina citri*'nin yayılış alanları

Asya Turunçgil Psillidi , *Diaphorina citri* Kuwayama'nın Konukçuları

Diaphorina citri yalnızca Rutaceae ailesine giren türlerde beslenir. Bu ailenin yaklaşık 100 türü ve melezi vardır. Başlıca konukçularının çoğu tatlı portakal (*Citrus sinensis*), limon (*C. limon*), pürüzlü limon (*C. jambhiri*), mandalina (*C. reticulata*), ekşi portakal (*C. aurantium*), greyfurt (*C. paradisi*) ve misket limonu (*C. aurantiifolia*) gibi ticari turunçgillerdir ([Aubert, 1990](#); [Tsai and Liu, 2000](#) ; [Westbrook et al., 2011](#) ; [Alves et al., 2014](#)). [Bununla birlikte, genellikle çitler için kullanılan bir rutaceum bitkisi olan portakal yasemini, *Murraya paniculata* gibi bazı süs türlerini de tercih etmektedir \(Anonim, 2025\).](#)

Asya Turunçgil Psillidi, *Diaphorina citri* Kuwayama'nın Doğal Düşmanları

Tamarixia radiata (Waterston) (Hymenoptera: Eulophidae)'nin Asya Turunçgil Psillidi'nin nimflerinin parazitoiti olduğu belirtilmiştir (Étienne et al., 2001). Ayrıca uğur böceklerinin (Coleoptera: Coccinellidae), sinir kanatlıların (Neuroptera: Chrysopidae) ve örümceklerin (Araneae) farklı türlerinin büyüme mevsimi boyunca *D.citri* üzerinde %100'e yakın ölümlere neden olduğu ifade edilmiştir (Michaud, [2004](#); [Qureshi and Stansly, 2009](#); [Monzo et al., 2014](#)). Ayrıca zararlının Coccinellidae, Neuroptera, Syrphidae, Hymenoptera ve Aracnida'ya bağlı çok sayıda doğal düşmanının olduğu saptanmıştır (Anonim, 2025).

Karantina Tedbirleri

Zararlı ile mücadelede bulaşık ülkelerden veya bölgelerden işlenmemiş yapraklı ve saplı halde turunçgil meyvesi ve süs bitkileri girişine müsaade edilmemelidir. Yalnızca yaprak ve dallardan arındırılmış paketli meyve ticaretine izin verilmelidir. Vektörün olası giriş noktalarında, fidanlıklarda ve özellikle sürgün döneminde bahçelerde kontroller ve sarı yapışkan tuzaklar ile takip yapılmalıdır. Ev bahçeleri ve yol kenarlarında süs amaçlı dikilen turunçgiller de kontrol edilmelidir (Anonim, 2025).

Huanglongbing'in ciddiyetini göz önünde bulundurarak EPPO, turunçgillerin ekim amaçlı ithalatının ve turunçgil Huanglongbing'inin (veya vektörlerinden herhangi birinin) bulunduğu bölgelerden veya ülkelerden turunçgil dallarının veya tomurcuklarının kesilmesinin yasaklanmasını önermiştir. AB topraklarında, pedinkülleri ve yapraklarıyla birlikte üçüncü ülkelerden meyve ithal etmek de yasaktır. Akdeniz bölgesi gibi hastalıktan ari ülkelerde farkındalık, izleme, gözetim, zararlı risk değerlendirmesi, karantina önlemleri ve eylem planları önerilmektedir (Duran-Vila et al., 2014). Huanglongbing ve vektörlerini tespit etme, kontrol altına alma ve ortadan kaldırma amacıyla resmi kontrol prosedürleri EPPO Standardı PM 9/27'de (EPPO, 2022) sağlanmıştır.

Zararlıının Tespiti

Sürvey amacıyla gelinen bahçe ve fidanlıklarda taze sürgünler gözle kontrol edilir. Ayrıca bahçe, fidanlık ve diğer kontrol noktalarında takip istasyonları oluşturularak asılan sarı yapışkan tuzaklarda gözle inceleme yapılmalıdır (Anonim, 2025).

Psillidin en yüksek aktivite dönemi, turuncgillerin taze sürgün verdiği ilkbahar ve sonbahar mevsimidir. Yeşil aksam kontrolleri bahçelerde ilkbahar ve sonbahar sürgün dönemlerinde, fidanlıklarda ise yıl boyunca yapılır. Obur sürgün veren tür veya çeşitlerde üretim sezonu boyunca yeşil aksam kontrolleri yapılır. Bahçe, fidanlık ve diğer yerlerde oluşturulan takip istasyonlarında sarı yapışkan tuzaklar yıl boyunca takip edilir. Tuzakla vektör takibi için 20x25 cm boyutlarında çift taraflı sarı yapışkan tablalar kullanılmalıdır. Tuzaklar yerden 1,5-2,0 metre yüksekçe taze sürgünlerin yakınına asılmalıdır (Anonim, 2025).

Gözle yapılan kontrollerde yumurta ve nimfler ile bulaşık olduğundan şüphelenilen sürgünler budama makası ile kesilerek teşhis için alınıp laboratuvara getirilmelidir (Anonim, 2025).

Ergin bireyleri yakalamak için emgi tüpü veya darbe yöntemi kullanılır. Darbe yönteminde, dallara vurarak ergin bireylerin toplama alanına düşmesi sağlanır ve kapalı bir kaba alınır. Sarı yapışkan tuzaklar birbirine ve başka herhangi bir yüzeye yapışmamış, üzerinde bulunan böceklerin morfolojik teşhislerini engellemeyecek şekilde olmalıdır. Alınan örnekler ve tuzaklar teşhis için ilgili enstitü/karantina müdürlüğüne gönderilmelidir (Anonim, 2025).

Biyolojik Kontrol

Biyolojik kontrol tek başına *D. citri* popülasyonunu, Huanglongbing'in yayılmasını önleyecek seviyelere düşürmek için yeterli değildir. Fakat ticari bahçelerde böcek ilacı uygulamalarının sıklığını azaltmaya yardımcı olup kentsel bahçelerdeki gibi ticari olmayan turuncgil ağaçları için en etkili yönetim stratejisidir (Kistner et al., 2016a,b; Monzo and Stansly, 2020).

Kimyasal mücadele

İnsektisit bazlı stratejiler en etkili olanlardır, ancak biyolojik kontrolle uyumlu olmalıdır (Qureshi et al., 2009; [Monzo et al., 2014](#)).

KAYNAKLAR

Anonim, (2025). Asya Turuncgil Psillidi (*Diaphorina citri*) Survey Talimatı. [https://www.tarimorman.gov.tr/GKGM/Belgeler/DB_Bitki_Sagligi/Survey/Asya_Turuncgil_Psilidi_\(Diaphorina_citri\)_Survey_Talimatı_\(2024\)](https://www.tarimorman.gov.tr/GKGM/Belgeler/DB_Bitki_Sagligi/Survey/Asya_Turuncgil_Psilidi_(Diaphorina_citri)_Survey_Talimatı_(2024)) (Erişim Tarihi: 24.02.2025).

Anonymous, (2025). *Diaphorina citri* (Asian citrus psyllid). <https://www.cabidigitallibrary.org/doi/10.1079/cabicompium.18615> (Accessed on 24 February 2025).

Aubert, B., (1987). Triozaerytreae Del Guercio ve *Diaphorina citri* Kuwayama (Homoptera: Psylloidea), two vectors of citrus greening disease: biological aspects and possible control strategies. *Fruits*, 42(3), 149-162.

Duran-Vila, N., Janse, J.D., Foissac, X., Melgarejo, P. & Bové, J.M., (2014). Addressing the threat of Huanglongbing in the Mediterranean region: the management of save the citrus industry. *Journal of Plant Pathology*, 96(4), 4-3.

EPPO/CABI (1996) Citrus greening bacterium. In: Quarantine pests for Europe. 2nd edition (Ed. By Smith, I.M.; McNamara, D.G.; Scott, P.R.; Holderness, M.). CAB INTERNATIONAL, Wallingford, UK.

EPPO, (2022). EPPO (European and Mediterranean Plant Protection Organization) Global Database. Database on quarantine pests (available online). *Diaphorina citri*. <https://gd.eppo.int> (Accessed on 17 February 2025).

FAOSTAT, (2023). Food Agriculture and Organization. <https://www.fao.org/faostat/en/#data/QCL> (Accessed on 22 February 2025).

GKGM, (2024). Turunçgil Yeşillenme Hastalığı. <https://dkib.org.tr/files/downloads/YaziSureti-TOB%20-%20Copy%202.pdf> (Erişim Tarihi: 17.02.2025).

Günaltın, İ., (2024). Salyangoz avcısı *Rumina decollata* (L.) (Gastropoda: Subulinidae) ile alternatif avlarından olan *Eobania vermiculata* (M.) (Gastropoda: Helicidae)'nın turunçgillerde popülasyon takibi ve biyolojisi üzerinde araştırmalar. Çukurova Üniversitesi Yüksek Lisans Tezi, Bitki Koruma Anabilim Dalı, Yüksek Lisans Tezi, Adana.

Gürbüz, T., (2024). Antalya İli Turunçgil Bahçelerinde Akdeniz Meyve Sineği, *Ceratitis capitata* (wiedemann) (Diptera: Tephritidae)'nekarşı farklı cezbedicilerin etkinliğinin araştırılması ve zarar oranının belirlenmesi, Hatay Mustafa Kemal Üniversitesi Fen Bilimleri Enstitüsü, Bitki Koruma Anabilim Dalı, Doktora Tezi, Hatay.

Jeppson, L.R., Baker, E.W. & Keifer, H.H., (1975). Mites Injurious to Economic Plants. University of California Press, Berkely, 614s.

Keçeci, M., (2022). Turunçgil Yetiştiriciliği için Potansiyel Tehdit: Asya Turunçgil Psillidi [*Diaphorina citri* Kuwayama (Hemiptera: Liviidae)]. İstilacı Zararlı Türler ve Mücadelesinde Yeni Yaklaşımlar (pp.251-277).

Kistner, E.J., Amrich, R., Castillo, M., Strode, V. & Hoddle, M.S., (2016). Phenology of the Asian citrus psyllid (Hemiptera: Liviidae) with special reference to biological control by *Tamarixia radiata* in residential areas of southern California. *Journal of Economic Entomology*, 109(3), 1047-1057.

Kistner, E.J., Melhem, N., Carpenter, E., Castillo, M. & Hoddle, M.S., (2016). Abiotic and biotic mortality factors affecting Asian citrus pest (Hemiptera: Liviidae) demography in southern California. *Annals of the Entomological Society of America*, 109(6), 860-871.

Kumru Özcan, G., (2024). Akdeniz Meyve Sineği, *Ceratitis capitata* (wiedemann) (Diptera: Tephritidae)'nin Antalya İli Nar ve Turunçgil Bahçelerinde Popülasyon Gelişimi, Isparta Uygulamalı Bilimler Üniversitesi, Lisansüstü Eğitim Enstitüsü, Bitki Koruma Anabilim Dalı, Yüksek Lisans Tezi, Isparta.

Kureshi, J.A., Kostyk, B.C. & Stansly, P.A., (2014). Insecticide suppression of the Asian citrus psyllid *Diaphorina citri* (Hemiptera: Liviidae) vector of Huanglongbing pathogens. *PLoS ONE*, 9(12) e112331.

Liu, Y.H. & Tsai, J.H., (2000). Effects of temperature on the biology and life table parameters of the Asian citrus Psyllid *Diaphorina citri* Kuwayama (Homoptera: Psyllidae). *Annals of Applied Biology*, 137(3), 201-206.

Michaud, J.P., (2004). Natural mortality of the Asian citrus psyllid (Homoptera: Psyllidae) in central Florida. *Biological Control*, 29(2), 260-269.

Monzo, C., Qureshi, J.A. & Stansly, P.A., (2014). Insecticide sprays, natural enemy communities and predation on the Asian citrus weed *Diaphorina citri* (Hemiptera: Psyllidae). *Bulletin Entomological Research*, 104(5), 576-585.

Monzo, C. & Stansly, P.A., (2020). Sampling and economic thresholds for the Asian Citrus Psyllid. In: Asian Citrus Psyllid. Biology Ecology and Management of the Huanglongbing Vector , [edited by Kureshi, JA, Stansly, PA]. Wallingford, England: CABI. 156-165.

SAG, (2003). Servicio Agrícola y Ganadero de Chile. Plagas cuarentenarias, Karar N° 3.080/2003. <https://normativa.sag.gob.cl/Publico/Normas/DetalleNorma.aspx?id=216948> (Accessed on 22 February 2025).

Önelge, N., Berigel, G. & Kazak, M., (2024). Doğu Akdeniz Bölgesi Turunçgil Süs Bitkilerinde *Citrus cachexia* viroid (ccavd)'nin Araştırılması.

Qureshi, J.A. & Stansly, P.A., (2009). Exclusion techniques reveal the significant biotic mortality suffered by populations of the Asian citrus psyllid *Diaphorina citri* (Hemiptera: Psyllidae) in Florida citrus. *Biological Control*, 50(2), 129-136.

TEPGE, (2023). Turunçgil Ürün Raporu. Tarımsal Ekonomi ve Politika Geliştirme Enstitüsü, TEPGE Yayın No: 382; ISBN: 978-605-7599-26-1.

Tuzcu, Ö., (1998). Trunçgiller lisans ders notları. Ç.Ü.Z.F. Bahçe Bitkileri Bölümü, Adana.

TÜİK, (2024). Türkiye İstatistik Kurumu. Bitkisel Üretim Verileri. <https://data.tuik.gov.tr/Bulten/Index?p=Bitkisel-Uretim-Istatistikleri-2022-45504> (Erişim tarihi: 22.02.2025).

EPIGENETIC INHERITANCE OF ENVIRONMENTAL CHANGES IN PLANTS: MECHANISMS AND IMPLICATIONS

İbrahim YILDIZHAN

Department of Agricultural Biotechnology, Faculty of Agriculture, Iğdir University, Iğdir,
Turkey

ORCID ID: <https://orcid.org/0009-0009-6611-088X>

Dr. Öğr. Üyesi Baris EREN

Department of Agricultural Biotechnology, Faculty of Agriculture, Iğdir University,
Iğdir, Turkey

ORCID ID: <https://orcid.org/0000-0002-3852-6476>

ABSTRACT

Epigenetics is a critical biological process that shapes the developmental processes and environmental adaptations of plants through mechanisms that regulate gene expression without altering the DNA sequence. When plants are exposed to environmental stresses such as drought, salinity, temperature changes and pathogen attacks, they can adapt to these challenges by regulating their genetic information through epigenetic modifications. Nowadays, it is suggested that some epigenetic modifications can be transmitted across generations, which may increase the long-term resilience of plants against environmental stresses. The aim of this study was to investigate the heritability of environmental epigenetic changes in plants, to elucidate the mechanisms of these processes and to evaluate their potential applications for agricultural sustainability. This study included a systematic analysis of the current literature on the mechanisms of epigenetic inheritance. Experimental studies investigating epigenetic modifications in plants were evaluated and focused on the role of DNA methylation, histone modifications and small RNAs. In particular, the effects of specific environmental stressors on epigenetic regulation and how these changes are transmitted to the next generation are discussed. Epigenetic changes have been observed in plants, particularly in response to abiotic and biotic stress conditions. Studies show that some epigenetic marks are maintained over generations, creating transgenic epigenetic memory. For example, plants exposed to drought stress were found to respond faster to the same stress conditions in the next generation. However, the persistence of epigenetic changes varies among species, and some of them appear to be deleted in subsequent generations. The findings indicate that environmental stresses in plants can be inherited through epigenetic modifications. This reveals the potential of developing epigenetic-based stress tolerance without the need for genetic modifications in plant breeding. In particular, in terms of agricultural sustainability, selecting and breeding plants with epigenetic memory may contribute to the creation of more resilient agricultural systems against global threats such as climate change. Future research should focus on the extent to which epigenetic marks are permanent, under which conditions they can be erased and the molecular mechanisms of these processes. In this way, epigenetic information can be integrated into agricultural Technologies to develop more productive and resistant plant species.

Key Words: Epigenetics, Histone modification, DNA methylation.

INTRODUCTION

Epigenetics represents one of the most profound and rapidly evolving fields in modern biology. The term "epigenetics" was first introduced by Waddington in 2012 to describe the complex interactions between genes and the environment that influence an organism's phenotype without altering the underlying DNA sequence (Waddington, 2012). Over the past few decades, the scope of epigenetics has expanded significantly, encompassing a wide range of molecular mechanisms such as DNA methylation, histone modifications, and RNA-mediated silencing. These mechanisms regulate gene expression patterns and contribute to phenotypic plasticity and adaptation in response to environmental cues (Berger et al., 2009; Zhang et al., 2018).

In plants, epigenetic mechanisms are particularly crucial due to their sessile nature. Unlike animals, plants cannot escape from environmental stressors such as drought, extreme temperatures, UV radiation, and pathogen attacks. Consequently, plants have developed sophisticated epigenetic regulatory networks to modulate gene expression in response to environmental challenges (Bräutigam & Cronk, 2018). These regulatory systems allow plants to adjust their growth, development, and reproductive strategies in real time, enhancing their ability to survive and thrive under diverse environmental conditions (Richards et al., 2017).

Epigenetic changes in plants can be transient or stable, with stable modifications often being transmitted across generations through transgenerational epigenetic inheritance (Quadrana & Colot, 2016). For example, DNA methylation patterns and histone modifications can be maintained through mitosis and meiosis, ensuring that adaptive traits induced by environmental stimuli are inherited by subsequent generations (Law & Jacobsen, 2010). The ability of plants to retain a "memory" of past environmental experiences through epigenetic marks provides an evolutionary advantage, enabling them to rapidly adjust to changing environmental conditions without the need for genetic mutations (Hauser et al., 2011).

Mechanisms of Epigenetic Inheritance in Plants

The primary mechanisms of epigenetic inheritance in plants include DNA methylation, histone modifications, and RNA interference (RNAi) (Table 1).

DNA Methylation

DNA methylation is one of the most well-studied and stable forms of epigenetic modification in plants. It involves the addition of a methyl group (-CH₃) to the cytosine base in the context of CG, CHG, and CHH sequences (where H = A, T, or C) (Law & Jacobsen, 2010). DNA methylation is catalyzed by DNA methyltransferases (MET1, CMT3, and DRM2) and can repress gene expression by inhibiting transcription factor binding or recruiting repressive chromatin remodeling complexes (Zhang et al., 2018).

Table 1. Direct and indirect pathways of epigenetic mechanisms.

Direct Effects		
Category	Mechanism	Description
A. DNA Modifications	DNA Methylation	Addition of a methyl group to cytosine residues (in CG, CHG, and CHH contexts) to repress gene expression.
	Non-covalent DNA Modifications	Structural changes in DNA that affect gene expression without altering the sequence.
	Transcription Factors and Feedback Autoregulation	Regulatory proteins that bind to DNA to activate or repress transcription.

B. Chromatin Modifications	Covalent Modifications (Histone Modifications):	
	- Acetylation	Addition of an acetyl group to histone tails, promoting open chromatin and gene expression.
	- Methylation	Addition of a methyl group to histones, leading to either gene activation or repression.
	- Phosphorylation	Addition of a phosphate group, often linked to chromatin remodeling during stress responses.
	- Ubiquitination	Addition of ubiquitin molecules, influencing chromatin structure and protein degradation.
	- SUMOylation	Addition of SUMO proteins, affecting protein interactions and stability.
Non-covalent Modifications:	Histone Exchange	Replacement of histones with variants to modify chromatin structure.
	Histone Incorporation	Insertion of histones into nucleosomes, influencing chromatin accessibility.
	Chromatin Remodeling	Rearrangement of chromatin architecture to regulate gene expression.
	Non-coding RNA Interactions	Interaction of microRNAs and small RNAs with chromatin to silence genes.
	Interactions with Other Agents	Interaction of chromatin with proteins and other regulatory factors.
	Long-distance Chromosome Interactions	Chromosome folding and looping to bring distant regulatory elements into contact.
Indirect Effects (Post-transcriptional mechanisms)		
Mechanism	Description	
mRNA Silencing	Small RNAs (miRNAs, siRNAs) guide Argonaute proteins to degrade or block translation of target mRNAs.	
Alternative Splicing Regulation	Epigenetic changes influence splicing patterns, leading to different protein isoforms.	
RNA Editing	Modification of RNA molecules, altering codon sequences and protein products.	

* Adapted from Zhang et al. (2018) and other sources.

Histone Modifications

Histones are proteins that package and organize DNA into chromatin. Post-translational modifications of histone tails, such as acetylation, methylation, phosphorylation, and ubiquitination, influence chromatin structure and gene expression. For example, histone acetylation generally promotes gene activation, while histone methylation can lead to either gene activation or repression, depending on the specific lysine residue modified (Kouzarides, 2007). The Polycomb and Trithorax group proteins play critical roles in establishing and maintaining histone modification patterns in plants (He et al., 2011).

Small RNA-Mediated Gene Silencing

RNA interference (RNAi) involves small non-coding RNAs, including microRNAs (miRNAs) and small interfering RNAs (siRNAs), that guide gene silencing at both the transcriptional and post-transcriptional levels. In plants, siRNAs direct the establishment of DNA methylation patterns through the RNA-directed DNA methylation (RdDM) pathway, which reinforces gene silencing and heterochromatin formation (Hauser et al., 2011; Kong et al., 2017).

Epigenetic Responses to Environmental Changes

Plants exhibit remarkable plasticity in response to environmental stress, mediated by dynamic changes in epigenetic marks. Epigenetic modifications regulate plant responses to abiotic stressors such as drought, salt, temperature extremes, and nutrient availability (Figure 1).




Figure 1. Factors playing a role in epigenetic mechanism.

Vernalization – A well-known example of epigenetic adaptation in plants is vernalization, where exposure to prolonged cold triggers flowering in species like *Arabidopsis thaliana*. This process involves stable changes in histone methylation at the FLOWERING LOCUS C (FLC) gene, which are maintained through cell divisions but reset in the next generation (Andrés and Coupland, 2012).

Drought Tolerance – Plants exposed to mild drought stress exhibit stable changes in DNA methylation and histone modification patterns, which enhance their ability to withstand future drought episodes. These epigenetic changes are thought to be transmitted to offspring, contributing to transgenerational drought tolerance (Ashapkin et al., 2020).

UV Radiation and Pathogen Defense – Epigenetic responses to biotic stress, such as pathogen infection and UV radiation, are also well documented. For example, exposure of *Arabidopsis* to UV-B radiation induces increased somatic recombination rates and changes in DNA methylation, which are maintained across generations (Boyko et al., 2010).

Transgenerational Epigenetic Memory and Adaptation

Transgenerational epigenetic inheritance provides plants with a powerful mechanism to "remember" past environmental conditions and pass on adaptive traits to future generations. This memory is thought to arise from the stable maintenance of DNA methylation and histone modification patterns during DNA replication and cell division (Richards et al., 2017). Evidence of transgenerational epigenetic inheritance has been observed in both model and natural plant populations, highlighting the evolutionary significance of epigenetic plasticity in plants (Schmitz et al., 2013).

One approach to distinguish genetic from epigenetic variation is to analyze cloned or genetically identical plant populations grown under different environmental conditions. Studies have shown that epigenetic variation in these systems directly reflects environmental influences rather than genetic differences (Savoie et al., 2015). Statistical approaches and genome-wide epigenomic profiling have further refined our ability to identify and quantify environmentally induced epigenetic changes in plants (Richards et al., 2017).

DISCUSSION and CONCLUSION

Epigenetic inheritance in plants has emerged as a crucial mechanism for regulating gene expression and facilitating adaptation to changing environmental conditions. The ability of plants to modify their gene expression patterns in response to environmental cues through DNA methylation, histone modifications, and small RNA-mediated gene silencing reflects a sophisticated and dynamic regulatory network. This discussion will explore the functional significance of epigenetic inheritance in plants, its evolutionary implications, and the current challenges and future directions in this rapidly evolving field.

Functional Significance of Epigenetic Inheritance in Plants

Epigenetic mechanisms allow plants to respond rapidly to environmental stresses without the need for changes in the underlying DNA sequence. DNA methylation, histone modifications, and small RNA activity work together to fine-tune gene expression, enabling plants to adapt to abiotic and biotic stresses such as drought, extreme temperatures, pathogen attacks, and nutrient deficiency (Bräutigam & Cronk, 2018). The role of vernalization, in which histone methylation at the FLOWERING LOCUS C (FLC) gene promotes flowering in response to cold temperatures, demonstrates the functional importance of epigenetic modifications in regulating developmental processes (Andrés and Coupland, 2012).

Similarly, the transmission of drought tolerance across generations through stable DNA methylation patterns highlights the adaptive potential of epigenetic inheritance. Plants subjected to mild drought stress have been shown to accumulate specific methylation marks that are maintained through mitosis and meiosis, allowing progeny to better withstand drought conditions (Ashapkin et al., 2020). This indicates that epigenetic inheritance serves as a mechanism for stress memory, which is essential for plant survival in fluctuating environmental conditions.

Evolutionary Implications

Epigenetic inheritance challenges the traditional view of evolution based solely on genetic variation and natural selection. The ability of plants to pass on epigenetic modifications across generations suggests that epigenetic plasticity plays a critical role in evolutionary processes. Unlike genetic mutations, which occur randomly and are often irreversible, epigenetic changes are reversible and can be fine-tuned in response to environmental pressures (Richards et al., 2017).

This adaptive flexibility allows plants to respond rapidly to environmental changes while maintaining the potential for reversibility if conditions change. For example, epigenetic variation in genetically identical populations of *Arabidopsis* and *Quercus lobata* has been linked to environmental factors such as temperature and water availability (Schmitz et al., 2013; Savoie et al., 2015). The ability to rapidly establish and erase epigenetic marks provides plants with a powerful mechanism for evolutionary adaptation that operates independently of genetic mutations.

Challenges and Limitations

Despite the growing body of evidence supporting the role of epigenetic inheritance in plants, several challenges remain. One key challenge is distinguishing between genetic and epigenetic variation, particularly in complex natural environments where both factors are likely to interact. Controlled experimental designs involving cloned or genetically identical plant populations have provided insights into the role of epigenetics, but extending these findings to natural populations remains difficult (Richards et al., 2017).

Furthermore, the molecular mechanisms underlying the stable transmission of epigenetic marks across generations are not fully understood. While DNA methylation and histone modifications have been implicated in transgenerational inheritance, the specific proteins and pathways involved remain to be fully characterized (Zhang et al., 2018). The dynamic nature of epigenetic marks also raises questions about the long-term stability of epigenetic inheritance and its impact on evolutionary fitness.

Future Directions

Future research on plant epigenetics should focus on expanding our understanding of the molecular mechanisms governing epigenetic inheritance and their evolutionary significance. Advances in high-throughput sequencing and epigenomic profiling have provided powerful tools for mapping epigenetic marks and linking them to specific environmental cues (Schmitz et al., 2013). Integrating these techniques with functional genomics and gene editing approaches (e.g., CRISPR-based epigenome editing) will enable researchers to explore the causal relationships between epigenetic modifications and adaptive traits.

Additionally, exploring the role of small RNAs in transgenerational inheritance and environmental adaptation represents a promising avenue for future research. Small RNAs have been shown to guide DNA methylation and gene silencing, but their role in maintaining epigenetic memory across generations remains poorly understood (Hauser et al., 2011; Kong et al., 2017). Investigating how small RNAs interact with chromatin remodeling complexes and transcription factors will provide valuable insights into the regulatory networks underlying epigenetic inheritance.

CONCLUSION

Epigenetic inheritance represents a powerful mechanism for regulating gene expression and facilitating plant adaptation to environmental changes. DNA methylation, histone modifications, and small RNA activity form a complex and dynamic regulatory network that allows plants to respond to abiotic and biotic stresses. The evolutionary implications of epigenetic plasticity challenge traditional views of genetic inheritance and natural selection, highlighting the role of epigenetics in shaping plant adaptation and evolution. Overcoming the challenges associated with distinguishing genetic and epigenetic variation and elucidating the molecular mechanisms governing transgenerational inheritance will be critical for advancing our understanding of plant epigenetics.

REFERENCES

- Andrés, F., & Coupland, G. (2012). The genetic basis of flowering responses to seasonal cues. *Nature Reviews Genetics*, 13(9), 627-639.
- Ashapkin, V. V., Kutueva, L. I., Aleksandrushkina, N. I., & Vanyushin, B. F. (2020). Epigenetic mechanisms of plant adaptation to biotic and abiotic stresses. *International journal of molecular sciences*, 21(20), 7457.
- Berger, S. L., Kouzarides, T., Shiekhatar, R., & Shilatifard, A. (2009). An operational definition of epigenetics. *Genes & development*, 23(7), 781-783.
- Boyko, A., Blevins, T., Yao, Y., Golubov, A., Bilichak, A., Ilnytsky, Y., ... & Kovalchuk, I. (2010). Transgenerational adaptation of *Arabidopsis* to stress requires DNA methylation and the function of Dicer-like proteins. *PloS one*, 5(3), e9514.
- Bräutigam, K., & Cronk, Q. (2018). DNA methylation and the evolution of developmental complexity in plants. *Frontiers in Plant Science*, 9, 1447.
- Hauser, M. T., Aufsatz, W., Jonak, C., & Luschnig, C. (2011). Transgenerational epigenetic inheritance in plants. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, 1809(8), 459-468.
- He, G., Elling, A. A., & Deng, X. W. (2011). The epigenome and plant development. *Annual review of plant biology*, 62(1), 411-435.
- Kong, X., Luo, X., Qu, G. P., Liu, P., & Jin, J. B. (2017). *Arabidopsis* SUMO protease ASP1 positively regulates flowering time partially through regulating FLC stability. *Journal of Integrative Plant Biology*, 59(1), 15-29.
- Kouzarides, T. (2007). Chromatin modifications and their function. *Cell*, 128(4), 693-705.
- Law, J. A., & Jacobsen, S. E. (2010). Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nature Reviews Genetics*, 11(3), 204-220.
- Quadrana, L., & Colot, V. (2016). Plant transgenerational epigenetics. *Annual Review of Genetics*, 50(1), 467-491.
- Richards, C. L., Alonso, C., Becker, C., Bossdorf, O., Bucher, E., Colomé-Tatché, M., ... & Verhoeven, K. J. (2017). Ecological plant epigenetics: Evidence from model and non-model species, and the way forward. *Ecology letters*, 20(12), 1576-1590.
- Savoie, A. M., & Saunders, G. W. (2015). Evidence for the introduction of the Asian red alga *Neosiphonia japonica* and its introgression with *Neosiphonia harveyi* (Ceramiales, Rhodophyta) in the Northwest Atlantic. *Molecular Ecology*, 24(23), 5927-5937.
- Schmitz, R. J., Schultz, M. D., Urich, M. A., Nery, J. R., Pelizzola, M., Libiger, O., ... & Ecker, J. R. (2013). Patterns of population epigenomic diversity. *Nature*, 495(7440), 193-198.
- Waddington, C. H. (2012). The epigenotype. *International journal of epidemiology*, 41(1), 10-13.
- Zhang, H., Lang, Z., & Zhu, J. K. (2018). Dynamics and function of DNA methylation in plants. *Nature reviews Molecular cell biology*, 19(8), 489-506.

REVIEW OF *IGF-IGENE* VARIABILITY AND ITS IMPACT ON SHEEP BREEDING IN TURKEY**TÜRKİYE'DE *IGF-IGEN* ÇEŞİTLİLİĞİ VE KOYUN YETİŞTİRİCİLİĞİ ÜZERİNDEKİ ETKİLERİ ÜZERİNE BİR DERLEME****Öğr. Gör. Dr. Hasret ÖZTÜRK**

Department of Animal Sciences, Unit of of Biometry and Genetics, Faculty of Agriculture, Iğdır University, Iğdır, Turkey

ORCID ID: <https://orcid.org/0000-0002-1419-1050>**ÖZET**

İnsülin Benzeri Büyüme Faktörü 1 (*IGF-I*) geni, büyüme, kas gelişimi, metabolizma ve üreme gibi temel biyolojik süreçlerde hayati rol oynayan hormon benzeri bir genetik faktördür. Hipofiz bezi tarafından salgılanan büyüme hormonu (GH) ile çalışarak hücre büyümesini, farklılaşmasını ve doku yenilenmesini düzenler. *IGF-I*, özellikle iskelet kası, karaciğer, yağ dokusu ve bağ dokusunda aktif olarak eksprese edilir ve organizmanın gelişim sürecinde anabolik (yapıcı) etkiler gösterir. Türkiye'deki koyun ırklarında *IGF-I* polimorfizmleri, canlı ağırlık, karkas kompozisyonu ve üreme performansı gibi ekonomik açıdan önemli özelliklerle ilişkilendirilmiştir. Bu derleme, Türk koyun ırklarındaki *IGF-I* gen çeşitliliği üzerine gerçekleştirilen mevcut ancak sınırlı sayıda araştırma incelemekte ve bu çeşitliliğin büyüme performansı ile seçici yetiştirme üzerindeki etkilerini değerlendirmektedir. Ayrıca, çalışmalar *IGF-I* mutasyonlarının kuzu doğum ağırlığı, karkas kalitesi ve üreme verimliliği üzerindeki genetik etkilerini ele almaktadır. Bununla birlikte, küresel çalışmalarla yapılan karşılaştırmalar, *IGF-I* polimorfizmlerinin hayvancılık açısından taşıdığı önemi ortaya koymaktadır. Marker Destekli Seleksiyon (MAS) ve genomik seleksiyon yaklaşımlarının *IGF-I* genetik belirteçlerini içerecek şekilde kullanılması, Türk koyunlarında genetik kazançları artırmak için önerilmektedir. Potansiyel faydalara rağmen, mevcut çalışmaların örneklem büyüklüğü ve kapsam açısından sınırlı olması, geniş ölçekli genomik araştırmalara duyulan ihtiyacı ortaya koymaktadır. Gelecekte, GWAS ve QTL haritalama yöntemlerinin birlikte kullanılması, *IGF-I* polimorfizmlerinin koyun verimliliği üzerindeki genetik etkilerini daha kapsamlı bir şekilde ortaya koyabilir. GWAS, küçük etkili genetik varyantları belirlerken, QTL haritalama büyük etkili gen bölgelerini tespit ederek seçici yetiştirme programlarını destekleyebilir. Bu yaklaşımlar, *IGF-I* verilerinin ulusal hayvancılık veri tabanlarına entegrasyonu ile birleştiğinde, Türk koyunculunun sürdürülebilirliği ve verimliliğine önemli katkılar sağlayacaktır.

Anahtar Kelimeler: *IGF-I*, Koyun Irkları, Genetik Polimorfizm, MAS**ABSTRACT**

The Insulin-like Growth Factor 1 (*IGF-I*) gene is a hormone-like genetic factor that plays a crucial role in growth, muscle development, metabolism, and reproduction. It works in conjunction with growth hormone (GH), secreted by the pituitary gland, to regulate cell growth, differentiation, and tissue regeneration. *IGF-I* is highly expressed in skeletal muscle, liver, adipose tissue, and connective tissue, exerting anabolic (constructive) effects during the organism's developmental process. In Turkish sheep breeds, *IGF-I* polymorphisms have been associated with economically

significant traits, such as live weight, carcass composition, and reproductive performance. This review examines the existing but limited studies on *IGF-I* gene variability in Turkish sheep breeds and evaluates its impact on growth performance and selective breeding. Additionally, this study investigates the genetic effects of *IGF-I* mutations on lamb birth weight, carcass quality, and reproductive efficiency. The comparisons with global studies highlight the significance of *IGF-I* polymorphisms in livestock production. The use of Marker-Assisted Selection (MAS) and genomic selection approaches incorporating *IGF-I* genetic markers is recommended to enhance genetic gains in Turkish sheep populations. Despite its potential benefits, the small sample sizes and limited scope of current studies emphasize the need for large-scale genomic research. In the future, the combined use of GWAS and QTL mapping methods may provide a more comprehensive understanding of the genetic effects of *IGF-I* polymorphisms on sheep productivity. GWAS identifies small-effect genetic variants, while QTL mapping detects major-effect genomic regions, supporting selective breeding programs. When integrated with national livestock databases, these approaches will make significant contributions to the sustainability and efficiency of sheep farming in Turkey.

Keywords: *IGF-I*, Sheep Breeds, Genetic Polymorphism, MAS

GİRİŞ

Bu kısa derleme, Türk koyun ırklarındaki *IGF-I* gen değişkenliği üzerine gerçekleştirilmiş olan ancak sayıca sınırlı olan mevcut çalışmalarını incelemekte ve bu genetik değişkenliğin büyüme performansı ve seçici yetiştirme üzerindeki potansiyel etkilerini değerlendirmektedir.

Koyun yetiştiriciliği

Koyun yetiştiriciliği, dünya çapında hayvancılık sektörünün kritik bir bileşeni olarak hem ekonomik hem de çevresel anlamda önemli katkılarda bulunur. Bu sektör, et, süt ve yün gibi çok çeşitli ürünler sunar; bu ürünler gıda sanayisinden tekstil sektörüne kadar geniş bir kullanım alanına sahiptir (Çiçek vd., 2022). Koyunlar, özellikle kırsal ve açık arazilere sahip bölgelerde ekonomik açıdan hayati bir faaliyet olarak görülür ve birçok aile için ana gelir kaynağıdır. Bu hayvanlar, farklı iklim ve arazi koşullarına kolayca uyum sağlayabilir, bu da onları çeşitli çevresel şartlarda yetiştirmeyi mümkün kılar. Ayrıca, yüksek kaliteli koyun eti ve yünü gibi ürünler uluslararası pazarlarda büyük değere sahiptir ve önemli ihracat kalemleri arasında yer alır (Günaydın, 2009). Koyun yetiştiriciliği, aynı zamanda çevresel etkileri düşük olan sürdürülebilir tarım uygulamaları ile de uyumludur, bu da onu ekolojik açıdan tercih edilen bir faaliyet haline getirir.

Türkiye’de Koyun yetiştiriciliği

Türkiye, coğrafi konumu ve iklim çeşitliliği sayesinde birçok yerli koyun ırkına ev sahipliği yapmaktadır. Ülkenin tarım sektöründe önemli bir yere sahip olan koyun yetiştiriciliği hem geleneksel aile işletmelerinde hem de büyük ölçekli ticari çiftliklerde yaygın bir şekilde uygulanmaktadır (Çiçek vd., 2022). Türkiye'nin zengin mera alanları bu hayvanlar için ideal koşulları sağlar. Özellikle İç Anadolu, Doğu Anadolu ve Güneydoğu Anadolu bölgelerinde, geniş mera alanları sayesinde koyun yetiştiriciliği özellikle yoğundur. Bu bölgelerdeki koyun yetiştiriciliği, ülkenin tarımsal üretim modelinin koyuncululuğun gelişimine uygun koşullar sağlaması ve bazı yerel koyun ırklarının iklim ve çevresel faktörlere yüksek adaptasyon yetenekleriyle yetiştirilmesi açısından avantaj sunmaktadır. Koyun yetiştiriciliği, Türkiye ekonomisi için büyük önem taşıyarak et, süt, yapağı ve deri gibi çeşitli değerli ürünler sunmakta ve insanların temel ihtiyaçlarını uzun yıllardır karşılamaktadır. Türkiye’de koyunlar, dünya genelinde olduğu gibi, çeşitli iklim ve arazi koşullarında başarıyla yetiştirilmekte ve bu alanda yapılan bilimsel çalışmalar sektörün daha da ileriye taşınmasına katkıda bulunmaktadır (Kaymakçı vd., 2000).

Türkiye'de koyun yetiştiriciliği sektöründe uygulanan yenilikler ve teşvikler, 2024 yılı itibarıyla toplam 44,080,584 koyunun yetiştirildiği bir popülasyon artışını tetiklemiştir (Ertürk vd., 2015; TÜİK, 2025). Ülkenin çeşitli illerinde koyun popülasyonu büyük farklılıklar göstermekte olup, özellikle Van 3,040,146 koyun ile en yüksek popülasyona sahip iken; Konya 2,931,323 ve Ankara 1,811,254 koyun ile takip etmektedir. Bu iller geniş mera alanları ve uygun çevresel koşulları ile koyun yetiştiriciliği için idealdir. Diğer yandan, Şanlıurfa ve Diyarbakır, sırasıyla 2,364,954 ve 1,894,610 koyun ile Güneydoğu Anadolu Bölgesi'nde; Balıkesir ise Marmara Bölgesi'nde 1,222,213 koyun ile dikkat çekmektedir. Karadeniz Bölgesi'nde ise Bartın, Rize ve Zonguldak illeri, sırasıyla 4,053, 8,028 ve 14,520 koyun ile en az koyun popülasyonuna sahip illerdir (TÜİK, 2025).

Türkiye'deki koyun ırklarının ekonomik katkısı ve yetiştirilme özellikleri, her bir ırkın genel yüzdesine dayanarak değerlendirilebilir. Ülkede en yaygın yetiştirilen koyun ırkı %48 ile Akkaraman'dır ve İç Anadolu, Doğu Anadolu, ve Güneydoğu Anadolu bölgelerinde popülerdir; adaptasyon kabiliyeti yüksek ve et ile yün verimi oldukça değerlidir. Morkaraman %13 ile ikinci sırada olup, soğuk iklimlere uygun, yüksek yün ve et verimi ile bilinir. Kıvırcık ırkı %9 ile batıda, özellikle Marmara ve Ege'de, kaliteli yünü ile öne çıkar. İvesi ve Pırıt/Pırlak her biri %7 ile sıralanır; İvesi Güneydoğu Anadolu'da süt verimi, Pırıt/Pırlak Ege'de et ve süt verimi ile tanınır. Anadolu Merinosu ve Hamdani her biri %5 ile az yaygınken; Merinos yün kalitesi, Hamdani dayanıklılığı ve et-süt verimi ile dikkat çeker. Karayaka %3 ile Karadeniz'de, zorlu iklim koşullarına uyumu ile bilinirken, Merinos ve Sakız sırasıyla %3 ve %2 ile nadir yetiştirilen ırklar arasında yer alır; Merinos yüksek yün kalitesi, Sakız ise Ege'de süt ve et kalitesi ile değerlidir (Taşkın & Kandemir, 2022a, 2022b, 2022c, 2022d; Kandemir & Taşkın, 2022a, 2022b, 2022c). Bu ırklar, Türkiye'nin zengin hayvancılık kültürünü ve tarımsal çeşitliliğini yansıtarak, çeşitli üretim amaçlarına uyum sağlar.

Koyun Yetiştiriciliğinde Büyüme Özelliklerinin Önemi

Büyüme özellikleri, koyun yetiştiriciliğinde hem verimliliği hem de yetiştiricinin karlılığını doğrudan etkileyen temel unsurlardır. Kuzu vücut ağırlığı gibi büyüme parametreleri, et üretimi başta olmak üzere süt ve yapağı verimi gibi çeşitli üretim faktörlerini de doğrudan etkiler. Bu nedenle, büyüme performansının iyileştirilmesi hem ürün kalitesini artırmak hem de ekonomik getiriyi maksimize etmek açısından koyun yetiştiriciliğinde öncelikli hedeflerden biri haline gelmiştir. Gelişmiş büyüme özellikleri, sadece daha fazla et ve süt sağlamakla kalmaz, aynı zamanda yüksek kaliteli yapağı üretimi için de kritik öneme sahiptir, bu da yetiştiricilere ekonomik olarak daha çekici seçenekler sunar Armstrong vd. 2018; Ghasemi vd., 2019; Ramos vd., 2023). Dolayısıyla, koyun yetiştiriciliğinde genetik iyileştirmeler ve yönetim stratejilerinin optimize edilmesi, genel verimliliği ve kârlılığı artırmanın yanı sıra, ürün çeşitliliğini ve pazar fırsatlarını da genişletmektedir.

Koyun Yetiştiriciliğinde Moleküler Hayvan Islahının Önemi

Hayvan ıslahı, koyun yetiştiriciliğinin verimliliğini, hastalık direncini ve adaptasyon kabiliyetini iyileştirme yoluyla önemli katkılarda bulunur. Moleküler hayvan ıslahı, hayvanların genetik yapısını anlamak ve iyileştirmek için genetik markörlerin kullanıldığı bir süreçtir. Bu alanda, özellikle Tek Nükleotid Polimorfizmleri (SNP) gibi genetik varyasyonlar, belirli özelliklerle güçlü ilişkiler gösterdiği için büyük önem taşır (Dekkers, 2004). Marker Assisted Selection (MAS) yöntemi, genetik markörlerle ilişkilendirilen özellikleri taşıyan bireylerin seçilmesi sürecidir ve üretim, hastalık direnci, adaptasyon, büyüme hızı ve üreme gibi özelliklerle ilişkili genetik lokusları tanımlamak için kullanılır. Moleküler ıslah ve polimorfizm çalışmaları sayesinde, belirli genetik markörler doğrudan belirlenerek, genetik olarak üstün hayvanların seçiminde rehber olarak kullanılır (Goddard ve Hayes, 2009). Bu yöntemler, koyun yetiştiriciliğinin verimliliğini ve kalitesini artırma potansiyeline sahiptir.

IGF-IGenin Hayvancılıkta Önemi ve Potansiyeli

İnsülin Benzeri Büyüme Faktörü 1 (*IGF-I*), büyüme, kas gelişimi, metabolizma ve üreme gibi temel biyolojik süreçlerde kritik bir rol oynayan hormon benzeri bir genetik faktördür. Hipofiz bezi tarafından salgılanan büyüme hormonu (GH) ile çalışarak hücre büyümesini, farklılaşmasını ve doku yenilenmesini teşvik eder. Özellikle iskelet kası, karaciğer, yağ dokusu ve bağ dokusu gibi dokularda aktif olarak eksprese edilen *IGF-I*, organizmanın gelişim sürecinde anabolik (yapıcı) etkiler gösterir *IGF-I*'nin özellikle sığırlarda, domuzlarda ve keçilerde büyüme performansı ve canlı kilo alımı üzerindeki etkileri incelenmiştir. Örneğin, null mutant ve yabancı tip fareler arasında yapılan karşılaştırmalarda, null mutant farelerin yabancı tiplere göre daha düşük iskelet büyüme hızı gösterdiği gözlemlenmiştir (Baker vd., 1993). Sığırlarda, *IGF-IGenin* büyüme performansı ve canlı kilo alımı ile ilişkisi belirlenmiş (Siadkowska vd., 2006), domuzlarda kas kütlesi ve canlı kilo alımı üzerinde önemli bir rol oynadığı bildirilmiştir (Casas-Carrillo vd., 1997), keçilerde ise et verimi ve kilo artışı üzerinde etkili olduğu bulunmuştur (Zhang vd., 2008). Makui koyun ırkında büyüme ve et verimi üzerinde potansiyel bir etkisi olduğu tespit edilmiş ve bu gen aday bir gen olarak önerilmiştir (Hajihosseino vd., 2013). Bu bulgular, *IGF-IGenin* hayvancılıkta verimliliği artırmak ve genetik iyileştirme hedeflerine ulaşmak için markör destekli seçim programlarında umut verici bir aday gen olarak değerlendirilmesini desteklemektedir.

Türkiye'de Yapılan IGF-IPolimorfizmi Çalışmaları

IGF-I/Bsp143II Polimorfizmi

Türkiye'deki koyun ırkları üzerinde yapılan *IGF-I/Bsp143II* gen polimorfizmi çalışmaları, yalnızca Akkaraman ve Kıvırcık ırklarında gerçekleştirilmiştir. Akkaraman ırkında BB genotipi monomorfik olarak baskınken (Bayram vd., 2019), Kıvırcık ırkında en yüksek oranda AA genotipi (%80 civarında) gözlemlenmiş, BB ve BA genotipleri ise çok daha düşük frekansta bulunmuştur (Kaplan ve Atalay, 2018). Akkaraman ırkında, sahip olunan genotipin kuzuların doğumdan itibaren 30, 60 ve 90. günlerdeki kilo alımları ile istatistiksel bir ilişkisi araştırılmak istenmiş fakat genin monomorfik olması nedeniyle istatistiksel bir analiz uygulanamamıştır. Bu bulgular, iki ırkın genetik yapısında *IGF-IGenine* dair farklı kalıplar sergilediğini ve bu genin fonotipik özelliklerle olan potansiyel ilişkilerini araştırmak için farklı yaklaşımlar gerektirdiğini işaret etmektedir.

2010 yılından sonra yapılan *IGF-I/Bsp143II* polimorfizmi çalışmaları, dünya genelinde birçok farklı ülkede ve koyun ırkında gerçekleştirilmiştir. Çin'de Small Tail Han, Hu, Texel ve Dorset ırkları üzerinde yapılan çalışmalarda (He vd., 2012); Kolombiya'da OPC x Pelibuey melez ırkı üzerinde (Lenis-Valencia vd., 2021); Polonya'da Renkli Polonya Merinosunda (Grochowska vd., 2017); İran'da Sanjabi ırkında (Bakhtiar vd., 2017); Hindistan'da Munjal koyunlarında (Kumar vd., 2023); Irak'ta Hamdani koyunlarında (Bayraktar ve Shoshin, 2021); Nijerya'da Yankasa ve aynı zamanda Balami, Uda, Yankasa koyunlarında (Umego vd., 2018; Raji vd., 2017) bu polimorfizm incelenmiştir. Grochowska vd.,(2017) Renkli Polonya Merinos koyun popülasyonunda AA ve AB genotiplerinin gözlemlendiğini, ancak BB genotipinin bulunamadığını belirtmiştir. Çalışma, *IGF-IGenotiplerinin* koyunlarda ön shaft ağırlığına ($p=0.006$), böbrek yağ sınıfına ($p=0.002$) ve EUROP yağ sınıfına ($p=0.005$) anlamlı bir etkisi olduğunu göstermiştir. Ayrıca, bu genotipler karkas sınıfının dış yağlılığını ($p=0.038$), damlama kaybını ($P=0.049$) ve et renginin öznel değerlendirmesini ($p=0.043$) önemli ölçüde etkilemiştir. Kumar vd., (2023), *IGF-I/Bsp143II* polimorfizminde A alelinin, daha yüksek vücut ağırlıkları ile ilişkilendirildiğini ve Munjal koyunlarının performansını artırmak için seçim kriterlerinde kullanılabileceğini ortaya koymuştur. Öte yandan, Bayraktar ve Shoshin (2021) tarafından yapılan çalışmada, Hamdani koyunlarında *IGF-I*'in AA genotipinin popülasyonda AB ve BB genotiplerinden daha sık rastlanıldığı belirtilmiştir. Ayrıca, AB genotipine sahip kuzuların daha yüksek ADG2'ye sahip olduğu gözlemlenmiştir. Umego vd., (2018) Yankasa popülasyonunda AA genotipinin büyüme özellikleri açısından avantajlı olduğunu ve bu genotipin büyüme performansını artırmak için ıslah programlarında kullanılabileceğini önermektedir. Raji vd., (2017) *IGF-IGen* polimorfizmi üzerine

yaptığı çalışmada, Balami popülasyonunda BB genotipi, AB genotipine kıyasla daha yüksek vücut ağırlığıyla (56 kg vs. 45.80 kg) ilişkilendirmiştir.

Bu bulgular, *IGF-I*geninin, koyunlarda karkas ve et kalitesi özellikleri için önemli bir aday gen olarak değerlendirilebileceğini ortaya koymaktadır. Bu geniş çaplı araştırmalar, *IGF-I/Bsp143II* polimorfizminin farklı coğrafi koşullar ve genetik arka planlar altında nasıl değişebileceğini ve bu değişikliklerin koyun ırklarının büyüme, sağlık ve üretim özellikleri üzerindeki etkilerini anlamak için önemli bilgiler sağlamaktadır.

IGF-I/ HaeII Polimorfizmi

Türkiye'deki koyun ırkları arasında Dağlıç, Karayaka, Akkaraman, Morkaraman, Malya, Sakız, Kıvırcık ve İvesi üzerinde yapılan *IGF-I/HaeII* polimorfizmi çalışmaları tek bir çalışmada incelenmiştir (Tawfeeq vd., 2020). Karayaka, Kıvırcık, Dağlıç, Malya ve Akkaraman ırklarında AA genotipi oldukça yüksek frekansta bulunur ve bu ırklarda baskın genotip olarak öne çıkar; diğer genotiplere kıyasla belirgin bir şekilde daha yaygındır. İvesi ve Morkaraman ırklarında ise BA genotipi diğerlerine göre daha yüksek frekansta bulunmakta ve bu iki ırk için baskın genotip durumundadır. Sakız ırkında en yaygın genotip AA iken, BA genotipi de yüksek frekansa sahip olarak dikkat çeker. Buna karşın, BB genotipi tüm ırklarda en düşük frekansa sahiptir ve diğer genotiplere göre oldukça seyrek görülmektedir.

2010 yılından sonra yapılan *IGF-I/HaeII* polimorfizmi çalışmaları, dünya genelinde birçok farklı ülkede ve çeşitli koyun ırklarında gerçekleştirilmiştir. Mısır'da Barki koyunlarında (Darwish vd., 2017), Kolombiya'da OPC koyunlarında (Valencia vd., 2022), Polonya'daki Renkli Merinos koyunlarında (Grochowska vd., 2021), İran'da Zandi (Nazari vd., 2016; Kazemi vd., 2011), Mehraban (Bahrami vd., 2013), Zel (Gharahveysi vd., 2015) ve Makui (Hajihosseino vd., 2013) koyunlarında çalışmalar yapılmıştır. Darwish vd., (2017) tarafından yapılan *IGF-I*gen polimorfizmi çalışmasında, GG genotipi, temiz yapağı ağırlığı (CFW) özelliği ile pozitif ve anlamlı bir ilişki göstermiştir ($p < 0.001$, Odds Ratio = 2.83). Buna karşın, AA genotipi bu özellikte negatif ve anlamlı bir ilişki göstermiştir ($p < 0.05$, Odds Ratio = 0.15). Öte yandan, lif çapı (FD) ölçümleri, farklı *IGF-I*genotipleri ile anlamlı bir ilişki göstermemiştir ($p > 0.05$). Bu çalışma, Mısır'ın Barki koyunlarında *IGF-I*gen polimorfizmi ile yünün CFW özelliği arasındaki ilişkiye dair kanıtlar eklemektedir. Nazari vd., (2016), Zandi koyunlarında AA genotipinin canlı doğum ağırlığı ile ilişkili olduğunu bulmuşlardır. Hajihosseino vd. (2013), Makui koyunlarında bu polimorfizmi incelemiş ve AA genotipinin, diğer genotiplere göre üstün doğum ağırlığı gösterdiğini belirtmiştir.

Bu geniş çaplı araştırmalar, *IGF-I/HaeII* polimorfizminin farklı coğrafyalarda ve genetik arka planlar altında nasıl varyasyonlar gösterdiğini ve bu değişikliklerin koyun ırklarının büyüme, sağlık ve üretim özellikleri üzerindeki etkilerini anlamak için önemli bilgiler sağlamaktadır.

IGF-15' UTR polimorfizmleri

Kadir Esen & Esen (2022) Türkiye'de yetiştirilen etlik koyun ırklarında *IGF-15'UTR* polimorfizmi ile büyüme ve karkas özellikleri arasındaki ilişkiyi incelemek amacıyla bir çalışma gerçekleştirilmiştir. Bu çalışmada, Kıvırcık, Karacabey Merin, Ramlıç, Alman Siyah Başlı Mutton × Kıvırcık, Hampshire Down × Merino ırkları üzerinde araştırmalar yapılmıştır. Araştırma, vücut ölçümleri (CW ve LC), MLD gelişimi, karkas dışı bileşenler (testis ağırlığı) ve perakende karkas yüzdesi (sırt yüzdesi) gibi çeşitli özellikleri kapsamaktadır. PCR-SSCP yöntemi kullanılarak *IGF-15'UTR*'nin üç varyantında sekiz nükleotid değişikliği tespit edilmiştir: P1 varyantlarında benzersiz bir silme (g.171328230 delT), P2 varyantları rs401028781, rs422604851 ve g.171328404C > Y SNP'leri ile tanımlanmış, P3 varyantları ise P1 veya P2'de gözlemlenmeyen bir heterozigot substitüsyon (g.171328260G > R) ve üç homozigot substitüsyon (g.171328246T > A, g.171328257T > G, g.171328265T > C) içermektedir. Büyüme ve üretim özellikleri açısından istatistiksel olarak anlamlı bir fark, süttan kesimde göğüs genişliğinde ($p < 0.01$) ve bir yaşında bacak çevresinde ($p < 0.05$) bulunmuştur. P1 varyantları daha zayıf bir profil gösterirken Musculus

longissimus dorsi oranı daha yüksek olmasına rağmen farklar istatistiksel olarak anlamlı değildir ($p > 0.05$). P2 varyantlarında sırt ($p < 0.01$) ve fileto oranı ($p > 0.05$) daha yüksektir. Ayrıca, P3 varyantlarının boyun ve bacak oranı, P1 varyantlarının omuz oranından daha yüksek olmasına rağmen, varyantlar arasında belirgin bir fark bulunmamıştır. Sonuçlar, *IGF-15*'UTR'deki nükleotid değişikliklerinin, MAS stratejisi kullanılarak koyun karkas kalitesi özelliklerinin iyileştirilmesinde kullanılabileceğini ve farklı varyantlar arasındaki farkların tanınabileceğini göstermektedir.

SONUÇ VE DEĞERLENDİRME

Koyun yetiştiriciliği, küresel ölçekte çeşitli faydalar sağlayarak gelişmiş ve gelişmekte olan ülkeler için stratejik bir öneme sahiptir. Bu önem, kırsal kalkınma ve tarımsal gelişim politikaları çerçevesinde koyun yetiştiriciliğinin desteklenmesini zorunlu kılar. Özellikle hayvan ıslahı, koyun yetiştiriciliğinde verimliliği artırma, hastalık direncini güçlendirme ve adaptasyon kabiliyetini iyileştirme gibi önemli katkılar sunar. Türkiye'de koyun çiftlikleri, yerel ırkların genetik potansiyelini maksimize etmek için genetik seçim ve çaprazlama tekniklerini aktif olarak kullanmaktadır. Bu teknikler, yerel koşullara daha uyumlu, yüksek verim ve sağlık özelliklerine sahip koyunların yetiştirilmesini sağlar, böylece ülkenin hayvancılık sektörü sürekli olarak iyileştirilir. Bu stratejik yaklaşım, sürdürülebilir tarım uygulamaları ve kırsal ekonomik gelişim için temel bir yapı taşı oluşturmaktadır.

IGF-1 geni, dünya çapında çeşitli koyun ırkları üzerinde detaylı bir şekilde incelenmiş ve fenotipik özelliklerle olan bağlantısı istatistiksel olarak gösterilmiştir. Araştırmalar, *IGF-1* geninin büyüme hızı, kas gelişimi ve et verimi gibi önemli özelliklerle nasıl ilişkili olduğunu ve bu genetik varyasyonların koyun yetiştiriciliğinde ekonomik verimliliği nasıl etkileyebileceğini açığa çıkarmaktadır. Ancak Türkiye'de, *IGF-1* geninin polimorfizm çalışmaları koyun ırklarının genetik çeşitliliğinin incelenmesi konusunda yeterli çalışma yapılmamaktadır.

Genome-Wide Association Study (GWAS) ise, genetik varyasyonları genom çapında analiz ederek bu varyasyonların belirli hastalık veya özelliklerle olan ilişkilerini araştıran bir yöntemdir. GWAS, genetik farklılıkları tespit ederek bu farklılıkların özelliklerle olan bağlantılarını belirlemeye çalışır (Zhang vd., 2013; Ghasemi vd., 2019). Quantitative Trait Loci (QTL), kromozomlar üzerinde belirli noktalarda bulunan ve miktar niteliklerini (boy, ağırlık gibi sürekli özellikler) etkileyen gen veya gen gruplarını tanımlar. QTL haritalama, bu genlerin kromozomal konumlarını saptamayı amaçlar ve seleksiyonda hayvanın genetik değerini daha yüksek doğrulukla tahmin etmeyi ve böylece genetik kazancı artırmayı hedefler. Fenotiple ilgili fizyolojik olayları etkileyen genlerdeki varyasyonların, ilgili fenotipteki kantitatif varyasyonları etkileyebileceği bildirilmiştir (Tambasco vd., 2003; Margawati vd., 2006). Türkiye'de yapılan bazı çalışmalara rağmen (Yılmaz vd., 2022; Arzik vd., 2023; Argun Karlı vd., 2024; Yaman vd., 2024), bu çalışmaların artırılması ve birçok ırkta yapılması, Türkiye'de bulunan koyun ırklarından elde edilen et, süt, yapağı gibi özellikleri iyileştirmek için faydalı olacaktır. Bundan dolayı, gelecekte, GWAS ve QTL haritalama yöntemlerinin birlikte kullanılması, *IGF-1* polimorfizmlerinin koyun verimliliği üzerindeki genetik etkilerini daha kapsamlı bir şekilde ortaya koyabilir.

Türkiye'deki araştırmaların gerek koyun sayısı gerek koyun ırkı çeşitliliği göz önüne alındığında sınırlı olması, bu genetik faktörün hayvancılıktaki potansiyelinin tam olarak ortaya konulamamasına neden olmaktadır. Bu durum, genin yetiştiricilikteki rolünün ve potansiyel faydalarının tam olarak anlaşılmasını engellemekte ve sektördeki ilerlemeyi kısıtlamaktadır. Bu konuda daha fazla bilimsel araştırma yapılması, Türkiye'nin koyun ıslah çalışmalarına büyük katkılar sağlayacak ve özellikle *IGF-1* genindeki polimorfizmlerin detaylı incelenmesi, yerli koyun ırklarının genetik çeşitliliğini anlamamızı ve ıslah stratejilerinin etkinliğini artırmamızı sağlayacaktır. *IGF-1*'nin koyunların et verimi, büyüme hızları ve genel sağlık durumları üzerindeki etkilerinin daha iyi anlaşılması, verimli koyun yetiştiriciliği için genetik seçim programlarının geliştirilmesine olanak tanıyacak ve bu da koyun yetiştiriciliğinin verimliliğini artırarak

Türkiye'nin hayvancılık sektöründeki ekonomik büyümesine ve sürdürülebilirliğine önemli katkılarda bulunacaktır. Dolayısıyla, *IGF-I* geni üzerindeki polimorfizm araştırmalarına yoğunlaşmak, ülkenin koyun yetiştiriciliği alanındaki rekabet gücünü artırmak için kritik bir adımdır. Bu çabalar, aynı zamanda tarım sektöründeki yenilikçi yaklaşımları teşvik edecek ve ekonomik büyümeye katkıda bulunacaktır.

Genel olarak, Türkiye'deki koyun yetiştiriciliği, hem ekonomik hem de ekolojik açıdan büyük bir sürdürülebilirlik potansiyeline sahiptir. Gelecekte, genetik seleksiyon çalışmalarının yanı sıra sürdürülebilir koyun yetiştiriciliği politikaları, Türkiye'nin küçükbaş hayvancılık sektörünü daha rekabetçi ve verimli bir hale getirilmesinde kritik bir rol oynayacak. Bu gelişmeler, ülkenin hayvancılık endüstrisinin uluslararası pazarda daha güçlü bir konum kazanmasına yardımcı olabilir ve aynı zamanda yerel ekonomilere de önemli katkılarda bulunabilir.

KAYNAKÇA

- Armstrong, E., Ciappesoni, G., Iriarte, W., Da Silva, C., Macedo, F., Navajas, E. A., ... & Postiglioni, A. (2018). Novel genetic polymorphisms associated with carcass traits in grazing Texel sheep. *Meat science*, *145*, 202-208.
- Arzik, Y., Kizilaslan, M., Behrem, S., White, S. N., Piel, L. M., & Cinar, M. U. (2023). Genome-wide scan of wool production traits in Akkaraman sheep. *Genes*, *14*(3), 713.
- Argun Karsli, B., Demir, E., Bilginer, U., Dogru, H., Karsli, T., & Kaya, S. (2024). Genome-wide discovery of selection signatures in four Anatolian sheep breeds revealed by ddRADseq. *Scientific Reports*, *14*(1), 20518.
- Bahrami, A., Behzadi, S., Miraei-Ashtiani, S. R., Roh, S. G., & Katoh, K. (2013). Genetic polymorphisms and protein structures in growth hormone, growth hormone receptor, ghrelin, insulin-like growth factor 1 and leptin in Mehraban sheep. *Gene*, *527*(1), 397-404.
- Baker, J., Liu, J. P., Robertson, E. J., Efstratiadis, A. 1993. Role of insulin-like growth factors in embryonic and postnatal growth. *Cell*, *75*:73-82.
- Bakhtiar, R., Abdolmohammadi, A., Hajarian, H., Nikousefat, Z., & Kalantar-Neyestanaki, D. (2017). Investigation of the 5' flanking region and exon 3 polymorphisms of IGF-1 gene showed moderate association with semen quality in Sanjabi breed rams. *Theriogenology*, *104*, 186-191.
- Bayraktar, M., & Shoshin, O. (2021). Estimate of the association of *IGF-I* and *IGFALS* genes with growth traits in Hamdani sheep. *Brazilian Archives of Biology and Technology*, *64*, e21210262
- Bayram, D., Akyüz, B., Arslan, K., Ozdemir, F., Aksel, E., & Çınar, M. (2019). *DGAT1*, *CAST* and *IGF-I* gene polymorphisms in Akkaraman lambs and their effects on live weights up to weaning age. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, *25*(1).
- Casas-Carrillo, E., Kirkpatrick, B.W., Prill-Adams, A., Price, S.G., Clutter, A.C. 1997. Relationship of growth hormone and insulin-like growth factor-1 genotypes with growth and carcass traits in swine. *Animal Genetics*, *28* (2): 88-93.
- Çiçek, A., Ayyıldız, M., Erdal, G., & Erdal, H. (2022). Türkiye'de koyun yetiştiriciliğinin önemi ve ekonomik analizi. *MAS Journal of Applied Sciences*, *7*(Özel Sayı), 1303-1322.
- Darwish, H. R., El-Shorbagy, H. M., Abou-Eisha, A. M., El-Din, A. E., & Farag, I. M. (2017). New polymorphism in the 5' flanking region of IGF-1 gene and its association with wool traits in Egyptian Barki sheep. *Journal of Genetic Engineering and Biotechnology*, *15*(2), 437-441.
- Dekkers, J. C. (2004). Commercial application of marker-and gene-assisted selection in livestock: strategies and lessons. *Journal of animal science*, *82*(suppl_13), E313-E328.
- Ghasemi, M., Zamani, P., Vatankhah, M., & Abdoli, R. (2019). Genome-wide association study of birth weight in sheep. *Animal*, *13*(9), 1797-1803.

- Goddard, M. E., & Hayes, B. J. (2009). Mapping genes for complex traits in domestic animals and their use in breeding programmes. *Nature Reviews Genetics*, 10(6), 381-391.
- Ertuğrul, M., Dellal, G., Soysal, İ., Elmacı, C., Akın, O., Arat, S., ... & Yılmaz, O. (2009). Türkiye yerli koyun ırklarının korunması. *Uludağ Üniversitesi Ziraat Fakültesi Dergisi*, 23(2), 97-119.
- Ertürk, Y. E., Yalcin, S., and Yılmaz, O. (2015). Livestock support policies in Turkey since 2000. Vi. Sci. Agric. Symposium” Agrosym 2015, 2019–2027.
- Gharahveysi, S., & Reykandeh, H. R. B. (2015). The genetic study of polymorphism of IGF-1 gene in Zel sheep. *International Journal of Advanced Life Sciences*. 8(1).
- Günaydın, G. (2009). Koyun yetiştiriciliğinin ekonomi politiği. *Uludağ Üniversitesi Ziraat Fakültesi Dergisi*, 23(2), 15-32.
- Grochowska, E., Borys, B., Janiszewski, P., Knapik, J., & Mroczkowski, S. (2017). Effect of the *IGF-I* gene polymorphism on growth, body size, carcass and meat quality traits in Coloured Polish Merino sheep. *Archives animal breeding*, 60(2), 161-173.
- Hajihosseini, A., Hashemi, A., Razavi-Sheshdeh, S.A., Pirany, N. 2013. Association of the polymorphism in the 5' flanking region of the ovine *IGF-I* gene with growth and development traits in Makui sheep of Iran. *European Journal of Zoological Research*, 2 (4): 19-24.
- He, J. N., Zhang, B. Y., Chu, M. X., Wang, P. Q., Feng, T., Cao, G. L., ... & Li, N. (2012). Polymorphism of insulin-like growth factor 1 gene and its association with litter size in Small Tail Han sheep. *Molecular biology reports*, 39, 9801-9807.
- Kader Esen, V., & Esen, S. (2023). Association of the IGF1 5' UTR polymorphism in meat-type sheep breeds considering growth, body size, slaughter, and meat quality traits in Turkey. *Veterinary Sciences*, 10(4), 270.
- Kandemir, Ç., & Taşkın, T. (2022a). Türkiye’de Koyun Irklarının Bölgelere Göre Durumu: İç Anadolu Bölgesi. *Uluslararası Anadolu Ziraat Mühendisliği Bilimleri Dergisi*, 4(4), 97-105.
- Kandemir, Ç., & Taşkın, T. (2022b). Türkiye’de Koyun Irklarının Mevcut Durumu ve Geleceği: Karadeniz Bölgesi. *Yüzüncü Yıl Üniversitesi Fen Bilimleri Enstitüsü Dergisi*, 27(1), 101-112.
- Kandemir, Ç., & Taşkın, T. (2022c). Türkiye’de Koyun Irklarının Mevcut Durumu ve Geleceği: Akdeniz Bölgesi. *Türkiye Tarımsal Araştırmalar Dergisi*, 9(1), 97-106.
- Kaplan, S., & Atalay, S. (2018). Single nucleotide polymorphism of ovine leptin and insulin-like growth factor 1 gene in Kivircik crossbred ewes. *Pakistan Journal of Zoology*. vol. 50(3), pp 851-856, 2018.
- Kaymakçı, M., Eliçin, A., Tuncel, E., Pekel, E., Karaca, O., Işın, F., ... & Sönmez, R. (2000). Türkiye’de küçükbaş hayvan yetiştiriciliği. *Türkiye Ziraat Mühendisliği V. Teknik Kongresi*, 2, 765-793.
- Kumar, S., Dahiya, S. P., Magotra, A., Ratwan, P., & Bangar, Y. (2023). Influence of single nucleotide polymorphism in the IGF-1 gene on performance and conformation traits in Munjal sheep. *Zygote*, 31(1), 70-77.
- Kazemi, S. M., Amirinia, C., Emrani, H., & Gharahveysi, S. (2011). Study and identification of Insulin-like Growth Factor-I gene polymorphisms in Zel sheep population. *American Journal of Animal and Veterinary Sciences*. 6 (4): 176-179.
- Lenis-Valencia, C., Hernández-Herrera, D., & Álvarez-Franco, L. (2021). Polymorphisms of the CAPN, CAST, LEP, GH, GHR, IGF-1 and MSTN loci of Colombian Creole hair x Pelibuey sheep sheep crossbreeds. *Veterinarska stanica*, 52(1), 35-44.
- Margawati, E. T., Raadsma, H. W., & Martojo, H. (2006). Quantitative trait loci (QTL) analysis for production traits of birth weight and weight 360 days in backcross sheep. *HAYATI Journal of Biosciences*, 13(1), 31-35.

- Nazari, F., Noshary, A., & Hemati, B. (2016). Association between insulin-like growth factor I polymorphism and early growth traits in Iranian Zandi sheep, found Polymerase Chain Reaction Restriction Fragment Length Polymorphism (PCR-RFLP). *Iranian Journal of Applied Animal Science*, 6(3), 665-669.
- Raji, A. O., Mohammed, A., Igwebuike, J. U., & Alphonsus, C. (2017). Association of IGF 1 gene polymorphisms with some morphometric traits of Nigerian indigenous sheep breeds. *Nigerian Journal of Biotechnology*, 34, 97-104.
- Ramos, Z., Garrick, D. J., Blair, H. T., Vera, B., Ciappesoni, G., & Kenyon, P. R. (2023). Genomic regions associated with wool, growth and reproduction traits in Uruguayan Merino sheep. *Genes*, 14(1), 167.
- Siadkowska, E., Zwierzchowski, L., Oprzadek, J., Strzalkowska, N., Bagnicka, E., Krzyzewski, J. 2016. Effect of polymorphism in IGF-1 gene on production traits in Polish Holstein-Friesian cattle. *Animal Science Papers and Reports*, 24 (3): 225-237.
- Tambasco, D.D., Paz, C.C.P., Pereira, AP., Alencar, MM ., Freitas, A.R., Coutinho, L.L., Packer, I.U. and Regitano, L.C.A., 2003. Candidate genes for growth traits in beef cattle crosses. *J. Anim. Breed. Genet.* 120, 51–56.
- Taşkın, T., & Kandemir, Ç. (2022a). Doğu Anadolu Bölgesinde Yetiştirilen Koyun Irklarının Mevcut Durumu. *Journal of Animal Production/Hayvansal Üretim*, 63(1).
- Taşkın, T., & Kandemir, Ç. (2022b). Türkiye’de koyun ırklarının mevcut durumu ve geleceği: Ege Bölgesi. *Ege Üniversitesi Ziraat Fakültesi Dergisi*, 59(3), 485-498.
- Taşkın, T., & Kandemir, Ç. (2022c). Marmara bölgesinde yerli ve kültür koyun ırklarının mevcut durumu. *Doğanın Sesi*, 5(9), 17-33.
- Taşkın, T., & Kandemir, C. (2022d). Türkiye’de Koyun Irklarının Mevcut Durumu ve Geleceği: Güneydoğu Anadolu Bölgesi. *Kadirli Uygulamalı Bilimler Fakültesi Dergisi*, 2(1), 93-105.
- Tawfeeq, M., Keskin, İ., & İlhan, F. (2020). The Identification of Genetic Variation in Insulin-Like Growth Factors-I (*IGF-I*) Gene Region in Some Turkish Shee. *Selcuk Journal of Agriculture and Food Sciences*, 34(3), 189-192.
- (TÜİK, 2025). Türkiye İstatistik Kurumu Bitkisel Üretim ve Hayvancılık İstatistikleri. www.tuik.gov.tr (Erişim 10 Şubat 2025).
- Umego, C. M., Kabir, M., Adeyinka, I. A., Alao, R. O., Mallam, I., Ibrahim, O. A., & Jinadu, L. A. (2018). Single nucleotide polymorphism in the insulin-like growth factor-1 gene and its effects on growth traits in Yankasa sheep. *Nigerian Journal of Animal Science*, 20(4), 323-332.
- Valencia, C. P. L., Franco, L. Á. Á., & Herrera, D. H. (2022). Association of single nucleotide polymorphisms in the CAPN, CAST, LEP, GH, and IGF-1 genes with growth parameters and ultrasound characteristics of the Longissimus dorsi muscle in Colombian hair sheep. *Tropical animal health and production*, 54(1), 82.
- Yaman, Y., Doğan, Ş., Kirbaş, M., Önalı, A. T., & Kal, Y. (2024). Investigation of growth traits in Turkish Merino lambs using multi-locus GWAS approaches: Middle Anatolian Merino. *BMC Veterinary Research*, 20(1), 567.
- Yılmaz, O., Kizilaslan, M., Arzik, Y., Behrem, S., Ata, N., Karaca, O., ... & Cemal, I. (2022). Genome-wide association studies of preweaning growth and in vivo carcass composition traits in Esmé sheep. *Journal of Animal Breeding and Genetics*, 139(1), 26-39.
- Zhang, L., Liu, J., Zhao, F., Ren, H., Xu, L., Lu, J., ... & Du, L. (2013). Genome-wide association studies for growth and meat production traits in sheep. *PloS one*, 8(6), e66569.

**SUSTAINABLE AGRICULTURE AS A REGULATORY AND SUPPORTIVE
ACTIVITY OF PUBLIC ADMINISTRATION****İDARENİN DÜZENLEYİCİ VE DESTEKLEYİCİ BİR FAALİYETİ OLARAK
SÜRDÜRÜLEBİLİR TARIM****Doç. Dr. Asuman ÇAPAR**

Sivas Cumhuriyet Üniversitesi Hukuk Fakültesi

ÖZET

Son yıllarda dünya üzerindeki yaşamsal kaynakların sınırlılığının daha da fark edilmeye başlaması ile sürdürülebilirlik kavramı ön plana çıkmaya başlamıştır. Çok yönlü bir boyutu olan sürdürülebilirlik kavramı genel itibariyle, insan türünün günlük ihtiyaçlarını karşılarken, gelecek kuşakların ve doğanın kendi ihtiyaçlarını gelecekte karşılayabilme yeteneğine zarar vermemesi olarak ifade edilebilir. Sürdürülebilirliğin önemli bir boyutunu tarım oluşturmaktadır. İnsanın mevcut ve gelecekteki doğal yaşamını devam ettirebilmesinde, tarımın hayati bir yeri bulunmaktadır. Bu nedenle devletler, sürdürülebilir tarım politikaları geliştirmektedirler. Tarımsal faaliyetlerin desteklenmesi, tarımın korunması, güvenliğin ve sürdürülebilirliğin sağlanması, temelde kendisinin ve milletin varlığını korumak yükümlülüğü olarak devletin asli görevleri arasındadır. Türkiye’de devlet idaresi altında bu alanda faaliyette bulunan Cumhurbaşkanı, Tarım ve Orman Bakanlığı gibi organlar olduğu gibi diğer kamu tüzel kişileri arasında da çeşitli idari organlar bulunmaktadır. Temel olarak bu kurum ve kuruluşlar, kanuni yetkileri çerçevesinde düzenleyici, destekleyici ve denetleyici idari faaliyetlerde bulunmaktadırlar. Bu çerçevede sürdürülebilir tarımı destekleyici ve teşvik edici devlet politikaları güdülmektedir. Kalkınma planlarında ve stratejik planlarda sürdürülebilir tarım konusuna yer verilerek planlamalar yapılmaktadır. Belirlenen bu politikalarda yalnızca günün ihtiyaçlarını karşılamak hedeflenmemekte, aynı zamanda insan sağlığının ve ekolojinin korunması sağlanarak, ilerleyen süreçte de ülke ihtiyaçlarının karşılanabilir olması hedeflenmektedir. Coğrafi özellikler olarak tarım ülkesi olmasına rağmen, Türkiye’de sürdürülebilir tarımı tehdit eden bazı faktörler bulunmaktadır. Kuraklık ve yanlış tarım uygulamaları bunların başında gelmektedir. Bu konuda öngörülen planlamaların hayata geçirilmesi, sürdürülebilir tarımı destekleyici politikaların uygulamaya konulması, istikrarlı şekilde yürütülmesi ve eksiklerin tamamlanması önem taşımaktadır.

Anahtar Kelimeler: Tarım, Sürdürülebilirlik, Su yönetimi, Düzenleyici ve destekleyici idari faaliyetler.

ABSTRACT

In recent years, the concept of sustainability has come to the forefront as the limited nature of vital resources on Earth has become more apparent. Sustainability, which has a multidimensional aspect, can generally be defined as meeting the daily needs of the human species without compromising the ability of future generations and nature to meet their own needs. Agriculture constitutes an important aspect of sustainability. It plays an important role in ensuring the continuation of human life, both at this moment and in the future. For this reason, governments develop sustainable agricultural policies. Supporting agricultural activities, protecting agriculture, ensuring food security, and maintaining sustainability are fundamental responsibilities of the state, originating from its duty to preserve both itself and the nation. In Türkiye, state administrative

bodies such as the Presidency and the Ministry of Agriculture and Forestry, as well as other public legal entities, include various administrative organs operating in this field. These institutions and organizations primarily engage in regulatory, supportive, and supervisory administrative activities within the framework of their legal authority. Within this scope, state policies are implemented to support and promote sustainable agriculture. Sustainable agriculture is incorporated into development plans and strategic plans, ensuring that necessary measures are taken. These policies not only aim to meet immediate needs but also protect human health and ecology in order to ensure that the country's agricultural needs can continue to be met in the future. Even though Türkiye is an agricultural country in terms of its geographical characteristics, certain factors threaten the sustainability of agriculture. Drought and improper agricultural practices are among the most important ones. The implementation of planned strategies, the enforcement and consistent execution of policies that support sustainable agriculture, and the rectification of deficiencies are of great importance in this regard.

Keywords: Agriculture, Sustainability, Water Management, Regulatory and supportive administrative activities

GİRİŞ

Biyolojik çeşitlilik ve doğal kaynaklar üzerindeki dünyadaki baskılar ve iklim değişikliğinin etkileri artarken, gıda sistemlerinin sosyal, ekonomik ve çevresel açıdan daha üretken ve sürdürülebilir olması gerekliliği ön plana çıkmaktadır. Bu çerçevede, bir taraftan çevre dostu üretimin artması, yeni uygulama teknikleriyle verimli ve etkin üretimin geliştirilmesi önem arz ederken, diğer taraftan israfın ve gıda kayıplarının azaltılması gerekmektedir (Türkiye Tarımsal Kuraklıkla Mücadele Stratejisi ve Eylem Planı 2023-2027 [TTKMEP], 2022: 30).

Türkiye ekonomisinin temel taşlarından birisini teşkil eden tarımın önemi ülkemiz ve dünya için her geçen gün artmaktadır. Tüketim alışkanlıklarının değişmesi, nüfus artışı, küresel ısınma sebebiyle gıdaya erişimin önümüzdeki dönemlerde daha pahalı ve zor hale geleceği, böylelikle 2050'ye kadar mevcut gıda tüketiminin %60 oranında artma ihtimali, tarımın önemini daha da artırmaktadır. Tarım alanları sınırlı olduğu için bu konuda tek çözüm verimliliği artırmak olarak görülmektedir. Bununla birlikte, Türkiye'de tarımsal üretim giderek artmasına ve çeşitlenmesine rağmen, tarımın ekonomimizdeki yeri giderek azalmaktadır (Akbaş, 2019: 35). Bu noktada sürdürülebilir tarım konusu, ülkede çok yönlü boyutu ile önemli gündem başlıkları arasında yerini korumaktadır.

Bu çalışma ile sürdürülebilir tarım konusunda idarenin düzenleme ve destekleme faaliyetlerine, kuşbakışı ve bütüncül bir şekilde bakılması hedeflenmektedir. Konu incelenirken hem devlet faaliyetleri hem de diğer kamu düzel kişileri bakımından sürdürülebilir tarım konusundaki düzenleyici ve destekleyici faaliyetler genel olarak değerlendirilecektir. Sürdürülebilir tarım konusunda faaliyette bulunan uluslararası kuruluşlar bu çalışma kapsamına girmemektedir.

SÜRDÜRÜLEBİLİRLİK KAVRAMI VE TARIM

Sürdürülebilirlik kavramı esasen çok boyutlu bir kavram olup insanın ve doğadaki diğer canlıların yaşamsal faaliyetlerinin devamlılığı bakımından etki edebilecek pek çok faktörü kapsamına almaktadır. Bu bağlamda sürdürülebilirlik kavramına ve sürdürülebilirliğin tarımsal boyutuna değinilecektir.

Kavramsal Olarak Sürdürülebilirlik

Bir şeyin sürdürülebilir olması, onun mevcut durumunu devam ettirebiliyor olması veya kendini yenileyebiliyor olması anlamına gelmektedir. Bu minvalde sürdürülebilirlik kavramı gelecek

nesillere ekolojik, sosyal ve ekonomik şartları devam ettirebilir bir dünya bırakmak anlamında kullanılmaya başlanmıştır. Bu yönüyle sürdürülebilirlik, sadece yaşayanları değil gelecek nesilleri de ilgilendirmektedir. Sürdürülebilirlik kelimesi bu anlamıyla ilk kez 1983 yılında Birleşmiş Milletlerin yayınlamış olduğu *Ortak Geleceğimiz* isimli raporda kullanılmıştır. Buna göre sürdürülebilirlik gelecek kuşakların ve doğanın kendi gereksinimlerini karşılayabilme yeteneklerini tehlikeye düşürmeden, günlük ihtiyaçlarımızı gidermek ve kalkınmak olarak ifade edilmiştir. Sürdürülebilirlik kavramı ekonomik, toplumsal ve ekolojik boyutları içeren bütünsel bir yaklaşımdır (Güner, 2020: 18).

Bir şeyin sürdürülebilmesi için o şeyin korunması ve ona değer verilmesi ve o şeyin belirli bir yaşam döngüsünde devam ettiriliyor olması gerekir. Bu, senelerce süren bir devamlılığın yanı sıra amaçlara ulaşmada da doğru kaynakların ve yolların kullanılması anlamına gelir. Bu süreçte tüm yaşam döngüsü bir bütün olarak değerlendirilmelidir. Küresel sistemi, ekoloji, enerji, ekonomi gibi yaşamsal faktörleri korumak için toplumsal ve fiziksel bir vizyon oluşturulmalıdır. Bunu sağlamak için toplumsal ve bireysel olarak sosyal açıdan sorumluluklarımızın bilincinde olmamız gerekir (Güner, 2020: 19-20).

Sürdürülebilirlik gelecek nesillerin genel refahını ve gıda güvenliğini tehlikeye düşürmeden çevresel bir maliyetle gıda ürünlerinin ve diğer tarımsal ürünlerin üretilebilmesidir. Sürdürülebilirlik, gelecek nesillerin ihtiyaçlarının karşılanabilmesi amacıyla bugünün kaynaklarının gereksiz bir şekilde kullanıma izin verilmemesi ilkesine dayanır. Bu nedenle sürdürülebilirlik konusunda doğal kaynakların ve insan kaynaklarının yönetimi birinci derecede önem taşımaktadır. İklim değişikliği insanları ve dünyamızı tehdit ederken sürdürülebilir tarıma geçiş oldukça önem taşımaktadır. Kırsal kalkınma çalışmaları yoksulluğu azaltır, güçlü bir tarım sektörünü destekler ve yanında sürdürülebilir ekonomik kalkınmayı sağlar (Kubaş ve Aydın, 2022: 56-60).

Tarımın Sürdürülebilirlik Boyutu

Tarım konusu hem ekonomik boyutu olan, hem toplumsal boyutu olan hem de ekolojik boyutu olan bir konudur. Sürdürülebilirlik kavramının, tarımın bütün boyutlarıyla ilgili olduğu ifade edilebilir.

Hayvanların, insanların ve hatta doğada yaşayan bütün canlıların hayati döngülerini sağlayan en önemli temel dayanaklardan birisini tarımsal üretim oluşturmaktadır. Tarımsal üretim bakımından vazgeçilmez unsur ise sürdürülebilir tarımdır (Baran vd., 2021: 1). Sürdürülebilir tarım kavramı aslında endüstriyel tarımın neden olduğu sorunları çözebilmek amacıyla ileri sürülmüş çözüm yöntemlerinin altında toplandığı bir başlık olarak da görülebilir. Burada öncelikle doğal dengeyi bozarak istenilen ölçüde ve sürekli bir gelişmenin sağlanamayacağı ve tarımsal üretim bakımından dünyada gerekli olan kaynakların sınırsız olmadığı kabul edilmelidir (Turhan, 2005: 14).

Sürdürülebilir tarım, uzun vadede doğal kaynakların korunması yanında çevreye zarar vermeyen teknolojik metodların kullanıldığı bir tarımsal yapının oluşturulmasıdır (Turhan, 2005:13). Sürdürülebilir tarım uygulamaları aynı zamanda doğal yaşama katkı sunmaktadır. Toprağın verimliliğini artırarak nesli tükenme tehlikesi altındaki hayvanların ve bitkilerin varlıklarına önemli katkıda bulunur. Bundan dolayı sürdürülebilir tarım uygulamaları verimi ve karlılığı artırır (Kubaş ve Aydın, 2022: 60).

Tarımın sürdürülebilir olma potansiyeli, bugün itibariyle yeterli gıdayı ve diğer tarımsal ürünleri, çevre ve insan refahını teşvik edecektir. Gelecek nesillerin bunu yapabileceğini koruyacak biçimde üretebilme potansiyeli, sürdürülebilir tarım bakımından kuvvetlidir. Daha çok yoğunlaşmanın, iklim değişikliğinin, biyolojik çeşitlilik kaybının ve diğer çevresel değişikliklerin sürdürülebilirlik zorluklarını karşılamak zor olacaktır. Fakat doğru teşvikler, yenilikçi araştırmalar ve siyasi iradeyle bu gerçekleşebilir (Kubaş ve Aydın, 2022: 60).

Sürdürülebilir tarımın desteklemesi amacıyla getirilebilecek bazı öneriler bulunmaktadır. Öncelikle üreticilerin kullandıkları ilaç ve gübre miktarı tam olarak belirlenerek ihtiyaçlara göre kullanılması sağlanmalı, gereksiz kullanımlardan kaçınılmalıdır. Kimyasalların ve gübrelerin uygulama vakitlerinin de çok iyi ayarlanması gerekmektedir. Zamansız, düzensiz veya aşırı yapılan işlemler yarardan çok zarar getirebilecektir. Bitki kalıntıları, hayvan gübresi ve besin maddeleri işleme artıkları gibi birtakım organik maddelerin kullanımları yaygınlaştırılmalıdır. Tarım topraklarının verimliliklerinin sürdürülebilmesi için gerekli kanuni düzenlemeler yapılmalıdır. Zararlılarla mücadelede mümkün olduğunca biyolojik mücadele teknikleri kullanılmaya çalışılmalıdır. Ayrıca biyoteknolojik gelişmeler takip edilerek bu konularda pratiğe aktarılacak faydalı çalışmalar emsal alınabilir (Karadavut vd., 2004: 65).

Tarımsal üretimin doğaya zarar vermeden yapılması gereği bugün artık iyice ortaya çıkmıştır. Bu alanda hedefe ulaşabilmek için öncelikle mevcut problemlerin doğru tespit edilmesi gerekmektedir. Bunu sağlayabilmek için su kaynaklarının kirlenmesini, toprağın tuzlaşmasını, erozyonu ve diğer zararları en aşağıya çekebilen sürdürülebilir tarım tekniklerinin geliştirilmesi gerekmektedir. Dünyada besin üretimini arttırmanın yolları aranırken, tarımda kullanılan doğal kaynakları da güvence altına alacak yeni yöntemler geliştirme zorunluluğu doğmuştur. Özellikle, uygulanan yoğun ve yanlış girdili yetiştirme teknikleri tarım bölgelerinde geri dönüşü olmayan zararlara neden olabilmektedir (Turhan, 2005: 22).

Sürdürülebilir tarım, tarımsal üretimde çevresel, sosyal, agronomik ve ekonomik boyutları dengelemeyi amaçlamaktadır. Tarımda verimliliği muhafaza ederken diğer taraftan da çevreye verilen zararı azaltarak, kısa ve uzun vadede ekonomiyi canlı tutarak, tarımla uğraşanların yaşam kalitesini yükseltmek önemlidir. Ayrıca sürdürülebilir bir yaşam açısından mutlaka geleceğe ilişkin planlamalar yapılmalı ve bunların kişilerin yaşam standartlarını düşürmeyecek, bireysel hürriyetlerini sınırlamayacak nitelik taşımasına özen gösterilmelidir. Doğal kaynakların korunması sürdürülebilir bir tarım açısından zorunluluktur. Aksi taktirde bir dönem bakımından geleceği düşünmeden ve aşırı şekilde yapılan etkinlikler neticesinde insanların uzun dönemde güçlüklerle karşılaşması kaçınılmazdır. Bundan dolayı ilk olarak bu konudaki hedefler iyi belirlenerek açıkça ortaya konulmalı, sürdürülebilir olmayan tarımsal etkinliklere destek verilmemelidir. Çevreye zarar verebilecek faaliyetlere karşı devlet önlemler almalıdır. Örneğin toprağın aşırı şekilde sulanmasına, yanlış ilaçlanmasına ve gübrenmesine karşı eğitim yapılmalı ve kontrol önlemleri alınmalıdır. Bu bakımdan sürdürülebilir tarım konusunda bir veri tabanı oluşturulmalı, bunun için gerekli sosyal, ekonomik ve teknik veri bankalarının uygulamaları geliştirilmelidir (Turhan, 2005: 14).

Sürdürülebilir bir şekilde üretim yapmak, gerekli sosyal ve yerel şartları sağlamak, yüksek seviyede gıda kalitesini teşvik etmek bakımından çiftçilik alternatiflerinden birisi organik tarımdır (Kubaş ve Aydın, 2022: 60). Sürdürülebilir tarım anlayışında toprak ve su kaynaklarının korunması, entegre ilaç yönetimi gibi bazı uygulamalar yer almakla birlikte, sentetik gübre, ilaç gibi doğal olmayan yöntemlerin kullanılmasından kaçınılarak kalite, sağlık ve çevresel standartlarla buluşan organik tarım teknikleri önemli rol oynamaktadır (Turhan, 2005:14).

Birleşmiş Milletler 2030 Sürdürülebilir Kalkınma Amaçları arasında “2030’a kadar sürdürülebilir gıda üretimi sistemlerinin güvence altına alınması ve üretimi ve üretkenliği artıran, ekosistemlerin sürdürülmesine yardımcı olan, iklim değişikliğine, aşırı hava koşullarına, kuraklığa, sellere ve diğer felaketlere uyum sağlama kapasitesini güçlendiren ve toprak kalitesini devamlı olarak artıran dayanıklı tarım uygulamalarının hayata geçirilmesi” bulunmaktadır (<https://turkiye.un.org/tr/sdgs/2>, 05.02.2025).

Kırsal alanlar, kaynakların bilinçsiz kullanımı ve nüfus artışı gibi nedenlerle baskı altındadır. Bu durum, ülkemizde olduğu gibi gelişmekte olan ülkelerde daha belirgindir. Özellikle, kırsal alanlarda tarımdan kaynaklanan su ve toprak kirliliği, hatalı ilaçlama ve gübreleme ile tarım topraklarının yanlış kullanımı gibi sebeplerle doğal kaynaklara zarar verilmektedir. Sürdürülebilir

tarım, yalnızca insan ve doğal kaynaklar arasındaki dengenin sağlanması ve kırsal çevrenin korunmasıyla mümkündür (Karadavut vd., 2004: 61).

SÜRDÜRÜLEBİLİR TARIM KONUSUNDA TÜRKİYE’DEKİ GENEL ÇERÇEVE

Türkiye’de sürdürülebilir tarım konusunda devreye giren idari organlar ve bunların görev ve faaliyet alanlarına genel esaslarıyla değinilecektir.

Sürdürülebilir Tarım Konusunda Sorumlu İdari Organlar

Devletin ve milletin varlığının devamı için gerekli tedbirleri almak devletin temel anayasal görevleri arasındadır. Bu bakımdan yaşamsal önem taşıyan tarımsal faaliyetlerde yetkili ve görevli organlar konusunda ilk olarak devlet idaresini görüyoruz. Tarımın ve bu alanda çalışanların korunması, desteklenmesi Anayasa’nın 45. maddesi¹ ile devlete görev olarak yüklenmiştir. Bu kapsamda ilk olarak merkezi idarenin başında bulunan Cumhurbaşkanının Anayasal ve kanuni yetkileri çerçevesinde düzenleme ve denetleme yetkisi bulunmaktadır.

Bakanlıklar arasında temel olarak tarım ve ormancılık alanında faaliyette bulunmak üzere Tarım ve Orman Bakanlığı (TOB) bulunmaktadır. Bu alanda yalnızca TOB değil, bunun koordinatörlüğünde ilgili bakanlıklar, valilikler, başkanlıklar, üniversiteler, yerel yönetimler ve sivil toplum kuruluşları da faaliyette bulunmaktadır. Başkent teşkilatında konularına ve ihtiyaçlara göre çeşitli müdürlükler şeklinde bir örgütlenme görülmektedir. Başkentteki bakanlık örgütünün taşrada uzantıları bulunmakta olup, başta Tarım ve Orman İl Müdürlükleri olmak üzere, İlçe Müdürlükleri bulunmaktadır. İl ve ilçe müdürlükleri haricinde taşrada da konularına ve ihtiyaca göre merkeze bağlı, onun hiyerarşisi altında faaliyet gösteren çeşitli müdürlükler bulunmaktadır.

Tarım alanında faaliyette bulunan idari organlar bunlardan ibaret olmayıp, Orman Genel Müdürlüğü ve Devlet Su İşleri Müdürlüğü gibi devletten ayrı bazı kamu tüzel kişilikleri de görev yürütmektedir. Bunların haricinde Toprak Mahsulleri Ofisi Genel Müdürlüğü, Tarım ve Kırsal Kalkınmayı Destekleme Kurumu ve koordineli çalışılan çeşitli ilgili genel müdürlükler bulunmaktadır. Bu alanda çok çeşitli idari organlar faaliyette bulunmakla birlikte, amaçlara ulaşabilmek için bunların koordineli, eş zamanlı ve uyumlu bir politika gütmesi gerekmektedir.

Bu alanda görevli kurum ve kuruluşların çokluğu aynı zamanda mevzuatta da çokluğu ve dağınıklığı beraberinde getirmiştir. Bu nedenle konuya ilişkin pek çok kanunu görmek mümkündür. Bunlara örnek olarak 5488 Sayılı Tarım Kanunu, 2872 Sayılı Çevre Kanunu, 5363 Sayılı Tarım Sigortaları Kanunu, 4342 Sayılı Mera Kanunu, Gıda ve Yem Kanunu, 6831 Sayılı Orman Kanunu, 1380 Sayılı Su Ürünleri Kanunu, 5442 Sayılı İller İdaresi Kanunu, 5996 sayılı Veteriner Hizmetleri, Bitki Sağlığı, 5403 Sayılı Toprak Koruma ve Arazi Kullanma Kanunu, 167 Sayılı Yeraltı Suları Hakkında Kanun, 2090 Sayılı Tabii Afetlerden Zarar Gören Çiftçilere Yapılacak Yardımlar Hakkındaki Kanun, 831 Sayılı Sular Hakkında Kanun gibi kanunlar örnek gösterilebilir.

Sürdürülebilir Tarımı Düzenleyici ve Destekleyici İdari Faaliyetler

Yukarıda belirtilen kurum ve kuruluşlar, Anayasa’dan ve ilgili kanunlardan aldıkları yetkiler çerçevesinde, sürdürülebilir tarım konusunda düzenleyici ve destekleyici faaliyetlerde bulunma

¹ Anayasa madde 45, “Tarım, hayvancılık ve bu üretim dallarında çalışanların korunması”: “Devlet, tarım arazileri ile çayır ve mer’aların amaç dışı kullanılmasını ve tahribini önlemek, tarımsal üretim planlaması ilkelerine uygun olarak bitkisel ve hayvansal üretimi artırmak maksadıyla, tarım ve hayvancılıkla uğraşanların işletme araç ve gereçlerinin ve diğer girdilerinin sağlanmasını kolaylaştırır. Devlet, bitkisel ve hayvansal ürünlerin değerlendirilmesi ve gerçek değerlerinin üreticinin eline geçmesi için gereken tedbirleri alır.”

yetkisine sahiptirler. Bu amaçla ilgili idarelerin gerektiği hallerde düzenleyici ve bireysel nitelik taşıyan işlemler yapma yetkisi bulunmaktadır.

TOB'un 2024-2028 Stratejik Planı arasında “*Tarım, orman ve su kaynakları ile doğal ekosistemleri koruyarak, verimli ve sürdürülebilir tarımsal üretimi, yeterli ve güvenilir gıdaya erişimi ve kırsal kalkınmayı sağlamak amacıyla politikalar belirlemek ve uygulamak*” misyonu ve “*tarım, orman ve su kaynaklarında sürdürülebilir ve güvenli bir gelecek*” elde etmek vizyonu görülmektedir. Ayrıca stratejik planda amaçlar arasında erişilebilir, yeterli ve sürdürülebilir tarımsal ürün arzını, su ve toprak kaynaklarıyla biyolojik çeşitliliğin sürdürülebilir yönetimini sağlamak da bulunmaktadır. Bu kapsamda genetik kaynakların ve biyolojik çeşitliliğin sürdürülebilirliğini, korunan alan sayısının çoğalmasını ve bunların etkin yönetimini, av ve yaban hayatı yönetiminde sürdürülebilirliği, toprağı koruyarak onun verimli kullanılmasını, su kaynaklarının kalite ve miktar bakımından korunmasını ve verimli kullanılmasını sağlamak hedeflenmektedir (TOB Stratejik Plan 2024-2028).

Bu çerçevede sürdürülebilir tarım konusundaki düzenleyici ve destekleyici idari faaliyetler şu başlıklar şeklinde örneklendirilebilir:

Sigortalama Faaliyetleri

Ortaya çıkabilecek çeşitli risklere karşı bazı tarım ürünleri ve tarımsal faaliyetlerin sigortalanması öngörülmüştür. Bu kapsamda tarım sigortasının kapsamının genişletilmesi hedeflenmektedir. Bu konuda Hazine ve Maliye Bakanlığı, TOB, Tarım Sigortaları Havuzu, Hazine Müsteşarlığı çeşitli çalışmalar yürütmektedir. Her yıl çıkarılan Cumhurbaşkanlığı Kararı ile Devlet Destekli Tarım Sigortaları uygulamalarının kapsamının genişletilmesi ve yaygınlaştırılmasına yönelik çalışmalar devam etmektedir (TTKMEP, 2022: 58).

Gıda Güvenliğine İlişkin Faaliyetler

Gıda güvenliği, doğru üretim ve zararlılarla mücadele sürdürülebilir tarım konusunun diğer bir boyutu olarak ayrı bir önem taşımaktadır. Bu konuda TOB ile birlikte bazı kurum ve kuruluşların da düzenleme, denetleme faaliyetleri bulunmaktadır.

Bu konuda TOB'un kabul ettiği stratejik planın yanısıra, Ticaret Bakanlığının belirlemiş olduğu Yeşil Mutabakat Eylem Planı mevcuttur. Buna göre sürdürülebilir tarım hedefi doğrultusunda; Avrupa Birliği'nin (AB) anti-mikrobiyal ve pestisitlerin azaltılmasına dair hedefleriyle uyumlu olarak Türkiye'de anti-mikrobiyal ve pestisitlerin kullanımının azaltılmasına yönelik yönünde çalışmalar gerçekleştirilecektir. Pestisitlerin azaltılmasına dair çalışmalar kapsamında, biyoteknik ve biyolojik mücadele metotlarının kullanımının yaygınlaştırılması amaçlanmaktadır. Kimyasal gübre kullanımının azaltılmasına ilişkin AB'nin amaç ve politika değişiklikleri göz önünde tutularak çalışmalar yürütülecektir. Arazi toplulaştırma tescil faaliyetleri yürütülecektir. Organik tarım üretiminin geliştirilmesi kapsamında çalışmalar yapılacaktır. AB organik tarım mevzuatının uyumlaştırma çalışmalarının bitirilmesi ve devamında AB ile organik tarım alanında karşılıklı tanıma için Komisyon nezdinde girişimler yapılması hedeflenmektedir. Aydın, İzmir, Denizli ve Ağrı İllerinde kurmuş olan Tarıma Dayalı (jeotermal sera) İhtisas Organize Sanayi Bölgelerinde jeotermal kaynaklardan yararlanılarak bitkisel üretim yapılacaktır. Yenilenebilir enerji kullanan üretim tesisleri ve seralar desteklenecektir. Tarımsal üretimde artıkların yeniden değerlendirilmesi konusunda AR-GE çalışmaları yapılacaktır. tüketicinin bilinçlendirilmesi ve gıda artıklarının geri dönüşümünün sağlanmasına dair farkındalık yaratma çalışmaları gerçekleştirilecektir. Avrupa Komisyonunca açıklanan Tarladan Sofraya ve Biyoçeşitlilik Stratejilerine ilişkin bilgilendirme çalışmaları düzenlenecektir (TOB 2024-2028 Stratejik Plan; Yeşil Mutabakat Eylem Planı, 2021: 30).

Su Yönetimi Faaliyetleri

Sürdürülebilir tarım konusunda önem arz eden hususlardan bir diğeri su yönetimidir. Su yönetimini tarımdan ayrı düşünmek mümkün değildir. Bu nedenle tarıma ilişkin planlamalar yaparken aynı zamanda su yönetimi de birlikte düşünülmelidir. Bu kapsamda TOB'ın tarım ve su yönetimine ilişkin bazı tespitleri bulunmaktadır. Buna göre taşkın ve kuraklık yönetim planları tamamlanarak etkin tedbirler alınması sağlanmalıdır. Tarımsal kuraklıkla mücadele ve kuraklık yönetimi konusunda alınan 5140 sayılı Cumhurbaşkanı kararında kırsal kesimde, sulu ve kuru ziraat alanlarında yatırımlar, su yönetimi tarım ve sulama teknikleri, tohum ve bitki çeşitliliği, haşerat ve hastalıklarla mücadele, sosyal ve ekonomik destekler, arazi kullanım planları, mera otlatma planları ve acil eylemin uygulanmasına ilişkin her türlü tedbirleri kapsayan karar alınmıştır.²

Su kaynaklarının kalitesini artırıcı ve kirlenmesini önleyici önlemler uygulanmalıdır. Uluslararası gelişmeler de dikkate alınarak, havzalarda yönetim ve koruma planları hazırlanmalı ve uygulanmalıdır. Sularda tarımsal faaliyetlerden doğan kirliliği önlemeye yönelik uygulamalar artırılmalıdır. Teferruatlı toprak etütleriyle arazi kullanım planlamaları tamamlanmalı, üretimde stratejik ürünlere öncelik veren üretim planlamaları gerçekleştirilmelidir. Bilinçsiz ilaç ve gübre kullanımı başta olmak üzere tarım topraklarını kirlletici etmenlerin takibi yapılmalı ve kirliliğin önlenmesine yönelik faaliyetlerin geliştirilmesi sağlanmalıdır (TOB, Stratejik Plan 2024-2028: 38).

Türkiye'de acilen Su Kanunu'na ihtiyaç duyulmaktadır. Mevzuatımızda dağınık şekilde kanuni düzenlemeler bulunmakla birlikte, bunlar su stresi altında bulunan ülkemizin gelecekteki ihtiyaçlarına cevap verebilecek nitelik taşımamaktadır. Genel su, özel su kavramlarında olduğu gibi suyun hukuki durumunda karmaşalar bulunmaktadır. Mineralli suların tanımı ve tahsisinde karmaşa olması, kullanılmış suların tekrar kullanımı ve su tahsisine ilişkin ortaya çıkan bazı sorunların bulunması, mükerrer uygulamalardan kaynaklanan problemlerin, bir diğer deyişle kurumlararası çakışmanın giderilmesi ihtiyacı, AB müktesebatına uyum süreci gibi bazı etmenler Su Kanunu'na olan ihtiyacı göstermektedir (Doğu Kırkıcı, 2019:1-22). Bu nedenle sürdürülebilir tarım konusunda aynı zamanda yasama organının da sorumluluğu bulunmaktadır. Hatta idari organlar yetkilerini kanundan aldığı için bu alanda temel çıkış noktası ilgili kanuni düzenlemelerdir. Bundan dolayı kanun koyucunun sürdürülebilir tarım konusundaki sorumluluğu, idari organların sorumluluğu kadar önem taşımaktadır.

Türkiye'de Su Kanunu Taslağı, 2021 yılında düzenlenen I. Su Şurası kapsamında oluşturulan geniş katılımlı Su Hukuku Çalışma Grubu çalışmaları kapsamında ele alınmış ve ardından TOB'un merkez birimleri ile bağlı ve ilgili birimlerin görüşlerine sunulmuştur. Gelen görüşlerin değerlendirilmesi ve uygun görülenlerin taslağa alınmasının ardından nihai halini alan Su Kanunu Taslağının kanunlaşma süreci devam etmektedir (TTKMEP, 2022: 58). Bu noktada sürdürülebilir tarım konusunun en önemli ayaklarından birisini teşkil eden su sorununa bir yön verilecektir. Taslağın kanunlaşması ile su yönetimi konusunda devlet politikasının yön bulması kolaylaşacaktır ve daha hızlı mesafe kat edilecektir.

Kuraklıkla Mücadele Faaliyetleri

Kuraklıkla mücadele konusunda 2022 yılında alınan 5140 sayılı Cumhurbaşkanı Kararı ile ilgili tüm ziraat alanlarında su yönetimi, tarım teknikleri, hastalık ve zararlılarla mücadele, yatırımlar, sulama teknikleri, bitki ve tohum çeşitliliği, mera otlatma planları, sosyal ve ekonomik destekler, kısıtlamalar ve acil eylemin uygulanması konusunda, idari organlar tarafından her türlü tedbir alınabileceğine ilişkin karar alınmıştır. Tarımsal Kuraklık Yönetimi biriminin kurularak, TOB'un buna ilişkin yönetmelik çıkaracağı belirtilmiştir. TOB ilerleyen dönemde konuya ilişkin

² RG, Sayı: 31738, 02.02.2022.

yönetmelik çıkarmıştır. Bu noktada tarım üretiminin sürdürülebilir tarım kapsamında yapılması için gerekli denetimler yerinde ve zamanında yapılmalıdır.

Su ve toprak kaynaklarının, sürdürülebilir ve ekonomik tarımsal kullanımına yönelik araştırma-geliştirme çalışmalarına hız verilmelidir. 2023-2027 stratejik kalkınma hedefleri doğrultusunda TOB, Üniversiteler ve Uluslararası Araştırma Kuruluşları gibi kuruluşlar bu konuda çalışmalar yürütmektedir. Doğal kaynakların sürdürülebilirliği açısından; toprak ve su kaynaklarımızın mevcut durumuna yönelik tarımsal kuraklıkla mücadele izleme araç ve yöntemlerinin geliştirilmesi, toprak kaynaklarının korunması amacıyla risk alanlarında iklim dostu tarım uygulamalarının yaygınlaştırılması, su kaynaklarının geliştirilmesi, korunması ve sürdürülebilir kullanımı çerçevesinde havza bazında su tasarrufu sağlanması hedeflenmektedir (TTKMEP, 2022: 53).

Yangınlarla Mücadele Faaliyetleri

Sürdürülebilir tarım konusunda etkili olan faktörlerden bir diğeri ekolojik dengeye zarar veren yangınlarla mücadeledir. Bu konuda orman yangınlarının önlenmesi ve orman yangınlarıyla mücadele konusunda çeşitli faaliyetler yapılmaktadır. Ayrıca orman dışı yangınlarla mücadelede gerekli kurumsal kapasitenin geliştirilmesi yönünde çalışmalar yapılmaktadır. Bu konuda TOB ile birlikte Orman Genel Müdürlüğü ve yerel yönetimler, kurumsal koordinasyonun sağlanmasına, orman dışı yangınlara müdahalede gerçekleşen sorunların giderilmesine ve kurumsal kapasitenin geliştirilmesine yönelik çalışmaların artırılmasını hedeflemektedir (TTKMEP, 2022: 58).

SONUÇ

Gerek devletin gerekse diğer bazı kamu tüzel kişilerinin sürdürülebilir tarımsal faaliyetler konusunda çok çeşitli görev ve yetkileri bulunmaktadır. İşin özelliği gereği sürdürülebilir tarım konusunda çok yönlü bir çalışma sistemi gereklidir. Sürdürülebilir tarım, tek yönlü bir yaklaşımı benimsemeyip su yönetimi, gıda güvenliği, çevrenin ve toprağın korunması gibi pek çok faktörle birlikte değerlendirilmesi gereken bir konudur. Zira hedeflenen başarıya tek bir yoldan ulaşmak mümkün olmayıp, bunu etkileyen pek çok faktör bulunmaktadır.

Bu nedenle sürdürülebilir tarım konusunda birden çok idari organın devreye girdiği görülmektedir. Tüm bu idari organlar, kıt olan kaynakların daha iyi ve etkin bir şekilde kullanımının sağlanarak, yarınlara daha iyi, yaşanabilir bir gelecek bırakma hedefine yönelmiştir. Bu alanda faaliyette olan idarelerin koordineli, eş zamanlı ve uyumlu bir politika gütmesi gerekmektedir. Bu konuda anayasal ve kanuni yetkiler çerçevesinde idari organların düzenleyici ve bireysel işlemler yapma yetkisi bulunmaktadır. İdare bu konudaki planlama, düzenleme ve destekleme yetkisini, bilimsel verilere ve öngörülere göre kullanmak durumundadır. Ancak bu sayede hedefe ulaşmak kolaylaşacak ve mümkün olacaktır.

Sürdürülebilir tarım konusunda en büyük sorumluluğun devlete düştüğünü ifade etmek yanlış olmayacaktır. Çünkü ülkenin kaynaklarının korunarak, halka uygun yaşam alanlarının sağlanması, devletin temel yükümlülükleri arasında bulunmaktadır. Burada kurumsal çeşitliliğin yanı sıra mevzuatta da çeşitlilik göze çarpmaktadır. Bu alandaki mevzuat eksikliğinin giderilmesi, etkin tedbirlerin alınması ayrıca önem taşımaktadır. Sürdürülebilir tarımı sağlamak hususunda gerekli kanuni düzenlemelerin hayata geçirilmesi, durumun gereğine göre mevzuatın güncellenmesi konusunda temel görev esasında yasama organınınındır. Bu nedenle sadece idari organların değil, yasama organının da bu konuda sorumluluğu bulunmaktadır.

KAYNAKÇA

- Akbaş, B. (2019). Sürdürülebilir Tarımda Entegre Mücadele Çalışmalarının Ülkemiz Açısından Değerlendirilmesi. *Yalvaç Akademi Dergisi*, 4 (1), 32-40.
- Baran, M. F. & Bellitürk, K. & Çelik, A. (2021). Türkiye’de Sürdürülebilir Tarım Uygulamaları: Zorluklar ve Potansiyeller, Ankara.
- Doğu Kırkıcı, D. Tarım ve Orman Bakanlığı Su Yönetimi Genel Müdürlüğü Su Kanunu Taslağı, <https://www.tarimorman.gov.tr/SYGM/Belgeler/Havza%20H%C4%B0E-Sunumlar/Su%20Kanunu%20Tasla%C4%9F%C4%B1.pdf>, 05.02.2025.
- Güner, U. (2020). Çevresel Sürdürülebilirlik, <https://books.googleusercontent.com/books/content?>, 05.02.2025.
- Karadavut, U. & Aksoyak, Ş. & Palta, Ç. (2004). Kırsal Çevre Sorunları ve Sürdürülebilir Tarım. *Bitkisel Araştırma Dergisi*, 1: 61-66.
- Kubaş, A., & Aydın, N. (2022). Sürdürülebilir tarım algısı. *Balkan and Near Eastern Journal of Social Sciences*, 8 (04), 56-62.
- TOB 2024-2028 Stratejik Plan, <https://www.tarimorman.gov.tr/SGB/Belgeler/stratejikplan.pdf>, 05.02.2025
- Turhan, Ş. (2005). Tarımda sürdürülebilirlik ve organik tarım. *Tarım Ekonomisi Dergisi*, 11(1 ve 2), 13-24.
- Türkiye Tarımsal Kuraklıkla Mücadele Stratejisi ve Eylem Planı 2023-2027, Ankara, 2022.
- Yeşil Mutabakat Eylem Planı, 2021, <https://ticaret.gov.tr/data/60f1200013b876eb28421b23/MUTABAKAT%20YE%C5%9E%C4%B0L.pdf>, 05.02.2025.
- <https://turkiye.un.org/tr/sdgs/2>, 05.02.2025.

USE OF FLAXSEED (*Linum usitatissimum* L.) WITH HIGH ADDED VALUE**KETEN (*Linum usitatissimum* L.) TOHUMUNUN KATMA DEĞERİ YÜKSEK ÜRÜN OLARAK KULLANILMASI****Doç. Dr. Mehmet Zeki KOÇAK**

İğdır Üniversitesi, Ziraat Fakültesi, Tarla Bitkileri Bölümü, İğdır, Türkiye

ORCID ID: 0000-0002-8368-2478

Prof. Dr. Bünyamin YILDIRIM

İğdır Üniversitesi, Ziraat Fakültesi, Tarla Bitkileri Bölümü, İğdır, Türkiye

ORCID ID: 0000-0003-2463-6989

ÖZET

Keten (*Linum usitatissimum* L.), yağ ve lif özellikleri nedeniyle önemli endüstriyel ürünlerden biridir ve çok sayıda kullanım alanına sahiptir. Ayrıca, tüketiciler ve üreticiler için son yıllarda keten, sadece temel beslenme aracı olarak değil, aynı zamanda sağlıktaki önemli kullanım ve endüstrinin birçok alanları bulunmaktadır. Keten temelli ürünlerin karakterizasyonu ve tanıtımı/adaptasyonu büyük ilgi görmektedir. Keten bitkisi, yapısında %35-65 oranında yağ içermektedir. Özellikle yağlık keten çeşitleri gıda sektöründe geniş bir kullanımı olduğu kadar aynı zamanda keten yağının hızlı kuruma özellikleri nedeniyle de boya, vernik, sabun, macun ve polimerlerin hazırlanması dâhil olmak üzere çok büyük endüstriyel uygulamalara sahiptir. Yağlık olarak kullanılmasının yanı sıra; keten bitkisinin saplarından elde edilen liflerin; doğal, yumuşak, emici, dayanıklılık özellikleriyle tekstil sanayisinde, otomobil parçalarının, gemi halatı, yelken, çadır bezi, hortum yapımı, gaz maskesi, muşamba üretimi, kâğıt, paket ve ısı yalıtım malzemesi yapımında geniş kullanım alanına sahiptir. Keten bitkisi tohumu zengin omega-3, omega-6 ve omega-9 yağ asitleri (α -linolenik asit, linoleik asit, oleik asit), lignan, secoisolariciresinol diglukosid (SDG). Bu bileşiklerin, anti-kanser etkileri (meme, kolon ve cilt vb.), anti-inflamatuvar etkileri, anti-oksidatif kapasiteleri, kardiyovasküler hastalıklar, anti-diabetik, anti-bakterial, anti-viral, anti-fungal, ruhsal-psikolojik hastalıklar olması ve biyoaktivite özellikte olmalarından kaynaklı insan sağlığı açısından tıbbi olarak birçok kullanım alanı mevcuttur.

Anahtar kelimeler: *Linum usitatissimum*, Keten tohumu, Yağ asitleri, Keten lifi**ABSTRACT**

Flax (*Linum usitatissimum* L.) is one of the important industrial crops due to its oil and fiber properties, and it has a wide range of applications. In recent years, flax has gained significant attention not only as a basic nutritional resource but also for its important uses in health and in various industrial fields for both consumers and producers. The characterization and promotion/adaptation of flax-based products have attracted great interest. The flax plant contains 35-65% oil. Particularly, oilseed flax varieties are widely used in the food industry and have extensive industrial applications due to the rapid drying properties of flax oil, including the preparation of paints, varnishes, soaps, putties, and polymers. In addition to its use as an oilseed,

the fibers obtained from the stalks of the flax plant have wide applications in industries such as textiles, thanks to their natural, soft, absorbent, and durable properties. These fibers are used in the production of automobile parts, ship ropes, sails, tent fabrics, hoses, gas masks, oilcloth, paper, packaging, and thermal insulation materials. Flaxseeds are rich in omega-3, omega-6, and omega-9 fatty acids (α -linolenic acid, linoleic acid, oleic acid), lignan, and secoisolariciresinol diglucoside (SDG). These compounds are known for their anti-cancer effects (breast, colon, skin, etc.), anti-inflammatory properties, antioxidant capacities, and their benefits in cardiovascular diseases, anti-diabetic, antibacterial, antiviral, antifungal, and psychological-mental health disorders. They also have bioactive properties, making flax a plant with numerous medicinal applications for human health.

Key words: *Linum usitatissimum*, Flaxseed, fatty acid, Flax fiber

GİRİŞ

Keten (*Linum usitatissimum* L.), tarih boyunca tarımı yapılan ve çok yönlü kullanımıyla öne çıkan önemli bir bitkidir. Genel olarak lif üretimi için yetiştirilen keten bitkisi, son yıllarda keten tohumunun sağlık, gıda, kozmetik ve endüstriyel alanlarda katma değeri yüksek bileşenler içermesinden dolayı zamanla daha fazla ilgi görmektedir. Keten tohumunun içerdiği omega-3 ve onega-6 gibi yağ asitleri, lignanlar, protein ve diyet lifi sayesinde insan sağlığına sunduğu faydalarla fonksiyonel bir gıda olarak kullanımı öne çıkmaktadır. Aynı zamanda farmasötik ve kozmetik endüstrisinde antioksidan, anti-inflamatuvar ve cilt yenileyici özellikleri ile değerlendirilmektedir (Koçak, 2023; Jahan ve ark. 2024; Shahada ve ark., 2024).

Bunun yanı sıra keten tohumu, biyoplastik üretimi, hayvan yemi katkısı ve biyoyakıt gibi çevre dostu ve sürdürülebilir alternatif ürünlerin geliştirilmesinde de önemli bir ham madde kaynağı olarak önemli bir tohumdur. Birçok araştırmada, keten tohumunun işlenerek fonksiyonel bileşenlerinin izole edilmesi ve çeşitli sektörlerde daha etkin kullanılması üzerine yoğunlaşmaktadır. Bu bağlamda, keten tohumunun katma değeri yüksek ürünlere dönüştürülmesi, hem ekonomik hem de çevresel açıdan büyük bir potansiyel taşımaktadır (Pokharel, 2023; Rani ve ark., 2023; Chand ve ark., 2024; Jahan ve ark. 2024).

Bunlara ek olarak, keten tohumu farmakolojik etkileriyle de dikkat çekmektedir. Kardiyovasküler hastalıklara karşı koruyucu etkileri, kan şekeri seviyelerini düzenleyici özellikleri ve bağırsak sağlığı üzerindeki olumlu etkileri birçok bilimsel çalışma ile desteklenmektedir. Ayrıca keten tohumunun anti-kanserojen özellikleri, içerdiği yüksek orandaki lignanlar sayesinde araştırmalara konu olmaktadır. Bağışıklık sistemini destekleyici bileşenler içermesi, keten tohumunun fonksiyonel gıda ve takviye edici ürünlerde kullanımını artırmaktadır (Goyal ve ark., 2014; Shahada ve ark., 2024). Endüstriyel anlamda, keten tohumunun ekstraksiyon yöntemiyle elde edilen keten yağı, boya, vernik ve tekstil endüstrisinde de geniş bir kullanım alanına sahiptir. Yüksek oranda α -linolenik asit içermesi nedeniyle besinsel açıdan da önemli bir kaynak olup, bitkisel yağ endüstrisinde özel bir yere sahiptir. Ayrıca, tohumdan elde edilen protein, hayvancılık sektöründe yem katkısı olarak kullanıldığı gibi, vegan ve vejetaryen beslenme düzenlerinde de protein kaynağı olarak değerlendirilmektedir (Gutte ve ark., 2015; Soni ve ark., 2016; Koçak ve ark., 2023).

Genel olarak bu çalışmada, keten tohumunun farklı endüstri alanlardaki kullanım potansiyeli ele alınarak, gıda, sağlık ve endüstriyel sektörlerde katma değerli bir ürün olarak değerlendirilmesi üzerinde kayda değer bilgiler verilmiştir. Ayrıca, keten tohumundan elde edilen bileşenlerin işleme teknikleri ve bu bileşenlerin verdiği avantajlar belirtilmiştir.

Keten Tohumunda Bileşenler;

Keten tohumunda, zengin bir kimyasal içeriğe sahiptir (Rubilar ve ark., 2010);

- **Yağ Asitleri:** Keten tohumu, %35-55 oranında yağ içerir. Bu yağın büyük kısmını omega-3 (α -linolenik asit), omega-6 (linoleik asit) ve omega-9 (oleik asit) yağ asitleri oluşturur.
- **Lignanlar:** Keten tohumunda bulunan lignanlar, özellikle secoisolariciresinol diglukozit (SDG), antioksidan ve östrojen benzeri etkiler gösterir.
- **Protein:** Keten tohumu, yaklaşık %20-25 oranında yüksek kaliteli bitkisel protein içerir.
- **Diyet Lifi:** Çözünür ve çözünmez lifler açısından zengin olan keten tohumu, sindirim sağlığına katkıda bulunur.
- **Vitamin ve Mineral:** Keten tohumu, E vitamini, B vitaminleri, magnezyum, potasyum ve çinko gibi önemli mikro besin öğeleri içerir.

Sağlık Alanındaki Kullanım;

Keten tohumu, içerdiği biyoaktif bileşikler sayesinde önemli oranda sağlık alanında kullanılmaktadır.

Kardiyovasküler etkisi;

Keten tohumunun kardiyovasküler sağlığı destekleyen birçok faydalı bileşiğe sahip olmasıyla, içerdiği yüksek miktarda omega-3 yağ asidi, özellikle α -linolenik asit (ALA), kalp-damar hastalıklarına karşı koruyucu etkiler göstermektedir. Ayrıca, mevcut yağ asidi, kötü kolesterol (LDL) seviyelerini düşürürken iyi kolesterol (HDL) seviyelerini artırdığı ve kan basıncını düzenleyerek hipertansiyon riskini azalttığı bilinmektedir (Parikh ve ark., 2018; Al-Madhy ve ark., 2023).

Keten tohumundaki lignanlar ve antioksidanlar, damar sertliği oluşumunu engellediği ve damar duvarlarının elastikiyetini artırabilir. Anti-inflamatuar özellikleri sayesinde, kalp-damar sisteminde inflamasyona bağlı hasarları önleyerek kalp krizi ve inme riskini düşürür. Bunun yanı sıra, kanın pıhtılaşma eğilimini azaltarak tromboz gibi ciddi sağlık sorunlarına karşı koruyucu bir rol oynamaktadır. Aynı zamanda, yapılan klinik çalışmalarda, yapılan düzenli keten tohumu tüketiminin trigliserit seviyelerini düşürdüğü ve kalp ritim bozukluklarını önlediği görülmüştür. (Al-Madhy ve ark., 2023; Noreen ve ark., 2023).

Anti-kanser etkisi;

Keten tohumu, özellikle lignanlar ve omega-3 yağ asitleri gibi biyoaktif bileşenler sayesinde anti-kanser özellikler göstermektedir. Lignanlar, fitoöstrojenik etkileriyle hormon ilişkili kanser türlerine, özellikle meme ve prostat kanserine karşı koruma sağlayabilir. Bu bileşikler, östrojen metabolizmasını düzenleyerek kanser gelişimini engelleyebildiği bilinmektedir. Ayrıca, mevcut lignanlar, antioksidan özellikleriyle serbest radikallerin neden olduğu hücre hasarı önler ve DNA koruması sağlayarak tümör oluşumunu engelleyebilir. Omega-3 yağ asitleri ise inflamasyon azaltıcı etkileriyle kanser hücrelerinin büyümesini yavaşlatabilir ve metastaz riskini düşürebilir. Bunların yanı sıra, yapılan klinik çalışmalarda, keten tohumu tüketiminin kolon, cilt ve akciğer kanseri gibi farklı kanser türlerinin önlenmesinde de etkili olabileceği gösterilmiştir (Mueed ve ark., 2023; Stepien ve ark., 2025).

Anti-diabetik etkisi;

Mevcut tohumun içerdiği yüksek diyet lifi, kan şekerinin düzenlenmesine katkı sağlamakta olup çözünür lifler, bağırsakta viskoz bir yapı oluşturarak glikoz emilimini yavaşlatır ve yemek sonrası kan şekerindeki ani yükselmeleri engellemektedir. Ayrıca, omega-3 yağ asitleri, inflamasyonu azaltarak insülin direncini düşürmeye yardımcı olmaktadır. Klinik çalışmalarda, keten tohumu tüketiminin hem tip 1 hem de tip 2 diyabet hastalarında glisemik kontrolü iyileştirdiği gösterilmiştir. Bu etkiler, insülin duyarlılığını artırarak ve glikoz metabolizmasını düzenleyerek gerçekleşmektedir (Alawlaqi ve ark., 2023).

Anti-inflamatuar ve Antioksidan etkisi;

Tohumun içerdiği omega-3 yağ asitleri (α -linolenik asit), lignanlar ve diğer antioksidan bileşenler sayesinde güçlü anti-inflamatuar ve antioksidan özelliklere sahiptir. Bu özellikler, vücutta inflamasyonun azaltılmasına ve serbest radikallerin neden olduğu oksidatif stresin engellenmesine katkı sağlamaktadır. Anti-inflamatuar etkiler, özellikle kronik hastalıkların önlenmesinde ve yönetiminde önemlidir. Keten tohumu, bağışıklık sistemini dengeleyerek romatoid artrit, astım, inflamatuvar bağırsak hastalıkları (IBD) ve diğer inflamasyon kaynaklı rahatsızlıkların semptomlarını hafifletebilmektedir. Omega-3 yağ asitleri, inflamatuvar sinyalleri baskılayarak hücresel düzeyde koruma sağlar (Coklar ve Akbulut, 2019).

Antioksidan etkiler ise hücrelerin DNA, protein ve lipid yapılarında meydana gelebilecek hasarları önler. Lignanlar, fenolik bileşikler ve tokoferoller, serbest radikallerle savaşarak yaşlanma belirtilerini geciktirir ve kanser gibi dejeneratif hastalıkların riskini azaltır. Yapılan çalışmalar, keten tohumu tüketiminin hücre yenilenmesini desteklediğini ve oksidatif stresin azalmasına yardımcı olduğunu göstermektedir. Bu özellikler, keten tohumunun hem koruyucu hem de tedavi edici bir fonksiyonel gıda olarak değerini artırmaktadır (Touré ve Xueming, 2010).

Gıda Sektöründeki Kullanımı;

Keten tohumunun, zengin besin içeriği ve fonksiyonel özellikleri hem bir gıda ürünü hem de gıda katkı maddesi olarak geniş bir kullanım alanına sahip olması ve tercih edilen bir bileşen olmasını sağlamaktadır. Birçok fonksiyonel gıdaların (bisküvi, ekmek, mısır gevreği, granola, enerji barları) üretiminde yaygın olarak kullanılmaktadır (Shahada ve ark., 2024).

Keten Tohumu Yağı

Keten tohumu yağı, soğuk sıkım yöntemiyle elde edilerek gıda sektöründe sıklıkla kullanılmaktadır. Yüksek omega-3 içeriği sayesinde sağlıklı yemeklerde, salatalarda ve soslarda tercih edilir. Özellikle omega-3, lignan ve diyet lifi içeriği nedeniyle keten tohumu, besin takviyelerinin üretiminde kullanılırlar. Ayrıca, bebek mamaları ve özel diyet ürünlerinde de kullanılmaktadır. Doğal yağ asitleri ve protein içeriği, hem büyüme çağındaki çocuklar hem de özel diyet gereksinimi olan bireyler için değerli bir katkıdır (Yawale, 2022; Koçak ve ark., 2023; Shahada ve ark., 2024).

Endüstriyel Kullanımı

Keten tohumu ve yağı, aynı zamanda farklı birçok sektörde ve çeşitli uygulamalara kullanılmaktadır; Keten yağının hızlı kuruma özelliği ve dayanıklılığı sayesinde boya ve vernik endüstrisinde yaygın olarak kullanılmaktadır. Özellikle ahşap yüzeyler için üretilen doğal verniklerde, keten yağı hem parlaklık hem de koruyuculuk sağlar (Yawale, 2022; Koçak ve ark., 2023). Çevre dostu materyallere olan talebin artmasıyla birlikte, keten yağı biyoplastik üretiminde önemli bir hammadde olarak değerlendirilmektedir. Keten yağı bazlı polimerler, geri dönüştürülebilir ve biyolojik olarak parçalanabilir özelliklere sahip olup, ambalaj sektöründe kullanılmaktadır (Gutiérrez ve ark., 2010; Dzuovor ve ark., 2018).

Keten tohumu yağı, nemlendirici ve antioksidan özellikleri nedeniyle kozmetik sektöründe tercih edilmektedir. Cilt bariyerini güçlendiren, kuruluğu önleyen ve yaşlanma karşıtı etkiler sunan bu yağ, krem, losyon, şampuan ve sabun gibi ürünlerin formülasyonunda kullanılmaktadır. Ayrıca, keten yağındaki E vitamini ve diğer antioksidanlar, cildi çevresel hasarlara karşı koruyarak ürünlere katma değer sağlamaktadır (Kolodziejczyk ve ark., 2012; Sarma ve Begum, 2024). Genel olarak lif üretimi için yetiştirilen keten bitkisinin liflerinden üretilen tekstil ürünleri, dayanıklılığı ve çevre dostu özellikleri nedeniyle popülerdir. Özellikle doğal kumaş üretiminde keten lifi, sürdürülebilir moda sektöründe uyum sağlamaktadır. Keten tohumu yağı ise tekstil yüzeylerinde su itici kaplamalar için kullanılmaktadır. Ayrıca, keten tohumu ve yağı, hayvan yemlerinde değerli bir katkı maddesi olarak yer almaktadır. Özellikle hayvanların et, süt ve yumurta kalitesini

artırmak için yemlere keten tohumu eklenmektedir. Bu uygulama, ürünlerin omega-3 yağ asitleri açısından zenginleşmesini sağlamaktadır.

Sonuç

Keten (*Linum usitatissimum* L.), gerek içerdiği besin bileşenleri gerekse endüstriyel kullanım potansiyeli ile çok yönlü bir bitkidir. Keten tohumu, omega-3 ve omega-6 gibi yağ asitleri, lignanlar, protein ve diyet lifi açısından zengin içeriğe sahip olup, kardiyovasküler hastalıklar, diyabet, kanser ve enflamatuar hastalıklar gibi çeşitli sağlık sorunlarına karşı koruyucu ve tedavi edici özellikleri olan bir bitkidir. Ayrıca, gıda sektöründe, fonksiyonel gıdalar ve besin takviyeleri için önemli bir bileşen olan keten tohumu, ekmek, bisküvi ve diğer unlu mamuller gibi çeşitli ürünlerde yaygın olarak kullanılmaktadır. Endüstriyel alanda keten tohumu yağı, boya, vernik, tekstil ve kozmetik gibi çok sayıda sektörde kıymetli bir ham madde olarak değerlendirilmektedir. Aynı zamanda, keten bitkisi lifleri dayanıklılığı, doğal yapısı ve çevre dostu özellikleri sayesinde tekstil, ambalaj, gemi halatı ve inşaat gibi alanlarda geniş bir kullanım alanına sahiptir. Bunlara ek olarak, keten tohumunun hayvan yemlerine katılması da besin değerini zenginleştirerek hayvansal ürünlerin kalitesini artırmaktadır. Genel olarak, keten tohumu ve liflerinin çok yönlü kullanım potansiyeli hem ekonomik hem de ekolojik açıdan önemli avantajlar sunmaktadır. Sürdürülebilir tarım uygulamaları ve geliştirilen yeni teknolojiler sayesinde keten bitkisinin önemi daha da artacak ve farklı sektörlerdeki katma değeri yüksek uygulamaları yaygınlaşacaktır.

KAYNAKLAR

- Alawlaqi, M. M., Al-Rajhi, A. M., Abdelghany, T. M., Ganash, M., & Moawad, H. (2023). Evaluation of biomedical applications for linseed extract: antimicrobial, antioxidant, anti-diabetic, and anti-inflammatory activities in vitro. *Journal of Functional Biomaterials*, 14(6), 300.
- Al-Madhagy, S., Ashmawy, N. S., Mamdouh, A., Eldahshan, O. A., & Farag, M. A. (2023). A comprehensive review of the health benefits of flaxseed oil in relation to its chemical composition and comparison with other omega-3-rich oils. *European journal of medical research*, 28(1), 240.
- Chand, M., Chopra, R., Talwar, B., Homroy, S., Singh, P. K., Dhiman, A., & Payyuni, A. W. (2024). Unveiling the potential of linseed mucilage, its health benefits, and applications in food packaging. *Frontiers in Nutrition*, 11, 1334247.
- Coklar, H., & Akbulut, M. (2019). Bioactive compounds, antioxidant activity and some physicochemical properties of the seed and seed-oil of *Mahonia aquifolium* berries. *Journal of Food Measurement and Characterization*, 13, 1269-1278.
- Dzuvor, C. K. O., Taylor, J. T., Acquah, C., Pan, S., & Agyei, D. (2018). Bioprocessing of functional ingredients from flaxseed. *Molecules*, 23(10), 2444.
- Goyal, A., Sharma, V., Upadhyay, N., Gill, S., & Sihag, M. (2014). Flax and flaxseed oil: an ancient medicine & modern functional food. *Journal of food science and technology*, 51, 1633-1653.
- Gutiérrez, C., Rubilar, M., Jara, C., Verdugo, M., Sineiro, J., & Shene, C. (2010). Flaxseed and flaxseed cake as a source of compounds for food industry. *Journal of soil science and plant nutrition*, 10(4), 454-463.
- Gutte, K. B., Sahoo, A. K., & Ranveer, R. C. (2015). Bioactive components of flaxseed and its health benefits. *International Journal of Pharmaceutical Sciences Review and Research*, 31(1), 42-51.
- Jahan, R., Malik, S., Shoaib, S., Alomary, M. N., Ansari, M. A., Islam, N., & Khan, S. (2024). Medicinal and Nutritional Importance of *Linum usitatissimum* L. in Human Health. In *Medicinal*

Plants and their Bioactive Compounds in Human Health: Volume 1 (pp. 25-45). Singapore: Springer Nature Singapore.

Koçak, M. Z., Kumlay, A. M., & Alma, M. H. (2023). Morphological and molecular characterization of flax (*Linum usitatissimum* L.) accessions obtained from different locations in Turkey. *Genetic Resources and Crop Evolution*, 70(8), 2235-2261.

Koçak, M. Z. (2023). How growing conditions are influential on the agronomic attributes and fiber-related quality parameters of flax (*Linum usitatissimum* L.) fibers: A seismomorphogenesis approach. *International Journal of Agriculture Environment and Food Sciences*, 8(1), 220-234.

Kolodziejczyk, P., Ozimek, L., & Kozłowska, J. (2012). The application of flax and hemp seeds in food, animal feed and cosmetics production. In *Handbook of natural fibres* (pp. 329-366). Woodhead Publishing.

Mueed, A., Deng, Z., Korma, S. A., Shibli, S., & Jahangir, M. (2023). Anticancer potential of flaxseed lignans, their metabolites and synthetic counterparts in relation with molecular targets: current challenges and future perspectives. *Food & Function*, 14(5), 2286-2303.

Noreen, S., Tufail, T., Ul Ain, H. B., & Awuchi, C. G. (2023). Pharmacological, nutraceutical, and nutritional properties of flaxseed (*Linum usitatissimum*): An insight into its functionality and disease mitigation. *Food Science & Nutrition*, 11(11), 6820-6829.

Parikh, M., Netticadan, T., & Pierce, G. N. (2018). Flaxseed: Its bioactive components and their cardiovascular benefits. *American Journal of Physiology-Heart and Circulatory Physiology*.

Pokharel, A. (2023). Development of Flax-and Hemp-Based Polylactic Acid Films for Bioplastic Applications (Doctoral dissertation, University of Saskatchewan).

Rani, R., Gosh, T., & Badwaik, L. S. (2023). Optimization of mustard, soybean and flaxseed meal blend formulation for development of biopolymeric film and its characterization. *Sustainable Chemistry and Pharmacy*, 33, 101147.

Rubilar, M., Gutiérrez, C., Verdugo, M., Shene, C., & Sineiro, J. (2010). Flaxseed as a source of functional ingredients. *Journal of soil science and plant nutrition*, 10(3), 373-377.

Touré, A., & Xueming, X. (2010). Flaxseed lignans: source, biosynthesis, metabolism, antioxidant activity, bio-active components, and health benefits. *Comprehensive reviews in food science and food safety*, 9(3), 261-269.

Sarma, S. D., Begum, T., & Lal, M. (2024). Application of linseed in cosmetic and personal care products development. In *Linseed* (pp. 193-203). Academic Press.

Shahada, C., Morya, S., & Awuchi, C. G. (2024). A narrative review on nutraceutical, food and industrial applications of flaxseed (*Linum usitatissimum*. L). *Cogent Food & Agriculture*, 10(1), 2306017.

Soni, R. P., Katoch, M., Kumar, A., & Verma, P. (2016). Flaxseed Composition and its health benefits. *Res. Environ. Life Sci*, 9, 310-316.

Stepień, A. E., Trojniak, J., & Tabarkiewicz, J. (2025). Anti-Oxidant and Anti-Cancer Properties of Flaxseed. *International Journal of Molecular Sciences*, 26(3), 1226.

Yawale, P. (2022). A comprehensive review on recent novel food and industrial applications of flaxseed: 2014 onwards. *Food and Feed Research*, 49(1), 67-95.

PORTABLE MINI-SCALE VERTICAL FARMING SYSTEM DESIGN: GROWING MICROSPROUT IN SOIL-LESS MEDIA

Alev ALDEMİR

Marmara University, Institute of Pure and Applied Sciences, Department of Electrical and Electronics Engineering, İstanbul, Turkey

ORCID ID: <https://orcid.org/0009-0006-6363-1565>

İsmail KIYAK

Marmara University, Faculty of Technology, Department of Electrical and Electronics Engineering, İstanbul, Turkey

ORCID ID: <https://orcid.org/0000-0002-5061-6378>

Faruk AYDIN

Marmara University, Faculty of Technology, Department of Electrical and Electronics Engineering, İstanbul, Turkey

ORCID ID: <https://orcid.org/0000-0002-6738-6775>

Ahmet KIRIK

Filizhane Biotechnology R&D Agricultural Applications Tic.Ltd. Şti, İstanbul, Turkey

ORCID ID: <https://orcid.org/0009-0002-7103-7268>

Abstract

Introduction and Objective: Commercial or individual efforts to grow microgreens/microflora indoors have gained momentum with the increasing awareness of access to healthy and safe food. With this awareness, microgreens/microflora seem to be the rising trend of recent years, but they have been at the center of life since the very beginning of life with the germination of each seed. In countries such as Turkey, where the rate of urban life is higher than rural life in terms of population density, various projects and studies are carried out to increase small-scale growing environments. The aim of this study is to design an environment that will enable people to access fresh greenery easily and economically with portable systems in places such as residential areas and shopping centers where there is no possibility of soil farming. **Materials and Methods:** In this study, a small-scale portable vertical agriculture production center was designed. Containers are used as growing media and the structure, which is divided into germination, production and control rooms, can be designed as grid-dependent/independent for energy and water needs. In order to fully meet the light spectrum needs of the plant for rooting, germination, sprouting and greening periods, LED- based artificial light sources covering 400-700 nm wavelengths were designed. In addition, a sensor-dependent control algorithm was used to support growth performance in germination and production environments. In the study, white radish was used as microgreens and no macro or micro elements were used in solid culture to understand the effect of the designed luminaire.

Results: White radish microgreens were observed from the sprouting stage to the harvest stage and plant height, Ph and Ec values were recorded. The effects of photosynthetic photon flux density (PPFD), wavelength and color temperature parameters of the artificial light used during harvesting on the plant growth process were comparatively analyzed. It was observed that the designed LED-based artificial light effectively provides the entire PAR region light spectrum required for plant growth stages. **Discussion and Conclusion:** At the end of the study, it was determined that the white radish microgreens grown reached harvest size in approximately 20% shorter time than the commonly used production elemental growing media products.

Keywords: Microgreens/microsprouts in solid culture media, microsprout cultivation armature design, sustainable agriculture, portable vertical farming system.

Introduction

The goal of vertical farming is to produce a large amount of crops in small areas in inverse proportion to the small size of the area. [1] These farms, called vertical farms, are also known as indoor farms. These farms are closed plant production organizations where environmental factors such as temperature, humidity, light, CO₂ concentration are observed and communication with external factors is minimized. [2]

The aim of the vertical farming system is to create a controlled agriculture system with plants grown in closed areas by keeping the plant growing conditions at optimum levels. It is possible to grow different types of plants such as vegetables, fruits or ornamental plants in vertical farming. The biggest reason why vertical farming is especially preferred is to grow plants in limited areas due to the lack of necessary land in urban areas. Likewise, it is preferred for growing plants in areas with unfavorable climatic conditions. In addition to these reasons, vertical farming will provide fresh, stable and reliable products all year round.[3]

One of the most important reasons for the growth and development of a plant is light. The time that any plant spends under the light is called the insolation period. Insolation periods for plants vary according to the growing season, year, and day of the plant. When calculating the optimal insolation time for a plant, the photosynthesis performed by the plant and the photoperiod illumination time are taken into account. [4]

UV and IR values are extremely important in the growth and development stages of a plant. At the same time, the spectrum values of the light source during the growth and development phase should be in the range of 400-700nm. [5]

In a study conducted in the Asian region, the effect of LED colors projected on the potato plant on the growth of the plant was examined. In this study, it was observed that the blue LED light spectrum as well as the combined red blue and green LED light spectrum had a positive effect on the potato plant. [6]

In a study for horticultural applications, a lighting module with 550/350mm dimensions was designed. The number of red and blue LEDs was determined as 50 and 40 respectively. LED selection is extremely important for the correct and effective operation of the luminaire and the selected LED was selected depending on the desired luminous flux, spectral distribution of the radiation, photosynthetic photon flux PPF (mol/s), operation control capabilities. SAMSUNG LED was selected within these required specifications. With the designed luminaire, it was concluded that an efficient luminaire was designed for plant cultivation. It was concluded that an effective, cheap and reliable system was established with this armature. [7]

In an experiment for vertical farming, a soybean plant was used. This study was divided into 2 parts. In the first part, light was placed for a period of 24 hours and in the second part, the effect of lighting on the different developmental stages of the plant was compared with the crop yield. The first part of the experiment resulted in a high number of flowering and production from the seeds. In the second experiment, as a result of partial illumination in different developmental stages, it was observed that illumination can also have an effect if the other conditions for the development of the plant are complete. As a result, it was reached as a result of the experiment that effective solutions can be produced in additional and correct lighting system in horticultural culture. [8]

Photosynthetically active radiation (PAR) distributions of LED light sources with different characteristics (A, B, C, D) designed by different lighting companies, which are widely preferred in greenhouse lighting systems, were evaluated. According to the measurement results of the study, the implicit PAR distributions of the LED light distribution in different distribution ranges (400-700nm

wavelength) were determined and different distribution uniformity evaluation criteria were examined in detail. The results are as follows; When analyzed on the daily lighting side, type A and B lighting scopes can be used as both full artificial and complementary photosynthesis lighting fixtures according to the designed layout, while type C and D lighting scopes gave results as complementary photosynthesis lighting fixtures when applied according to the designed layout. [9]

In an LED study on growing basil plants, hyper red and dark blue LED light combinations were applied and no difference was observed. In addition, different combinations of LED light were applied to the plant in question and were found to be beneficial for the germination and growth of the basil plant compared to daylight. Height increased by 30% and total wet weight increased by 50%. [10]

As a result of research and applications, it was concluded that with the application of one or two rows of intermediate lighting LEDs, mini cucumber yield increased by 22.3% and 30.8% compared to no intermediate interior lighting. [11] In another study conducted in this regard, it was observed that mini cucumber yield increased with intermediate lighting only in the early production stage and gradually decreased towards the late production period. [12]

The cucumber plant was subjected to a research process of diversion. As the amount of blue light increases, the amount of dry matter decreases. In addition, it was observed that as the amount of blue light increased, chlorophyll content per leaf area increased along with net photosynthetic rate and stomatal conductance. [13]

Microgreens, also known as micro sprouts, are young and tender plant varieties produced using seeds of various vegetable species, herbaceous plants, different aromatic plants and edible wild plants. Germination takes between 7-21 days depending on the selected seedling. Harvesting time is ready for harvesting when the first real leaves appear after the leaves emerging from the seed complete their development after germination is over. [14]

Microgreens are a new class of specialty crops produced from immature greens produced from the seeds of vegetables, herbs or grains, including wild species. [15]

Microgreens are good sources of nutrients and antioxidants, mainly minerals, carotenoids, and phenolic compounds. Due to their richness in vitamins and phytochemicals, microgreens have a strong antioxidant capacity and are effective in regulating plasma lipoprotein and cholesterol metabolism. These properties also indicate a potential value in the prevention and treatment of chronic diseases. Many studies have shown that microgreens have higher nutritional values than mature plants. [16]

As a result of a research, it was proved that the vitamin C and protein ratio increased in lettuce and spinach plants by adjusting the ratio of LED light quality well. [17] In another study, when the pepper plant was exposed to LED supplementation and fluorescent light under the same conditions in a dark environment in the process after grafting and comparing the two different types of illumination, it was observed that there was a higher survival rate and recovery efficiency with LED. [18]

The darkness level was set as the controllable factor to elucidate whether LED supplementation applied in plant cultivation affects the rate of necrotic base formation in the plant and to know the light intensity required. Inoculated cucumber seedlings were exposed to four different LED light intensities for three consecutive days. The light intensities were 25 $\mu\text{mol}/(\text{m}^2\text{s})$, 50 $\mu\text{mol}/(\text{m}^2\text{s})$, 75 $\mu\text{mol}/(\text{m}^2\text{s})$, and 100 $\mu\text{mol}/(\text{m}^2\text{s})$. As a result of this experiment, the white light intensity applied at an intensity of 50 $\mu\text{mol}/(\text{m}^2\text{s})$ was found to be the ideal light level for the formation of the necrotic layer of cucumber seedlings. [19]

The spinach plant was simultaneously illuminated with a warm white LED lamp, a cool white LED lamp and a garden LED lamp. A table showing the dry mass of the plants illuminated with each LED lamp compared to the dry mass of the plant illuminated with daylight was created. The observed results show that the dry mass of spinach illuminated with LED grow lamp is 15% higher than the dry mass illuminated with cold white lamp and 87.5% higher than the dry mass illuminated with warm white lamp.[20]

LED lighting was applied to the purple leaf lettuce plant, which has high nutritional value, before harvesting and the result of this application was observed in terms of the formation of nutritional quality and flavor of the plant. Red blue color LED light was applied in a ratio of 4 to 1 as a light source. The application method was carried out at different intensities and durations. As a result of this application, the optimum continuous lighting (CL) intensity and duration were found. It was an example for theoretical and technical guidance of artificial lighting (PFAL) for lettuce production. 2 different experiments were conducted. In these experiments, lighting was provided at different stages. According to the results of Experiment 1, it was observed that there was no change in lettuce sprout and fresh root weights as a result of CL treatment, but higher light intensity and CL duration significantly increased both sprout and root dry weights. In Experiment 2, ascorbic acid (ASA) content was not significantly different. However, it was concluded that the antioxidant levels in the leaves increased as the duration of CL increased. [21]

In this study, a portable vertical agriculture system was designed based on a container physical environment. In the designed environment, an LED artificial light source arrangement, air conditioning system, irrigation system and control system projects were made for the purpose of growing micro greens / micro sprouts. In the small-scale equivalent of the designed system, white radish microgreens/micro sprouts application was carried out under the same climate and LED source artificial light. The results were evaluated by comparative analysis of the application.

Portable Vertical Farming System Design Project

Within the scope of the study, it was found appropriate to use a 40' container for the portable scale vertical farming system project. The design system was realized with Dialux Evo program. The design stages are as follows:

- The total container area is divided into control room, production room and germination room.

- Vertical agricultural shelves, plant trays, irrigation system, air conditioning system and doors were added to these spaces.
- A water tank and solar panels are positioned on top of the container for places without mains connection.
- The colors and surface materials of the objects were determined.
- Finally, artificial light sources were added and the planning and calculations of the sections to grow plants with horticulture LEDs were completed.

Portable Vertical Farming

Within the scope of the project, a 40' container (length 12 m, width 2.35 m, height 2.4 m) was preferred as a portable production system space for microfilter/microgreens cultivation. The main reason for the

preference of the container is that it is more advantageous in terms of investment and installation cost compared to other alternatives as well as being an easily transportable structure

The container designed in Dialux evo program is shown in Figure 2.1.




Figure 2.1. Portable Vertical Farming System

The production room, sprouting room and control room of the vertical agricultural container designed in Dialux Evo program are shown in Figure 2.2, Figure 2.3, Figure 2.4.



Figure 2.2. Portable Vertical Farming System Production




Figure 2.3. Portable Vertical Farming System Sprouting Room



Figure 2.4. Portable Vertical Farming System Control Room

Portable micro sprout/micro greenery growing media control room is designed with dimensions of approximately 2,35X2X2,4 meters. Production room is designed approximately 2,35X7,5X2,4 meters. Germination room is designed approximately 2,35X2,5X2,4 meters.

The design of the shelves where the plants are placed in the vertical agriculture container designed in Dialux Evo program is shown in Figure 2.5.




Figure 2.5. Vertical Farming System production racks

The dimensions of the shelf selected for the cultivation of micro sprout plants were drawn in the dialux evo program and shown as follows width: 120cm, length: 60cm, height: 200cm.

The size of the trays used for growing micro sprouts will be 30x50cm.

There are 12 shelves in the portable micro sprout/micro greens growing environment and each shelf has 5 tiers. The distance between the shelves is 35 cm and the lowest floor of the shelf is 10 cm above the ground. There are 4 trays on each floor of the shelf. 10 of the 12 shelves are in the production room and 2 shelves are in the sprouting room. There is one inverter air conditioner each in the sprouting and production room. Air filter is used to balance the air circulation in the room. In addition, small fans were used and a thermometer was used to control the temperature inside the room.

The portable structure is divided into control, germination and production rooms, with the addition of a water tank and solar panels for locations where grid connection may be difficult.

The exterior view of the micro sprout growing container designed in Dialux evo program is shown in Figure 2.6




Figure 2.6 Microfilis Growing Container

Artificial Light Design

It is known that plants have different morphologies under different wavelengths of light and this is studied under the name of photomorphogenesis. For example, it is known that the formation of leaves and roots is supported more in plants fed with blue light, while fruit formation is triggered more in plants fed with red light [5,9].

The wavelength plot of the photon and energy weights of the PAR range of the designed light is shown in Figure 2.8




Figure 2.8 Wavelength dependence of the photon and energy weight of the PAR range[22-24]

PAR defined by PPF does not distinguish between different wavelengths between 400 and 700 nm and assumes that wavelengths outside this range have zero photosynthetic effect. If the exact spectrum of light is known, photosynthetic photon flux density (PPFD) values ($\mu\text{mol s}^{-1}\text{m}^{-2}$) can be varied by applying different weighting factors to different wavelengths. This results in a quantity called the yield photon flux (YPF).[22] The red curve in the graph shows that photons around 610 nm (orange-red) have the highest amount of photosynthesis per photon. However, since shorter wavelength photons

carry more energy per photon, the maximum amount of photosynthesis per event unit is at a longer wavelength, around 650 nm (dark red).[23,24]

It is seen that red and blue light are generally emphasized as effective wavelengths for growth in artificial light sources in plant growing environments. Along with these wavelengths, green light has also been shown to play an important role in the direction and efficiency of photosynthesis. It is understood that artificial light sources containing warm or cold white wavelengths are used to obtain green light.[20] The PAR region, which is the photosynthetic wavelength range for plants, is shown in Figure 2.9.




Figure 2.9 Yield photon flux curve [20]

The photon flux spectral distribution curve of the designed LED-based artificial light source is shown in Figure 2.10. SAMSUNG_4A4B0 Blue LED chips containing approximately 5% blue light at 400 nm wavelengths were preferred in the artificial light design of the microgreens/microphilis growing environment to support the root formation of the plant.

SAMSUNG vertical agricultural LED chips containing intense red light at effective wavelengths of 650-660 nm were preferred for the plant's stem growth, leaf number and leaf area. SAMSUNG_F2RNYZ WHITE LED chips with 572.8 nm effective wavelength at 6500 K⁰ color temperature and 572.8 nm effective wavelength containing green light wavelengths were preferred to help the plant use PAR region light with high energy for photosynthesis. Samsung brand LED chips produced for vertical agriculture were preferred to provide light as an artificial light source. The spectral distribution curve of the PAR region, the photosynthetic wavelength range for plants, is shown in Figure 2.10.




Figure 2.10 Photon flux spectral distribution curve of LED-based artificial light source




Figure 2.11 Vertical axis PAR value variation of LED based luminaire

Photosynthetically active radiation (PAR) variation values of the designed vertical farming LED-based artificial light source as a function of height are shown in Figure 2.11.

The cultivation process was carried out by illuminating the tray 25 cm above the tray where 4 microfilis were grown using 2 lights of 120cm. During this process, PPF measurements were made with a 10x15 grid and these measurement patterns are shown in Figure 2.12




Figure 2.12 Growing area and PPF measurement

The illumination of the skylight angle form of the LED-based light source in the dialux evo program is shown in Figure 2.13




Figure 2.13 Skylight Angle Form of LED Luminaire shown in Dialux program

The arrangement of the LED-based light source on a PCB is shown in Figure 2.14.




Figure 2.14: LED PCB Board

There are 72 LEDs on the PCB. Of these LEDs, 6 blue, 6 red and 60 white LEDs were used. 2835 Samsung brand LED chips produced for vertical agriculture were used. The visuals of the LED-based artificial light source are shown in Figure 2.15



Figure 2.15 Micro Sprout Plant Growth Luminaire

The technical specifications of the LED-based artificial light source are given in Table 1.

Table 1: Specifications of the Microphilisis Magnification Armature Used

Input Voltage/Current	230 V, 0.12 A
Input Power	28 WATT
LED LIGHT ANGLE	120°
LED Chip	SAMSUNG White-Blue-Red LED
Light source PAR region	400nm-700nm
Light source effective wavelength	660 nm
Photosynthetic Photon Flux (PPF)	67.5 $\mu\text{mol/s}$
Heat generated	95.54 BTU
Length	1200 mm
Weight	0.5 kg

A driver model known as DOB LED (Driver On Board), which is expressed as on the driver by adding it to the modules (driver), was used.



Figure 2.16 Micro Sprout Plant Growth Armature Driver

In this luminaire, there is no need for an external driver. With this driver mounted under the PCB, a useful luminaire design has emerged in order to create a ready finished product design. This constant current driver is 27.6 Volt input 220-230V output voltage value for the driver driven with 120 mA. The visualization of the driver with the specified features is shown in Figure 2.16.




Figure 2.17 DC /DC Driver circuit diagram with active suppression

The DOB drive first converts the AC input energy into DC unregulated signal by AC/DC converter. In order to convert the unregulated DC signal into a constant DC voltage signal in a controlled manner, an isolated type forward DC-DC converter is used. In the active suppression forward converters shown in Figure 2.17, where the suppression switch and capacitor are connected upwards, the S2 switch and the Cc suppression capacitor are connected in parallel to the primary windings of the TR transformer.

Here, in addition, the circuit works as a classical step-down converter. Thus, the voltage stress of the switch is equal to the sum of the voltages V_{di} and V_{Cc} .

Air Conditioning and Irrigation System

In the portable vertical farming system, an air conditioning system was designed for two spaces: germination room and production room. The germination room is designed as approximately 6 m². For the heating, cooling and ventilation needs of the germination room, it was found appropriate to position a 9000 btu inverter wall type air conditioner. The production/growing room is designed as approximately 19 m². For the heating, cooling and ventilation needs of the production/growing room, an inverter wall type air conditioner with 18000 btu power was found suitable.

For use in vertical farming applications, a solution tank in the control room and an irrigation system for liquid circulation from the production trays on the shelves were added to the project. According to the number of trays and the length of the pipe, the liquid circulation pump was found to be suitable with a power of approximately 150 W. It is planned to have a water tank with a capacity of 1 ton on the roof for places without mains water connection and to feed the irrigation system.

Energy and Control System

Portable mini-scale vertical farming system is designed as a system with energy management system. Designed as a hybrid energy system of interconnected grid and photovoltaic solar panels, the system can be fed only by interconnected grid connection for projects where low investment cost is desired. Approximately 20 PV panels of approximately 2000X1000 mm in size can be positioned on approximately 26 m² roof area. Depending on the grid-dependent or -independent connection status, the number of PV panels, charging control unit with MPPT feature, inverter power for AC conversion of DC energy and battery capacity required for storage can be integrated into the project portable mini-scale vertical agriculture system.

The design of the micro sprout growing container, which was designed in Dialux evo program, is shown in Figure 2.18, where PV panel and water tank can be positioned on the top of the container




Figure 2.18 Microfiltration Container

Air conditioning automation has been realized in the production and germination rooms so that temperature/humidity sensors control the heating/cooling inverter air conditioners. In addition, ambient measurements will be made with carbon dioxide sensors and carbon dioxide cylinders will be checked at specified intervals.

White Radish Microgreens Growing Results

In order to understand the effects of the LED-based artificial light source of the portable vertical farming system designed within the scope of the project on the growth parameters of the plant, Nitrogen/Nitrogen (N), which are the main macro elements that support growth in solid culture (cocopeat), Secondary microelements such as Phosphorus (P), Potassium (K) and Calcium (Ca), Magnesium (Mg), Sulphur (S), Iron (Fe), Zinc (Zn), Boron (B), Manganese (Mn), Copper (Cu), Molybdenum (Mo), Chlorine (Cl) were also not used.

The white radish microgreen plant has 4 different main stages. These are; rooting, germination period, sprouting period, greening period. These stages are shown in figure 3.1, figure 3.2, figure 3.3



Figure 3.1 White Radish Microgreens Germination-Rooting Stage



Figure 3.2 White Radish Microgreens Sprouting Stage



Figure 3.3 White Radish Microgreens Greening Stage

Some important values were recorded day by day during the period from rooting to the end of the greening period of white radish microgreens. The table of values is shown in table 2.

Table 2. Table of Values of White Radish Microphilis During the Growing Process of White Radish Under Plant Growth Lighting Fixture

	FERTILIZER	IRRIGATION	PLANT		LIGHT
History	pH	Method	Boy	Period	Distance
5.02.2025	7,5	Spraying	0cm	Rooting	25cm
6.02.2025	7,5	Spraying	0cm	Rooting	25cm
7.02.2025	6	Spraying	1cm	Germination	25cm
8.02.2025	6	Spraying	3cm	Germination	25cm
9.02.2025	6	Bottom irrigation	5cm	Germination	25cm
10.02.2025	6	Bottom irrigation	8cm	Sprouting	25cm
11.02.2025	6	Bottom irrigation	9cm	Sprouting	25cm
12.02.2025	6	Bottom irrigation	10cm	Sprouting	25cm
13.02.2025	6	Bottom irrigation	12cm	Greening	25cm
14.02.2025	6	Bottom irrigation	15cm	Greening	25cm
15.02.2025	6	Bottom irrigation	15cm	Greening	25cm

The period from seed to harvest time was 10 days for white radish microphilis plant. After 4 main periods of rooting, germination, germination, sprouting and greening, the microphilis plant

completed its growth as of the 10th day and reached a height of 15 cm. And harvested. Ph values were generally around 6 and spraying and bottom irrigation method was used

Conclusion

The results of the PAR region measurement, which is the photosynthetic wavelength range of the light used during the design and white radish cultivation process, are shown in table 3.

Table 3: PPF Measurement Results for the cultivation area

28,6	33,9	43,3	43,2	43,2	43,3	34	28,7	0,50 m	
36,7	42,1	51,4	52,6	52,6	51,4	42,2	36,8		
36,8	42,5	51,7	52,8	52,8	51,7	42,6	36,9		
36,7	42,1	51,4	52,6	52,6	51,4	42,2	36,8		
28,6	33,9	43,3	43,2	43,2	43,3	34,1	28,7	0,00 m	
0,00 m								1,20 m	

When other methods on the market for the cultivation of microphilis/microgreens were investigated, it was concluded that mostly chemical fertilizers are used. Elements containing heavy metals such as Nitrogen/Nitrogen (N), Phosphorus (P), Potassium (K) are used to accelerate plant growth. However, in our production technique, 100% plant-based growing media (coconut fiber (cocopeat), 100% plant-based fertilizer (seaweed fertilizer) and 100% organic matter (beneficial bacteria and enzymes) are used. Our microfilizers are grown with tap water and we use organic fertilizers, beneficial bacteria and enzymes in a way that is most suitable for nature and human health.

When grown in normal daylight, microphyllis/microgreens complete their development in 30 days. When these plants are grown with special plant growth lighting fixtures with special cultivation methods, their development is completed in about 12-13 days. The white radish microphilis, which was designed within the scope of the project, completed the development process in about 10 days with Samsung vertical agriculture LED chip light source luminaires. When the project product is compared with its counterparts in the market, it has been determined that the harvesting time of microphilis/microgreens is shortened by approximately 20%. Since climate, weather conditions, humidity, temperature, wind, etc. conditions are uncontrolled, it is concluded that products produced with vertical farming techniques can be produced with the same operation every day of the year through air conditioning, air conditioning equipment and artificial lights.



Figure 4.1 White Radish Microphyllis Plant Just Before Harvest

The last image of the white radish microgreen plant before it completes its development and is harvested is shown in Figure 4.1. White radish, which does not contain Nitrogen (N), Phosphorus (P), Potassium (K) macro-elements, or desired microgreens / microfoliage applications are very suitable for use in sectors such as the pharmaceutical industry by dehydration in the mini-scale portable vertical agriculture system designed. At the same time, thanks to its additive-free growth feature, it is easy to integrate into biotechnology products and is also suitable for use as plant-based protein food supplements

Bibliography

- [1] Beacham, A. M., Vickers, L. H., & Monaghan, J. M. (2019). Vertical farming: a summary of approaches to growing skywards. *The Journal of Horticultural Science and Biotechnology*, 94(3), Pp 277-283.
- [2] Şahin, M., & Akkoy, S. (2022) *Alternative Agriculture Practice: Vertical Farming*.
- [3] Samur-Çelebi, Elif. A research on vertical farming practices in the reuse of existing buildings in terms of sustainability: The case of Istanbul porcelain factory. Diss. Master Thesis. Gebze Technical University. Institute of Science and Technology. Gebze, 2019.
- [4] Dayioğlu, Mehmet Ali, and Hasan Silleli. "Artificial Lighting System Design for Greenhouses: Daily Light Integral Method." *Journal of Agricultural Machinery Science* 8.2 (2012): Pp 233-240.
- [5] Seyhan, Temuçin Göktürk, et al. "Principles of Artificial Lighting in Vertical Farming Facilities." *Agricultural Engineering* 376 (2022): Pp 97-106.
- [6] Chen, Li-li, et al. "Effects of different LEDs light spectrum on the growth, leaf anatomy, and chloroplast ultrastructure of potato plantlets in vitro and minituber production after transplanting in the greenhouse." *Journal of Integrative Agriculture* 19.1 (2020): Pp 108-119.
- [7] MASHKOV, Petko Hristov, et al. "LED horticulture lighting for high-voltage AC operation." In: 2018 IEEE XXVII International Scientific Conference Electronics-ET. IEEE, 2018. Pp 1-4.
- [8] VEGNER, Igor Gass, et al. "Horticulture LEDs in Soybean Outdoor Light Supplementation-a Case Study." In: 2021 Joint Conference-11th International Conference on Energy Efficiency in Domestic Appliances and Lighting & 17th International Symposium on the Science and Technology of Lighting (EEDAL/LS:17). IEEE, 2022. p. 1-4.
- [9] DEMIR, Vedat, et al. "Determination of FAR distribution levels of different LED light sources used for photosynthesis in greenhouses." *Journal of Ege University Faculty of Agriculture*, 2021, 58.3: Pp 385-397.
- [10] BARBI, Silvia, et al. "Effects of different LED light recipes and NPK fertilizers on basil cultivation for automated and integrated horticulture methods." *Applied Sciences*, 2021, 11.6: Pp 2497.
- [11] HAO, X.; LITTLE, C.; KHOSLA, S. "LED inter-lighting in year-round greenhouse mini-cucumber production." In: VII International Symposium on Light in Horticultural Systems 956. 2012. Pp. 335-340.
- [12] KUMAR, K. G. S., et al. "Comparison of HPS lighting and hybrid lighting with top HPS and intra-canopy LED lighting for high-wire mini-cucumber production". In: VIII International Symposium on Light in Horticulture 1134. 2016. Pp. 111-118.
- [13] Hogewoning, Sander W., et al. "Blue light dose-responses of leaf photosynthesis, morphology, and chemical composition of *Cucumis sativus* grown under different combinations of red and blue light." *Journal of experimental botany* 61.11 (2010): Pp 3107-3117.

- [14] Gioia, F. D., and P. Santamaria. "Microgreens, agrobiodiversity and food security.(eds) Microgreens." *Eco- logicaeditore, Bari* (2015): Pp.7-24.
- [15] Kyriacou, M. C., Roupael, Y., Di Gioia, F., Kyratzis, A., Serio, F., Renna, M., & Santamaria, P. (2016). Micro- scale vegetable production and the rise of microgreens. *Trends in food science & technology*, V.57, Pp.103-115.
- [16] Zhang, Y., Xiao, Z., Ager, E., Kong, L., & Tan, L. (2021). Nutritional quality and health benefits of microgreens, a crop of modern agriculture. *Journal of Future Foods*, V.1(1), 58-66.
- [17] Ohashi, K., Takase, M., Kon, N, Fujiware.,K., Kurata, K., 2007. Effect of light quality growth and vegetable quality in leaf lettuce, spinach and komatsuna. *Environ. Control Biol.* Pp.189-198.
- [18] Huyen , N. Pham, T., Gioi, D., Ngoc Thang, V., Kim, I., 2014. Effect of light -emitting diode irradiation during healing and acclimatization period on the survival rate and seedling quality of grafted pepper. *J Agric. Life Environ. Sci* Pp.39-47.
- [19] Zhong, L., Zhai, T., Hao, S., Lin, B., Chu, Y., Shen, B., 2022Scanning the lights and examining their proportions during recovery after cucumber transfer. *J. Henan Agricultural Science.* 07, Pp 123-133.
- [20] Madias, Evangelos-Nikolaos D., et al. "The effect of artificial lighting on both biophilic and human-centric design." *Journal of Building Engineering* 76 (2023): 107292.
- [21] Shen, Weiyang, et al. "Pre-harvest short-term continuous LED lighting improves the nutritional quality and flavor of hydroponic purple-leaf lettuce." *Scientia Horticulturae* 334 (2024): 113304.
- [22] Barnes, Charles, et al. "Accuracy of quantum sensors measuring yield photon flux and photosynthetic photon flux." *HortScience* 28.12 (1993): Pp 1197-1200.
- [23] Nelson, Jacob A., and Bruce Bugbee. "Economic analysis of greenhouse lighting: light emitting diodes vs. high intensity discharge fixtures." *PloS one* 9.6 (2014): e99010.
- [24] Cope, Kevin R., M. Chase Snowden, and Bruce Bugbee. "Photobiological interactions of blue light and photosynthetic photon flux: Effects of monochromatic and broad-spectrum light sources." *Photochemistry and photobiology* 90.3 (2014): Pp 574-584.

**THE EFFECT OF ESSENTIAL OILS FROM MEDICINAL AND AROMATIC
PLANTS ON PLANT GROWTH**

**TIBBİ VE AROMATİK BİTKİLERDEN ELDE EDİLEN UÇUCU YAĞLARIN BİTKİ
GELİŞİMİ ÜZERİNE ETKİSİ**

Doç. Dr. Mehmet Zeki KOÇAK

Iğdır Üniversitesi, Ziraat Fakültesi, Tarla Bitkileri Bölümü, Iğdır, Türkiye.

ORCID ID: 0000-0002-8368-2478

Prof. Dr. Bünyamin YILDIRIM

Iğdır Üniversitesi, Ziraat Fakültesi, Tarla Bitkileri Bölümü, Iğdır, Türkiye.

ORCID ID: 0000-0003-2463-6989

ÖZET

Tıbbi ve aromatik bitkilerden elde edilen uçucu yağlar, esas olarak terpenler, fenolikler ve diğer sekonder metabolitlerden oluşan uçucu organik bileşiklerdir. Bunlar, antimikrobiyal, antioksidan ve allelopatik özellikleri de dâhil olmak üzere çeşitli biyolojik aktiviteleri nedeniyle tarımsal araştırmalarda büyük ilgi görmüştür. Bu derleme, uçucu yağların bitki büyümesi üzerindeki hem uyarıcı hem de engelleyici rollerini kapsayan çok yönlü etkilerini ortaya koymakta ve sürdürülebilir tarımdaki potansiyel uygulamalarını vurgulamaktadır. Uçucu yağlar bitki büyümesini doğrudan ve dolaylı mekanizmalarla etkileyebilir. Düşük konsantrasyonlarda, "lavanta (*Lavandula* spp.), biberiye (*Rosmarinus officinalis*) ve nane (*Mentha piperita*)" gibi bazı uçucu yağların tohum çimlenmesini teşvik ettiği, kök ve sürgün uzamasını artırdığı ve fotosentetik verimliliği geliştirdiği gösterilmiştir. Bu büyümeyi teşvik edici etkiler genellikle oksinler ve sitokininler de dâhil olmak üzere bitki hormon seviyelerinin modülasyonuna, besin alımının ve metabolik süreçlerin geliştirilmesine bağlanmaktadır. Aynı zamanda, daha yüksek konsantrasyonlarda uçucu yağlar, allelopatik özelliklerinden dolayı tohum çimlenmesini ve kök büyümesini engelleyerek fitotoksik etkiler gösterebilir. Uçucu yağlar ayrıca bitki savunma mekanizmalarında önemli bir rol oynarlar. Antimikrobiyal özellikleri bitkileri patojenik mantarlardan, bakterilerden ve nematodlardan koruyabilmektedir. Böylece hastalık insidansını azaltır ve daha sağlıklı büyümeyi teşvik eder. Bunlara ek olarak, uçucu yağlar antioksidan enzim aktivitelerini artırarak ve oksidatif hasarı azaltarak bitkinin kuraklık, tuzluluk ve ağır metal toksisitesi gibi abiyotik streslere karşı direncini artırabilir. Genel anlamda, tıbbi ve aromatik bitkilerden elde edilen uçucu yağlar, etkileri konsantrasyona bağlı, türe özgü olmak üzere bitki büyümesinin düzenlenmesinde ikili bir rol sergiler. Düşük konsantrasyonlar büyümeyi uyarabilir ve stres toleransını artırabilirken, yüksek konsantrasyonlar fitotoksositeye yol açabilir. Uçucu yağların tarımsal uygulamalara entegrasyonu, sürdürülebilir mahsul yönetimi için umut vaat etmekte ve haşere kontrolü, yabancı ot yönetimi ve büyümenin artırılması için sentetik kimyasallara doğal alternatifler sunmaktadır. Uçucu yağların uygulama yöntemlerini ve konsantrasyonlarını optimize etmek, çeşitli tarım sistemlerinde etkinliklerini ve güvenliklerini sağlamak için daha fazla araştırma yapılması gerektiği öngörülmektedir.

Anahtar kelimeler: Uçucu yağlar, bitki büyümesi, allelopati, abiyotik stres, sürdürülebilir tarım.

ABSTRACT

The effects of essential oils derived from medicinal and aromatic plants on plant growth. Essential oils, consisting mainly of terpenes, phenolics and other secondary metabolites, are volatile organic compounds that have attracted considerable interest in agricultural research due to their diverse biological activities, including antimicrobial, antioxidant and allelopathic properties. This review demonstrates the versatility of supporting oils, covering both their supportive and inhibitory roles on vegetation, and highlights their potential introduction in sustainable agriculture. Essential oils can affect plant growth through both direct and indirect mechanisms. At low concentrations, essential oils from plants such as lavender (*Lavandula* spp.), rosemary (*Rosmarinus officinalis*) and peppermint (*Mentha piperita*) have been shown to promote seed germination, increase root and shoot elongation and improve photosynthetic efficiency. These growth-promoting effects are often attributed to the modulation of plant hormone levels, including auxins and cytokinins, and the enhancement of nutrient uptake and metabolic processes. Conversely, at higher concentrations, essential oils can exert phytotoxic effects by inhibiting seed germination and root growth due to their allelopathic properties. Additionally, essential oils play an important role in plant defence mechanisms. Their antimicrobial properties help protect plants against pathogenic fungi, bacteria and nematodes, thereby reducing disease incidence and promoting healthier growth. In addition, essential oils can enhance antioxidant enzyme activities and mitigate oxidative damage, thereby increasing plant tolerance to abiotic stresses such as drought, salinity and heavy metal toxicity. Overall, essential oils from medicinal and aromatic plants have a dual role in regulating plant growth, depending on their concentration and species-specific effects. While low concentrations can stimulate growth and improve stress tolerance, high concentrations can lead to phytotoxicity. The integration of essential oils into agricultural practices holds great promise for sustainable crop management by providing natural alternatives to synthetic chemicals for pest control, weed management and growth enhancement. Further research is needed to optimise the application methods and concentrations of essential oils to ensure their efficacy and safety in different agricultural systems.

Key words: Essential oils, plant growth, allelopathy, abiotic stress, sustainable agriculture

GİRİŞ

Tıbbi ve aromatik bitkiler, içerdikleri aktif bileşenler sayesinde hem geleneksel hem de modern tarımda ve özellikle sağlık alanlarında önemli bir yere sahiptir. Bu bitkilerden elde edilen uçucu yağlar, kimyasal bileşimlerine bağlı olarak farklı biyolojik aktivitelere sahip olup; antimikrobiyal, antifungal, insektisit ve allelopatik özellikler göstermektedirler (Giannenas ve ark., 2020; Akademic ve ark. 2023).

Uçucu yağlar, bitkilerin sekonder metabolitleri arasında yer alır ve çeşitli monoterpenler ve fenolik bileşikler içerir. Timol, karvakrol, sineol ve linalool gibi bileşenler, hem bitki gelişimini teşvik edici hem de engelleyici allelopatik etkilere sahip olabilir. Bu etkiler, uçucu yağların bitkiler arası rekabeti düzenleme, hastalık ve zararlılara karşı koruma sağlama ve bitkilerin büyüme düzenleyici mekanizmalarını değiştirme potansiyelinden kaynaklanmaktadır (Saad ve ark., 2013; Sharifi-Rad ve ark., 2017).

Aynı zamanda, bitkisel uçucu yağların tarımda kullanımları kimyasal pestisitlere alternatif doğal biyopestisitlerin geliştirilmesi açısından büyük bir önem taşımaktadır. Özellikle kekik (*Thymus vulgaris*), lavanta (*Lavandula angustifolia*), biberiye (*Rosmarinus officinalis*) ve nane (*Mentha* spp.) gibi bitkilerden elde edilen uçucu yağların, tarım ekosistemlerinde biyolojik mücadele ve bitki gelişim düzenleyicileri olarak kullanımı üzerine yapılan çalışmalar giderek artmaktadır. Ayrıca, uçucu yağ bileşenlerinin fitotoksik etkileri de dikkate alınarak uygun dozajlarda kullanımı önem arz etmektedir (Ngegba ve ark., 2022; Gupta ve ark., 2023).

Bunlara ek olarak, son yıllarda bu alanda yapılan çalışmalar, uçucu yağların tarımsal üretimde kullanılmasının, sentetik kimyasallara olan bağımlılığı azaltabileceğini ve ekolojik sürdürülebilirliği artırabileceğini göstermektedir. Özellikle biyopestisit olarak kullanım alanları genişletilerek, organik tarımda zararlı böceklerin, patojenlerin ve yabancı otların kontrol altına alınmasında etkili bir yöntem olarak değerlendirilmektedir (Koul ve ark., 2008).

Uçucu Yağların Bitki Gelişimine Etkileri

❖ **Allelopatik Etki:** Uçucu yağlar, diğer bitkilerin çimlenmesini, kök ve sürgün gelişimini inhibe edebilir. Özellikle bazı monoterpenler (timol, karvakrol) bitkiler arası rekabeti etkileyerek ekolojik avantaj sağlayabilir. Ayrıca, allelopati, bir bitkinin salgıladığı kimyasal bileşikler aracılığıyla diğer bitkilerin büyümesini ve gelişimini olumlu veya olumsuz yönde etkilemesi olarak bilinmektedir. Bu kimyasal bileşikler, genellikle sekonder metabolitler olup bitki tarafından kök salgıları, yaprak döküntüleri veya uçucu yağlar yoluyla çevreye bırakılabilir. Uçucu yağlar, bitkilerin allelopatik etkileşimlerinde önemli bir rol oynayan doğal bileşikler arasında yer alır (Semerdjieva ve ark., 2022).

Çimlenme Üzerindeki Etkiler:

Uçucu yağlar, hedef bitkinin tohum çimlenme oranını azaltabilir veya tamamen engelleyebilir. Özellikle Timol ve Karvakrol gibi fenolik bileşenler, çimlenme sürecini baskılayarak rekabetçi avantaj sağlarlar.

Kök ve Sürgün Gelişimi Üzerindeki Etkiler:

Uçucu yağların bileşenleri, kök gelişimini inhibe ederek su ve besin alımını zorlaştırabilir. Örneğin, Nane ve Kekik uçucu yağları yüksek konsantrasyonlarda kök uzamasını engelleyebilirler.

Fotosentez ve Metabolizma Üzerindeki Etkiler:

Linalool ve Sineol gibi bazı bileşikler, bitkinin fotosentetik pigmentlerini bozarak fotosentezi azaltabilirler.

Biyokimyasal ve Hücresel Etkiler:

Allelopatik bileşikler, hücre zarının geçirgenliğini değiştirebilir, enzim aktivitelerini bozabilir ve hücre bölünmesini engelleyerek bitkide strese neden olabilirler.

❖ **Tohum Çimlenmesi Üzerine Etkiler:** Çeşitli araştırmalar, bazı uçucu yağların tohum çimlenmesini engellediğini, bazılarının ise teşvik ettiğini göstermektedir. Bitkiler arası etkileşimde önemli rol oynayan bu bileşenler, toprak mikrofloraları ve diğer bitkiler üzerinde de çeşitli biyokimyasal ve fizyolojik etkiler oluşturabilmektedir. Bu etkileşimlerden biri de tohum çimlenmesi üzerine olan etkileridir (Fattahi ve ark., 2019).

Allelopatik Etki: Uçucu yağların içerdiği bileşikler (timol, karvakrol, sineol, linalool vb.) toprak ortamına salındığında, tohum kabuğu üzerinde etki göstererek su ve gaz alışverişini engelleyebilir. Bu durum, tohum çimlenme oranını azaltabilir ya da geciktirebilir.

Hormonel Etkiler: Bazı uçucu yağlar, absisik asit (ABA) gibi çimlenmeyi baskılayan hormonların seviyelerini artırabilirken, gibberellin (GA) gibi çimlenmeyi destekleyen hormonların sentezini engelleyebilir.

Membran Geçirgenliği ve Oksidatif Stres: Uçucu yağlar, tohum zarının yapısını bozarak, su ve besin maddesi alımını olumsuz etkileyebilir. Aynı zamanda serbest radikal oluşumu ve oksidatif stres artışı nedeniyle tohumda metabolik bozukluklara yol açabilir.

Enzim Aktivitesi ve Solunum: Uçucu yağlar, tohum çimlenmesi için gerekli olan enzimlerin (amilaz, proteaz vb.) aktivitesini baskılayabilir veya solunumu azaltarak enerji üretimini düşürebilir.

❖ **Hastalık ve Zararlılara Karşı Koruma:** Uçucu yağlar, bitkilerin doğal savunma mekanizmalarının bir parçası olarak sentezlediği, çeşitli kimyasal bileşiklerden oluşan kompleks karışımlardır. Bu bileşikler, bitki gelişimi üzerinde olumlu etkiler sağlarken, aynı zamanda hastalık ve zararlılara karşı da koruyucu bir rol oynar. Kekik, lavanta, biberiye ve nane gibi bitkilerden elde edilen uçucu yağlar; timol, karvakrol, sineol ve linalool gibi bileşenler içerir. Bu bileşenler antifungal, antibakteriyel ve insektisit özellikleriyle bitkileri çeşitli patojenlerden ve zararlılardan koruyabilir (Raveau ve ark., 2020).

IAntifungal Etkiler: Mantar hastalıkları, bitkiler için önemli bir tehdit oluşturur ve ürün kayıplarına neden olabilir. Uçucu yağlar, mantar patojenlerinin gelişimini engelleyerek biyolojik koruma sağlayabilir.

Antibakteriyel Etkiler: Bakteriyel enfeksiyonlar, bitkilerde ciddi hastalıklara neden olabilir. Uçucu yağlar, bakteriyel patojenlerin hücre duvarlarını bozarak onları etkisiz hale getirebilir.

İnsektisit ve Akarisit Etkiler: Zararlı böcekler ve akarlar, bitki gelişimini olumsuz etkileyerek verim kaybına yol açar. Uçucu yağların böcek öldürücü ve kovucu özellikleri sayesinde biyopestisit olarak kullanımı mümkündür.

Nematisit Etkiler: Toprak kökenli zararlılardan biri olan nematodlar, bitki köklerine zarar vererek büyümeyi engellemektedirler. Uçucu yağlar biyonematisit olarak ta kullanılabilir.

❖ **Büyüme Düzenleyici Etkiler:** Uçucu yağlar, bitki hormonlarıyla etkileşime girerek büyüme hızını değiştirebilir. Özellikle oksin ve sitokin gibi hormonlar üzerindeki etkileri nedeniyle kök ve sürgün gelişimini değiştirebilirler (Gaspar ve ark., 1996).

Oksin, bitkilerde hücre uzaması ve büyümesinin kontrol edilmesinde anahtar rol oynayan bir büyüme hormonudur. Bitkilerde oksin konsantrasyonları, kök gelişimini, sürgün uzamasını ve tohum çimlenmesini etkiler.

Oksin ve Uçucu Yağlar: Bazı uçucu yağ bileşenleri, oksin seviyelerini artırabilir veya azaltabilir. Örneğin, **timol** ve **karvakrol** gibi fenolik bileşikler, oksinlerin biyolojik aktivitesini modüle edebilir. Bu, kök gelişimini uyarabilir veya inhibe edebilir. Özellikle bazı uçucu yağlar, oksin düzeylerini artırarak köklerin daha fazla uzamasını sağlayabilir.

Sitokininler, hücre bölünmesi, büyüme ve organ gelişimini destekleyen bir diğer önemli büyüme hormonudur. Sitokininlerin etkisi, özellikle sürgün gelişimi, yaprak büyümesi ve hücre bölünmesi üzerinde belirgindir.

Sitokinin ve Uçucu Yağlar: Uçucu yağlar, sitokinin seviyelerini etkileyerek bitkilerin organ gelişimini ve büyüme hızını değiştirebilir. Özellikle **linalool** ve **sineol** gibi terpenoid bileşenler, sitokinin üretimini teşvik edebilir ve bitkilerin hücre bölünmesini artırabilir. Bu da sürgün uzamasını hızlandırabilir ve genç bitkilerin sağlıklı gelişmesini destekleyebilir.

Tarımsal Uygulamalar ve Kullanım Alanları

Uçucu yağların tarımda ve bitki koruma alanında kullanımı, sürdürülebilir tarım uygulamalarının gelişmesine katkı sağlamaktadır. Özellikle kimyasal pestisitler ve herbisitlere alternatif olarak kullanılan bu doğal bileşikler, çevre dostu ve insan sağlığına zarar vermeyen çözümler sunmaktadır. Aşağıda, uçucu yağların doğal herbisit olarak, biyolojik mücadelede ve bitki büyümesini teşvik eden maddeler olarak kullanımlarına dair detaylı bir açıklama yapılmıştır (Isman ve ark., 2011; Kesraoui ve ark., 2022).

❖ **Doğal Herbisit Olarak Kullanım:**

Uçucu yağlar, yabancı otlarla mücadelede etkili ve çevre dostu bir alternatif olarak kullanılabilir. Yabancı otların büyümesini inhibe etme kabiliyetleri, özellikle **kekik** (*Thymus vulgaris*), **nane** (*Mentha piperita*) ve **ardıç** (*Juniperus communis*) gibi bitkilerden elde edilen uçucu yağlarla öne çıkmaktadır.

❖ **Biyolojik Mücadelede Kullanım:**

Uçucu yağlar, zararlı böcekler, mantar ve diğer patojenlerle mücadelede etkili doğal pestisitler olarak kullanılabilir. Uçucu yağlar, özellikle **insektisit** ve **fungisit** özellikleri nedeniyle tarımsal zararlılara karşı potansiyel bir alternatiftir.

❖ **Bitki Büyümesini Teşvik Eden Doğal Maddeler:**

Uçucu yağlar, stres koşullarında (kuraklık, tuzluluk gibi) bitkilerin büyümesini teşvik etmek için de kullanılabilir. Özellikle belirli dozlarda uygulandığında, uçucu yağlar bitkilerin stresle başa çıkmalarına yardımcı olabilir ve büyümelerini destekleyebilir.

Sonuç

Tıbbi ve aromatik bitkilerden elde edilen uçucu yağlar, bitki gelişimi üzerinde hem olumlu hem de olumsuz etkilere sahip olabilir. Bu etkiler, uçucu yağın türüne, konsantrasyonuna ve uygulama yöntemine bağlıdır. Doğru kullanıldığında, uçucu yağlar bitkilerin büyümesini teşvik edebilir, stres faktörlerine karşı direncini artırabilir ve zararlılara karşı koruma sağlayabilir. Ancak, yüksek konsantrasyonlarda toksik etkiler gösterebileceği unutulmamalıdır. Bu nedenle, uçucu yağların tarımda kullanımını dikkatli bir şekilde planlanmalıdır. Bu nedenle, tarımsal üretimde uçucu yağların kontrollü kullanımı, sürdürülebilir tarım ve doğal mücadele yöntemleri açısından önemli bir potansiyel taşımaktadır.

KAYNAKLAR

- Academic Press. Greff, B., Sáhó, A., Lakatos, E., & Varga, L. (2023). Biocontrol activity of aromatic and medicinal plants and their bioactive components against soil-borne pathogens. *Plants*, 12(4), 706.
- Fattahi, B., Arzani, K., Sourı, M. K., & Barzegar, M. (2019). Effects of cadmium and lead on seed germination, morphological traits, and essential oil composition of sweet basil (*Ocimum basilicum* L.). *Industrial Crops and Products*, 138, 111584.
- Gaspar, T., Kevers, C., Penel, C., Greppin, H., Reid, D. M., & Thorpe, T. A. (1996). Plant hormones and plant growth regulators in plant tissue culture. *In vitro Cellular & Developmental Biology-Plant*, 32, 272-289.
- Giannenas, I., Sidiropoulou, E., Bonos, E., Christaki, E., & Florou-Paneri, P. (2020). The history of herbs, medicinal and aromatic plants, and their extracts: Past, current situation and future perspectives. *In Feed additives* (pp. 1-18).
- Gupta, I., Singh, R., Muthusamy, S., Sharma, M., Grewal, K., Singh, H. P., & Batish, D. R. (2023). Plant essential oils as biopesticides: Applications, mechanisms, innovations, and constraints. *Plants*, 12(16), 2916.
- Isman, M. B., Miresmailli, S., & Machial, C. (2011). Commercial opportunities for pesticides based on plant essential oils in agriculture, industry and consumer products. *Phytochemistry reviews*, 10, 197-204.
- Kesraoui, S., Andrés, M. F., Berrocal-Lobo, M., Soudani, S., & Gonzalez-Coloma, A. (2022). Direct and indirect effects of essential oils for sustainable crop protection. *Plants*, 11(16), 2144.

- Koul, O., Walia, S., & Dhaliwal, G. S. (2008). Essential oils as green pesticides: potential and constraints. *Biopestic. Int*, 4(1), 63-84.
- Ngegba, P. M., Cui, G., Khalid, M. Z., & Zhong, G. (2022). Use of botanical pesticides in agriculture as an alternative to synthetic pesticides. *Agriculture*, 12(5), 600.
- Raveau, R., Fontaine, J., & Lounès-Hadj Sahraoui, A. (2020). Essential oils as potential alternative biocontrol products against plant pathogens and weeds: A review. *Foods*, 9(3), 365.
- Saad, N. Y., Muller, C. D., & Lobstein, A. (2013). Major bioactivities and mechanism of action of essential oils and their components. *Flavour and Fragrance Journal*, 28(5), 269-279.
- Semerdjieva, I., Atanasova, D., Maneva, V., Zheljazkov, V., Radoukova, T., Astatkie, T., & Dincheva, I. (2022). Allelopathic effects of Juniper essential oils on seed germination and seedling growth of some weed seeds. *Industrial Crops and Products*, 180, 114768.
- Sharifi-Rad, J., Sureda, A., Tenore, G. C., Daglia, M., Sharifi-Rad, M., Valussi, M., ... & Iriti, M. (2017). Biological activities of essential oils: From plant chemoecology to traditional healing systems. *Molecules*, 22(1), 70.

COTTON LEAFWORM, *Spodoptera littoralis* (Boisduval) (LEPIDOPTERA: NOCTUIDAE) AN IMPORTANT QUARANTINE PEST CAUSING HIGH LOSSES IN VEGETABLES

SEBZELERDE BÜYÜK KAYIPLARA YOL AÇAN ÖNEMLİ BİR KARANTİNA ZARARLISI PAMUK YAPRAKKURDU, *Spodoptera littoralis* (Boisduval) (LEPIDOPTERA: NOCTUIDAE)

Eda ÖZÇALICI

Ağrı Ziraî Karantina Müdürlüğü, Ağrı, Türkiye

ORCID ID: <https://orcid.org/0009-0006-7822-4030>

İsmail ALASERHAT

Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, 76000, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6929-0179>

ÖZET

Pamuk yaprakkurdu, *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae), dünya genelinde birçok sebze ve meyveye zarar veren en önemli zararlılardan biridir. Bu zararlı, Akdeniz ve Asya ülkelerinde pamuk (*Gossypium hirsutum* L.), tütün (*Nicotiana tabacum* L.) ve mısır (*Zea mays* L.) bitkilerinin en önemli zararlılarından biri olarak kabul edilmektedir. Larvalar polifag olup, seralarda ve açık alanlarda yetiştirilen çeşitli süs, endüstri ve sebze bitkilerinde ciddi ekonomik kayıplara yol açmaktadır. Zararlı European and Mediterranean Plant Protection Organization (EPPO)'in A2 listesinde yer almaktadır. Türkiye'de Bitki Karantinası Yönetmeliği'nin Ek-1/B listesinde yer almaktadır. *Spodoptera littoralis* uygun koşullarda yılda 4-6 döl verir. Polifag larvaların konukçularına ciddi zararlar vermesi nedeniyle, zararlının kontrol altına alınması büyük önem taşımaktadır. Yumurtadan çıkan larva yaprağın alt yüzeyini (epidermisini) kemirerek zararlı olur. Larva dönemi ilerledikçe larvalar bitkinin diğer kısımlarına yayılırlar ve zarar oranları da artar. Zararlı, yıl boyunca ekonomik değeri yüksek birçok bitkide ciddi zararlar meydana getirebilmektedir. Sebzelerin yaprak, çiçek ve meyvelerinde beslenen zararlının larvaları, meyveleri delerek iç kısımlarına ulaşır ve bu tahribat sonucunda kalite ve verim kaybı ortaya çıkar.

Bu çalışmada, *Spodoptera littoralis* ile ilgili şüana kadar yürütülmüş çalışmalar incelenerek, zararlının morfolojisi, biyolojisi, zarar şekli, yayılışı, konukçuları, doğal düşmanları ve mücadelesi ile ilgili bilgiler derlenmiştir. Böylelikle zararlının dünyadaki mevcut durumu ortaya çıkarılmıştır.

Anahtar Kelimeler: Pamuk yaprakkurdu, *Spodoptera littoralis* (Boisduval), Zararlı, Karantina, Mücadele

ABSTRACT

The cotton leafworm, *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae), is one of the most important pests damaging many vegetables and fruits worldwide. This pest is considered one of the most important pests of cotton (*Gossypium hirsutum*), tobacco (*Nicotiana tabacum*) and maize (*Zea mays*) plants in Mediterranean and Asian countries. The larvae are polyphagous and cause severe economic losses in various ornamental, industrial and vegetable crops grown in greenhouses and outdoors. The pest is on the A2 list of the European and Mediterranean Plant Protection Organization (EPPO). It is listed in Annex-1/B of the Plant Quarantine Regulation in Türkiye. *Spodoptera littoralis* produces 4-6 generations per year under suitable conditions. Because polyphagous larvae cause severe damage to their hosts, control of the pest is important. The larva hatches from the egg and causes damage by feeding at the lower surface (epidermis) of the leaf. As the larval period progresses, the larvae spread to other parts of the host and the damage rates increase. The pest can cause serious damage to many crops of high economic value throughout the year. The larvae of the pest feed on the leaves, flowers and fruits of vegetables and penetrate into the inner parts of the fruit, causing damage and loss of quality and yield.

This study reviewed previous studies on *S. littoralis* and compiled information on the pest's morphology, biology, damage type, distribution, hosts, natural enemies and management. Thus, the current status of the pest in the world was revealed.

Key Words: Cotton leafworm; *Spodoptera littoralis* (Boisduval); Pest; Quarantine; Management

GİRİŞ

Pamuk yaprak kurdu, *Spodoptera littoralis* (Boisduval) Lepidoptera takımı Noctuidae familyasına ait oldukça geniş coğrafik alanlara yayılmış polifag bir zararlıdır. Ana konukçusu pamuk olup soya, tütün, şeker pancarı, patates, mısır, domates, biber, marul, kabak, üzüm, elma ve şeftali gibi sebze ve meyvelerin de içinde bulunduğu yaklaşık 40 familyadan 87 bitki türünde zarar yaptığı bilinmektedir (Özyardımcı, 2002; Hadim, 2008; Anonymous, 2025a). Afrika, Avrupa'nın Akdeniz ülkelerinde ve Ortadoğu'da yaygın olarak bulunduğu bilinmektedir (Pineda et al., 2007; Shairra and Nouh, 2014). Zararlı ile mücadele yapılmadığında %100'e varan kayıplara neden olabilmektedir (Aydın, 2002; Ünlü ve Kornoşor, 2003; Hadim ve Gürkan, 2007; Anonim, 2008; Yıldırım ve Başpınar, 2008). Akdeniz havzasında bulunan ülkelerde yaygın olarak bulunan Pamuk Yaprak kurdu EPPO tarafından A2 karantina listesine tabii tutulmuştur. Türkiye'de ise Akdeniz, Ege, Marmara ve Güneydoğu Anadolu Bölgeleri'nde bulunmaktadır (Anonim, 2025a,b). Pamuk yaprak kurdu ile mücadelede kültürel, kimyasal ve biyolojik yöntemler kullanılmasına rağmen, zararlının tam olarak kontrol altına alınmadığı ve konukçularında yol açtığı zararların günümüzde de devam ettiği görülmektedir. Ekonomik açıdan önemli tarımsal ürünlerin zararlı tarafından ciddi şekilde tahrip edilmesi, insektisit kullanımının ve uygulama dozlarının artmasına yol açmaktadır. Ancak insektisitlerin sürekli ve yüksek dozlarda uygulanması, zararlının kullanılan ilaçlara karşı direnç geliştirmesine neden olmaktadır (Saidy et al., 1989; Delen ve ark., 2005). Kimyasal mücadele, tarımsal ürünlerde kalıntıya neden olmakla kalmaz aynı zamanda çevre, hayvan, insan sağlığı açısından da riskler içerir. Bu sebeple daha çevre dostu olan entomopatojen mikroorganizmaların kullanımı gittikçe artmaktadır.

Pamuk yaprak kurdu, *Spodoptera littoralis* (Boisduval)'in Morfolojisi, Biyolojisi ve Zarar Şekli

Ergin kelebeklerin kanat açıklığı 35-40 mm olup kanatlarda gri kahverengi zeminin üzerinde karışık açık sarı çizgiler bulunmaktadır (Şekil 1a). Kelebekler gündüz kuytu yerlerde saklanır, gece aktiftirler. Yumurtalar yaklaşık 0.4 mm çapında şekilleri yuvarlak olup genelde ikili gruplar halinde yaprağın alt ve üst yüzüne yerleştirilirler. Dişi bireyler tarafından bırakılmalarının ardından üstleri deve tüyü renginde tüylerle kaplanır (Şekil 1b) (Anonymous 2025b). Gelişmesini tamamlamış olan larva 45-50 mm boyunda koyu kahverengi veya siyahımsı kadife görünüştür. Başının üzerinde üçgen şeklinde kahverengi bir leke bulunur. *Spodoptera littoralis*'in geç evrelerinde, ikinci ve üçüncü toraks segmentindeki siyah lekelerin taban kısmında küçük sarı ila beyaz noktalar görülmektedir. Dikkat çekici özelliklerden bir diğeri ise sırttaki koyu lekelerdir, en belirgin olarak 1 ve 8. abdominal segmentte bulunur (Şekil 1c). Larvalar yaprakta meyve üzerinde bulunur. Pupa kahverengidir ve yaklaşık 15-20 mm uzunluğundadır (Şekil 1c). Pupa yaklaşık 0,5 mm uzunluğunda iki dikenden oluşan bir kremastera sahiptir. Zararlı yılda 4-6 döl verir (Anonim, 2008).




Şekil 1. Pamuk yaprak kurdu, *Spodoptera littoralis* (Boisduval)'in a) ergini, b) yumurtaları, c) larvası, d) pupası

Birçok sebzenin çiçeklerinde, meyvelerinde ve yapraklarında beslenirler. Larvalar, sebzelerin yaprak, çiçek ve meyveleriyle beslenir. Meyveleri delerek iç kısımlarına girerken, yaprakları yiyerek elek benzeri bir yapı meydana getirirler. Zarar verdikleri yıllarda önemli ürün kayıplarına yol açabilirler. Domates, biber, patlıcan gibi sebzelerde kalite ve verim kaybına neden olurlar. Bunun sonucunda da pazar değerini azaltabilirler (Anonymous, 2025b).

Yayılışı

Pamuk yaprak kurdu, *S. littoralis* Avrupa, Asya ve Afrika'nın çeşitli bölgelerinde zarara neden olan bir türdür. Avrupa'da Arnavutluk, İtalya, İngiltere ve Portekiz; Asya'da Kıbrıs, Türkiye, Irak, İran, İsrail, Birleşik Arap Emirlikleri, Lübnan, Bahreyn Yemen; Afrika'da Mısır, Libya, Etiyopya, Sudan, Uganda, Zaire, Zambia ve Senegal'de yayılış gösterirler. Ülkemizde ise zararlı özellikle güney bölgelerinde yaygındır (Şekil 2) (Anonim, 2025d).



Şekil 2. Pamuk yaprak kurdu, *Spodoptera littoralis* (Boisduval)'in dünyadaki yayılışı (EPPO, 2024).

Konukçuları

Konukçu sayısı çok fazla olan *S. littoralis*, başta pamuk olmak üzere pirinç, tütün, ayçiçeği, mısır, yonca, domates, yerfıstığı, patlıcan, lahana, biber, şeker kamışı, börülce ve birçok bitki türünde ciddi zararlara neden olabilen bir zararlıdır (Anonim, 2025d).

Doğal Düşmanları

Ülkemizde saptanmış doğal düşmanları arasında, *Chrysoperla carnea* (Stephens, 1836) (Neuroptera: Chrysopidae), *Nabis pseudoferus* (Remane, 1949) (Hemiptera: Nabidae) ve *Microplitis rufiventris* (Kokujev, 1914) (Braconidae: Microgastrinae) bulunmaktadır (Anonim, 2008).

Mücadele Yöntemleri

Pamuk yaprak kurduna karşı kültürel, kimyasal ve biyolojik mücadele yöntemleri uygulanmasına rağmen, zararlının tamamen kontrol altına alınamadığı ve konukçuları üzerinde oluşturduğu zararların günümüzde de sürdüğü gözlemlenmektedir (Özkan ve ark., 2011).

Zararlının mücadelesinde kültürel mücadele olarak zararlının yoğun popülasyonlar oluşturduğu nemli ve loş ortamları ortadan kaldırmak gerekir. Bu yüzden aşırı sulama, fazla gübreleme ve sık ekimden kaçınılmalıdır. Ekim ve dikim usulüne uygun olarak yapılmalıdır (Anonim, 2008).

Biyoteknik mücadelede kullanılan feromon tuzaklar, yetişkin Lepidoptera türlerini tespit etmek için kullanılabilen ve bu türlerin izlenmesinde birincil yöntem olan bir tekniktir. Larvalar, çoğunlukla toprakta pupa haline gelmesi nedeniyle, pupa evresi hariç tüm gelişim aşamaları konak bitkileri veya ürünlerinde görsel olarak tespit edilebilir. Yetişkinlerin veya larvaların bir bitki veya üründe tespiti için çeşitli tanımlama yöntemleri mevcuttur.

Pamukta, 10.000 yumurta kütesi / hektarlık bir ekonomik eşik, pestisit müdahalesinin planlanması için güvenilir ve pratik olarak kabul edilmiştir (Hosny et al., 1986). *Spodoptera littoralis*'in tarla popülasyonları için çeşitli insektisit sınıflarına karşı insektisit direncinin geliştiği bildirilmiştir ve yönetim stratejilerinin, farklı etki biçimlerine sahip uygulanan insektisitlerin rotasyonuna dayandırılması önerilmektedir (Issa et al., 1984a,b; Sawicki, 1986; Temerak, 2002; Elghar et al., 2005).

Spodoptera littoralis'in yumurtaları ve larvalarının doğal kontrolüne yardımcı olan çeşitli parazitoit türleri belirlenmiştir (Gerling, 1971; Vojtech et al., 2005; Depalo et al., 2010; Hatem

et al., 2016; Agbodzavu et al., 2018). Pamuk yaprakkurdunun bulunduğu bölgelerde *Bacillus thuringiensis*'e dayalı ticari ürünlerin kullanımı yaygın olarak gerçekleştirilmektedir (Navon et al., 1983; Magholifard et al., 2020). Ayrıca, entomopatojen funguslar, entomopatojen nematodlar ve ticari olarak temin edilebilen özel bir nükleopolihedrovirüs, *S. littoralis* popülasyonlarının kontrolü için sahada kullanılmaktadır (Resquín-Romero et al., 2016; Sutanto et al., 2017; Sobhy et al., 2020).

Sonuç olarak *S. littoralis* polifag bir zararlı olup konukçu dizini oldukça fazladır. Ayrıca zararlının bir karantina zararlısı olması önemini daha da artırmaktadır. Günümüzde globalleşen dünya şartlarında *Spodoptera littoralis*'in uluslararası ticaret yoluyla EPPO bölgesine daha fazla getirilip yayılabileceği düşünülmektedir. Bitki sağlığı önlemleri, zararlılardan arı alanlara veya zararlılardan arı üretim alanlarına dikim amacıyla bitki yetiştirilmesini ve ihracat öncesi ürünlerin denetlenmesini gerekli kılar.

KAYNAKLAR

- Agbodzavu, M.K., Lagat, Z.O., Gikungu, M., Rwomushana, I., Ekesi, S. & Fiaboe, K.K.M., (2018). Performance of the newly described endoparasitoid *Cotesia icipe* Fernandez-Triana & Fiaboe on *Spodoptera littoralis* (Boisduval). *Journal of Applied Entomology*, 142 (7), 646–653.
- Anonim, (2008). Ziraî Mücadele Teknik Talimatları Cilt 2. Gıda Tarım ve Hayvancılık Bakanlığı Tarımsal Araştırmalar ve Politikalar Genel Müdürlüğü Yayınları, Ankara, 260 s.
- Anonim, (2025a). Pamuk yaprak kurdu, *Spodoptera littoralis*. <https://agrobasesapp.com/turkey/pest/pamuk-yaprak-kurdu> (Erişim Tarihi: 16.02.2025).
- Anonim, (2025b). Pamuk yaprak kurdu, *Spodoptera littoralis*. https://www.tarimziraat.com/hastalik_ve_zararlılar/endustri_bitkileri_zararlılari/%20soyada_pamuk_yaprak_kurdu/ (Erişim Tarihi: 16.02.2025).
- Anonim, (2025c). Pamuk yaprak kurdu. <https://www.koppert.com.tr/bitki-zararlılari/tirtillar/pamuk-yaprak-kurdu/> (Erişim Tarihi: 17.02.2025).
- Anonim, (2025d). Çizgili Pamuk Yaprak Kurdu (*Spodoptera* spp.). https://www.progenseed.com/images/upload/4493_4975.pdf (Erişim Tarihi: 17.02.2025).
- Anonymous, (2025a). EPPO (European and Mediterranean Plant Protection Organisation). https://www.eppo.int/ACTIVITIES/plant_quarantine/A2_list (Accessed on 16 February 2025).
- Anonymous, (2025b). *Spodoptera littoralis*. https://en.wikipedia.org/wiki/Spodoptera_littoralis (Accessed on 17 February 2025).
- Aydın, H., (2002). *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae)'e Karşı Bazı *Tanacetum* spp. (Compositae) Ekstraktlarının Biyolojik Aktivitelerinin Belirlenmesi. Ankara Üniversitesi, Fen Bilimleri Enstitüsü, (Basılmamış) Yüksek Lisans Tezi, Ankara, 100 s.
- Delen, N., Durmuşoğlu, E., Günçan, A., Güngör, N., Turgut, C. & Burçak, A., (2005). Türkiye'de pestisit kullanımı, kalıntı ve organizmalarda duyarlılık azalışı 123 sorunları. Türkiye Ziraat Mühendisliği 6. Teknik Kongre (3-7 Ocak Ankara).
- Depalo, L., Marchetti, E., Baronio, P., Martini, A., & Dindo, M.L., (2010). Location, acceptance and suitability of *Spodoptera littoralis* and *Galleria mellonella* as hosts for the parasitoid *Exorista larvarum*. *Bulletin Entomology*, 63, 65–69.
- Dündar, M.B., (2024). Bakulovirüs Enfeksiyonunun *Spodoptera littoralis* (Lepidoptera: Noctuidae) Üzerinde Sublethal Etkilerinin Belirlenmesi. Yüksek Lisans Tezi. Ankara Üniversitesi Fen Bilimleri Enstitüsü Bitki Koruma Anabilim Dalı, Ankara.

Elghar, G.E.A., Elbermawy, Z.A., Yousef, A.G. & Elhady, H.K.A., (2005). Monitoring and characterization of insecticide resistance in the cotton leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *Journal of Asia-Pacific Entomology*, 8(4), 397–410.

EPPO, (2025). EPPO (European and Mediterranean Plant Protection Organization) Global Database. Database on quarantine pests (available online). *Spodoptera littoralis*. <https://gd.eppo.int> (Accessed on 17 February 2025).

Gerling, D., (1971). Occurrence, abundance, and activity of some local parasitoids attacking *Spodoptera littoralis* [Lepidoptera: Noctuidae] in selected cotton fields in Israel. *Annals of the Entomological Society of America*, 64, 492–499.

Gördebak, S., (2024). Bazı Bitki Ekstraktları ve Nano Partiküllerinin *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) Larvaları Üzerine İsektisidal Etkisinin Araştırılması. Yüksek Lisans Tezi. Tokat Gaziosmanpaşa Üniversitesi Lisansüsü Eğitim Enstitüsü Bitki Koruma Anabilim Dalı, Tokat.

Hadim, N. & Gürkan, M.O., (2007). Pamuk yaprak kurdu (*Spodoptera littoralis* Boisd.) (Lepidoptera: Noctuidae)'nda sentetik pyretroidlere karşı ortaya çıkan direncin biyokimyasal mekanizmaları, 60. Türkiye II. Bitki Koruma Kongresi (27-29 Ağustos 2007) Bildirileri, Isparta, 342 s.

Hadim, N., (2008). Pamuk yaprak kurdu *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae)'te insektisitlere karşı oluşan direncin biyokimyasal ve moleküler karakterizasyonu, Ankara Üniversitesi Biyoteknoloji Enstitüsü Doktora Tezi, Ankara, 146 s.

Hatem, A., Shaver, D. & Vargas-Osuna, E. (2016). Parasitism and optimization of *Hyposoter didymator* (Hymenoptera: Ichneumonidae) reared on *Spodoptera littoralis* and *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Journal of Economic Entomology*, 109(3), 1058–1063.

Hosny, M.M., Topper, C.P., Moawad, G.M. & El-Saadany, G.B., (1986). Economic damage thresholds of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) on cotton in Egypt. *Plant Protection*, 5(2), 100–104.

Issa, Y.H., Keddis, M.E., Abdel-Sattar, M.A., Ayad, F.A. & El-Guindy, M.A. (1984a). Investigation of resistance to organophosphorus insecticides in field species of cotton leafworm during the 1980-1984 cotton growing seasons. *Bulletin of the Egyptian Entomological Society, Economic Series*, 14, 399-404.

Issa, Y.H., Keddis, M.E., Abdel-Sattar, M.A., Ayad, F.A. & El-Guindy, M.A. (1984b). Investigation of resistance to pyrethrins in cotton leafworm field strains 1980-1984 cotton growing seasons. *Bulletin of the Egyptian Entomological Society, Economic Series*, 14, 405-411.

Magholifard, Z., Hesami, S., Marzban, R. & Salehi Jouzani, G., (2020). Individual and combined biological effects of *Bacillus thuringiensis* and multicapsid nucleopolyhedrovirus on biological stages of the Egyptian cotton leafworm *Spodoptera littoralis* (B.) (Lep.: Noctuidae). *JAST*, 22(2), 465-476.

Navon, A., Wysoki, M. & Keren, S., (1983). The power and effect of *Bacillus thuringiensis* preparations against *Spodoptera littoralis* and *Boarmia (Ascotis) selenaria* larvae. *Fitopasitika*, 11, 3-11.

Özkan, F., Demirbağ, Z. & Demir, İ., (2011). Bakterilerin *Spodoptera littoralis* (Lepidoptera: Noctuidae)'e karşı mikrobiyal mücadele etmeni olarak araştırılması, Türkiye IV. Bitki Koruma Kongresi (28-30 Haziran 2011) Bildirileri, Kahramanmaraş, 496 s.

Özyardımcı, B., (2002). *Spodoptera littoralis* (Boisd.)'in değişik popülasyonlarında chlorpyrifosa duyarlılık farklarının radyoizotop izleme tekniği ile saptanması üzerinde araştırmalar. Doktora Tezi. Ankara Üniversitesi Bitki Koruma Anabilim Dalı, s. 1-7. Ankara

- Partal, E., (2023). Bazı Entomopatojen Nematodların *Spodoptera littoralis* (Boisduval, 1833) (Lepidoptera: Noctuidae)'e Etkinliklerinin Laboratuvar Koşullarında Belirlenmesi. Yüksek Lisans Tezi. Ege Üniversitesi Fen Bilimleri Enstitüsü Bitki Koruma Anabilim Dalı, İzmir.
- Pineda, S., Schneider, M.I., Smagghe, G., Martinez, A.M., Del Estal, P., Vinuela, E., Valle, J. & Budia, F., (2007). Lethal and sub lethal effects of Methoxyfenozide and Spinosad on *Spodoptera littoralis* (Lepidoptera: Noctuidae). *J. Econ. Entomol.*, 100: 773-780.
- Resquín-Romero, G., Garrido-Jurado, I. & Quesada-Moraga, E., (2016). *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae) kontrolü için entomopatojenik mantarlar ve bunların ekstraktlarının kombine kullanımı. *Biyolojik Kontrol* 92, 101–110.
- Saidy, M.F.E., Auda, M. & Degheele, D., (1989). Detoxification mechanisms of diflubenzuron and teflubenzuron in the larvae of *Spodoptera littoralis* (Boisd.). *Pesticide Biochemistry and Physiology*, 35, 211-222.
- Sawicki, R.M., (1986). Successfully preventing resistance to synthetic pyrethrins. *Agribusiness Worldwide*, 8(20), 22-25.
- Shairra, S.A. & Nouh, G.M., (2014). Efficacy of entomopathogenic nematodes and fungi as biological control agent against the cotton leaf worm, *Spodoptera littoralis* (Boisd.). *Egypt. J. Biol. Pest Control.*, 24(1), 247-253.
- Sobhy, H.M., Abdel-Bary, N.A., Harras, F.A., Faragalla, F.H. & Hussein, H.I., (2020). Efficacy of entomopathogenic nematodes against *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (H.) (Lepidoptera: Noctuidae). *Egypt. J. Biol. Pest Control.*, 30 (73), 1-8.
- Sutanto Koko, D., El Salamouny, S., Tufail, M., Khawaja, G.R., Sukirno, S., Shepard, M., Shapiro, M. & Saad Aldawood, A., (2017). Evaluation of natural additives to enhance the persistence of *Spodoptera littoralis* (Lepidoptera: Noctuidae) Nucleopolyhedrovirus (SpliMNPV) under field conditions in Saudi Arabia. *Journal of Economic Entomology*, 10 (3), 924–930.
- Temerak, S.A., (2002). Historical records of resistance of field cotton leafworm (*Spodoptera littoralis*) to conventional insecticides as influenced by resistance programs in maize. *Resistant Pest Management Bulletin*, 12, 7-10.
- Ünlü, L., & Kornoşor, S., (2003). Şanlıurfa ilinde saptanan Noctuidae (Lepidoptera) familyası türleri ve morfolojik özellikleri. *Harran Üniversitesi Ziraat Fakültesi Derg.*, 7, 19-28.
- Vojtech, E., Meissle, M. & Poppy, G.M., (2005). Effects of Bt maize on herbivorous *Spodoptera littoralis* (Lepidoptera: Noctuidae) and parasitoid *Cotesia marginiventris* (Hymenoptera: Braconidae). *Transgenic Research*, 14 (2), 133–144.
- Yıldırım, E.M. & Başpınar, H., (2008). Aydın ili çilek alanlarında saptanan Noctuidae (Lepidoptera) familyası türleri, yayılışı, zararı ve popülasyon dalgalanmaları üzerinde çalışmalar. *ADÜ Ziraat Fakültesi Dergisi*, 5(2), 115-121.

**WASTE MANAGEMENT IN AGRICULTURE: RECYCLING OF ORGANIC
WASTE AND COMPOSTING TECHNIQUES**

**TARIMDA ATIK YÖNETİMİ: ORGANİK ATIKLARIN GERİ DÖNÜŞÜMÜ VE
KOMPOSTLAMA TEKNİKLERİ**

Hasine KÜÇÜKYILDIRIM

Harran University, Faculty of Agriculture, Department of Soil Science and Plant Nutrition,
Şanlıurfa, Turkey

ORCID ID: <https://orcid.org/0000-0001-5822-9439>

Abdulkadir BÜYÜK

Iğdır University Faculty of Agriculture, Department of Field Crops, Iğdır/ Turkey²

ORCID ID: <https://orcid.org/0009-0002-8323-8032>

ÖZET

Tarım sektöründe organik atık yönetimi, çevresel sürdürülebilirlik ve tarımsal verimlilik açısından büyük önem taşımaktadır. Tarımsal faaliyetler sırasında oluşan bitkisel, hayvansal ve gıda işleme atıkları doğru şekilde geri dönüştürüldüğünde, hem çevreye olan olumsuz etkiler azaltılmakta hem de tarımsal üretimde verimli kullanım sağlanmaktadır. Kompostlama, organik atıkların yeniden değerlendirilmesi için en etkili yöntemlerden biridir. Geleneksel, vermikompost, hızlı termofilik ve anaerobik kompostlama gibi çeşitli teknikler, atıkların biyolojik ayrışmasını hızlandırarak, toprağın besin değerini artıran ve su tutma kapasitesini yükselten kompost ürünlerinin elde edilmesini sağlar. Kompost kullanımı, kimyasal gübre ihtiyacını azaltarak sürdürülebilir tarım uygulamalarını destekler ve tarımsal üretimde doğal döngülerin korunmasına katkı sağlar, tarımsal toprakların biyolojik aktivitesini destekleyerek mikrobiyal çeşitliliği artırır ve toprak sağlığını iyileştirir. Ayrıca, organik madde içeriğinin yükseltilmesi sayesinde, toprak erozyonunun önlenmesine katkı sağlar. Kompostlanan organik atıkların gübre olarak kullanılması, kimyasal gübre ihtiyacını azaltarak çiftçilerin üretim maliyetlerini düşürmesine yardımcı olur. Bununla birlikte, organik gübrelerin toprağa uzun vadeli faydalar sağladığı ve kimyasal gübrelere kıyasla daha dengeli bir besin kaynağı sunduğu bilinmektedir. Bu nedenle, organik atık yönetimi stratejilerinin yaygınlaştırılması ve çiftçilerin bu konuda bilinçlendirilmesi, çevre dostu ve sürdürülebilir tarım uygulamalarının başarısı için kritik bir adımdır. Devlet teşvikleri, eğitim programları ve yerel yönetimlerin destekleri ile kompost kullanımının artırılması, uzun vadede tarım sektöründe daha verimli ve ekolojik olarak dengeli bir üretim modeline geçişi mümkün kılacaktır.

Anahtar kelimeler: Organik atık yönetimi, Çevresel sürdürülebilirlik, Tarımsal verimlilik, Kompostlama

ABSTRACT

Organic waste management in the agricultural sector is of great importance in terms of environmental sustainability and agricultural productivity. When plant-based, animal-based,

and food processing wastes generated during agricultural activities are properly recycled, both the negative environmental impacts are reduced, and efficient use in agricultural production is ensured. Composting is one of the most effective methods for reutilizing organic waste. Various techniques such as traditional composting, vermicomposting, rapid thermophilic composting, and anaerobic composting accelerate the biological decomposition of waste, resulting in compost products that enhance soil nutrient content and increase water retention capacity. The use of compost supports sustainable agricultural practices by reducing the need for chemical fertilizers and contributes to the preservation of natural cycles in agricultural production. Additionally, it enhances the biological activity of agricultural soils, increases microbial diversity, and improves soil health. By increasing the organic matter content, it also helps prevent soil erosion. The use of composted organic waste as fertilizer reduces the need for chemical fertilizers, helping farmers lower production costs. Furthermore, organic fertilizers are known to provide long-term benefits to the soil and offer a more balanced nutrient source compared to chemical fertilizers. Therefore, the widespread implementation of organic waste management strategies and raising awareness among farmers are critical steps for the success of environmentally friendly and sustainable agricultural practices. Increasing the use of compost through government incentives, educational programs, and support from local administrations will enable the transition to a more efficient and ecologically balanced production model in the agricultural sector in the long term.

Keywords: Organic waste management, Environmental sustainability, Agricultural efficiency, Composting

GİRİŞ

Dünyada ve ülkemizde yıllar içerisinde artış gösteren tarımsal üretim sonucunda hem bitkisel atıklar hem de tarımsal endüstri atıkları da artan bir ivme göstermektedir. Bu atıklar önemli bir organik madde kaynağı olma özelliğindedir. Dünyada ve ülkemizde artarak devam eden tarımsal üretimin hem bitkisel üretim sonrası atıkların hem de tarımsal ürünlerin işlenmesi sonucu oluşan atıkların, çevresel kirliliğin önlenmesi ve atıkların değerlendirilmesi amacıyla geri dönüşümünün sağlanması oldukça önemlidir. (Güven, 2020).

Tarımsal faaliyetler sonucu oluşan organik atıklar, doğru şekilde geri dönüştürülmediğinde çevresel ve ekonomik sorunlara yol açabilir. Ancak, bu atıklar sürdürülebilir tarım uygulamaları ile verimli bir şekilde dönüştürülebilir. Organik atıkların geri dönüşümü, çevresel etkileri azaltmak ve tarımsal verimliliği artırmak için önemli bir stratejidir.

Arazi bozulması tarımsal üretkenliği, çevre ve gıda güvenliğini ciddi olarak etkileyen önemli sorunlarından biri olduğu düşünülmektedir. Arazi bozulmasını engellemenin en önemli yolu toprak organik maddesini arttırarak toprak sağlığını iyileştirmektir. Fakat günümüzde hayvansal gübre ve kompost kullanımından vazgeçilmiş olup kimyasal gübre kullanımı yaygınlaşmıştır. Fakat kimyasal gübreler giderek daha maliyetli ve güvensiz hale gelmiş bununla birlikte çevre kirlenmesine neden olmuş ve toprak organik karbonunu korumak açısından katkısı az olmuştur. Ticari gübrenin başlıca alternatifleri tarımsal atıklardan elde edilen kompost ve çiftlik gübresi olmuştur.

Kompost; biyokimyasal olarak ayrışabilir çok çeşitli organik maddelerin organizmalar tarafından stabilize edilmiş, mineralize olmuş halidir. Kompostlaştırma ise mikroorganizma adı verilen ve çoğunluğu gözle görülmeyen canlıların, ortamın oksijenini kullanarak çöp içerisindeki organik maddeleri biyokimyasal yollarla ayrıştırmasıdır (Erdin, 2009; Yaman 2012). Kompostlaştırma doğada, bitki yetişen her yerde doğal olarak yürüyen bir süreçtir. Bitkiler yapraklarını döktüğünde veya onların herhangi bir artığı toprağa düştüğünde, bu bitkisel artıklar doğada mikroorganizmalar tarafından ayrıştırılarak humus formuna dönüştürülür. Başka bir deyişle, mikroorganizmalar organik materyali ayrıştırıp daha stabil olan

organik son ürünlere dönüştürerek karbondioksit, su, ısıl enerji ve humusa çevirirler. Bu ekosistemdeki besin geri dönüşümünün göstergesidir. Kompost yapımı bu doğal olayın iyi bir şekilde incelenip taklit edilmesiyle oluşmuştur. Başka deyişle bu doğal ayrışma, ideal koşullar yaratılarak hızlandırılmıştır (Kaçar, 2018).

Kompost endüstrisi, 1980'lerde, atık suyun arıtılmasında kullanılan teknolojik gelişmelerle birlikte, tarım ve bahçecilik alanında kullanılan yüksek kaliteli kompost üretimi de gelişim göstermiştir (Goldstein, 2001; Güven, 2020). Kompost yapımında genellikle bitkisel materyaller (budama atıkları, yapraklar, buğday samanı, odunsu bitki materyali), gıda işleme atıkları (pirina, domates, yeşil ceviz kabuğu vb), hayvansal gübreler ve belediyelerin atık su arıtımından ortaya çıkan atık çamurlar kullanılır. Tüm bu kompostlama materyalleri organik maddece oldukça zengindir.

İklim değişikliği, kontrolsüz sanayileşme, bilinçsiz tarım faaliyetleri vb. sebepler nedeniyle tarım toprakları her geçen gün azalmakta olup, çevre kirliliği küresel bir problem haline gelmiştir. Hızla büyüyen dünya nüfusuna paralel olarak beslenme ihtiyacını karşılamak ise zorlaşmıştır. Bu sebeple toprakta sürdürülebilirlik kavramı önem arz etmeye başlamış ve fiziksel, kimyasal ve biyolojik özelliklerine etki ederek topraktan sürdürülebilir verim almayı sağlayan organik gübrelerin kullanımı ve çeşitliliği artış göstermiştir.

Tarımsal üretimde bitkinin toprakta gelişebilmesi, yetiştiği toprak ortamının fiziksel ve kimyasal özellikleri ile ilişkilidir. Kumlu topraklar genellikle düşük su tutma kapasitesi gibi kötü fiziksel ve zayıf kimyasal özelliklere sahiptir (Chesworth ve diğerleri 2008). Toprağın fiziksel özelliklerini düzeltmede ve sürekliliğini sağlamada en fazla başvurulan yöntem, toprağa organik kökenli materyallerin ilave edilmesidir. Kompost veya diğer organik düzenleyicilerin kullanılması, topraktaki mevcut besinlerin miktarını artırır ve besin ve su dinamikleri üzerinde uzun vadeli etkileri vardır (Srivastava ve diğerleri, 2016). Kumlu topraklarda, hatta organik maddenin ayrışmasının mikrobiyal aktivite için elverişsiz olduğu koşulların bulunduğu yarı kurak bölgelerde bile organik düzenleyicilerin etkileri daha kısadır.

Son zamanlarda ise toprağa bitki besin maddesi kazandırma, patojen baskılama ve atık değerlendirme gibi kompostlama kazanımları yanı sıra topraktaki mikrobiyal aktiviteyi daha da arttırmaya yönelik, daha hızlı ve daha etkili bir son ürün elde edilmesini sağlayan solucanlar ile kompostlama işlemi bir diğer deyişle vermikompostlama teknolojisi de önem kazanmıştır (Boran, 2023).

Vermikompostlama; belirli bir solucan türü ile atık dönüşüm sürecinin termofilik kompostlama işlemine göre daha hızlı gerçekleştiği biyoteknolojik bir kompostlama çeşididir (Edwards ve Bohlen 1996). Organik atıkların vermikompostlaması ile elde edilen ve vermikompost adı verilen son ürünün tarım topraklarına uygulanması halinde bitki gelişimi yanında toprağın fiziksel, kimyasal ve biyolojik özelliklerine olumlu etkileri bulunmaktadır.

Bununla birlikte tarım arazilerine doğru kompost uygulamaları; toprakların su tutma, su, hava ve ısı iletkenliği, agregasyon, hacim ağırlığı ve kıvam gibi bazı fiziksel ve mekaniksel özelliklerini, katyon değişim kapasitesi, bitki besin maddesi içeriği, toprak reaksiyonu ve tuzluluk gibi kimyasal özelliklerini ve yüksek karbon ve azot içeriği ile toprakların mikrobiyolojik özelliklerini olumlu yönde etkilemektedir (Jat ve Ahlawat 2006, Alam vd. 2007, Singh vd. 2008, Sarı vd. 2017, Temiz vd. 2021; Temiz, 2023).

TARIMSAL ATIKLAR VE GERİ DÖNÜŞÜM POTANSİYELİ

Tarımsal faaliyetler sırasında oluşan organik atıklar şu şekilde sınıflandırılabilir:

- Bitkisel atıklar (sap, saman, meyve-sebze artıkları, yaprak döküntüleri)
- Hayvansal atıklar (gübre, kesimhane artıkları)

- Gıda işleme atıkları (meyve ve sebze kabukları, kullanılmayan gıda artıkları)

Bu organik atıklar, uygun geri dönüşüm yöntemleri kullanılarak yeniden değerlendirilebilir ve tarım için faydalı bir kaynak haline getirilebilir.

Bitkisel Atıklar

Türkiye’de 1960’lı yıllarda üretilen toplam katı atık miktarı yılda 3-4 milyon ton iken, 2008 yılı itibarı ile toplam atık miktarı yaklaşık 24.36 milyon tona yükselmiş olup bunun yaklaşık 6.7 milyon tonu evsel nitelikli atıklardır. Kişi başı günlük katı atık üretim miktarı 1.15 kg’dır. Toplam atıkların ancak %1.1 (267.960 ton) geri kazanım tesislerine kompost üretimi amacıyla gönderilmiştir (TUİK, 2012). Katı atıkların %1.1’inin kompost üretiminde değerlendirilmesi yeterli değildir. Bu nedenle, katı atıklar içinde geri dönüşümünün sağlanması durumunda elde edilecek katma değeri en yüksek olan bitkisel atıkların yeniden değerlendirilmesi çok önemlidir. Öyle ki pazardan bir TL’ye alınan bir bitkisel ürünün atık kısımları iyi bir biçimde değerlendirildiği takdirde asıl fiyatının 5-10 misli değere ulaştığı görülmektedir. Özellikle ilaç, kozmetik, parfümeri ve boya sanayinde hiç değer verilmeyen bitkisel atık ve artıklar hammadde olarak kullanılmaktadır (Yaman,2012).

Geri dönüşüme konu olan bitkisel kökenli atıklar; meyve, sebze ve diğer bitkisel ürünlerin tüketim amaçlı olarak ev, lokanta ya da sanayi mutfağında kullanım öncesi atılan kabukları, çiçekleri, yaprakları, çekirdekleri, sap ve köklerini içeren evsel bitkisel atıklar ve gerek kentsel gerekse kırsal alanlarda yetişen ağaçların ve bitkilerin yaprakları, dalları, kabukları ve meyvelerini içeren evsel olmayan bitkisel atıklar olarak iki grupta incelenir.

Hayvansal Atıklar

Hayvan çiftliklerinde büyük ve küçükbaş hayvanlardan oluşan atıklar ve atık sular çevre kirliliğine neden olmaktadır. Hayvansal atıklar genel manada hayvan dışkıları, mezbaha atıkları ve hayvansal atıkların işlenmesi-paketlenmesi ve nakliyesi sırasında oluşan atıklar olarak sınıflandırılabilir. Bunun dışında işletmedeki gübre atıkları, hayvan kadavraları bunlara eklenebilir. Hayvansal atıklar genel olarak biyodizel, biyogaz, gübre ve yem olarak değerlendirilmektedir. Bu atıkların değerlendirilmesi hayvansal atığın türüne göre olmaktadır (Gökmen ve Sarıçoban; 2012).

Gıda İşleme Atıkları

Geri dönüşüm potansiyeli yüksek olan gıda işleme atıkları pirina, domates, yeşil ceviz kabuğu gibi meyve ve sebze kabukları ile kullanılmayan gıda atıkları kullanılarak elde edilen organik maddece zengin kompost tarım için önemli ve faydalı bir kaynak olarak kullanılabilir.

KOMPOSTLAMA TEKNİKLERİ

Kompostlama, organik atıkların kontrollü bir şekilde ayrışmasını sağlayarak besin değeri yüksek bir toprak düzenleyici elde etme yöntemidir. Başlıca kompostlama teknikleri şunlardır:

Geleneksel Kompostlama

- Organik atıklar katmanlar halinde yığılır ve belirli aralıklarla karıştırılarak havalandırılır.
- 3-6 ay içerisinde olgunlaşır.

Geleneksel kompostlama genellikle dört temel aşamadan oluşur:

- **Mezofilik Evre:** 15-40°C aralığında gerçekleşen bu evrede, mezofilik mikroorganizmalar organik maddenin parçalanmasına başlar (Tiquia ve Tam, 2002).
- **Termofilik Evre:** Sıcaklığın 45-70°C'ye ulaştığı bu aşamada, patojenlerin yok edilmesi ve organik maddelerin daha hızlı ayrışması sağlanır.

- **Soğuma Evresi:** Mikroorganizma aktivitesi azalırken kompost stabil hale gelir.
- **Olgunlaşma Evresi:** Besin açısından dengeli ve bitkiler için uygun bir kompost oluşur (Bernal ve ark., 2009).

Kompostlama sürecinde sıcaklık, nem, oksijen miktarı ve karbon/azot (C/N) oranı gibi faktörler büyük önem taşımaktadır. Optimum C/N oranı 25-30 arasında olmalıdır; düşük C/N oranları azot kaybına, yüksek oranlar ise ayrışma sürecinin yavaşlamasına neden olabilir (Kumar ve ark., 2010). Geleneksel kompostlama ile elde edilen kompost, toprak yapısını iyileştirir, su tutma kapasitesini artırır ve bitkiler için gerekli besin maddelerini sağlar. Aynı zamanda, sentetik gübre kullanımını azaltarak çevre dostu bir alternatif sunar (Zhu, 2007).

Vermikompost (Solucan Kompostu)

Solucan gübresi olarak da bilinen vermikompost, organik atıkların toprak solucanları aracılığıyla ayrışması sonucunda oluşan, humusa benzer bir maddeyi ifade eder. Sığır, koyun, at ve tavşan gibi hayvanların gübrelere, orman endüstrisinden kaynaklanan artıklar, mutfaktan çıkan organik atıklar, çim ve sebze-meyve bahçelerindeki budama ile hasat kalıntıları, kâğıt atıkları ve diğer pek çok organik materyal solucanların besin kaynağı olarak değerlendirilerek vermikompost üretimi gerçekleştirilebilir. Solucanların sindirim sisteminde bulunan çeşitli mikroorganizmalar sayesinde bu atıklar parçalanarak, hoş olmayan kokular içermeyen ve organik madde açısından zengin bir gübreye dönüşür. Bu sürecin temel hedefi, organik atıkları en hızlı ve verimli şekilde işleyerek değerlendirmektir.

Vermikompost, besin maddelerinin yavaş salınımını sağlayarak tarımsal üretimi artıran ve besin kayıplarını en aza indiren bir özellik göstermesiyle güvenilir bir organik gübre olarak kabul edilir. Kullanıldığı topraklarda fiziksel, kimyasal, biyolojik ve mikrobiyolojik açıdan iyileştirmeler yapması, onu tarımsal uygulamalarda önemli bir unsur haline getirir. Vermikompostun bilinen başlıca avantajları arasında toprak yapısını düzenleyici etkisi, yeterli miktarda bitkiler için faydalı besin maddeleri içermesi, bazı zararlı organizmaları ve bitki hastalıklarını kontrol etme yeteneği, toprak kalitesini yükselterek verimi artırması, çevre dostu olması ve uzun vadede ekonomik bir seçenek sunması sayılabilir. Solucanların organik maddeleri sindirmesiyle elde edilen vermikompost, bitki gelişimi, toprak iyileştirme, bitki sağlığı ve çevresel faydalar açısından normal komposttan daha üstün olduğu belirtilen bir materyaldir (Fritz ve ark., 2012; Bellitürk ve ark., 2013; Bellitürk ve ark., 2015). Vermikompostlama süreci, organik atıkların solucanlar tarafından biyogübreye dönüştürülmesi olarak tanımlanır ve günümüzde organik katı atık yönetiminde yaygın olarak uygulanmaktadır (Manyuchi ve ark., 2013). Organik atıkların vermikompostta dönüşümünde en sık kullanılan altı solucan türü bulunmaktadır. Ilıman iklimlerde *Eisenia fetida* (*Eisenia andrei* ile benzer tür), *Dendrobaena veneta* ve *Lumbricus rubellus* yaygınken, tropikal bölgelerde *Eudrilus eugeniae*, *Perionyx excavatus* ve *Perionyx hawayana* tercih edilmektedir. Bunların dışında farklı solucan türleri de kullanılabilir de, bahsedilen altı tür kadar yaygın olarak tercih edilmemektedir (Edwards, 2004).

Vermikompostun içeriğini en çok etkileyen faktörlerden birinin, solucanlara verilen yem olduğu belirlenmiştir. Bunun yanı sıra, kullanılan solucan türü, üretim ortamının koşulları ve uygulanan uzmanlık seviyesi de önemli etkenler arasında yer almaktadır. Sadece sığır gübresinin kullanılmasıyla üretilen vermikompostun organik madde oranının diğer türlere kıyasla %44 daha yüksek olduğu tespit edilmiştir. Bu durum, sığır gübrelere "vermikompost teknolojisi" ile işlenerek daha verimli ve değerli bir solucan gübresine dönüştürülmesi gerektiğini göstermektedir. Bu gübrenin N, P ve K gibi temel besin elementlerini yeterli seviyede içermesinin yanı sıra, özellikle ülkemiz tarım topraklarında eksikliği yaygın olarak görülen Zn (çinko) ve B (bor) gibi mikro elementler açısından da zengin olması dikkate değer bir özelliktir. Organik madde içeriğine ek olarak, çinko ve bor eksikliği bulunan tarım alanlarında vermikompost gibi organik gübrelere kullanımının yaygınlaştırılması büyük önem taşımaktadır. Vermikompost, yalnızca toprak için gerekli olan makro ve mikro besin

elementleri bakımından değerli olmakla kalmaz, aynı zamanda toprak yapısını iyileştirici etkisiyle de önemli avantajlar sunar.

Çizelge 1. Sığır gübresi, öğütülmüş kağıt atığı, zeytin budama atığı, % 50 zeytin budama atığı+% 50 sığır gübresi atığından elde edilen 4 çeşit vermikompostun analiz sonuçları.

Kimyasal İçerik	Sığır Gübresi Vermikompostu (Arancon ve Edwards, 2011)	Gıda+Bahçe Atığı Vermikompostu (Lange, 2005)	Kağıt Atıkları Vermikompostu (Bellitürk ve ark., 2015a)	Zeytin Budama Atığı Vermikompostu (Bellitürk ve ark., 2014)	Sığır Gübresi+ Zeytin Budama Atığı Vermikompostu (Bellitürk ve ark., 2014)
N, %	1.90	1.81	1.11	1.80	1.62
P, %	4.70	1.01	0.14	0.23	0.44
K, %	1.40	1.04	0.32	1.74	1.77
B, ppm	58	-	-	64	45
Ca, ppm	23	0.28	26	4.15	3.75
Fe, ppm	3454	1440	-	15451	12653
Mg, ppm	5802	2100	0.33	0.54	0.47
Mn, ppm	160	346	132	586	525
Zn, ppm	516	387	49	70	104
Org. Mad.%	44	30	39	48	48

Hızlı (Termofilik) Kompostlama


Yüksek sıcaklıkta mikroorganizma faaliyetleri hızlandırılarak kompostlama süreci kısaltılır. 4-8 hafta gibi kısa sürede kompost elde edilir. Sıcak kompostun dezavantajlarından biri, yığın oluşturulduktan sonra yeni organik madde eklenemeyecek olmasıdır. Yığını hazırlamak için bir kafes içine su, odun külü, saman, taş, talaş, tahıl, yaprak gibi maddeleri hayvan gübresi ile yığıp karıştırabilirsiniz. Yığın hazırlandıktan sonra, sıcaklık 30 dereceye ulaşana kadar küf mantarları, bakteriler, protozoalar ve nematodlar aktif rol oynar. 20–40 derece arasındaki bu evreye” mezoflik evre” adı verilir. 30–40 derece arasında, asıl humuslaştırıcı organizmalar olarak bilinen aktinomisetler egemen olmaya başlar. Sıcaklık 40–50 dereceye ulaştığında kompostlaşmayı başlatan organizmaların neredeyse tamamı ölür ve bunların yerini 70 dereceye kadar dayanabilen ve ısı üreten termofilik bakteriler alır. 60–70 derece aralığında birkaç spor dışında temel olarak bütün patojenik organizmalar birkaç saat içinde ölür. Termofilik bakteriler kendileri için mevcut besini tükettiklerinde ısı üretmeyi durdurur ve kompost soğumaya başlar. Sıcaklık, yeniden termofilik evre eşiğinin altına düşünce, yığının içinde kalan organik maddeyle beslenen, mantar ve aktinomiset ağırlıklı yeni bir grup mikroorganizma çoğalır. Bunlar, ölü bakterileri de besin olarak kullanır. Tüm bu aşamalar tamamlandıktan sonra, çok sayıda solucan ve böcek larvası oluşur. Elimizde kalan malzeme, artık daha fazla parçalanamayan maddelerden oluşan organik bir kütledir.

Anaerobik Kompostlama (Biyogaz Tesisleri)

Biyogaz, organik kökenli atık ve artıkların oksijensiz bir ortamda parçalanması (anaerobik fermantasyon) sonucunda oluşan, renksiz, kokusuz, havadan daha hafif ve parlak mavi bir alevle yanan bir gaz karışımıdır. İçeriğinde yaklaşık olarak %40-70 oranında metan, %30-100 arasında karbondioksit, %0-3 oranında hidrojen sülfür ile eser miktarda azot ve hidrojen bulunur (Öztürk, 2008).

Biyogaz, birçok farklı organik atık türünden üretilen ve hammadde sorunu bulunmayan bir enerji kaynağıdır. Şeker pancarı, ayçiçeği, kanola gibi enerji bitkileri, bahçe atıkları, kâğıt sanayi atıkları, yemek artıkları, sebze, meyve, tahıl ve yağ sanayisi atıkları, hayvansal gübreler, deri ve tekstil sektörü atıkları, evsel ve kentsel organik atıklar ile atık su arıtma tesisi çamurları biyogaz üretiminde yaygın olarak kullanılan kaynaklar arasındadır (Gürel ve Şenel, 2010). Biyogaz üretim sürecinde kullanılan hammaddeler ile enerji akışları Şekil 1'de gösterilmektedir (Berglund ve Börjesson, 2006).

Şekil 1: Biyogaz Üretiminde Kullanılan Hammaddeler ve Enerji Akışları



Kompost Kullanımının Faydaları

Kompost kullanımının tarımsal ve çevresel faydaları şunlardır:

- En başta katı atık depolama alanlarında tasarruf sağlar. Bu alanlarda birikerek çevreye zararlı metan gazlarının üretilmesini yol açması engeller, faydalı ve değerli bir kaynağa dönüşmesi sağlar.
- Kompostlama, toprak yapısını iyileştirir. Yağış olmadığı dönemlerde nemi toprakta tutar ve buharlaşmayı azaltır. Aynı zamanda kompost sonucunda ortaya çıkan humus hafif bünyeli topraklarda besin maddesi ve su tutma yeteneğinin artmasını sağlar.
- Toprağın hava almasına yardımcı olur. Toprağın üst kısmında havadar bir tabaka oluşturur ve topraktaki çeşitli canlıların hareket etmesine uygun bir ortam hazırlar. Böylece toprağın hava almasını kolaylaştırır.
- Bitkilerin ihtiyacı olan zamanda besin sağlar. Hasat edilerek topraktan uzaklaştırılan organik maddenin yerini alır ve topraktaki humus oranını dengeler. Bitkilerin istediği zamanda ve formda alabilecekleri besinleri kapsar.
- Ayrıca yüksek mineralli gübrelemeye karşı bir nevi tampon görevi görür ve bu sayede bitkilerin zarar görmesini engeller.

- Kompostlama sonucu elde edilen kompost tarımda, bahçecilikte ve benzeri alanlarda kullanılabilir. Böylece gübre tüketiminin azaltılması, yapay gübrenin az kullanılması ise çevre kirliliğinin azaltılmasını ve ekonomik açıdan tasarruf edilmesini sağlar.
- Toprakta bulunan toksit maddeler ve ağır metaller, bitkilerin köklerinin ememeceği bir forma dönüşür ve sabitlenir. Toprağa kompost eklenmesi kirlenmiş toprakların iyileşmesini sağlar.
- İşlenmesinde zorluk yaşanan toprakların daha kolay işlenmesini sağlar.
- Asidik toprakların pH'sini artırır.
- Bitkilerin büyümesini hızlandırır ve güçlenmelerini sağlar. Toprağa sağladıkları hümitik asit sayesinde bitkilerin gelişimini hızlandırır.

Faydaları sıralanan kompostlamanın sonucunda elde edilen ürün tarımda, çeşitli spor ve oyun tesislerinin kurulmasında, çöp toplama alanlarının yeşillendirilmesinde, ormancılıkta ve bahçecilikte yaygın bir şekilde değerlendirme imkânı bulunmaktadır.

SONUÇ

Organik atıkların geri dönüşümü ve kompostlama teknikleri, sürdürülebilir tarımın temel unsurlarından biridir. Bu yöntemlerin yaygınlaştırılması, hem çevresel etkileri azaltacak hem de tarımsal verimliliği artıracaktır. Çiftçilerin ve tarımsal işletmelerin bu konuda bilinçlendirilmesi, etkili bir atık yönetimi stratejisi oluşturulması için kritik bir öneme sahiptir.

KAYNAKÇA

- Bellitürk, K., Görres, J. H., Turan, H. S., Göçmez, S., Bağdatlı, M. C., Eker, M. ve Aslan, S. (2014). Zeytin Bitki Artıkları-Ahır Gübresi-Kum Karışımı İle Yapılacak Olan Vermikompostun Tarımda Kullanılabilirliğinin Araştırılması. Namık Kemal Üniversitesi Bilimsel Araştırma Projesi, Proje No: NKUBAP.00.24.AR.13.15.
- Bellitürk, K., Aslan, S., Eker, M. (2013). Ekosistem Mühendisleri Diye Adlandırılan Toprak Solucanlarından Elde Edilen Vermikompostun Bitkisel Üretim Açısından Önemi. Hasad (Bitkisel Üretim) Aylık Tarım Dergisi, Eylül, İstanbul, Yıl: 29 (340): 84- 87.
- Bellitürk, K., Shrestha, P. ve Görres, J. H. (2015). The Importance of Phytoremediation of Heavy Metal Contaminated Soil Using Vermicompost for Sustainable Agriculture. Rice Journal 3:2, 6: e114, doi: 10.4172/2375-4338.1000e114
- Bellitürk, K., Zahmacıoğlu, A., Şerif, E., Top, M. (2015a). Kağıt Atıklarının Vermikompost (Solucan Gübresi) Yapılarak Değerlendirilmesi Projesi. Trakya Toprak (Verimli Toprakların Dergisi), Yıl: 1, Sayı: 3, Sayfa: 21-24, Tekirdağ.
- Berglund, M. & Börjesson, P. (2006), "Assessment of Energy Performance in the Life-Cycle of Biogas Production", Biomass & Bioenergy, 30 (3), 254–266.
- Bernal, M. P., Albuquerque, J. A., & Moral, R. (2009). Composting of animal manures and chemical criteria for compost maturity assessment: A review. Bioresource Technology, 100(22), 5444-5453.
- Boran, D. (2023). Farklı Materyallerden Elde Edilen Kompost ve Vermikompostun Biyokimyasal Özellikleri ve Bakteriyel Çeşitliliğinin Belirlenmesi. Ankara Üniversitesi Fen Bilimleri Enstitüsü. Doktora Tezi. Ankara
- Edwards, C. A., Bohlen, P. J. 1996. Biology and ecology of earthworms, Springer Science & Business Media
- Edwards, C.A. (2004). Earthworm Ecology (2nd Edition). CRC Press. Boca Raton, FL, London, New York, Washington. 448 pp.

- Fritz, J. I., Franke-White, I. H., Haindl, S., Insam, H., Braun, R. (2012). Microbiological Community Analysis of Vermicompost Tea and its Influence on the Growth of Vegetables and Cereals. *Canadian Journal of Microbiology*, 58:836-847.
- Goldstein, N. (2001). The industry in the United States: past, present, and future. *Compost Utilization in Horticultural Cropping Systems*, 3-16).
- GÖKMEN, S., & SARIÇOBAN, C. HAYVANSAL ATIKLARIN GERİ DÖNÜŞÜMÜ. 1. ULUSAL GERİ KAZANIM KONGRE ve SERGİSİ, 121.
- Gürel, A. & Şenel, Z. (2010), "Organik Atıklardan Biyogaz Üretimi", II.Trakya Bölgesi Kalkınma Girişimcilik Sempozyumu, 1-2 Ekim, İğneada-Kırklareli, 123-133.
- Güven, O.B. (2020). Farklı Tarımsal Atık ve Kompost Uygulamalarının Toprak Özellikleri Üzerine Etkileri. Çanakkale Onsekiz Mart Üniversitesi Fen Bilimleri Enstitüsü. Yüksek Lisans Tezi. Çanakkale
- Jat, R.S. and Ahlawat, I.P.S. 2006. Direct and residual effect of vermicompost, biofertilizers phosphorus on soil nutrient dynamics and productivity of chickpea- fodder maize. *Journal of Sustainable Agriculture*, 28, 41-54.
- Kaçar, H. (2018). Elma İşleme (Ön Eleme-Ayırma) Katı Atıklarının Kompostlaştırılmasında Farklı Sıcaklık Uygulamalarının Süreç Performansına Etkisi. Süleyman DEMİREL Üniversitesi Fen Bilimleri Enstitüsü. Yüksek Lisans Tezi. Isparta
- Kumar, M., Ou, Y. L., & Lin, J. G. (2010). Co-composting of green waste and food waste at low C/N ratio. *Waste Management*, 30(4), 602-609.
- Lange, M. G. (2005). A Comparison Analysis of Vermicomposting Strategies in Food Substrates with an Emphasis on Nutrient Values and Reproduction (Research Advisor: Dr. Christopher Baxter and Dr. Ken Killian). Pioneer Undergraduate Research Fellowship, May 27, 2005. pp. 1- 15.
- Manyuchi, M. M., Phiri, A., Muredzi, P., Chitambwe, T. (2013). Comparison of Vermicompost and Vermiwash BioFertilizers from Vermicomposting Waste Corn Pulp. *International Scholarly and Scientific Research & Innovation* 7 (6): 389- 392.
- Öztürk, H.H. (2008), "Yenilenebilir Enerji Kaynakları ve Kullanımı", Teknik Yayınevi, Ankara, 367.
- Srivastava, V., De Araujo, A. S. F., Vaish, B., Bartelt-Hunt, S., Singh, P., ve Singh, R. P. (2016). Biological response of using municipal solid waste compost in agriculture as fertilizer supplement. *Reviews in Environmental Science and Bio/Technology*, 15(4), 677-696.
- Temiz, Ç. (2023). Farklı Kompostlama Yöntemleriyle Elde Edilmiş Organik Materyallerin Bazı Toprak Fiziksel Özellikleri, Organik Karbon Formları ve Mısır Bitkisinin Verimi Üzerine Etkileri. Ankara Üniversitesi Fen Bilimleri Enstitüsü. Doktora Tezi. Ankara
- Tiquia, S. M., & Tam, N. F. Y. (2002). Characterization and composting of poultry litter in forced-aeration piles. *Process Biochemistry*, 37(8), 869-880.
- Zhu, N. (2007). Effect of low initial C/N ratio on aerobic composting of swine manure with rice straw. *Bioresource Technology*, 98(1), 9-13.

DEVELOPMENT OF FOOD FROM PAST TO PRESENT AND FOOD SECURITY

GEÇMİŞTEN GÜNÜMÜZE GIDANIN GELİŞİMİ VE GIDA GÜVENCESİ

Dr. Osman İNAN

Bağımsız araştırmacı, İstanbul, Türkiye

ORCID ID: <https://orcid.org/0000-0001-7306-5516>

Dr.Öğr.Üyesi Sema KONYALI

Tekirdağ Namık Kemal University, Faculty of Agriculture, Department of Agricultural Economics, Tekirdağ, Türkiye

ORCID ID: <https://orcid.org/0000-0002-6049-495X>

ÖZET

Giriş ve Amaç: Günümüzde nüfus artışı, sanayileşme, kentleşme, turizm, tarım gibi sebeplerle oluşan insan kaynaklı sorunlar tarihte daha önce olmadığı kadar gıda güvencesine tehdit oluşturmaktadır. Ek olarak, nüfus artışından kaynaklanan çevre sorunları, kuraklık, savaşlar, nükleer enerji, asit yağmurları gibi sebepler ekonomik, çevresel ve sosyal sürdürülebilirliği olumsuz yönde etkilemekte ve gıda güvencesini kuşkuya düşürmektedir. Gıda güvencesinin; uluslararası antlaşmalarda da tanımlanan insani, vicdani ve toplumsal vd. nedenleriyle devletlerin sorumluluğu altında sağlanması elzemdir. Bu çalışmada, geçmişten günümüze gıdanın gelişimini incelemek ve gıda güvencesinin mevcut durumunu ortaya koyarak sorunlara çözüm önerileri getirebilmek amaçlanmıştır.

Gereç ve Yöntem: Bu çalışma, konu ile ilgili literatürün ve raporların incelenmesine dayanmaktadır. Çalışmanın ilk bölümünde, gıdanın insanlığın ortaya çıkışından günümüze değin geçirdiği süreçler olarak gelişimi ayrıntılı olarak açıklanmıştır. Sonrasında, günümüzde gıda güvencesinin genel durumu ortaya konulmaya çalışılmıştır. Bu amaçla, ilk olarak gıda güvencesi kavramı hakkında bilgiler verilmiştir. İkinci olarak gıda güvencesinin sağlanmasının zorunluluğu detaylı bir şekilde açıklanmıştır. Üçüncü olarak, gıda güvencesinin mevcut durumu hakkında uluslararası kuruluşların raporlarına dayanarak bilgiler verilmiş ve gıda güvencesinin ölçülmesi konusu incelenmiştir.

Bulgular: Ülkeler, vatandaşlarının gıda hakkını sağlamaya çalışmakta, bunun için günümüz sorunlarını çözmek, ulusal stratejiler belirlemek ve küresel ortak amaçları karşılamayı amaçlamaktadırlar. Gıda güvencesinin sağlanmasında ciddi zorluklar yaşanmaktadır. Ülkelerin bunu karşılamak için uygun strateji geliştirmelerine ve uluslararası işbirliklerine ihtiyaç bulunmaktadır. Bu doğrultuda gıda politikaları; gıda arz güvenliğinin sağlanması, gıda sistemlerinin dayanıklı yapıda olması, tüketicinin güvenilir gıdaya erişiminin tam ve istikrarlı bir şekilde güvence altına alınması ile sürdürülebilir olacaktır.

Tartışma ve Sonuç: Çalışmada, gıda politikasının sürdürülebilir başarısının, herkesin gıda güvencesinin sağlanabilmesi olduğu vurgulanmıştır. Çalışmanın, gıda güvencesi konusunda literatüre katkı sağlaması ve politika yapıcılara ışık tutması hedeflenmektedir.

Anahtar Kelimeler: Sürdürülebilirlik; Gıda; Tarım; Gıda Politikaları; Gıda Güvencesi

ABSTRACT

Introduction and Objective: Today, human-induced problems such as population growth, industrialisation, urbanisation, tourism and agriculture pose a threat to food security like never before in history. In addition, environmental problems arising from population growth, drought, wars, nuclear energy, acid rain, etc. adversely affect economic, environmental and social sustainability and put food security in doubt. It is essential to ensure food security under the responsibility of states for humanitarian, conscientious, social, etc. reasons defined in international treaties. In this research, it is aimed to examine the development of food from past to present and to bring solutions to the problems by revealing the current situation of food security.

Materials and Method: This study is based on a review of the literature and reports on the subject. In the first part of the study, the development of food as a process from the emergence of humanity to the present day is explained in detail. Afterwards, the general situation of food security today has been tried to be revealed. For this purpose, firstly, information about the concept of food security is given. Secondly, the necessity of ensuring food security is explained in detail. Thirdly, information on the current state of food security is given based on the reports of international organisations and the issue of measuring food security is examined.

Findings: Countries endeavour to ensure the right to food for their citizens, to solve contemporary problems, to define national strategies and to meet global common goals. There are serious challenges in ensuring food security. Countries need to develop appropriate strategies and international co-operation to meet this challenge. In this direction, food policies will be sustainable by ensuring the security of food supply, ensuring that food systems are resilient, and ensuring that consumers' access to safe food is fully and consistently secured.

Discussion and Conclusion: The study emphasized that the sustainable success of food policy is to ensure food security for all. The study is expected to contribute to the literature on food security and shed light on policy makers.

Keywords: Sustainability; Food; Agriculture; Food Policies; Food Security

GİRİŞ

Gıda ihtiyacının ve güvenilir gıdaya olan talebinin karşılanması insanın öncelikli ve en temel olan gereksinimidir. İnsanlar tarih boyunca mücadelelerini, içgüdüsel olarak ilk başta günlük gıdalarını elde edebilmek ve hayat sürecinde gıdasız kalmamak üzerine kurmuşlardır. Bireyler ve toplumlar gıdaya daha kolay ulaştıkça yaşamın diğer temel gereksinimlerine ve daha gelişmiş değerlere daha hızlı ulaşmışlardır. İnsanlığa büyük ilerleme kaydettiren yerleşik düzene geçmek, medeniyetlerin kurulması, ve varlıklarını uzun süre devam ettirebilmeleri, tarihte ilk imparatorlarının ortaya çıkması, geniş bozkırları aşabilme becerisi, bilimsel gelişmelerin temellerinin atılması, ticaretin gelişebilmesi, kalabalık yerleşimler kurabilmesi, uzak coğrafyalara keşiflerin yapılabilmesi, sanayi devrimi ile kentlere geniş kesimleri çekebilme becerisi, teknolojinin günümüzdeki seviyesine gelmesi gıda konusunda ilerleme kaydetmesiyle beraber yaşanan gelişmeler olmuştur.

Daha büyük topluluklar ve ülkelerin başarılarındaki gibi bireylerde kişisel mücadelesinde gıda sorunlarını hafiflettikçe, kendilerini daha iyi anlayabilmiş ve insanlığa daha yararlı olmuşlardır. Günümüze gelindiğinde de, gıdaya sahip olan ülkelerin daha refahta ve gıdaya kolay ulaşan bireylerin hayattan beklentilerinin yüksek ve yaşama katkılarının daha fazla olduğu görülebilmektedir. İnsan hayatının her döneminde pek çok yönüyle gelişebilmek için bireysel gıda güvencesinin tam olarak sağlanmasına muhtaçtır. Gıda güvencesinden yoksun olma durumu, daha çocukluk çağında olumsuz etkilerini göstermekte (Yanardağ, 2023), insanın yaşamı boyunca başarıyı engellemektedir. Gıda güvencesi hayattaki başarı ile beraber

koordineli olarak yükselmekte, gıda güvencesizliği ise yaşamın zor koşulları ile birleştiğinde olumsuz bir sarmala dönerek başarısızlığı beraberinde getirebilmektedir.

Bu çalışmada, öncelikle gıdanın tarihsel gelişimi ayrıntılı olarak açıklanmıştır. Sonrasında, günümüzde gıda güvencesinin genel durumu ortaya konulmaya çalışılmıştır. Bu amaçla, ilk olarak gıda güvencesi kavramı açıklanmış, ikinci olarak gıda güvencesinin bileşenleri ve gıda güvencesi için önemi ortaya konulmuştur. Üçüncü olarak, gıda güvencesinin mevcut durumu hakkında uluslararası kuruluşların raporlarına dayanarak bilgiler verilmiş ve gıda güvencesinin ölçülmesi konusu incelenmiştir. Sonuçta, gıda politikalarının sürdürülebilir başarısının, herkesin gıda güvencesinin sağlanabilmesi olduğu vurgulanmıştır. Çalışmanın, gıda güvencesi konusunda literatüre katkı sağlaması ve politika yapıcılara ışık tutması amaçlanmaktadır.

MATERYAL VE YÖNTEM

Bu çalışma, konu ile ilgili literatürün ve raporların incelenmesine dayanmaktadır. Çalışmada, öncelikle gıdanın insanlığın ortaya çıkışından günümüze değin geçirdiği süreçler olarak gelişimi ayrıntılı olarak açıklanmıştır. Sonrasında, günümüzde gıda güvencesinin genel durumu ortaya konulmaya çalışılmıştır. Bu amaçla, ilk olarak gıda güvencesi kavramı açıklanmıştır. İkinci olarak gıda güvencesinin sağlanmasının zorunluluğu detaylıca açıklanmıştır. Üçüncü olarak, gıda güvencesinin mevcut durumu hakkında uluslararası kuruluşların raporlarına dayanarak bilgiler verilmiş ve gıda güvencesinin ölçülmesi konusu incelenmiştir.

BULGULAR VE TARTIŞMA

GIDANIN GELİŞİMİ

İnsanlık tarihi incelendiğinde, homo sapiens'in ilk ortaya çıkışından bugüne değin insan ilk olarak gıdasını güvence etme peşinde koşmuştur. Bu büyük mücadelesinde çok büyük zorluklarla karşılaşmış ve ciddi aşamalar kaydetmiştir. Gıdayı elde edebilmeye ilk olarak toplayıcı olarak başlamıştır. Bu dönem aletleri çok ilkel olduğundan dönemin insanları da çok kolay elde edebilecekleri gıdalara ulaşabilmişlerdir. Toplanan gıdaların depolanması ya da takas edilmesi toplayıcılığın ilk dönemlerinde çok uzun süre düşünülmemiş (Özçelik, 2014), genellikle açlıkla mücadele etmek için gündelik beslenilmiştir. Gıdasını elde etme mücadelesinde bulunduğu coğrafyayı yeterli olarak anlayamamış ve gıdayı yeterince depolayabilme becerilerini henüz geliştirememiştir. Bu gibi uyum sağlama becerilerinin noksanlığından dolayı gıda temin etmek için yerini değiştirme zorunluluğu ve fiziksel mücadelesinin yüksek olması kendisine toplayıcılıkta güç, sezgi, faydalı olanı anlama gibi farklı anlamlarda beceriler kazandırmıştır (Aytan, 2021). İlk zamanlarda uzun bir süre toplayıcılık süreci devam ederken insanlar avlanarak gıda temin edebilmeyi de öğrenmiştir. Bundan sonra insan toplayıcı-avcı olarak uzun bir süre gıdasını sağlamaya çalışmıştır. Nitelikli bir gıda depolama anlayışının geliştiği ise halen uzunca bir süre için söylenememektedir. Ancak, toplayıcılıkta edinilen deneyimler ve gıda için daha büyük çapta avlanmayı beceribilmek insanlık için gıda temininde ciddi gelişmelerdir. Gıdaların toplanması olarak kabul edilen biriktirme iktisadı, çok uzun süre devam etmiş ve gelişmemiş toplumlarda yakın dönemlere kadar uygulaması sürmüştür.

Tarihsel süreçte insanların gruplar halinde avlanması, takası keşfetmeleri, ateşi bularak başta eti ısıtmaları, gıdaları ile ilgili diğer önemli gelişmelerdir. Gruplar halinde avlanabilme becerisi, büyük boyutlu canlıları avlamalarını sağlamış, insanları pek çok canlının önüne geçirmiş ve yaşam mücadelesinde güçlü canlılar arasında piramidin en üst basamaklarında yer almalarına katkı sağlamıştır. Yine takas becerisinin gelişmesi, insanları farklı besinler yemeye ve besin depolayabilme becerilerini geliştirmelerine imkân vermiştir. Bir diğer önemli adımda ateşin bulunması ile insanlar çeşitli hastalıklardan kurtulmuş ve gıdadan daha iyi faydalanabilmiştir.

Var oluşunun ilk yıllarında insanların bu becerilerini geliştirmeleri, beyin yapılarının daha iyi gelişmesini ve diğer canlılardan yüksek becerilere sahip olmalarını sağladığı anlaşılabilir. Diğer canlılara üstünlük kurabilen, coğrafyasındaki gıdalarından daha iyi faydalanabilen, etrafını algılama becerileri gelişen insanın gıda temelli ilerlemesinden memnun olduğu ve bununla gelişme kaydettiğini anladığı da anlaşılabilir.

Tarım devriminden hemen önceki yakın zamanlarda insanın el aletlerini kullanabilmesi, bunları geliştirmesi ve gıdasını edinebilmesinde bunlardan yararlanabilmesi, hem işbirliğini geliştirmiş hem de insanın kolayda gıda bulabilmek için uzak mesafelere gitmesini engellemiştir. İnsanın uzak mesafeler kat etmesini, gıda bulmak amacıyla mevsim değişikliklerinde farklı coğrafyalara taşınmasını, göçe bağlı yaşamasını da engelleyen gıda becerileri geliştirmiş olması aynı zamanda onu fiziksel anlamda yavaşlatmıştır. Bunun neticesinde, daha büyük grupların oluştuğu, gruplar arasında işbirliklerinin arttığı, gıda işbirliklerinin artmasının diğer materyallerin de işbirliğini arttırdığı tahmin edilebilir. Yine M.Ö. 9.500-8.000’li yıllar olarak hesaplanan dünyanın farklı yerlerinde farklı zamanlarda ortaya çıktığı tahmin edilen tarım devrimi ile, başlıca tarım ürünlerinde verim ve üretim artışı sağlandığı, tarımda bir nevi daha başlangıç seviyesinde bir uzmanlaşmanın sağlanması ile geçmişten gelen gıda yeniliklerinden insanların oldukça faydalanmaları sebebiyle göçebelikten kurtularak tarım şehirlerine daha hızlı bir geçiş yapıldığı söylenebilir (Direk, 2012). Tarım devrimi sonrasında yine bireyler ve toplumlar arasında gıda işbirliklerinin arttığı ve dünyada o güne kadar olmayan iletişimin geliştiği bilinmektedir.

Gıdaya olan görece bu daha kolay olan erişim neticesinde ilk yerleşmeler, bir kaç yüzyıl sonra ilk medeniyetleri meydana getirmiştir. Tarımın keşfedildiği ilk yıllarda ve sonraki yüzyıllarda daha büyük işbirliklerinin gelişmesi, gıdanın depolanması, hava durumunun tahmin edilmeye çalışılması, tohumun saklanması gibi olaylar sayısız tarihçi tarafından günümüz tarım teknolojilerinin ilk ayak izleri sayılmaktadır. İlk şehirlerin kuruluşu olarak kabul edilen M.Ö. 4.500 - 4000 'li yıllara kadar gıdanın temini, yaklaşık olarak bu gibi gıdayı ortak alanlarda depolama, ekim yerinin hayvanların evcilleştirilerek sürülmesi ile devam etmiş, ilk homo sapienslerden beri gelen toplayıcılık ise daha yakın ancak daha dar alanlarda sonraları da devam etmiştir. Tarım devriminin başlangıcı sayılan M.Ö. 9.500’li yıllar ile ilk şehirlerin ortaya çıktığı M.Ö. 4.500’li yılları arasındaki dönemde insanları zorlayan en önemli üretim sorunlarından bazıları; ani mevsim değişimleri ve küresel hava olayları olmuştur. Mevsimsel değişimleri öngöremeyen, hava olaylarını kestiremeyen ve başına gelenleri de yorumlayamayan insan her ne kadar bulunduğu coğrafya uygun gıda üretmeyi denese dahi mevcut bilgisinin yetersiz olması sonucu olarak meydana gelen olumsuz hava olayları sonrasında yerini değiştirmek zorunda kalmıştır. Ancak, yine de ilk yerleşimler incelendiğinde insanların her zaman yerleşim için verimli arazilerin yakınlarını seçtikleri anlaşılmaktadır. İnsanlar, olumsuz hava olaylarına uğrayıp yerleşimini terk etse de insan toplulukları bir zaman sonra yıkıma uğrayan bu yerleşim yerleri üzerinde tekrar yeni topluluklar kurabilmişlerdir. Bu durum özellikle ilk dönem insanların, doğaya olan büyük bağlılığı göstermektedir.

Tarım, kentlerin beslenmelerini desteklemek için çok büyük öneme sahip olduğundan kent yerleşmeleri ilk önce tarım ekonomisinin yoğun olduğu bu yerlerde meydana gelmiştir. Tarım alanlarının sulanması sonucu elde edilen artı ürün ve bu artı ürünü korumak için kentlerin etrafının surlarla çevrilmesi, surların kentlerin etrafına çevrilme iradesini ortaya koyan bir örgütsel yapının olduğunu göstermektedir (Işık, 2018). Sümer medeniyeti günümüzde çoğu tarihçi tarafından insanlığın üzerinde geliştirdiği ilk önemli medeniyetlerden biri olarak kabul edilmektedir. Sümerlerin, yaşadıkları Mezopotamya ve civarında kuruldukları coğrafyada, ilk yıllardan itibaren tarımı o zamanın şartlarına göre ileri seviyede yaptıkları anlaşılmaktadır. Gündelik hayatı gıda üretim bilgileri de dahil olmak üzere, M.Ö. 3200’lu yıllarda yazıyı geliştirmeleri sonrasında tabletlere işlemişlerdir. Tabletlerde tuttukları vergi kayıtları ve harcamalarında gıda konularına değinmeleri ile gıda bilgisi arşivi mirası da bırakmışlardır. Sümerler ile aynı dönemlerde gelişen Asya, Indus vadisi ve daha kuzey uygarlıklarının da aynı

veya benzer tarımsal ilerlemeler geliştirdikleri bilinmektedir. Bu gibi daha büyük yerleşmeler, karmaşık gıda sistemlerinin gelişmesini, üretimin kaydının tutulmasını, ilk vergilerin gıda üzerinden alınmasını, sulama ile ilgili sistemlerin gelişmelerini sağlamıştır. Sümer, Mısır, İran gibi dönemin daha verimli topraklar üzerinde kurulu medeniyetlerinde gıda üretimi gelişirken tarımdaki gelişmeler dünyanın her yerinde aynı düzeyde olmadığı bilinmektedir. Toplayıcılık ile daha geniş coğrafyalara gıda bulmak için yayılmak ihtiyacı, yakın zamana kadar devam etmiştir. Sümer ve dönemin diğer devletlerinin uzunca bir zaman varlıklarını devam ettirebilmiş olmalarında tarım sistemlerini geliştirmiş olmalarının payından sıklıkla bahsedilmektedir. Sümer kayıtları, Mezopotamya bölgesindeki verimli topraklarda kurulu zengin şehir devletlerinin dağlık bölgelerde ve çöllerde yaşayan göçebelere sürekli saldırı tehdidi altında olduğu yazılıdır. Saldırı ve savaşlar neticesinde bu vadileri işgal ederek vadi devletleri ve imparatorluklar kurmuş fakat imparatorluklar merkezi yönetim ve kontrol özelliği taşımamaktadır. Fethedilen şehirler işgalcilerin egemenlik haklarını kabul ederek haraç ödemekte, buna karşın mahalli yönetimde bağımsızlıklarını korumalarına izin verilmektedir. İstilalar ve savaşlar, Sümer hayat tarzını fazla etkilememiş, çünkü istilacılar kısa zamanda şehir uygarlığını benimsemişlerdir (Doğruyol, 2021; Güran, 2003; Childe, 1995).

Tarihin ilk yıllarında bazı başarılı olanları dışında medeniyetlerin ömürleri nispeten kısa olmuştur. Büyük imparatorlukların dağılması sonralarında küçük işbirlikleriyle kurulan toplumlar tarımın grupları yönetmenin önemini bilincine vardıklarından suya yakın, bereketli olarak görülen, ova yerlerde konuşlanmışlardır. Daha çok batıda yerleşik tarımı ana eksenine alarak kurulan toplumlar bulunmakta iken batıya kıyasen doğudaki toplumlar toplayıcı-avcı ağırlıklı olmuşlardır. Ancak hayvan evcilleştirmekte başarılı olmaları da bu düzende olmalarında etkili olmuştur. Batılı toplumların daha çok inek, koyun, keçi başta olmak üzere hayvanları tarımda kullanmak üzere evcilleştirip eğittikleri, doğudaki toplumların ise göçebe hayatın da etkisiyle başta at, köpek, kurt gibi hayvanları diğer coğrafyalara kolay taşınmak ve savaşmak amacıyla daha fazla eğittikleri anlaşılmaktadır. Roma İmparatorluğu'nun yıkılışı sonrası ve sonrası yaşanan gelişmeler, özellikle kavimler göçü nedeniyle Avrupa'ya taşınan nüfus çok büyük kentleri meydana getirmiştir. Bu yerleşimlerin beslenmesi için ekilmeyen arazilerin tarıma kazandırılması, toprak işlemede hayvan gücünden yararlanılması, o zamana kadar ekimi yapılmamış ürünlerin ekilmesi, ekim nöbeti yapılması gibi daha gelişmiş gıda çözümlerini gerektirmiştir (Direk, 2012). Avrupa'da geçmişi kimilerine göre yıkılan feodal beylikler sonrasında ortaya çıktığı döneme bir kısım için ise iki bin yıl öncelerine götürülebilen toplu tarım sistemi (açık tarım sistemi), 8. Yüzyılda artan nüfusun ihtiyaçlarının karşılanması için önemli bir çözüm sunmuştur. Bu sistemin temelinde, köylülere ait topraklar dağıtık haldedir ve dönüşümlü ekim, ortak otlatma usulü benimsenmiştir. Ortak toprakların bir bölümü nadasa bırakılır. Bu sistem, çok büyük gruplarda ve uzun yıllar devam etmiştir. Sonraları, toplumsal ilişkilerin karmaşık bir hal alması ve pazar ekonomisinin gelişmesi ile bu sistemin uygulanmasını zorlaştırmış, bu sistem yerini bireysel çiftçiliğe bırakmıştır.

M.Ö. 1000 ile M.S. 1.000 yılları arası döneme denk gelen dönemde yerleşik sayılabilecek batılı toplumlar ile daha az yerleşik olan doğulu toplumlar kıyaslandıklarında, tarımın rolü daha büyük nüfuslu yerleşmeleri, daha önemli bilimsel gelişmelerin ortaya çıkması vb. ilerlemelerde görülmektedir. Batının ilerleme çağı olan bu dönem, bugünkü pek çok bilim dalının temelini atılması ile, tarım ve gıda sistemlerinin gelişmesine katkı sağlamıştır. Bu dönem batıda zenginliğin oluşması, zengin kesimlerin ortaya çıkması, küçük köylü sınıfının ve büyük toprak sahiplerinin oluşması, birkaç yüzyıl sonra bunların soylu sınıf adına ülkeyi yönetmeleri ile doğuda ise büyük hükümdarlık varlığı ile devam etmiştir. Gelişmelerinde tarımın da önemli katkısı olan tarihteki büyük yerleşimlerin sadece hep iyi zamanları olmamıştır. Bu büyük yerleşimler hastalık, salgın, ganimet, çatışmalar ve savaşlar gibi büyük sorunlarla boğuşmuşlardır. Doğu ile Batının yıllarca uzak olması, birbirinden çeşitli alışkanlıklar ve beğeniler geliştirmiş olmaları ile farklılaşmaları ticarete zenginlikler getirmiştir. Doğuda üretimi yaygın ürünler dönemlerinin tüccarları ile daha çok batıya taşınmıştır. Tarihi çok daha

eskilere dayanan İpek Yolu ve Baharat Yolu gibi geçişlerde gıda ürünleri ve fikirler taşınmıştır. Doğudan sulama konusunda, batıdan ise gen kaynakları ile ilgili bilgiler transfer olmuştur (Kalan, 2014; Durukan, 2015; Koçak, 2019; Şahin, 2020).

14. yüzyılda batıda hastalıklar, savaşlar ve Avrupa coğrafyasının politik karmaşasında, halkın gıdasını düşünmek, bunun için üretim sahalarını yeniden tasarlamak, gıdaya müdahale etmek ve gıdanın temini için yeni arayışlara girmek daha çok kaçınılmaz olmuştur. 15. yüzyıla gelindiğinde gıda konusunda devam eden ilerlemeler özellikle coğrafi keşifler çağında oldukça artmıştır. Keşifler ile o zamana kadar bilinmeyen gıdalar bilinen dünyaya taşınmış, buralarda üretilmiş, o günün dünyasında bilinen yerlerde pazarlanmıştır. Aynı şekilde, bilgi, bitki örnekleri ve gıdanın kendisi taşınmıştır. Daha uzak bölgeler (yenidünya dedikleri Amerika kıtası ve diğer yeni coğrafyalar), özellikle batılılar için kendi topraklarında emek ve sermaye (ganimet) kaynağı olmuştur. Yenidünyadan gelen pek çok ürün; o zamana kadar varlığı bilinmeyen gıdalar, Uzak Doğudan gelen bol ve yeni ürünler batılıların kentlerinde insan sağlığına uygun ve besleyici gıdalar olarak hastalıklara ve salgılara direnç oluşturmuştur. Coğrafi keşiflerin bir diğer önemli etkisi, 16. yüzyıla gelindiğinde deniz aşırı ve döneme göre oldukça büyük şirketlerin ortaya çıkması olmuştur. Bu şirketler dönemin tarım ve gıda ürünlerinin ticaretinde büyük rol edinmişlerdir. Batının bu dönemde güçlenmesi, ilk şirketlerde gıdanın payı, bunun en önemli nedenlerinden sayılabilir. Batılılar, dolaylı olarak gıdanın medeniyetlerine ilk kez ciddi faydalarını gördükleri M.Ö. 600'lü yıllarda gelişen bilim adımlarını, M.S. 600'lü yıllarda İslam'ın yükselişi ile Müslümanlarının bilimi ileri taşımalarından faydalanarak daha kalıcı olarak hayatın içerisine alarak geliştirmişlerdir. Keşifler, ilk şirketler ve bunların medeniyete etkilerine bilimi ekleyerek başta gıdada olmak üzere insanlık tarihinde önemli bir gelişim kaydedilmiştir. Gıda üretiminde dönemin bilimsel gelişmeleriyle beraber hızlı gelişmeler yaşanmış; daha büyük kentlerin oluşturulması, dış ticaretin ileri seviyelere getirilmesiyle daha büyük sermayeler gıda piyasalarında yer almıştır. Görüldüğü üzere, bir kez daha gıda temelli gelişmeler doğrudan ya da dolaylı olarak insanlığa büyük bir basamak daha atlatmıştır.

19. yüzyıla gelindiğinde yeni kıta Amerika'nın gelişmesi, üzerindeki devletlerin yüzyılın başlarında birer birer bağımsızlıklarını elde etmeleri ve artık yavaş yavaş günümüzde de en güçlü ülke olduğu varsayılan Amerika Birleşik Devletleri (ABD)'nin bu kıtada yer alması gıda sistemlerini çokça karmaşıklaştırmıştır. Coğrafi keşiflere kadar Avrupa ve Asya kıtalarında, kısmen bir parça da Afrika kıtasında devam eden gıda ticareti Kuzey Amerika ve Güney Amerika ile sonraki yıllarda Avustralya kıtasına da taşınmıştır. 1800'lü yıllardan sonrası için, geçmiş bin yıllardır yapılan üretimden çok daha fazlasının gerçekleştirildiği söylenebilir. İnsanlık tarihi boyunca bilinen coğrafyada devam eden gıda ilişkileri, Amerika kıtasının ekonomik yönden gelişmesiyle daha büyük alanlara ve çok daha hızlı bir halde taşınmıştır. Bu şekilde bir gıda transferini, keşfedilen ve bilinen kıtalarla daha sıkı gıda ilişkileri geliştiren Avusturya kıtası ve dünyanın diğer yerleri de izlemiştir. Bu dönemlerde, insanlığın gıda birikiminin katlanarak ilerlediği söylenebilir. 1760-1840 yılları arasında İngiltere merkezli olarak başlayan sanayi devrimi, kurulan fabrikalara işçi bulunması için çiftçilerin şehirlere göç etmeleriyle büyük şehirleri meydana getirmiş ancak kırsalda üretim yapacak insan sayısını azaltmıştır. Bunu takip eden yıllarda gıda üretimi için makine kullanımı, daha karmaşık planlar yapılması, emperyalizm gibi konular gıda arz fazlalarının oluşması, gıda fazlalarının eritilmesinin istenmesi, kırdan uzaklaşan insanların üretimden uzaklaşmaları ile daha çok başkasının üretmesine bağlı olmaları gıda sisteminin dönüşmesini zorunlu kılmıştır. Teknolojik gelişmelerin gıda üretimine sıçramasıyla ilk yıllarda üretim katlanarak artmış, oluşan arz fazlalarının değerlendirilmesi söz konusu olmuştur. 1840 yılında yapay gübrenin icat edilmesi beraberinde getirdiği verim artışı ve toprakların yeni yöntemler ile işlenmesi sonucunda, tarımsal faaliyetlerin pazar payı artmış ve tarım sektörü ekonomik bir güç olmaya başlamıştır (Doğan vd., 2015). Daha sonraları gelişmiş ülkelerin az gelişmiş ülkelere uygulayacakları gıda emperyalizminin temelleri böylece atılmış olmuştur.

20. yüzyılın başında ABD’de yaşanan büyük buhran ABD’de kırsal kesimin üretememesine ve küçük çiftçilerin üretim sahalarını kaybetmesine neden olmuştur. Buhranın etkisi ve sanayi devrimi sonrası ilk kez görülen sorunlar bütün dünyaya yayılmış ve küresel çapta gıdanın yoksulluğunun gıda krizlerine neden olabileceği ilk kez anlaşılmıştır. II. Dünya savaşından kazanarak çıkan ABD, ilk önemli süper güç olmuştur. Savaş sonraları kurulan, Avrupa Birliği (AB) ve diğer uluslararası işbirliklerinde tarım ve gıda konusu en önemli konulardan birisi olmuştur. Yüzyılın ortalarına gelmeden, gıdaya egemen olan milletler diğer ülkeler üzerinde hakim güçler olmuştur. ABD’de üretim fazlalarının görüldüğü “Yeşil Devrim” olarak adlandırılan dönemde, emperyalist ülkelerin gıda fazlası olurken daha fazla ülkede fakir kesimler gıda güvencesizliğinden muzdarip ve gıda yardımlarına bağımlı olmuşlardır. ABD Dışişleri eski bakanı Henry Kissinger bu konuda “Petrolü kontrol edersen ulusları kontrol edersin, yiyeceği kontrol edersen insanları kontrol edersin” ifadesini kullanmıştır. Bu ifade, günümüzde geçerliliğini korumakta ve içerdiği düşünce daha iyi anlaşılmaktadır. Son yıllarda güçlü ülkelerin gıda emperyalizmini tüm dünyaya yayma peşinde olmalarının olumsuz etkilerini yaşamış ve zamanla anlamış olmalı ki gelişmekte olan ülkeler de kendilerinin gıda arzını güvende tutmayı önemsemişlerdir. Bunun yanında, özellikle 1970’lerden sonra artan kentleşme, çevre kirliliği, gelir dağılımı eşitsizliği vb. konular ekonomik, çevresel ve kültürel konularda sürdürülebilirlik konusunu gündeme getirmiştir. Birbirinden farklı tanımları olsa da, bilimsel anlamda sürdürülebilirlik anlayışının ortak noktası olarak sürdürülebilirlik kavramı, doğal ve beşeri kaynakların ölçülü bir şekilde kullanılmasını hedefleyen süreçleri ifade etmektedir (Tıraş, 2002).

Sürdürülebilirlik kavramının 19. yüzyıl başlarında literatürde somut olarak kendini göstermeye başladığı belirtilmektedir (Bozdoğan, 2010). Ekonomik faaliyetlerin gerçekleştirilirken gelecek nesillerin hesaplanması, kaynakların korunması ve doğaya zarar verilmemesi hedeflenen düşüncenin küresel bir bilinç haline getirilmesi; devletlerin, işletmelerin ve toplumların sürdürülebilirlik temelli politikaları takip etmeleri geçmişten günümüze düzenlenmiş olan uluslararası toplantılar ve çalışmalar sonucunda olmuştur (Paker, 2018). Bu anlamda, çevrenin korunması ve geliştirilmesi konularının ilk defa ele alındığı platform olması özelliği ile 1972’de gerçekleştirilen Stockholm Çevre ve İnsan Konferansı, önemlidir (BM, 1972; Günal, 2019). Günümüzde kullanılan en geçerli sürdürülebilir kalkınma tanımı, 1987 yılında yayımlanan Ortak Geleceğimiz (Brundtland) Raporu ile yapılmıştır. Tanıma göre; “*sürdürülebilirlik, gelecek kuşakların kendi ihtiyaçlarını karşılama gücünü tehlikeye atmadan günümüz kuşağının ihtiyaçlarının karşılanmasıdır*”. Bu toplantıda sürdürülebilir kalkınmanın ana amaçları; büyümeyi canlandırmak, büyümenin şeklini değiştirmek, iş, yiyecek, enerji, su ve sağlık hizmetleri için gerekli temel ihtiyaçların karşılanması, sürdürülebilir bir nüfus düzeyi sağlamak, kaynak tabanını korumak ve arttırmak, teknolojiyi yeniden yönlendirmek ve riski yönetmek, karar sürecinde çevre ve ekonomiyi birleştirmek olarak belirlenmiştir (WCED, 1987). Bu tarihten günümüze çok sayıda toplantı yapılmış ve politikalar değişikliğe uğrayarak günün ihtiyaçlarına göre biçimlendirilmiştir. Sürdürülebilirlik çabalarının gıdaya yansması ise, gıda güvencesi sağlanırken ekonomik, sosyal ve çevre boyutuyla gıdanın etkilerinin azaltılması olarak belirlenmiş, gıda politikalarının sürdürülebilirliği bu şekilde ele alınmıştır.

GIDA GÜVENCESİ

Gıda Güvencesi Kavramı

1948 tarihli Birleşmiş Milletler (BM) İnsan Hakları Evrensel Beyanname’si, “*Gıdaya ulaşım hakkını temel insan hakkı*” olarak tanımlamıştır. Bu tarihten başlayarak insanın gıdası savunulan bir durum olmuştur. Daha sonraları 1966 yılında Ekonomik, Sosyal ve Kültürel Haklar Konvansiyonu ile yasal koruma altına alınan gıda hakkı, insan haysiyeti kavramından kaynaklanan temel bir hak olarak görülmüş, uluslararası sözleşmelere konu olmuştur. Devletler, vatandaşlarının gıda hakkını sağlamak için gıda güvencesi oluşturma çabasıdadır. Gıda hakkı,

bireyin sahip olduğu diğer insan hakları ile doğrudan ve dolaylı olarak güçlü bir ilişki içerisindedir. İnsan için bir varoluş meselesi olan gıda hakkı toplumların ve devletlerin ortaya çıkışında önemli rol oynamıştır (Tekir, 2016). İnsana, gıdasının güvence edilmesi bir hak olarak sağlanmalıdır. Gıda güvencesi, ilk zamanlarda “*gıda tüketiminin istikrarlı biçimde sürdürülebilmesi, üretim ve fiyatlardaki dalgalanmaların dengelenebilmesi için temel besin maddeleri bakımından yeterli dünya arzının her zaman var olması*” olarak tanımlanmıştır (BM, 1975).

Dünya Gıda Zirvesi-1996 raporunda (FAO, 1996) gıda güvencesi, “*bütün insanların her zaman aktif ve sağlıklı yaşamı için gerekli olan besin ihtiyaçlarını ve gıda önceliklerini karşılayabilmek amacıyla yeterli, sağlıklı, güvenilir ve besleyici gıdaya fiziksel ve ekonomik bakımdan sürekli erişebilmeleri*” olarak tarif edilmiştir. Gıda güvencesinin temel bileşenleri; “*yeterlilik, faydalanılabilirlik, erişilebilirlik ve istikrar*” üzerinde uzlaşmış temel boyutlar olmuş, ardından bunlar güncellenerek günümüzde anlaşılacağı halini almıştır (World Bank, 2023). *Yeterlilik bileşeni*, gıda arz güvenliğini kapsayacak şekilde gıdanın ihtiyaç duyulduğunda fiziksel olarak bulunabilmesini (*mevcudiyeti*) ifade etmektedir. *Faydalanılabilirlik bileşeni*, gıdanın güvenilir olması ve besinsel değerinin iyi olmasını tanımlamakta kullanılmaktadır. *Erişilebilirlik bileşeni*, güvenilir gıdadan ekonomik ve fiziksel olarak yararlanabilmeyi ifade etmektedir. Son olarak gıdanın *sürdürülebilirlik bileşeni*, gıda güvencesinin yeterlilik, faydalanılabilirlik ve erişilebilirlik koşullarının istikrarlı bir şekilde sürekliliğinin sağlanabilmesidir (FAO, 2010; World Bank (2023) .

Gıda Güvencesinin Zorunluluğu

Gıda hakkı kavramı, “*bireyin sağlıklı ve aktif bir hayat yaşaması için ihtiyaç duyduğu bütün besin öğelerine ve bunlara ulaşım araçlarına sahip olması*” olarak tanımlanmıştır (FAO ve OHCHR, 2010). Gıda herkes için olmalı, alınabilir olmalı ve ihtiyaç olduğunda bulunabilmelidir. Bireye hak olarak tanımlanan gıda güvencesinin “*yeterli, sağlıklı, güvenli ve besleyici gıdaya sürekli fiziksel ve ekonomik erişim*” olarak günümüzde anlaşıldığı görülmektedir (Keskin ve Demirbaş, 2022). Gıda güvencesi herkesin gıdaya tam, sağlıklı, uygun fiyatlı, süresiz şekilde hesap edildiğinde daha yapılması gereken çok fazla şey olduğu görülebilecektir. Günümüzde 700 milyona yakın insan her gün açlıkla mücadele etmektedir (FAO, IFAD, UNICEF, WFP and WHO, 2023).

Milyonlarca insan “*beslenme açısından yeterli ve güvenli gıdaların sınırlı veya belirsiz mevcudiyeti veya sosyal olarak kabul edilebilir şekillerde kabul edilebilir yiyecekler elde etme konusunda sınırlı veya belirsiz bir kabiliyet durumu*” olarak da tanımlanabilen gıda güvencesizliği sorununu derinden yaşamaktadır (USDA, 2000). 1.750 milyon insan fazla kilo sorunu yaşamaktadır. 800 milyon insan obezite ile mücadele etmektedir. 22.500 kişi ise açlıktan ölmektedir (Worldometer, 2024). Bir yandan insanlar açlık çekmekte gıda bulamamakta, çoğunlukla gıdaya erişememektedir. Bir tarafta ise tüketiciye bağlı yanlış beslenmek, gıdayı eritememek sebeplerinden yanlış beslenme sorunları yaşanmaktadır. Açlık gibi bilinçsiz beslenmek, yeterince beslenememek, obezlik de gıda güvencesi sorunudur.

Ulusal mevzuatlar ve uluslararası antlaşmalarda tanımlanan insani nedenler, gıda hakkının kutsallığını sayan dini nedenler, yaşadığımız yüzyılın varlığı ve yoksulluğu birlikte bulundurması hesap edildiğinde vicdani ve toplumsal nedenler ile uluslararası antlaşmalar ve devlet olmanın idareye verdiği sorumluluktan dolayı her kesimden vatandaşların gıda güvencesinin sağlanması gerekmektedir.

Devletler, vatandaşlarının gıda hakkını gözeterek bunu sağlamaya yönelik düzenlemeleri ile gıda hakkı ve gıda güvencesinin sağlanmasından ve sürdürülebilirliğinin korunmasından sorumludurlar. Anayasa, kanun ve diğer mevzuat düzenlemeleri ulusal düzeyde, bölgesel işbirlikleri ile ülkeler arasında antlaşmalarla ve daha üst insan haklarıyla ilgilenen uluslararası

hukuki metinlerde ve içtihatlarda gıda hakkı ve güvencesinin tanımı ve kapsamı, devletlerin ve diğer unsurların bu konudaki görev ve sorumlulukları tanımlanmıştır. Devletlerin anayasaları ile garanti edilmeye çalışılan gıda hakkı, yine ülkelerin bağlı oldukları uluslararası kuruluşlar ve uluslararası antlaşmalar ile de doğrudan veya dolaylı olarak gıda güvencesinin korunması çabalarına konu olmuştur (Peerzada, 2019).

Gıda güvencesinin bir hak olduğu ve bunun karşılanması ile ilgili sorunlar olduğu ülkeler ve uluslararası kurullar tarafından görülmüştür. Ülkeler, genelde Tarım Bakanlıkları ve diğer kuruluşları vb. ulusal çapta gıda güvencesini temin etmek için çalışmaktadır. BM Tarım ve Gıda Örgütü (FAO), Uluslararası Tarımsal Kalkınma Fonu (IFAD), Ekonomik Kalkınma ve İş Birliği Örgütü (OECD), Dünya Gıda Programı (WFP), Uluslararası Gıda Politikaları Araştırma Enstitüsü (IPFRI), Dünya Bankası (World Bank), Economic – Impact (EIT) gibi uluslararası kuruluşlar gıda güvencesi için çalışmaktadır. ABD Hastalık Kontrol ve Korunma Merkezleri (CDC), Avrupa İmar ve Kalkınma Bankası vb. kuruluşlar da, bölgesel ve uluslararası özellikte çalışmalar yapmakta, raporlar sunmaktadır.

Gıda Güvencesini Sağlamaya Yönelik Çalışmalar

Gıda güvencesi, ekonomik krizler, savaşlar, kıtlık ve kuraklık, küresel iklim olayları, plan/program/strateji/hesaplama hataları, uygulama hataları, vb. pek çok nedenden dolayı tehdit edilmektedir. Ancak en çok ve uzun süreli etkileyen nedenler işsizlik, gelir eşitsizliği vb. ekonomik nedenlerdir. Gıda güvencesi; insani, dini, vicdani ve toplumsal nedenleri ile devletlerin sorumluluğundadır. Ülkeler, günümüz sorunlarını çözmek için ortak amaçlar oluşturmak, ülkelerinde yapılabilecekleri belirlemek ve küresel bir mücadele vermek istemektedir. Bu anlamda Binyıl Kalkınma Hedefleri (BKH); Birleşmiş Milletler tarafından 2000 yılında ilan edilmiş, uluslararası kalkınmaya katkıda bulunmayı amaçlayan 193 ülke ve 23 uluslararası şirketin katkı sözü verdiği hedeflerdir. Bu hedefler;

- (1) Aşırı yoksulluğu ve açlığı ortadan kaldırmak,
- (2) Herkes için evrensel ilköğretim sağlamak,
- (3) Cinsiyet eşitliğini teşvik etmek ve kadınların güçlendirilmesini sağlamak,
- (4) Çocuk ölümlerini azaltmak,
- (5) Anne sağlığını iyileştirmek,
- (6) HIV/AIDS, sıtma ve diğer hastalıklarla mücadele,
- (7) Çevresel sürdürülebilirliğin sağlanması ve
- (8) Küresel ortaklık,

başlıklarında iyileştirmeler yapılmasını gözetiyordu. *Binyıl* ya da bir diğer ifadeyle *Milenyum Kalkınma Hedefleri*'nin süresi 2015 itibarıyla tamamlanmıştır.

BM üyesi 193 ülke tarafından yoksulluğu ortadan kaldırmak, dünya gezegenini korumak ve tüm insanların barış ve refah içinde yaşamasını sağlamak üzere 2030 yılı sonuna kadar ulaşılması için birbiri ile bağlantılı 17 amaç Ocak 2016'da yürürlüğe girmiştir. Sürdürülebilir Kalkınma Amaçları (SKA), diğer bir deyişle Küresel Amaçlar, Binyıl Kalkınma Hedefleri'nin ilerlemeleri üzerine inşa edilmekle birlikte iklim değişikliği, ekonomik eşitsizlik, yenilikçilik, sürdürülebilir tüketim, barış ve adalet gibi yeni alanları içermektedir. BM, 2030 SKA Gündemi'nin hedeflerine ulaşmak için gıda sistemlerini bütüncül ve koordineli bir yaklaşımla sürdürülebilir bir şekilde ele almak istemektedir. Sürdürülebilir Kalkınma Amaçları'nın süresi 2030 yılında dolacağı için 2020-2030 yılları arası "Eylem Onyılı" olarak adlandırılmaktadır. Bu amaçlara ulaşılmasında, (2) açlığa son ve (12) sorumlu tüketim ve üretim, (13) iklim eylemi gibi doğrudan ya da dolaylı olarak herkesin gıdaya ulaştırılmasını gerektiren amaçlar bulunmaktadır.

Bunlara çözüm bulmak amacıyla Gıda Sistemleri Zirvesi, 2030 Gündemi'nin hedefleri genelinde benimsenecek bir gıda sistemleri yaklaşımının ortak faydalarını en üst düzeye

çıkarmak ve iklim değişikliğinin zorluklarını karşılamak amacıyla BM tarafından toplanmıştır. Zirvede;

- (1) Herkes için besleyici ve güvenilir gıdaya erişiminin sağlanması,
- (2) Sürdürülebilir tüketim modeline geçilmesi,
- (3) Yeterli ölçekte doğaya uyumlu üretimin artırılması,
- (4) Adil geçim kaynaklarının geliştirilmesi,
- (5) Hassasiyetler, şoklar ve streslere karşı dayanıklılık sağlanması,

konu başlıklarında 5 ana eylem alanı belirlenmiştir. Türkiye Cumhuriyeti de bu eylem planlarını esas alarak ulusal yol haritasını belirlemiştir (TOB, 2021).

Gıda Güvencesinin Durumu ve Ölçülmesi

Gıda güvencesinin günümüzde üzerinde uzlaşılan dört bileşeninde de sorunlar bulunmaktadır. Bu sorunların belirlenmesi, analiz edilmesi, mevcut durumun görülmesi ve ülkeler arasında karşılaştırılma yapılması sorunların çözümünü sağlaması için amaçlanmaktadır. Uluslararası işbirliklerinin ve ülkelerin gıda güvencesi konusunda çalışan kuruluşlarının dönemlik raporlarıyla gıda güvencesi konusunda güncel gelişmelerden daha iyi bilgi edinilebilmektedir. Gıda sisteminde yer alan üreticiler, aracılar, tarıma ve gıdaya dayalı sanayi, gıda sektörü, tüketiciler, devletler ve uluslararası kuruluşlar/yükümlülükler sorunların iyileştirilmesinde bunlardan faydalanmaktadırlar (Koç vd., 2018).

Gıda güvencesine yönelik uluslar üstü çabalar, sorunun görülmesi, farkındalığın oluşması ve çözüm için işbirlikleri geliştirilmesi için oldukça faydalıdır. Ülkelerin bundan kendilerine özel sorunları anlamaları ve çözüm üretmeleri sağlanmalıdır. Küresel ölçekte gıda güvencesinin ölçülmesi ve ülkelerin gıda güvenliği ve güvencesini hangi faktörlerin etkilediğiyle ilgili olarak gıda güvence göstergelerini hesaplayan bazı kurumlar bulunmaktadır. Ancak genelde üç kurumun gıda güvencesi raporları hem kullanılan metotlar hem verdikleri önerilerle diğerlerinden daha dikkatle değerlendirilmektedir. Bunlar FAO, Uluslararası Gıda Politikaları Araştırma Enstitüsü (IFPRI), Ekonomik İstihbarat Birimi (EIU)'dir. Sayılan organizasyonların yayınlanan geçmiş dönem raporları, gıda güvencesinde güncel durum hakkında bilgiler vermektedir.

The State of Food Security and Nutrition in the World (SOFI), FAO tarafından 1999'dan beri güncellenen göstergelerle dünyanın farklı bölgelerini, farklı ülkelerini, bazı yıllar birbirinden farklı tür konularda kıyaslayabilen yıllık olarak yayınlanmaktadır. 2023 yılında yayımlanan rapor, FAO, IFAD, UNICEF, WFP and WHO tarafından ortaklaşa hazırlanmıştır. Önceki dönem SOFI raporunda kullanılan gıda güvencesi göstergeleri; availability (*bulunabilirlik*), access (*erişilebilirlik*), utilization (*kullanılabilirlik*) ve stability (*kararlılık/istikrar*) boyutları ve bunlara ait raporun metodoloji kısmında açıklanan göstergeler olmuştur. SOFI (2023) raporunda gıda güvencesinin daha iyi anlaşılmasını sağlaması düşünülen, ancak üzerinde yaygın olarak uzlaşa sağlanmasa da High Level Panel of Experts (HLPE) of the Committee on World Food Security (CFS) tarafından iki ek boyuta daha vurgu yapılmıştır. Bunlar (a) *Agency*; bireylerin veya grupların ne yediklerini, nasıl ürettiklerini, gıda sistemleriyle gıda üretme, süreç ve sistemlerini yönetmeleri ile gıda sistemlerini ve gıda politikalarını daha iyi nasıl iyileştirebilecekleri boyutudur. (b) *Sustainability*; gıda güvencesi ve beslenmenin geleceği için ekonomik, çevresel ve sosyal konularda temellendirilmiş uzun vadeli becerileri ifade etmektedir (HLPE, 2020).

SOFI (2023) raporunda küresel düzeyde gıda güvencesi konusunda bahsedilen en önemli konular; 2022 yılında bir önceki yılı göre küresel açlık düzeyinin önemli oranda daha da kötüleşmediğidir. Fakat COVID-19 pandemisine bağlı olarak ya da artan gıda fiyatlarına, artan üretici maliyetlerine ve iklim olaylarına bağlı olarak çok yerde arttığı belirtilmiştir. Genel olarak bakıldığında çocukların gıda güvencesi ile ilgili olumlu durumlar olduğu, buna karşın

kırsaldaki kadınlar için olumsuz durumlar olduğu belirtilmiştir. Yalnız bazı bölgeler için aşırı beslenme durumlarında artışlardan da söz edilebilir. Özellikle kentleşmeye gıda sistemlerini hızla dönüştürdüğü üzerinde durularak 2050 yılına kadar artması da hesap edildiğinde, daha fazla hesaplanması gereken bir konu olarak dikkat çekilmiştir. Hızlı kentleşmeye bağlı açlık, gıda güvensizliği ve yetersiz beslenme sorunlarına önlem alınması gerektiği vurgulanmıştır. Raporda, kır-kent dengesi ve dengesizliği vurgulanmış, doğrudan olmasa da kent tarımına önem verilmesi ve bu konuda bir şeyler yapılması önerilmiştir. Hükümetlere güncel gıda sisteminin sorunlarını görerek ve kentleşme konusunu da hesap ederek gıda sistemlerini dönüştürmesi ve yerel yönetimleri temel gıda sistemlerini şekillendirmede daha aktif rol almaları önerilmiştir.

Bir sonraki yıl SOFI (2024) raporunda ise bu sefer, sürdürülebilir gıda sistemleri için dönüştürücü finansman mekanizmalarının gerekliliği pek çok kez vurgulanmıştır. Küresel gıda sisteminde adil ve eşit bir dönüşüm sağlanmadan açlığı ve yetersiz beslenmeyi sona erdirmenin mümkün olmadığı altı çizilmiştir. Gıda maliyetlerinin ve gıda fiyatlarının artmakta olduğundan bahsedilerek gıdaya erişim konusunda özellikle kırılğan toplulukların karşılaştığı zorluklara dikkat çekilmiştir. Açlık kadar yetersiz ve dengesiz beslenmenin de çok önemli bir sorun oluşturduğu ve bir an evvel azaltılmasının gerektiği ifade edilmiştir. Gıda güvenliğinin temini için sadece kamu yatırımlarının değil, özel sektörün de aktif bir rol üstlenmesinin önemine atıflar yapılmıştır. Küresel gıda güvenliğini sağlama çabalarında sürdürülebilir başarı elde etmek için kamu-özel sektör iş birliğinin artırılmasının önemine dikkat çekilmiştir. Özel sektörün sermaye, istihdam, teknoloji ve pazarlara erişim konularındaki katkılarının, gıda sistemlerini güçlendirebileceği belirtilmektedir. Gıda güvencesinin ve beslenmenin finansmanı gibi, yeni bir kavramdan söz edilmektedir. Bununla, finansal kaynakların beslenme ve açlık sorununu iyileştirmesi hedeflenmektedir. Raporun sonuçları, küresel çapta başta iklim değişikliği ve ekonomik krizlerin tarımsal üretim üzerindeki olumsuz etkilerinin azaltılması ile daha etkin ve sürdürülebilir politikalar geliştirilmesi gerektiğini de göstermektedir (FAO, IFAD, UNICEF, WFP and WHO, 2024).

Global Food Policy Report (2023): Uluslararası Gıda Politikaları Araştırma Enstitüsü (IFPRI), tarafından yayımlanan raporda COVID-19 pandemisi şoku, savaşlar, doğal felaketlerle, artan gıda fiyatları gibi durumların gıda sistemlerini dönüştürdüğü ve gıda sistemlerinin yeniden gözden geçirilmesinin önemi vurgulanmıştır. Bu zamanların ileride görülmesi daha muhtemel bu durumların etkisinin azaltılması için politikalar geliştirilmesiyle geçirilmesi önerilmiştir. Raporda, daha çok yeni gelişen gıda sistemlerine odaklanılmış ve öneriler verilmiştir. Bunlardan bazıları; gıda sistemlerinin yapısal sorunlarının çözülmesi ve erken uyarı sistemlerinin geliştirilmesi ile en azından geçmişte yaşanan büyük şokların olumsuz etkilerini en aza indirmek önerilmiştir. Krizler tespit edilir edilmez ilk olarak büyük insani yardımların devreye alınması önerilmiştir.

Krizlerde gıda değer zincirlerinde özellikle alt basamaklarda olanların etkilendiklerine vurgu yapılmıştır. Bu sebeple hükümetlerin ilk başlarda gıda sistemlerini korumak için bu dezavantajlı grupları finansal olarak desteklemeleri önerilmiştir. Dezavantajlı grupların ve iklim dezavantajlı durumları güçlendirilmesi krizlerin etkisini azaltabilir ve gıda sistemlerini koruyabilir denilmektedir. Raporda, özellikle dezavantajlı kadınların desteklenmesine çokça vurgu yapılmış, kadınların ekonomik yönden güçlendirilmesinin gerektiği belirtilmiştir. Küresel ölçekte dünyada varlığı devam eden zorunlu göç durumunun özellikle üretimde değerlendirilmesi ve bu durumun olumlu yanlarının değerlendirilmesi tavsiye edilmiştir. Gıda sistemlerinin dönüşümü için özel sektör yatırımlarının desteklenmesi önerilmiştir. Yatırım fonlarının ve tarımsal desteklemelerin daha iyi kullanılması önerilmiştir. Raporda, en önemli konular olarak bunlar görülmüş, bölgesel olarak da gıda güvencesi konusunda yapılabileceklerin farklılıklarına değinilmiş ve bunlar için de öneriler sunulmuştur (IFPRI, 2023).

Bir sonraki yıl IFPRI (2024) tarafından hazırlanan *Global Food Policy Report (2024)*'unda da benzer sorunların devam ettiği görülmektedir. Açlık, gıda güvencesizliği ve yetersiz

beslenmenin, her türlü yetersiz beslenme ve diyetle ilişkili bulaşıcı olmayan hastalıklar dahil olmak üzere birçok kritik halk sağlığı sorununun temelini oluşturduğu belirtilmiştir. Sayılan sağlık sorunlarının dünya genelinde insanların refahı ve verimliliği için kısa ve uzun vadeli etkileri vurgulanarak açıklanmıştır. Düşük ve orta gelirli ülkelerde, açlık ve yetersiz beslenmenin düşürülmesindeki gelişmelerin yavaşlamasına karşın, aşırı kilo ve obezitenin dünya çapında hızla arttığı belirtilmiştir. Günümüzde birçok ülkenin birey, hane ve toplum düzeyinde çift yönlü olarak sorunları bulunmaktadır. Bir yandan yetersiz beslenme ve mikro besin eksikliği sorunu çekilirken, diğer yandan ise obezite ve yaşam boyunca diyetle ilişkili bulaşıcı olmayan hastalıkların bir arada bulunduğu savunulmaktadır. Bir diğer konu da, iklim değişikliğinin, gıdaların tedariki ve besin içeriği de dahil olmak üzere gıda sistemlerine yeni zorluklar getirdiğidir.

IFPRI (2024) ile diyet ve beslenme müdahalelerine daha bütüncül bir yaklaşım sağlamak için bazı öneriler getirilmiştir. Bunlardan ilki, beslenme için yönlendirilen düzenleyici/yatırım politikalarına ayrılan mali kaynakların etkilerinin ekonomik analizleri, kapasite ve politik ekonomi yönünden sürdürülebilir ve ölçeklenebilir olmasını sağlamak için yönetim değerlendirmeleri yapılmalıdır. Beslenme için basitçe ikinci en iyi seçenek, uygulanabilirlik açısından en iyi görülen birinci seçenektен daha faydalı olabilir. İkinci olarak, ilk en iyi uygulanabilir politika seçenekleri kümesini genişletmek için, bağlayıcı yönetim kısıtlarının tespit edilmesi ve iyileştirilmeleri gerekir. Üçüncü olarak, bilgiye erişimin iyileştirilmesi ve eşitlikçi hale getirilmesi, gıda sistemleri yönetişiminin iyileştirilmesinin temelindedir. Zaman içinde vatandaşlara karşı hesap verebilirliği artırmak için, kurumsal gıda endüstrisinin siyasi etkisinin karnesini tutmak; hükümetin beslenme hedeflerine yönelik taahhütlerini izlemek ve beslenme politikası yatırımlarının bütçelerini takip etmek gibi şeffaflık araçlarını geliştirme fırsatları vardır. Dördüncü olarak, tabandan gelen hareketler gıdanın durumunu katılımcı yolla yeniden şekillendirilmesinde kilit bir rol oynamakta ve vatandaşların dönüştürücü potansiyelini ortaya koymaktadır. Bu hareketlerin yeterli sivil alan ve kararlı siyasi liderlik mevcut olduğunda gelişmesi pek muhtemeldir. Çok paydaşlı platformlarda ve gıda politikası konseylerinde büyümeleri ve somutlaşmaları, ne zaman, neden ve nasıl daha iyi beslenme politikası kararlarına yol açtıklarını belirlemek için benzersiz bir doğal deney sağlar ve böylece giderek daha popüler hale gelen bu yöntemler hakkında daha fazla kümülatif öğrenme için temel sağlar (IFPRI, 2024).

Global Food Security Index (GFSI): The Economist Intelligence Unit (EIU) tarafından “Küresel Gıda Güvenlik Endeksi” ilk olarak 2012 yılında ortaya çıkarılmıştır. Oluşturulan endeks ile, dünyadaki gıda sistemlerinin incelenerek gıda güvencesizliğinin temel nedenleri ve affordability (*bulunabilirlik*), availability (*erişilebilirlik*), quality-safety (*kalite-güvenlik*) ve sustainability-adaptation (*sürdürülebilirlik-adaptasyon*) boyutları itibarıyla gıda güvencesizliğinden etkilenebilirliği düşük ve yüksek olan ülkelerin incelenmesi, politika önerileri verilmesi hesaplanmıştır. Endeks ile gıda güvencesinin boyutları da kendi içlerinde bazı göstergelerle önceden belirlenen metodla puanlanmaktadır. En sonunda bu 4 boyut birlikte değerlendirilerek ülkenin veya bölgenin gıda güvencesi konusunda ortalama puanına ulaşılmaktadır (Economist Impact, 2023).

GFSI’yi diğer hesaplama yöntemlerinden farklı kılan yönleri, gıda güvencesinin ölçülmesini açlıktan farklı bir şekilde ele alması, güvencesizliğin altında yatan faktörleri belirlemeye en yakın olması ve hükümetlerin politikalarını da gösterge olarak kabul etmesidir. Hükümetler, endeksi politikaların gözden geçirilmesi ve yatırımların belirlenmesi amacıyla, sivil toplum kuruluşları ve uluslararası örgütler ise odaklanılacak ülkelerin belirlenmesi için bir araştırma aracı olarak kullanmaktadır. Özel sektör ise, ülkelerdeki gıda tüketim trendlerini keşfetmek ve sosyal sorumluluk projelerini geliştirmek amacıyla endeksten yararlanmaktadır (Koç vd., 2018).

GFSI (2022) raporunda 113 ülke, Asya Pasifik, Avrupa, Latin Amerika, Güney Amerika, Orta Doğu ve Afrika ve Kuzey Amerika bölgeleri kendi aralarında incelenmiştir. 2019 ve 2022

yılları raporları kıyaslandığında gıda fiyatlarının artışı, COVID-19 pandemisi ve Ukrayna savaşı gibi nedenlerle endeksin genel satın alınabilirlik puanının %4 aşağı düştüğü görülmektedir. Bu küresel ölçekte gıdaya erişimde güçlükleri ortaya koymaktadır. Yine sosyal ve politika engeller gıdanın bulunabilirliğini düşürmüştür. Bu durumun kronik gıda yardımlarına olan bağımlılığı artırdığı vurgulanmıştır (Economist Impact, 2023). GFSI, çok fazla ülkelerin birbirleriyle karşılaştırılabilmesi, ülkelerin geçmiş yıllarıyla kıyas yapılabilmesi, gıda güvencesi bileşenlerinin anlaşılır göstergelerle her bir ülke için ayrı ayrı puanlanabiliyor olarak açıklanması sebebiyle kolay anlaşılabilir.

Gıda güvencesi indexleri, gıda güvencesi konusunda hazırlanmış bilimsel çalışmalar, deneyimlerle ortaya çıkmış diğer yayınlar, Türkiye'nin gıda güvencesindeki yeri, tehditler, geliştirilebilecekler konusunda imkan sağlamaktadır. Bu çalışmalardan yola çıkarak, Türkiye'de gıda güvencesi için genel bir yorum yapılabilir. Economist Impact (2023), 2022 yılı verilerine göre, Türkiye 49. sırada ve gıda güvencesi hakkında daha çok düşünmesi gereken 65.3 puanlık bir ortalama ile yer almıştır. Gıda güvencesinin dört boyutunun ayrı ayrı puanlandığı indexin dikkatle incelenmesi önemlidir. Söz konusu indexe genel olarak bakıldığında, Türkiye'nin avantajları, dezavantajları ve potansiyeli anlaşılabilir. Söz edilen gıda güvencesi indexi (GFSI), '*affordability (58.4), availability (65.3), quality and safety (78.5), sustainability and adaptation (61.2)*' bileşenlerinde Türkiye'yi puanlamış ve listelemiştir. Bu başlıklardan yola çıkarak, indekste kriterlerin puanlanması ve diğer ülkelerle kıyaslandığında Türkiye'nin hem sıralamadaki yeri hem puanı ilk olarak gıda güvencesinin tüm boyutlarında iyileştirilme yapılmasına ihtiyaç duyulduğunu göstermektedir (İnan, 2023).

Gıda güvencesinin ölçülmesine olanak sağlayan, farklı amaçlar ve çeşitli yöntemler gıda güvencesinin ölçülüp kıyaslanmasına olanak sağlayan diğer yaygın endeksler ise, Global Hunger Index (GHI) ve Hunger and Nutrition Reduction Commitment Index (HANCI) gibi örneklendirilebilir. Gıda güvencesi konusunda hesaplamalar yapan ve raporlar hazırlayan bu kurumların hesaplama yöntemleri detaylı incelendiğinde birbirlerinden farklıdır ve bu durumda farklı gösterilebilirler. Oysa, küresel boyutta alınması gereken tedbirler ile ülkeler arasında tutarlı olarak karşılaştırma yapmaya imkan sağlayacak gıda güvence parametrelerinin tespit edilmesi için, ülkeler arasındaki görüşmeler ve uzlaşmanın sağlanması gıda güvencesinin sürdürülmesi için önem arz etmektedir (Kulakoğlu, 2020).

Söz edilen raporlara detaylı bakıldığında, IPRFI'nin raporu daha çok küresel açlık göstergelerine ve açlığın takip edilmesine odaklanmaktadır. SOFI raporu ise çok daha konuda inceleme yapılmış olması sebebiyle çok fazla karşılaştırma yapılmasına imkân vermektedir. GFSI raporu ise ülkelerin karşılaştırılabilmesi ve açıklanabilmesi anlamında daha iyi kıyaslama yapmayı sağlamaktadır. Ancak gıda güvencesini ölçebilmek ve bu konuda bir karşılaştırma yapmak gerçekten de güçtür. Yine de bu raporların çıkarımları ve iyileştirmeler için öneriler vermeleri önemlidir.

SONUÇ VE ÖNERİLER

Gıdanın tarihi çok eskilere dayanmaktadır. İnsanlığın gıda sorunu, tarihte zenginler ve fakirler ayırımına uğramış bir kesim için gıdasını temin etmek her zaman sorunlu bir durumda olmuştur. Günümüzde, gelir dağılımındaki adaletsizlik, gıda paylaşımında da adaletsizlik oluşturmaya devam etmektedir. Son yıllarda ise bazı güçlü ülkeler gıda emperyalizmini tüm dünyaya yayma peşinde olmuşlardır. Bunun olumsuz etkilerini yaşamış ve zamanla anlamış olmalı ki, diğer ülkeler kendilerinin gıda güvencesini güvende tutmayı daha stratejik bir öncelikte ele almışlardır. Bu düşüncelerin sonucu olarak günümüzde gıda milliyetçiliği ve gıda egemenliği fikirleri kabul görmüştür. Gıda güvencesinin, vatandaşlarına devletlerin ve uluslararası kurumların çatısında garanti edilmeye çalışılmakta ancak gıda güvencesin herkesin gıdaya tam, sağlıklı, uygun fiyatlı, süreksiz şekilde hesap edildiğinde daha yapılması gereken çok fazla şey

olduğu görülebilmektedir. Gıdanın güvence edilmesi, tarihte insanlığa önemli basamaklar anlattırılmıştır. Gelecekte, herkesin gıdaya güvencesinin sağlanabilmesi insanlığı en yüksek mertebeye ulaştıracaktır. Gıda üretiminin sürdürülebilir hale getirilmesi, gıda israfının azaltılması, sağlıklı beslenme alışkanlıklarının oluşturulması ve gıdaya erişimde eşitsizliklerin giderilmesi gibi çok sayıda stratejinin bir arada uygulanması ile gıda güvencesi sağlanabilir. Bu yaklaşımla, hem çevreye duyarlı hem de toplumsal açıdan daha adil bir sistem oluşturulmasına katkı sağlanabilecektir.

KAYNAKLAR

Aytan, O. A. (2021). Hareketli avcı-toplayıcı grupların yaşam biçimiyle yerleşik çiftçi toplulukların yaşam biçim arasındaki insan-mekan ilişkisinin mukayesesi. *Kahramanmaraş Sütçü İmam Üniversitesi Sosyal Bilimler Dergisi*, 18(2), 979-1012.

BM (1972). UN Stockholm Environment Declaration. Stockholm: UN. Birleşmiş Milletler.

BM (1975). United Nations Report of the World Food Conference Rome 5-16 November 1974. New York 1975, Resolution XVII.

Bozdoğan, R. (2010). Sürdürülebilir gelişme düşüncesinin tarihsel arka planı. *Sosyal Siyaset Konferansları Dergisi*, 2010. Cilt: 0. Sayı 50.

Childe, G. (1995) Tarihte neler oldu?, (Çevirenler: Mete Tunçay ve Alaeddin Şenel) (6. Baskı). İstanbul: Alan Yayıncılık.

Direk, M. (2012). Tarım tarihi ve deontolojisi (2. Baskı). Konya: Eğitim Yayınevi. ISBN: 9786054392117.

Doğan, Z., Arslan, S. ve Berkman, A. (2015). Türkiye’de tarım sektörünün iktisadi gelişimi ve sorunları: tarihsel bir bakış. *Niğde Üniversitesi İktisadi Ve İdari Bilimler Fakültesi Dergisi*, 8(1), 29-41.

Doğruyol, A. (2021). Tarım devrimi ve zaman ölçümü. *Sakarya İktisat Dergisi*, 10(1), 103-114.

Durukan, M. (2015). Geç antik çağ’da Doğu Akdeniz’deki ekonomik gelişmenin nedenleri: İpek Yolu ve Baharat Yolu’nun rolü, LR1 amphoraları ve Kilikia’daki diğer kanıtlar. *Adalya*, no. 18 (June 2015): 237-58.

FAO (1996). The state of food security and nutrition in the world. Food and Agriculture Organization of the United Nations. World Food Summit, Rome.

FAO (2010). The Concept of the Right to Adequate Food. s.4. Food and Agriculture Organization of the United Nations.

FAO & OHCHR (2010). Right to Adequate Food. Fact Sheet No.34. Food and Agriculture Organization of the United Nations - Office of the United Nations High Commissioner for Human Rights. Geneva: United Nations.

FAO, IFAD, UNICEF, WFP and WHO. (2023). *The State of Food Security and Nutrition in the World (SOFI) 2023. Urbanization, agrifood systems transformation and healthy diets across the rural–urban continuum*. Rome, FAO. <https://doi.org/10.4060/cc3017en>

FAO, IFAD, UNICEF, WFP and WHO (2024). *The State of Food Security and Nutrition in the World 2024 – Financing to end hunger, food insecurity and malnutrition in all its forms*. Rome. <https://doi.org/10.4060/cd1254en>

Günel, D. (2019). *Aile işletmelerinin sürdürülebilirlik süreci üzerine bir Araştırma* (Yüksek Lisans Tezi). T.C. Marmara üniversitesi Sosyal Bilimler Enstitüsü İşletme Anabilim Dalı Sürdürülebilir Büyüme ve Kalite Yönetimi Bilim Dalı, İstanbul.

Güran, T. (2017). İktisat tarihi. İstanbul: Der Yayınları.

HLPE (2020). Food security and nutrition: building a global narrative towards 2030. A report by the High Level Panel of Experts on Food Security and Nutrition of the Committee on World Food Security. Rome. Access address: www.fao.org/3/ca9731en/ca9731en.pdf Access date: 21.07.2023.

IFPRI (2023). Global food policy report 2023: Rethinking food crisis responses. Washington, DC. International Food Policy Research Institute (IFPRI). <https://doi.org/10.2499/9780896294417>

IFPRI (2024). 2024 Global food policy report: food systems for healthy diets and nutrition. Washington, DC: International Food Policy Research Institute. <https://hdl.handle.net/10568/141760>

Işık, A. (2018). İlk şehirler ve yazılı medeniyete geçiş. Mavi Atlas, 6(2), 46-74.

İnan, O. (2023). Gıda arz güvenliği: (kavramlar - gerçekler - tehditler - hedefler). Aylık dergi (Mayıs, 2023). Avrasya Tarım Ekonomistleri Derneği (ATED).

Kalan, E. (2014). XIII-XIV. yüzyıllarda Kuzey İpek Yolu ve Altın Orda Hanları'nın ticaret politikaları. *Avrasya Etüdüleri*, 45(1), 43-62.

Keskin, B. ve Demirbaş, N. (2022). Is there a perfect way for food security measurement? Evaluation for Turkey. *Progress in Nutrition*, 24(1): e2022007.

Koç, G., Uzmay, A. ve Çınar, G. (2018). *Gıda güvencesini hesaplama yöntemleri ve son gelişmeler*. IX. IBANESS Kongreler Serisi – Edirne / Türkiye. 29-30 Eylül 2018. ss. 967-979.

Koçak, M. E. (2019). Moğollar zamanında ticaretin beşiği: İpek Yolu. *Genel Türk Tarihi Araştırmaları Dergisi* 1, sayı. 1 (Ocak 2019): 37-46.

Kulakoğlu, Ö. (2020). *Tarımda kendine yeterlilik ve gıda güvencesi* (Doktora Tezi). Tekirdağ Namık Kemal Üniversitesi Fen Bilimleri Enstitüsü Tarım Ekonomisi Anabilim Dalı, Tekirdağ.

Özçelik, A. (2014). *Tarım tarihi ve deontolojisi* (Ders notları). Ankara Üniversitesi Ziraat Fakültesi Tarım Ekonomisi Anabilim Dalı, Ankara.

Paker, Y (2018). *Çevresel sürdürülebilirlik ve tedarik zincirinde çevresel sürdürülebilirlik performansının ölçülmesi* (Yüksek Lisans Tezi). T. C. Dokuz Eylül Üniversitesi Sosyal Bilimler Enstitüsü İşletme Anabilim Dalı Üretim Yönetimi ve Endüstri İşletmeciliği Programı, İzmir.

Peerzada, R. İ. (2019). *Uluslararası hukukta gıda hakkı* (Doktora Tezi). İhsan Doğramacı Bilkent Üniversitesi Hukuk Bölümü, Ankara.

Şahin, T. (2020). İpek Yolu: tarihsel geçmişi, kültürü ve Türk dünyası için önemi. *Tarih Araştırmaları Dergisi*, 39(67), 71-97.

T. C. Tarım ve Orman Bakanlığı (2021). Sürdürülebilir gıda sistemlerine doğru Türkiye'nin ulusal yol haritası. T. C. Tarım ve Orman Bakanlığı. 2021.

Tekir, A. T. (2016). *Gıda Hakkı* (Yüksek Lisans Tezi). T.C. Marmara Üniversitesi Sosyal Bilimler Enstitüsü Kamu Hukuku Ana Bilim Dalı Kamu Hukuku Bilim Dalı, İstanbul.

Tıraş, H. (2002). Sürdürülebilir kalkınma ve Çevre: teorik Bir İnceleme. *Kahramanmaraş Sütçü İmam Üniversitesi İktisadi ve idari Bilimler Fakültesi Dergisi*, Cilt:2. Sayı:2. 2002. Ss 57-73.

USDA (2000). Guide to measuring household food security (Prepared by: Gary Bickel, Mark Nord, Cristofer Price, William Hamilton, John Cook). *Measuring Food Security in the United States*. USDA Food and Nutrition Service. Revised March, 2000.

Yanardağ, A. A. (2023). Bütünsel bir sosyal politika alanı olarak gıda güvencesi ve eğitim. *Politik Ekonomik Kuram*, 7(2), 516-528.

World Bank (2023). What is food security? Access address: <https://www.worldbank.org/en/topic/agriculture/brief/food-security-update/what-is-food-security> Access date: 21.07.2023.

Worldometers (2024). Worldometer – real time world statistics. Access address: <https://www.worldometers.info/tr/> Access date: 22.12.2024.

ANALYSING EXPERT OPINIONS ON AGRICULTURAL SUPPORTS IN RURAL AREAS IN KONYA CLOSED BASIN

Res. Assist. Tansu ALKAN

Niğde Ömer Halisdemir University, Faculty of Engineering, Department of Geomatics Engineering, Niğde, Turkey

Necmettin Erbakan University, Institute of Science, Konya, Turkey

ORCID ID: <https://orcid.org/0000-0001-8293-2765>

Prof. Dr. Süleyman Savaş DURDURAN

Necmettin Erbakan University, Faculty of Engineering, Department of Geomatics Engineering, Konya, Turkey

ORCID ID: <https://orcid.org/0000-0003-0509-4037>

ABSTRACT

Agricultural supports are of great importance in ensuring sustainability in the agricultural sector, in raising the income level of farmers, in increasing productivity with modern agricultural techniques and in protecting food security. When agricultural support policies are analysed in terms of both economic and social dimensions, they are necessary for strengthening the local economy and supporting rural development. In this study, opinions of experts on agricultural supports were taken in a survey conducted within the scope of a doctoral thesis to analyze the factors affecting the value of agricultural lands in the Konya Closed Basin. Experts were asked whether they were involved in agricultural activities or production and if so, their opinions on agricultural supports and their answers were analysed. Data were obtained from 472 experts belonging to the professions of geomatics engineering, agricultural engineering, forest engineering, civil engineering, urban planning, architecture and real estate appraiser by using a questionnaire form. 356 experts are not involved in agricultural production, while 116 experts are involved in agricultural production. While 24 of these experts did not express their opinions on agricultural subsidies, 92 of them expressed their opinions and suggestions. When the answers of the experts were examined, opinions and suggestions were obtained regarding the inadequacy of agricultural supports, the access of incentives to the real producer, the provision of machinery and grant support, the provision of support for fertilization and spraying that increase productivity, the planting of agricultural products that are suitable for soil structure and prevent erosion and desertification, the provision of training and support for organic agriculture, the provision of product and basin-based support, the provision of soil analysis support in fertilization, the provision of agricultural production that prevents environmental pollution, the provision of spraying support in the foresight of an engineer, the provision of training on agricultural production, the provision of farmer packages that will increase production activities, the removal of bureaucratic obstacles in agricultural support packages and the establishment of agricultural production policies. As a result of the study, it was determined that there are various problems regarding the effectiveness and adequacy of agricultural supports. The findings show that agricultural support policies should be re-evaluated, and a more efficient and sustainable agricultural support system should be established.

Keywords: Agricultural supports, Agricultural policies, Expert opinions

INTRODUCTION

Agriculture is one of the most time-honoured sectors, playing a pivotal role in addressing fundamental human needs and driving economic growth. Concurrently, agriculture stands as a pivotal sector regarding employment generation, income distribution, and the advancement of rural areas. Given its instrumental role in addressing numerous societal concerns, notably nutrition, the agricultural sector exhibits a structure that pervades all societal segments (Aktaş, Altıok & Songur, 2015). Agricultural production is an area that is carried out depending on biological, natural and human factors, has higher risks compared to other sectors and is considered less attractive in terms of investment (Topcu, 2008). Due to its disadvantageous characteristics, agriculture is one of the sectors that is protected with supportive policies in all countries, especially in developed countries (Tan, Hasdemir & Everest, 2015). Consequently, the agricultural sector requires state support to ensure its effective direction and encouragement.

Agricultural sector is susceptible to considerable volatility due to its exposure to unpredictable natural events. This exposure gives rise to a range of challenges, including those pertaining to market fluctuations, agricultural income, production and marketing (Narin, 2008). Consequently, agricultural investments are characterised by a high degree of risk, which in turn hinders producers' ability to generate a sustainable income. However, when the employment opportunities provided by the agricultural sector to the rural population are considered, in addition to its contributions to national income and foreign trade, it becomes evident that agriculture is an indispensable sector for the national economy (Yıldız, 2017). In this respect, the role of agricultural policies and supports is particularly significant in ensuring a sustainable agricultural sector.

The distinctive characteristics of agriculture necessitate the preservation of this sector in every nation, whether through direct state intervention or authorised institutions (Abay et al., 2017). To ensure the sustenance of its population, competitiveness in global markets, and the viability of its agricultural sector, a nation must implement and maintain support and incentive policies within the agricultural domain (Erdal, Erdal & Gürkan, 2013). This underscores the necessity to regard the agricultural sector not solely as an economic entity but also as a strategic domain, thereby ensuring its prioritization within the framework of long-term development plans.

Agricultural policies are designed to achieve a range of objectives, including increasing producer income and welfare, maintaining price stability, enhancing consumer welfare through affordable food provision, ensuring self-sufficiency, supporting rural development and preserving foreign currency reserves (Semerci, 2019). Agricultural policies should be oriented towards the economy of the countries and the solution of the problems encountered in the agricultural sector (Yüceer, Tan & Semerci, 2020). Agricultural supports play a pivotal role in agricultural policies. The primary objectives of implementing support policies in the agricultural sector include increasing income levels, safeguarding consumers against price volatility, balancing production and prices, encouraging production, diversifying product offerings, ensuring equitable income distribution, promoting productivity and quality enhancement (Gezer & Gezer, 2022; Oğul, 2022). In addition, the objectives of these policies include the promotion of highly competitive agricultural products, the reduction of foreign dependency through increased fertiliser production, and the reduction of production costs.

MATERIAL AND METHOD

The material of this study consists of experts specialized in geomatics engineering, agricultural engineering, forest engineering, civil engineering, urban and regional planning, architecture and real estate appraisers. In this study conducted within the scope of a doctoral thesis, a survey was conducted to analyse the factors affecting the value of agricultural lands in Konya Closed Basin. The opinions of experts on agricultural supports were also obtained in the survey form.

In this context, the participants were asked the question, “Are you involved in agricultural activity or production? If so, what are your opinions on supports in agricultural production?”.

A total of 472 experts participated in the survey. 356 of the participants were not involved in agricultural activity or production processes, while 116 were working in these areas. In this study, the opinions obtained from 116 experts involved in agricultural activities or production were analysed.

This survey study was conducted in the provinces of Aksaray, Ankara, Isparta, Konya, Karaman, Nevşehir and Niğde located in the Konya Closed Basin. The distribution of the provinces where the expert participants work is presented graphically in Figure 1.




Figure 1. Distribution of participants by province

The province with the highest number of participants was Konya with 67 experts. There were 15 participants in Karaman and 10 in Nevşehir. While Aksaray, Isparta and Niğde provinces were each represented by 7 participants, the province with the lowest number of participants was Ankara with 3 experts.

The distribution of the ages of the expert participants is presented graphically in Figure 2.




Figure 2. Distribution of participants by age

When the age distribution of the participants is examined, it is seen that the highest participation is in the 36-40 and 41-45 age groups; there are 26 participants in both groups. This is followed by the 46-50 and 26-30 age groups with 18 and 17 participants, respectively. There are 15 participants in the 31-35 age group and 3 participants in the 51-55 age group. It is observed that the number of participants decreases in the older age groups, while there are 7 participants in the 56-60 age group, while the lowest participation is in the 61-65 and over 65 age groups, with only 1 participant in each age group. Similarly, the participation is low in the 20-25 age group, with 2 participants in this age group.

The gender distribution of expert participants is presented graphically in Figure 3.




Figure 3. Distribution of participants by gender

40 females and 76 males participated in the survey.

The distribution of the educational level of expert participants is shown graphically in Figure 4.




Figure 4. Distribution of participants by educational level

According to the data on the education level of the survey participants, 50 of the participants had a bachelor's degree, 39 had a master's degree, and 27 had completed their doctorate education.

The occupational distribution of expert participants is presented graphically in Figure 5.




Figure 5. Distribution of participants by occupational

When the occupational distribution of the participants is examined, it is seen that the highest participation was provided by 65 agricultural engineers. This is followed by 13 geomatics engineers and 11 real estate appraisers. The participation in other occupational groups is lower, with 8 participants each for civil engineers and urban planners, 6 participants for architects and 5 participants for forest engineers.

The distribution of expert participants' years of experience is presented graphically in Figure 6.




Figure 6. Distribution of participants by years of experience

When the distribution of participants according to their years of experience is examined, the highest participation was provided by 47 people with 15 years of experience and the lowest participation was provided by 14 people with 1-4 years of experience. From other experience groups, there are 25 participants with 5-9 years of experience and 30 participants with 10-14 years of experience.

In addition, when the responses to the question "Have you worked in the field of real estate valuation? If so, please indicate how many years have you been working?" were examined, it was determined that 57 of the 116 participants who expressed their opinions on agricultural support worked in the field of real estate valuation, while 59 did not have any experience in this field. When the experience periods of the 57 participants working in the field of real estate valuation were examined, it was determined that 38 participants had 1-5 years of experience, 4 participants had 6-10 years of experience, 11 participants had 11-15 years of experience, and 4 participants had 16-20 years of experience.

RESULTS AND DISCUSSION

While 24 out of 116 experts engaged in agricultural activities did not express an opinion on agricultural supports, 92 experts did so. The opinions of the experts were compiled and listed as items:

- Incentives should reach real producers
- Machine and grant support should be provided
- Training for organic farming should be provided
- Fertilization and spraying support should be provided to increase productivity
- Agricultural products should be suitable for soil structure
- Agricultural production should be carried out to prevent environmental pollution

- Training should be provided on agricultural production
- The right agricultural products should be planted to prevent erosion and desertification
- Bureaucratic obstacles should be removed in agricultural support packages
- Agricultural lands should not be opened to settlement
- Organic farming should be supported
- Supports are insufficient, irregular and not suitable for the purpose
- Innovative agricultural machinery support should be provided
- Product and basin-based support should be provided
- Irrigation service should be provided
- Long-term support packages for production should be increased
- Supports should be removed, and product value should be increased
- Supports should be delivered to the actual users
- Agricultural production diversity should be increased
- Soil analysis support should be provided in fertilization
- Support for spraying should be provided according to the engineer's prediction
- Local seeds should be supported, and cooperatives should widespread
- Input costs should be reduced
- Farmer packages that will increase production activities should be created
- Agricultural production policies should be created

CONCLUSION

In this study, a survey was conducted to identify the factors affecting the value of agricultural land in the Konya Closed Basin, and the effectiveness of agricultural support policies was evaluated based on expert opinions. The findings indicate that the current agricultural supports are insufficient, inconsistent, and not aligned with their intended purpose, highlighting the need for a restructuring of the support system. It is emphasized that supports should reach actual producers, bureaucratic obstacles should be eliminated, and support mechanisms should be made more transparent.

To ensure the long-term development of the agricultural sector, it is recommended to increase machinery and grant support, promote innovative agricultural technologies, and provide fertilisation and spraying assistance based on soil analysis. Additionally, the widespread adoption of environmentally friendly farming practices, the selection of appropriate crops to prevent erosion and desertification, the expansion of incentives for organic farming, and the promotion of local seed use are considered essential measures.

Moreover, ensuring the continuity of agricultural production requires the implementation of product- and basin-based support models, the improvement of irrigation services, and the expansion of long-term incentive packages for producers. Establishing systems that ensure agricultural supports reach active producers and encouraging cooperative structures are also crucial steps in addressing structural issues within the sector.

In conclusion, improving farmers' economic conditions, enhancing production diversity, and reducing input costs are critical factors for the long-term viability of the agricultural sector.

Agricultural policies should be restructured with a holistic approach that considers economic, environmental, and social dimensions. Enhancing the efficiency and accessibility of support mechanisms will be one of the key steps in securing the future of agricultural production.

ACKNOWLEDGEMENT

This work has been supported by Necmettin Erbakan University Scientific Research Projects Coordination Unit under project number 23DR19001 and has been produced from the ongoing doctoral thesis titled "Investigation of Agricultural Land Valuation Methods and Analysis of Factors Affecting the Value of Agricultural Lands with Spatial Decision Support Systems in Konya Closed Basin".

REFERENCES

- Abay, C., Türkekul, B., Ören, M. N., Gürer, B. ve Özalp, B. (2017). Türkiye’de Üreticilerin Tarımsal Desteklerden Faydalanma Durumu Üzerine İnceleme. *Balkan ve Yakın Doğu Sosyal Bilimler Dergisi*, 3(1), 130–136.
- Aktaş, E., Altıok, M., & Songur, M. (2015). Farklı Ülkelerdeki Tarımsal Destekleme Politikalarının Tarımsal Üretim Üzerine Etkisinin Karşılaştırmalı Analizi. *Anadolu Üniversitesi Sosyal Bilimler Dergisi*, 15(4), 55-74. <https://doi.org/10.18037/ausbd.08912>
- Erdal, G., Erdal, H., & Gürkan, M. (2013). Türkiye’de Uygulanan Tarımsal Desteklerin Üretici Açısından Değerlendirilmesi (Kahramanmaraş İli Örneği). *International Journal of Social and Economic Sciences*, 3(2), 92-98.
- Gezer, T., & Gezer, M. A. (2022). Tarımsal Destek ve Kredilerin Tarımsal Üretim Üzerindeki Etkinliği. *Türk Tarım ve Doğa Bilimleri Dergisi*, 9(4), 1102-1113.
- Narin, M. (2008). Türkiye’de Uygulanan Tarımsal Destekleme Politikalarında Değişim. *Ekonomik Yaklaşım*, 19(Special), 183-225.
- Oğul, B. (2022). Tarımsal Destekler ve Tarımsal Üretim İlişkisi: Türkiye Ekonomisi Üzerine Ampirik Bulgular. *Tarım Ekonomisi Araştırmaları Dergisi*, 8(1), 44-56.
- Semerci, A. (2019). Türkiye’de Tarımsal Destekleme Uygulamaları. *ÇOMÜ Ziraat Fakültesi Dergisi*, 7(1), 181-186.
- Tan, S., Hasdemir, M., & Everest, B. (2015). Türkiye’de Tarımsal Destekleme Politikaları. *International Conference on Eurasian Economies*.
- Topcu, Y. (2008). Çiftçilerin Tarımsal Destekleme Politikalarından Faydalanma İstekliliğinde Etkili Faktörlerin Analizi: Erzurum İli Örneği. *Akdeniz University Journal of the Faculty of Agriculture*, 21(2), 205-212.
- Yıldız, F. (2017). Türkiye’de Merkezi Yönetim Bütçesinden Yapılan Tarımsal Destekleme Ödemelerinin Tarımsal Üretim Üzerindeki Etkisi: 2006–2016 Dönemi. *Sayıştay Dergisi*, (104), 45-63.
- Yüceer, S. E., Tan, S., & Semerci, A. (2020). Türkiye’de 2000-2020 Döneminde Tarımsal Destekleme Politikalarının Gelişiminin İncelenmesi. *Lapseki Meslek Yüksekokulu Uygulamalı Araştırmalar Dergisi*, 1(2), 36-46.

CLIMATE THREATS ON AGRICULTURAL PRODUCTION: EXTREME WEATHER EVENTS AND WATER MANAGEMENT**TARIMSAL ÜRETİMDE İKLİM TEHDİTLERİ: EKSTREM HAVA OLAYLARI VE SU YÖNETİMİ****Doç. Dr. Aysun ALTIKAT**

İğdır Üniversitesi, Mühendislik Fakültesi, İnşaat Mühendisliği Bölümü, İğdır, Türkiye

ÖZET

Giriş ve Amaç: Küresel iklim değişikliği, tarımsal üretim sistemleri üzerinde önemli etkiler yaratmakta ve gıda güvenliğini tehdit etmektedir. Artan sıcaklık, düzensiz yağış rejimleri ve ekstrem hava olayları, bitki gelişimi, verimlilik ve toprak sağlığı üzerinde olumsuz sonuçlar doğurmaktadır. Özellikle sıcaklık artışları, bitkilerin büyüme döngülerini değiştirerek su tüketimini artırmakta ve hastalık ile zararlı istilalarının yayılmasına neden olmaktadır. Yağış rejimlerindeki düzensizlikler ise kuraklık veya aşırı yağış gibi olaylarla üretimde dalgalanmalara sebep olmaktadır. Bu çalışmanın amacı, iklim değişikliğinin tarımsal üretim üzerindeki etkilerini bilimsel bulgular ışığında değerlendirmek ve adaptasyon stratejilerini ele almaktır. **Gereç ve Yöntem:** Bu çalışmada, iklim değişikliğinin tarımsal üretime etkilerini ele alan güncel akademik çalışmalar, raporlar ve politika belgeleri taranmıştır. Literatür taraması kapsamında farklı iklim senaryoları altında tarımsal verimlilik, toprak sağlığı, su yönetimi ve adaptasyon stratejilerine ilişkin veriler incelenmiştir. **Bulgular:** İklim değişikliği, tarımsal üretim süreçleri üzerinde çok yönlü etkiler yaratmaktadır. Artan sıcaklık, bazı bölgelerde büyüme sezonunu uzatırken, aşırı sıcaklık stresine bağlı verim kayıplarına da yol açabilmektedir. Kuraklık, su kaynaklarının azalmasına ve sulama ihtiyacının artmasına neden olurken, aşırı yağış ve sel gibi olaylar toprak erozyonunu artırarak bitkilerin kök gelişimini olumsuz etkilemektedir. Ekstrem hava olaylarının sıklığında ve şiddetinde yaşanan artış, tarımsal üretimin sürdürülebilirliği açısından büyük bir tehdittir. Tarım sektörünün bu değişimlere uyum sağlayabilmesi için sürdürülebilir tarım uygulamalarına, akıllı sulama sistemlerine ve iklim dostu üretim tekniklerine yönelmesi gerektiği bulgular arasında yer almaktadır. **Tartışma ve Sonuç:** İklim değişikliğinin tarımsal üretim üzerindeki etkileri giderek daha belirgin hale gelmekte ve gıda güvenliği açısından önemli riskler oluşturmaktadır. Tarım sektöründe bu riskleri azaltmak için adaptasyon stratejileri geliştirilmelidir. İklim dostu tarım uygulamalarının yaygınlaştırılması, hassas tarım tekniklerinin benimsenmesi ve yerel ölçekte su yönetim politikalarının geliştirilmesi kritik öneme sahiptir. Çiftçilerin değişen iklim koşullarına uyum sağlayabilmesi için eğitim programları ve mali teşvik mekanizmalarının artırılması gerekmektedir. Sonuç olarak, tarımsal üretimin sürdürülebilirliğini sağlamak ve küresel gıda krizlerinin önüne geçmek için iklim değişikliğine uyum stratejilerinin hızla uygulanması gerekmektedir.

Anahtar Kelimeler: İklim değişikliği; Tarımsal üretim; Ekstrem hava olayları; Adaptasyon stratejileri

ABSTRACT

Introduction and Purpose: Global climate change has significant impacts on agricultural production systems and threatens food security. Increasing temperatures, irregular precipitation regimes and extreme weather events have negative consequences on plant development,

productivity and soil health. In particular, temperature increases change the growth cycles of plants, increase water consumption and cause the spread of disease and pest infestations. Irregularities in precipitation regimes cause fluctuations in production with events such as drought or excessive precipitation. The aim of this study is to evaluate the effects of climate change on agricultural production in the light of scientific findings and to address adaptation strategies. **Materials and Methods:** In this study, current academic studies, reports and policy documents addressing the effects of climate change on agricultural production were reviewed. Within the scope of the literature review, data on agricultural productivity, soil health, water management and adaptation strategies under different climate scenarios were examined. **Results:** Climate change has multifaceted effects on agricultural production processes. While increasing temperatures extend the growing season in some regions, they can also lead to yield losses due to extreme heat stress. Drought causes water resources to decrease and irrigation needs to increase, while events such as excessive rainfall and floods increase soil erosion and negatively affect plant root development. The increase in the frequency and severity of extreme weather events is a major threat to the sustainability of agricultural production. Among the findings is that the agricultural sector should turn to sustainable agricultural practices, smart irrigation systems and climate-friendly production techniques in order to adapt to these changes. **Discussion and Conclusion:** The effects of climate change on agricultural production are becoming increasingly evident and pose significant risks in terms of food security. Adaptation strategies should be developed to reduce these risks in the agricultural sector. The dissemination of climate-friendly agricultural practices, the adoption of precision agriculture techniques and the development of water management policies at the local level are of critical importance. Education programs and financial incentive mechanisms should be increased for farmers to adapt to changing climate conditions. As a result, climate change adaptation strategies should be implemented rapidly to ensure the sustainability of agricultural production and prevent global food crises.

Key Words: Climate change; Agricultural production; Extreme weather events; Adaptation strategies

GİRİŞ


Küresel iklim değişikliği, tarım sektörü üzerinde büyük bir baskı oluşturmakta ve gıda üretimini etkileyen çeşitli riskleri ortaya çıkarmaktadır (IPCC, 2021). Tarımsal faaliyetler, çevresel koşullara duyarlı olduğundan, iklim değişikliğinin üretim üzerindeki etkilerini anlamak ve bu etkilere karşı çözümler üretmek büyük önem taşımaktadır (Lobell et al., 2011).

Son yıllarda sıcaklıkların kademeli olarak artması, tarım alanlarının üretkenliğini azaltmış ve bu durum, çiftçilerin geleneksel tarım yöntemlerini sorgulamalarına neden olmuştur. Ekstrem hava olaylarının daha sık ve şiddetli hale gelmesi, tarım üzerindeki baskıları daha da artırmaktadır. Kuraklıklar, sel felaketleri, dolu yağışları ve hortumlar gibi olaylar, tarlalardaki mahsullerin zarar görmesine ve ekonomik kayıplara yol açmaktadır. Aynı zamanda, toprak verimliliğinin azalması ve su kaynaklarının azalması, tarımsal üretimde ciddi sıkıntılara sebep olmaktadır (Rosenzweig et al., 2014).

Ayrıca, iklim değişikliği sadece tarımsal üretimi değil, aynı zamanda tüm tarım ekosistemlerini de etkilemektedir. Değişen iklim koşullarına bağlı olarak zararlı ve hastalıkların yayılma hızı artmakta, ürün verimliliği düşmektedir. Bütün bu gelişmeler, tarım sektöründe yeni adaptasyon stratejilerinin geliştirilmesini zorunlu hale getirmektedir (Skendžić et al., 2021). Bu makalede, iklim tehditlerinin tarımsal üretim üzerindeki etkileri bilimsel verilere dayanarak tartışılacak ve adaptasyon stratejileri sunulacaktır.

İKLİM DEĞİŞİKLİĞİ VE TARIMSAL ÜRETİME ETKİLERİ

İklim değışikliği, tarımsal üretim sistemlerini doğrudan etkileyen önemli bir faktördür. Sıcaklık artışları, yağış rejimlerindeki değışiklikler ve ekstrem hava olaylarının sıklığındaki artış, bitki gelişimi ve tarımsal verim üzerinde çeşitli olumsuz etkilere yol açmaktadır (Hatfield et al., 2011).



Şekil 1. İklim kaynaklı ekstrem çevresel olayların tarım, toprak ve mahsuller üzerindeki etkisi (Bibi and Rahman, 2023).

Sıcaklık Artışı ve Bitki Gelişimi

Küresel ısınma, bitki fizyolojisi ve gelişimi üzerinde doğrudan etkiler yaratmaktadır. Sıcaklık artışları, fotosentez oranını, solunum hızını ve su buharlaşmasını etkileyerek bitkilerin büyüme süreçlerini değıştirmektedir (Dusenge et al., 2019).

Fotosentez ve Solunum Üzerindeki Etkiler

Fotosentez, bitkilerin karbonhidrat üretimi için karbondioksit ve suyu kullanarak güneş enerjisini kimyasal enerjiye dönüştürdüğü temel bir süreçtir. Sıcaklık artışları, fotosentez enzimlerinin aktivitesini etkileyerek bu süreci olumsuz yönde etkileyebilir. Özellikle, optimal sıcaklık aralığının üzerine çıkan sıcaklıklar, fotosentez hızını azaltabilir ve bitki büyümesini sınırlayabilir. Ayrıca, yüksek sıcaklıklar bitki solunumunu artırarak, fotosentezle üretilen karbonhidratların daha hızlı tüketilmesine neden olur (Moore et al., 2021). Bu durum, bitkilerin net karbon kazancını azaltarak, biyokütle birikimini ve nihayetinde verimi düşürür.

Büyüme Dönemi ve Fenolojik Değışiklikler

Sıcaklık artışları, bitkilerin büyüme dönemlerini ve fenolojik aşamalarını da etkiler. Örneğin, artan sıcaklıklar, bitkilerin daha erken çiçeklenmesine ve olgunlaşmasına neden olabilir. Bu durum, bazı bölgelerde yetiştirme sezonlarının kısalmasına yol açabilir ve bu da verim kaybına neden olabilir. Özellikle, soğuk iklimlere adapte olmuş bitki türleri, sıcaklık artışlarından olumsuz etkilenebilir ve bu da tarımsal üretimde bölgesel kaymalara yol açabilir (Menzel and Sparks, 2006).

Sıcaklık artışlarının tarımsal verim üzerindeki etkileri, bitki türüne ve bölgesel koşullara bağlı olarak değışiklik gösterir. Örneğin, yapılan bir çalışmada, Filipinler'de sıcaklık artışının 1°C olduğu bir senaryoda, pirinç üretiminin %10 azalması beklenmektedir (Stuecker et al., 2018). Benzer şekilde, Avustralya'da buğday veriminde %50'lik bir düşüşün, iki derecelik ortalama

büyüme mevsim sıcaklık değişimi ve artan yaprak yaşlanması nedeniyle olacağı tahmin edilmektedir (Subedi et al., 2023).

Türkiye'de Durum

Türkiye'de de sıcaklık artışlarının tarımsal üretim üzerindeki etkileri gözlemlenmektedir (Chandio et al., 2021). Özellikle İç Anadolu Bölgesi'nde yaşanan kuraklık ve aşırı sıcaklar, hububat rekoltesinde önemli kayıplara yol açmıştır (Sen et al., 2012). Bu durum, gelecekte hava koşullarına bağlı olarak yaşanabilecek benzer sorunların tarımsal piyasalarda olumsuzluklar yaratacağı endişesini doğurmaktadır.

Yağış Rejimindeki Değişiklikler

İklim değişikliği, yağış rejimlerinde düzensizliklere ve aşırılıklara neden olarak tarımsal üretimi olumsuz yönde etkilemektedir (Trenberth, 2011). Hem kuraklık hem de aşırı yağış olaylarının artışı, toprak erozyonu, besin yetersizliği ve sel baskınları gibi sorunlara yol açmaktadır.

Kuraklık ve Su Kaynakları Üzerindeki Etkiler

Kuraklık, tarımsal üretim için gerekli olan suyun azalmasına neden olarak bitki gelişimini olumsuz etkiler. Su eksikliği, bitkilerin fotosentez kapasitesini düşürür, stomaların kapanmasına ve dolayısıyla karbon alımının azalmasına yol açar. Bu durum, bitki büyümesini ve verimini doğrudan etkiler (Farooq et al., 2009). Özellikle, suya duyarlı bitki türleri ve sulama imkânı kısıtlı olan bölgelerde, kuraklık nedeniyle ciddi verim kayıpları yaşanabilir.

Aşırı Yağışlar ve Toprak Erozyonu

Aşırı yağışlar, toprak erozyonunu artırarak tarımsal üretimi olumsuz etkiler. Yoğun yağışlar, toprağın üst tabakasının yıkanmasına ve besin maddelerinin kaybına neden olur. Bu durum, bitkilerin ihtiyaç duyduğu besin elementlerinin azalmasına ve dolayısıyla verim düşüşlerine yol açar (Cameron et al., 2013). Ayrıca, suya doymuş toprak koşulları, bitki köklerinin oksijen alımını engelleyerek kök çürüklüğüne ve bitki ölümlerine neden olabilir (Bhattarai et al., 2011).

Türkiye'de Yağış Rejimi Değişiklikleri

Türkiye'de yağış rejiminde görülecek değişikliklerin tarımsal üretimin azalmasına neden olacağı, kurak ve yarı kurak bölgelerdeki yağışlarda nispeten görülecek artışların ise ürün miktarında artışlara yol açabileceği belirtilmektedir (Kuzucu et al., 2016). Bu durum, bölgesel farklılıkların tarımsal üretim üzerindeki etkisini göstermektedir.

Ekstrem Hava Olaylarının Etkileri

İklim değişikliği, sıcaklık artışları ve yağış rejimindeki değişikliklerin yanı sıra, ekstrem hava olaylarının sıklığını ve şiddetini de artırmaktadır (Seneviratne et al., 2021). Kuraklık, sel, dolu fırtınaları, kasırgalar ve ani sıcaklık düşüşleri gibi ekstrem hava olayları, tarımsal üretim sistemleri üzerinde önemli olumsuz etkilere yol açmaktadır (Sikka et al., 2016). Bu olaylar mahsul kayıplarına, toprak erozyonuna, ürün kalitesinin bozulmasına ve çiftçilerin ekonomik kayıplarına neden olmaktadır.

Kuraklık ve Tarımsal Üretime Etkisi

Kuraklık, tarım sektörünü en fazla etkileyen ekstrem hava olaylarından biridir. Uzun süreli su kıtlığı, bitki büyümesini engeller ve fotosentez sürecini yavaşlatır. Özellikle suya duyarlı ürünler, kuraklık dönemlerinde ciddi verim kayıpları yaşar (Hussain et al., 2023).

FAO (2021) verilerine göre, 2000 yılından bu yana dünya genelinde yaşanan kuraklık olayları, tarımsal üretimde ortalama %30'luk bir düşüşe neden olmuştur (Bouabdelli et al., 2022). 2012 ABD kuraklığı, özellikle mısır ve soya fasulyesi üretiminde büyük kayıplara yol açmış ve mısır veriminde %26'lık bir azalmaya sebep olmuştur (Zipper et al., 2016). Türkiye'de 2021 yılı kuraklığı, özellikle İç Anadolu ve Güneydoğu Anadolu Bölgeleri'nde tahıl üretiminde %35 oranında düşüşe yol açmıştır (TÜİK, 2022). Kuraklığın etkileri, su kaynaklarına bağımlı olan

sulamalı tarım sistemlerinde daha belirgin hale gelmektedir. Sulama imkânı olmayan kuru tarım alanlarında ise kayıplar daha da fazla olmaktadır.

Kasırgalar ve Tarımsal Hasar

Kasırgalar, özellikle kıyı bölgelerinde tarımsal üretimi doğrudan etkileyen yıkıcı doğal afetlerdir. Şiddetli rüzgarlar, aşırı yağışlar ve fırtına dalgaları, tarım arazilerinde ciddi tahribata neden olmaktadır.

2017 yılında ABD'yi vuran Irma Kasırgası, Florida'daki tarım arazilerinde 2,5 milyar dolarlık zarara neden olmuştur (USDA, 2018). 2005 yılında Katrina Kasırgası, Louisiana'daki şeker kamışı tarlalarında %40'luk bir ürün kaybına yol açmıştır (Boyd, 2011). Asya-Pasifik bölgesinde yaşanan kasırgalar, özellikle pirinç üretimini olumsuz etkilemiş ve bazı yıllarda %20'ye varan verim kayıplarına sebep olmuştur (Panda, 2021). Kasırgaların etkileri sadece mahsul kaybı ile sınırlı kalmamakta, aynı zamanda tarım altyapısını da tahrip etmektedir. Seralar, sulama sistemleri ve tarımsal tesisler kasırga sonrası büyük zarar görmekte ve çiftçilerin toparlanması yıllar sürebilmektedir.

Dolu Fırtınalarının Tarımsal Üretime Zararları

Dolu yağışları, özellikle meyve ve sebze üretimi için büyük bir tehdit oluşturmaktadır. Fındık, üzüm, elma, domates gibi ürünler, dolu fırtınalarına karşı oldukça hassastır. Dolu tanelerinin doğrudan fiziksel zarara neden olması, mahsullerin satılabilir kalitesini ciddi şekilde düşürmektedir (Da and Halberstadt, 2010).

2018 İtalya dolu fırtınası, şaraplık üzüm üretiminde %30'luk bir kalite kaybına yol açmış ve şarap endüstrisinde büyük ekonomik zararlara sebep olmuştur (European Commission, 2019). 2021 yılında Fransa'da görülen dolu yağışları, elma ve şeftali üretiminde %25 oranında kayıpla sonuçlanmıştır (FAO, 2021). Özellikle seracılık yapılan bölgelerde dolu fırtınaları büyük maddi kayıplara yol açmaktadır. Bu tür olaylar nedeniyle sigorta maliyetleri artmakta ve çiftçiler ekonomik olarak zor durumda kalmaktadır.

Ani Sıcaklık Düşüşleri ve Don Olayları

Ani sıcaklık düşüşleri ve don olayları, özellikle ilkbahar aylarında çiçeklenme döneminde olan meyve ağaçları için büyük bir tehdit oluşturmaktadır. Don olayları, bitkilerin su içeriğini kristalize ederek hücre yapısına zarar verir ve verimi düşürür.

Fındık üretimi yapılan bütün illerde Mart 2014 tarihlerinde meydana gelen don olayı farklı oranlarda zarara sebep olmuştur (Balık ve Balık, 2015). 2012 yılında İspanya'da görülen don olayı, narenciye üretiminde %20 verim kaybına yol açmıştır (Gimeno et al., 2025). 2020 yılında ABD'nin Kaliforniya eyaletinde görülen don olayı, badem üretiminde 500 milyon dolarlık zarara neden olmuştur (Vuppapapati, 2023). Don olayları, özellikle yüksek ekonomik değere sahip tarım ürünlerinde büyük kayıplara sebep olmakta ve çiftçiler için ciddi ekonomik riskler oluşturmaktadır.

İklim değişikliği ile birlikte ekstrem hava olaylarının sıklığının ve şiddetinin artacağı öngörülmektedir (Zwiers et al., 2013). 2100 yılına kadar sıcak hava dalgalarının süresinin ve şiddetinin %50 oranında artması beklenmektedir (Yin et al., 2022). Bu da sıcak stresine bağlı tarımsal kayıpların daha da artacağı anlamına gelmektedir.

TARIMSAL SU YÖNETİMİ VE ADAPTASYON STRATEJİLERİ

İklim değişikliği, tarımsal üretim üzerinde önemli etkiler yaratmakta ve su kaynaklarının sürdürülebilir yönetimini zorunlu kılmaktadır. Bu bağlamda, akıllı sulama sistemleri, toprak iyileştirme ve organik tarım uygulamaları ile yerel ve ulusal politikaların entegrasyonu, tarımsal su yönetiminde adaptasyon stratejileri olarak öne çıkmaktadır (Sivakumar, 2011).

Akıllı Sulama Sistemleri

Su kaynaklarının etkin ve verimli kullanımı, tarımsal üretimin sürdürülebilirliği için kritik öneme sahiptir. Akıllı sulama sistemleri, bitkilerin su ihtiyacını doğru zamanda ve doğru miktarda karşılayarak su tasarrufu sağlamayı amaçlayan teknolojilerdir. Bu sistemler, sensörler, otomatik kontrol üniteleri ve veri analiz yazılımları ile entegre çalışarak sulama süreçlerini optimize eder (Obaideen et al., 2022). Su kaynaklarını verimli kullanmak için akıllı sulama sistemleri geliştirilmektedir. Damlama sulama ve hassas sulama teknolojileri ile su tasarrufu sağlanabilir (Ferreer and Soriano, 2007).

Damlama sulama, suyun doğrudan bitki kök bölgesine düşük debili damlalar halinde verilmesini sağlayan bir yöntemdir. Bu yöntem, suyun buharlaşma ve yüzey akışı ile kaybını minimize ederek su kullanım etkinliğini artırır (Mmolawa and Or, 2000). Özellikle su kıtlığı yaşayan bölgelerde, damlama sulama ile %30-50 oranında su tasarrufu sağlanabilmektedir (Narayanamoorthy, 2004).

Hassas sulama teknolojileri ise, bitki su tüketimini izleyen sensörler ve otomatik kontrol sistemleri kullanarak suyun optimal miktarda uygulanmasını sağlar (Bwambale et al., 2022). Bu teknolojiler, toprak nem sensörleri, hava durumu istasyonları ve bitki su stresi izleme cihazları ile entegre edilerek sulama zamanlaması ve miktarını hassas bir şekilde belirler. Bu sayede, su kullanımında %20-40 oranında tasarruf elde edilebilir (Lakhier et al., 2024).

Akıllı Sulama Sistemlerinin Avantajları

- ✓ Su Tasarrufu: Geleneksel sulama yöntemlerine kıyasla, akıllı sulama sistemleri su kullanımını optimize ederek önemli ölçüde su tasarrufu sağlar. Örneğin, otomatik damla sulama sistemleri ile %15,95 oranında su tasarrufu elde edilebilmektedir (Filintas et al., 2023).
- ✓ Enerji Verimliliği: Su pompalarının ve sulama ekipmanlarının etkin kullanımı sayesinde enerji tüketimi azalır. Bu durum, hem maliyetleri düşürür hem de çevresel faydalar sağlar (Tarjuelo et al., 2015).
- ✓ Verim Artışı: Bitkilerin su stresinin azaltılması, ürün verimini ve kalitesini artırır. Doğru sulama uygulamaları ile tarımsal verimde %10-20 oranında artış gözlemlenebilir (Ali and Talukder, 2008).
- ✓ İşgücü Tasarrufu: Otomatik sistemler sayesinde manuel sulama ihtiyacı azalır, bu da işgücü maliyetlerini düşürür ve insan hatasını minimize eder (Gallardo and Sauer, 2018).

Türkiye'de Akıllı Sulama Uygulamaları

Türkiye'de, su kaynaklarının etkin kullanımı amacıyla akıllı sulama teknolojilerine yönelik çalışmalar artmaktadır (Balta and Kulat, 2024). Özellikle İç Anadolu ve Güneydoğu Anadolu bölgelerinde, damlama sulama ve hassas sulama sistemlerinin kullanımı yaygınlaşmaktadır. Bu bölgelerde, akıllı sulama uygulamaları ile su kullanım etkinliğinde büyük oranda iyileşme sağlanmıştır (Çetin, 2025).

Akıllı sulama sistemlerinde, yapay zeka ve makine öğrenimi algoritmalarının entegrasyonu ile sulama süreçlerinin daha da optimize edilmesi hedeflenmektedir (Obaideen et al., 2022). Bu teknolojiler, sulama verilerini analiz ederek daha hassas ve öngörülebilir sulama kararları alınmasına olanak tanır. Ayrıca, Nesnelerin İnterneti (IoT) teknolojisi ile sensörlerin ve cihazların birbirine bağlanması, sulama sistemlerinin daha entegre ve koordineli çalışmasını sağlar (Adeyemi et al., 2017).

Toprak İyileştirme ve Organik Tarım

İklim değişikliğine karşı tarımsal üretim sistemlerinin dayanıklılığını artırmak için toprak iyileştirme ve organik tarım uygulamaları önemli stratejilerdir (Scialabba and Müller-Lindenlauf, 2010). Bu uygulamalar, toprağın fiziksel, kimyasal ve biyolojik özelliklerini iyileştirerek bitki sağlığını ve verimliliği artırır. Organik tarım ve toprak ıslah teknikleri ile

tarımsal üretim sistemleri iklim değişikliğine daha dayanıklı hale getirilebilir (Reganold and Wachter, 2016).

Toprak İyileştirme Teknikleri

- ✓ Organik Madde İlavesi: Kompost, yeşil gübreleme ve hayvan gübresi kullanımı ile toprağın organik madde içeriği artırılır. Bu uygulamalar, toprağın su tutma kapasitesini ve yapısını iyileştirir (Cooperband, 2002).
- ✓ Toprak Koruma Uygulamaları: Eğimli arazilerde teraslama, şeritvari ekim ve örtü bitkisi kullanımı ile erozyon kontrolü sağlanır. Bu yöntemler, toprağın verimli üst tabakasının korunmasına yardımcı olur (Lenka et al., 2017).
- ✓ Toprak pH Düzenlemesi: Kireçleme veya kükürt uygulamaları ile toprağın pH seviyesi bitki yetiştirme için optimal aralığa getirilir. Bu, bitki besin elementlerinin alımını kolaylaştırır (Kunhikrishnan et al., 2016).

Organik Tarımın Faydaları

Organik tarım, kimyasal gübreler ve pestisitler yerine doğal yöntemleri kullanarak tarımsal üretimi sürdürülebilir hale getirmeyi amaçlayan bir sistemdir (Gamage et al., 2023). İklim değişikliğine uyum sağlamak için organik tarımın birçok avantajı bulunmaktadır:

- ✓ Toprak Sağlığının Korunması: Organik tarım uygulamaları, toprak mikrobiyal aktivitesini artırarak biyolojik çeşitliliği destekler (Reganold and Wachter, 2016).
- ✓ Su Tutma Kapasitesinin Artışı: Organik madde içeriği yüksek topraklar, suyu daha iyi tutar ve kuraklık dönemlerinde bitkilerin su stresine karşı dayanıklılığını artırır (Bot and Benites, 2005).
- ✓ Karbon Tutma Potansiyeli: Organik tarım, atmosferdeki karbonun toprakta tutulmasını sağlayarak sera gazı emisyonlarının azalmasına katkıda bulunur. Araştırmalar, organik tarım uygulamalarının karbon depolama kapasitesini %20-30 oranında artırabileceğini göstermektedir (Lal, 2003).
- ✓ Ürün Kalitesinin Artışı: Organik tarımda yetiştirilen ürünlerde pestisit kalıntıları daha azdır ve besin içeriği açısından daha zengindir (Crinnion, 2010).

Organik Tarımın Yaygınlaşması ve Zorluklar

Organik tarım, birçok ülkede hızla yaygınlaşmasına rağmen bazı zorluklarla karşı karşıyadır:

- ✓ Düşük Verim: Konvansiyonel tarıma kıyasla organik tarımda verim %10-30 oranında daha düşük olabilir (Seufert et al., 2012). Bu durum, bazı çiftçilerin organik üretime geçişte tereddüt etmesine neden olmaktadır.
- ✓ Sertifikasyon Süreci: Organik tarım ürünlerinin sertifikalandırılması zaman alıcı ve maliyetli olabilir (Courville and ISEAL Alliance, 2006).
- ✓ Pazar Erişimi: Küçük ölçekli çiftçilerin organik ürünlerini satabilmesi için daha güçlü bir tedarik zinciri gereklidir (Benedek et al., 2018).

Türkiye’de organik tarım uygulamaları giderek yaygınlaşmaktadır. Bu, sürdürülebilir tarım politikalarının desteklenmesi ile daha da artırılabilir.

Yerel ve Ulusal Politikalar

Tarımsal su yönetimi ve adaptasyon stratejilerinin uygulanabilmesi için yerel ve ulusal politikaların desteklenmesi gerekmektedir (OECD, 2020). Tarımsal su yönetimi ve adaptasyon stratejileri, sadece çiftçilerin bireysel çabaları ile değil, aynı zamanda hükümetlerin ve uluslararası kuruluşların desteklediği politikalar ile etkin bir şekilde uygulanabilir. Tarımsal su yönetimi ile ilgili yerel ve ulusal politikalar, su tasarrufunu teşvik eden, çiftçilere mali destek sağlayan ve iklim değişikliğine uyumu artıran düzenlemeleri içermelidir.

Yerel Politikalar ve Uygulamaları

Yerel yönetimler, su kaynaklarının sürdürülebilir kullanımını sağlamak için çeşitli politikalar geliştirmektedir:

- ✓ Sulama Alt Yapısının Güçlendirilmesi: Modern sulama sistemlerine geçişin teşvik edilmesi ve açık kanal sistemlerinden kapalı boru sistemlerine dönüşüm projeleri.
- ✓ Eğitim ve Bilinçlendirme Programları: Çiftçilere su yönetimi ve akıllı sulama sistemleri hakkında eğitim verilmesi.
- ✓ Yerel Teşvikler ve Hibe Programları: Küçük ve orta ölçekli çiftçilere su tasarrufu sağlayan teknolojilere geçiş için destek verilmesi.

Türkiye’de Tarım ve Orman Bakanlığı’nın "Modern Basınçlı Sulama Sistemleri Hibe Destek Programı" çerçevesinde çiftçilere %50’ye kadar hibe desteği sağlanmaktadır (Yolal and Değirmenci, 2020).

Ulusal Politikalar ve Uluslararası İşbirlikleri

Tarımsal su yönetimi konusunda etkin bir ulusal politika oluşturulabilmesi için aşağıdaki adımlar önemlidir:

- ✓ Su Tasarrufunu Destekleyen Mevzuatlar: Sulama suyunun fiyatlandırılması, tarımsal su kullanımının düzenlenmesi ve aşırı su tüketimini önleyecek politikaların uygulanması.
- ✓ Araştırma ve Geliştirme (Ar-Ge) Çalışmaları: Akıllı sulama sistemleri ve su kullanım verimliliği üzerine bilimsel araştırmaların desteklenmesi.
- ✓ Uluslararası İşbirlikleri: Avrupa Birliği ve FAO gibi uluslararası kuruluşlar ile ortak projeler geliştirilerek, sürdürülebilir tarım ve su yönetimi konusunda küresel bilgi alışverişinin sağlanması.
- ✓

SONUÇ VE ÖNERİLER

İklim değişikliği, tarımsal üretim üzerinde giderek daha büyük etkiler yaratmaktadır. Artan sıcaklıklar, düzensiz yağış rejimleri, ekstrem hava olayları ve su kaynaklarının azalması, tarım sektörünü doğrudan tehdit etmektedir. Bu tehditler, sadece tarımsal üretimi değil, gıda güvenliği, ekonomik sürdürülebilirlik ve ekosistem dengesini de olumsuz yönde etkilemektedir. Tarım sektörü, bu değişimlere uyum sağlayabilmek için yenilikçi çözümler geliştirmek zorundadır. Bu bağlamda, akıllı sulama sistemleri, sürdürülebilir toprak yönetimi ve yerel ile ulusal politikaların güçlendirilmesi gibi stratejiler büyük önem taşımaktadır.

Ekstrem hava olaylarının tarımsal üretimdeki etkileri giderek artmaktadır. Özellikle kuraklık, sel, dolu fırtınaları ve ani sıcaklık değişimleri, mahsul kayıplarına yol açarak gıda üretiminde dalgalanmalara neden olmaktadır. Kuraklık, su kaynaklarının azalmasına ve sulama maliyetlerinin artmasına yol açarken, sel ve aşırı yağışlar toprak erozyonunu artırarak tarımsal üretimin sürdürülebilirliğini tehdit etmektedir. Kasırgalar ve dolu fırtınaları ise mahsullerin fiziksel olarak zarar görmesine neden olarak çiftçilerin ekonomik kayıplar yaşamasına sebep olmaktadır. Bu nedenle, tarım politikalarının ve yönetim stratejilerinin iklim değişikliğine uyum sağlayacak şekilde geliştirilmesi gerekmektedir.

Tarımsal su yönetimi, bu tehditlerle mücadelede kritik bir rol oynamaktadır. Akıllı sulama sistemleri, hassas tarım teknikleri ve sürdürülebilir su yönetimi yaklaşımları, suyun daha verimli kullanılmasını sağlamakta ve tarımsal üretimi korumaktadır. Damlama sulama, yağmurlama sulama ve hassas sulama teknolojileri, tarımsal üretimde su tasarrufu sağlayarak iklim değişikliğinin olumsuz etkilerini azaltabilir. Bu sistemlerin yaygınlaştırılması, özellikle su kıtlığı yaşayan bölgelerde tarımsal sürdürülebilirliği artıracaktır. Ayrıca, toprak iyileştirme uygulamaları ile organik tarım yöntemlerinin teşvik edilmesi, suyun toprakta daha uzun süre tutulmasını sağlayarak kuraklık etkisini azaltabilir.

İklim değişikliğine uyum sağlamak için yalnızca teknolojik çözümler yeterli değildir; yerel ve ulusal politikaların da etkin bir şekilde uygulanması gerekmektedir. Sulama altyapısının modernize edilmesi, çiftçilere yönelik teşvik programlarının artırılması ve eğitim programlarının yaygınlaştırılması, tarımsal üretimi daha dirençli hale getirecektir. Ayrıca, su yönetimi politikalarının geliştirilmesi, suyun aşırı tüketimini önleyecek ve tarımsal üretimde sürdürülebilirliği destekleyecektir. Tarım ve Orman Bakanlığı tarafından sunulan hibe destek programları, çiftçilerin modern tarım tekniklerine erişimini kolaylaştırabilir. Bununla birlikte, Avrupa Birliği, FAO ve diğer uluslararası kuruluşlarla iş birliği içinde geliştirilen projeler, küresel bilgi paylaşımını artırarak sürdürülebilir tarım uygulamalarının yaygınlaştırılmasına katkı sağlayacaktır.

Tarımsal üretimde iklim değişikliğinin olumsuz etkilerini azaltmak için aşağıdaki öneriler dikkate alınmalıdır:

- ✓ Akıllı Sulama Sistemlerinin Yaygınlaştırılması
- ✓ Toprak İyileştirme Çalışmalarının Artırılması
- ✓ İklim Dostu Tarım Uygulamalarının Geliştirilmesi
- ✓ Yerel ve Ulusal Politikaların Güçlendirilmesi
- ✓ Çiftçilerin Eğitimi ve Bilinçlendirilmesi
- ✓ Uluslararası İş Birliklerinin Desteklenmesi

Sonuç olarak, tarım sektörü için en büyük tehditlerden biri haline gelen iklim değişikliği ile mücadelede etkin adaptasyon stratejileri geliştirilmelidir. Tarımsal üretimin sürdürülebilirliği için yenilikçi çözümler, güçlü politikalar ve çiftçilerin bilinçlendirilmesi büyük önem taşımaktadır. Eğer bu önlemler zamanında alınmazsa, küresel gıda güvenliği ciddi şekilde tehlikeye girebilir. Bu nedenle, hükümetler, bilim insanları ve çiftçiler, iklim değişikliğine karşı dayanıklı tarım sistemleri oluşturmak için iş birliği içinde hareket etmelidir.

KAYNAKLAR

Adeyemi, O., Grove, I., Peets, S., & Norton, T. (2017). Advanced monitoring and management systems for improving sustainability in precision irrigation. *Sustainability*, 9(3), 353.

Ali, M. H., & Talukder, M. S. U. (2008). Increasing water productivity in crop production—A synthesis. *Agricultural water management*, 95(11), 1201-1213.

Balık, H. İ., & Balık, S. K. (2015). Fındıkta 2014 yılında meydana gelen don zararı üzerine bir araştırma. *GAP VII. Tarım Kongresi*, 28.

Balta, M. A., & Kulat, M. I. (2024). Transforming an irrigation system to a smart irrigation system: A case study from Türkiye (Turkey). *Irrigation and Drainage*, 73(5), 1799-1811.

Benedek, Z., Fertő, I., & Molnár, A. (2018). Off to market: but which one? Understanding the participation of small-scale farmers in short food supply chains—a Hungarian case study. *Agriculture and Human Values*, 35, 383-398.

Bhattarai, S. P., Midmore, D. J., & Su, N. (2011). Sustainable irrigation to balance supply of soil water, oxygen, nutrients and agro-chemicals. *Biodiversity, biofuels, agroforestry and conservation agriculture*, 253-286.

Bibi, F., & Rahman, A. (2023). An overview of climate change impacts on agriculture and their mitigation strategies. *Agriculture*, 13(8), 1508.

Bot, A., & Benites, J. (2005). *The importance of soil organic matter: Key to drought-resistant soil and sustained food production* (No. 80). Food & Agriculture Org..

- Bouabdelli, S., Zeroual, A., Meddi, M., & Assani, A. (2022). Impact of temperature on agricultural drought occurrence under the effects of climate change. *Theoretical and Applied Climatology*, 148(1), 191-209.
- Boyd, E. C. K. (2011). *Fatalities due to Hurricane Katrina's impacts in Louisiana*. Louisiana State University and Agricultural & Mechanical College.
- Bwambale, E., Abagale, F. K., & Anornu, G. K. (2022). Smart irrigation monitoring and control strategies for improving water use efficiency in precision agriculture: A review. *Agricultural Water Management*, 260, 107324.
- Cameron, K. C., Di, H. J., & Moir, J. L. (2013). Nitrogen losses from the soil/plant system: a review. *Annals of applied biology*, 162(2), 145-173.
- Chandio, A. A., Gokmenoglu, K. K., & Ahmad, F. (2021). Addressing the long-and short-run effects of climate change on major food crops production in Turkey. *Environmental Science and Pollution Research*, 28(37), 51657-51673.
- Cooperband, L. (2002). Building soil organic matter with organic amendments. *Center for Integrated Agricultural Systems (CIAS), College of Agricultural and Life Sciences, University of Wisconsin-Madison*.
- Courville, S., & ISEAL Alliance, U. K. (2006). Organic standards and certification. In *Organic Agriculture* (p. 201). Csiro Publishing.
- Crinnion, W. J. (2010). Organic foods contain higher levels of certain nutrients, lower levels of pesticides, and may provide health benefits for the consumer. *Alternative Medicine Review*, 15(1).
- Çetin, Ö. (2025). Irrigation Management and Innovative Approaches in Cotton Under Climate Change. In *Agriculture and Water Management Under Climate Change* (pp. 65-92). Cham: Springer Nature Switzerland.
- Da, R., & Halberstadt, L. (2010). Correlation between weather events and crop damages in northern Italy.
- Dusenge, M. E., Duarte, A. G., & Way, D. A. (2019). Plant carbon metabolism and climate change: elevated CO₂ and temperature impacts on photosynthesis, photorespiration and respiration. *New Phytologist*, 221(1), 32-49.
- European Commission. (2019). Climate Risks in Agriculture. European Climate Observatory
- FAO. (2021). Climate Change and Agriculture. Food and Agriculture Organization Report.
- Farooq, M., Wahid, A., Kobayashi, N. S. M. A., Fujita, D. B. S. M. A., & Basra, S. M. (2009). Plant drought stress: effects, mechanisms and management. In *Sustainable agriculture* (pp. 153-188). Dordrecht: Springer Netherlands.
- Fereres, E., & Soriano, M. A. (2007). Deficit irrigation for reducing agricultural water use. *Journal of Experimental Botany*, 58(2), 147-159.
- Filintas, A., Gougoulas, N., Kourgialas, N., & Hatzichristou, E. (2023). Management Soil zones, Irrigation, and Fertigation effects on Yield and Oil Content of *Coriandrum sativum* L. using Precision Agriculture with fuzzy k-Means clustering. *Sustainability*, 15(18), 13524.
- Gallardo, R. K., & Sauer, J. (2018). Adoption of labor-saving technologies in agriculture. *Annual Review of Resource Economics*, 10(1), 185-206.

- Gamage, A., Gangahagedara, R., Gamage, J., Jayasinghe, N., Kodikara, N., Suraweera, P., & Merah, O. (2023). Role of organic farming for achieving sustainability in agriculture. *Farming System, 1*(1), 100005.
- Gimeno, S., Crisafulli, V., Sobrino-Gómez, Á., & Sobrino, J. A. (2025). A Methodology for Estimating Frost Intensity and Damage in Orange Groves Using Meteosat Data: A Case Study in the Valencian Community. *Remote Sensing, 17*(4), 578.
- Hatfield, J. L., Boote, K. J., Kimball, B. A., Ziska, L. H., Izaurralde, R. C., Ort, D., ... & Wolfe, D. (2011). Climate impacts on agriculture: implications for crop production. *Agronomy journal, 103*(2), 351-370.
- Hussain, S. B., Karagiannis, E., Manzoor, M., & Ziogas, V. (2023). From stress to success: Harnessing technological advancements to overcome climate change impacts in citriculture. *Critical Reviews in Plant Sciences, 42*(6), 345-363.
- IPCC. (2021). *Climate Change 2021: The Physical Science Basis*. Cambridge University Press.
- Kunhikrishnan, A., Thangarajan, R., Bolan, N. S., Xu, Y., Mandal, S., Gleeson, D. B., ... & Naidu, R. (2016). Functional relationships of soil acidification, liming, and greenhouse gas flux. *Advances in agronomy, 139*, 1-71.
- Kuzucu, M., Dökmen, F., & Güneş, A. (2016). Effects of climate change on agriculture production under rain-fed condition. *International Journal of Electronics Mechanical and Mechatronics Engineering, 6*(1), 1057-1065.
- Lakhiar, I. A., Yan, H., Zhang, C., Wang, G., He, B., Hao, B., ... & Rakibuzzaman, M. (2024). A review of precision irrigation water-saving technology under changing climate for enhancing water use efficiency, crop yield, and environmental footprints. *Agriculture, 14*(7), 1141.
- Lal, R. (2003). Global potential of soil carbon sequestration to mitigate the greenhouse effect. *Critical reviews in plant sciences, 22*(2), 151-184.
- Lenka, N. K., Satapathy, K. K., Lal, R., Singh, R. K., Singh, N. A. K., Agrawal, P. K., ... & Rathore, A. (2017). Weed strip management for minimizing soil erosion and enhancing productivity in the sloping lands of north-eastern India. *Soil and Tillage Research, 170*, 104-113.
- Lobell, D. B., Schlenker, W., & Costa-Roberts, J. (2011). Climate trends and global crop production since 1980. *Science, 333*(6042), 616-620.
- Menzel, A., & Sparks, T. (2006). Temperature and plant development: phenology and seasonality. *Plant growth and climate change, 70-95*.
- Mmolawa, K., & Or, D. (2000). Root zone solute dynamics under drip irrigation: A review. *Plant and soil, 222*(1), 163-190.
- Moore, C. E., Meacham-Hensold, K., Lemonnier, P., Slattery, R. A., Benjamin, C., Bernacchi, C. J., ... & Cavanagh, A. P. (2021). The effect of increasing temperature on crop photosynthesis: from enzymes to ecosystems. *Journal of experimental botany, 72*(8), 2822-2844.
- Narayanamoorthy, A. (2004). Drip irrigation in India: can it solve water scarcity?. *Water policy, 6*(2), 117-130.
- Obaideen, K., Yousef, B. A., AlMallahi, M. N., Tan, Y. C., Mahmoud, M., Jaber, H., & Ramadan, M. (2022). An overview of smart irrigation systems using IoT. *Energy Nexus, 7*, 100124.
- OECD. (2020). *Agriculture and Water: Ensuring Sustainable Water Management*. OECD Publishing.

- Panda, A. (2021). Climate Change and Agricultural Insurance in the Asia and Pacific Region. *The Asian Development Bank: Mandaluyong, Philippines*.
- Reganold, J. P., & Wachter, J. M. (2016). Organic agriculture in the twenty-first century. *Nature Plants*, 2, 15221.
- Rosenzweig, C., Elliott, J., Deryng, D., et al. (2014). Assessing agricultural risks of climate change in the 21st century. *Proceedings of the National Academy of Sciences*, 111(9), 3268-3273.
- Scialabba, N. E. H., & Müller-Lindenlauf, M. (2010). Organic agriculture and climate change. *Renewable agriculture and food systems*, 25(2), 158-169.
- Sen, B., Topcu, S., Türkeş, M., Sen, B., & Warner, J. F. (2012). Projecting climate change, drought conditions and crop productivity in Turkey. *Climate Research*, 52, 175-191.
- Seneviratne, S. I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Luca, A. D., ... & Allan, R. (2021). Weather and climate extreme events in a changing climate.
- Seufert, V., Ramankutty, N., & Foley, J. A. (2012). Comparing the yields of organic and conventional agriculture. *Nature*, 485(7397), 229-232.
- Sikka, A. K., Rao, B. B., & Rao, V. U. M. (2016). Agricultural disaster management and contingency planning to meet the challenges of extreme weather events. *Mausam*, 67(1), 155-168.
- Sivakumar, B. (2011). Global climate change and its impacts on water resources planning and management: assessment and challenges. *Stochastic Environmental Research and Risk Assessment*, 25, 583-600.
- Skendžić, S., Zovko, M., Živković, I. P., Lešić, V., & Lemić, D. (2021). The impact of climate change on agricultural insect pests. *Insects*, 12(5), 440.
- Stuecker, M. F., Tigchelaar, M., & Kantar, M. B. (2018). Climate variability impacts on rice production in the Philippines. *PloS one*, 13(8), e0201426.
- Subedi, B., Poudel, A., & Aryal, S. (2023). The impact of climate change on insect pest biology and ecology: Implications for pest management strategies, crop production, and food security. *Journal of Agriculture and Food Research*, 14, 100733.
- Tarjuelo, J. M., Rodriguez-Diaz, J. A., Abadía, R., Camacho, E., Rocamora, C., & Moreno, M. A. (2015). Efficient water and energy use in irrigation modernization: Lessons from Spanish case studies. *Agricultural Water Management*, 162, 67-77.
- Trenberth, K. E. (2011). Changes in precipitation with climate change. *Climate research*, 47(1-2), 123-138.
- TÜİK. (2022). Türkiye’de Tarım Üretim İstatistikleri. Türkiye İstatistik Kurumu.
- USDA. (2018). Economic Analysis of Hurricane Irma’s Damage to Florida Agriculture. United States Department of Agriculture.
- Vuppalapati, C. (2023). Specialty crops almonds. In *Specialty crops for climate change adaptation: Strategies for enhanced food security by using machine learning and artificial intelligence* (pp. 627-741). Cham: Springer Nature Switzerland.
- Yin, C., Yang, Y., Chen, X., Yue, X., Liu, Y., & Xin, Y. (2022). Changes in global heat waves and its socioeconomic exposure in a warmer future. *Climate Risk Management*, 38, 100459.

Yolal, A. K., & Değirmenci, H. (2020). Basınçlı Sulama Sistemleri Hibe Destek Uygulamalarının Değerlendirilmesi: Yozgat İli Örneği. *Kahramanmaraş Sütçü İmam Üniversitesi Tarım ve Doğa Dergisi*, 23(5), 1175-1183.

Zipper, S. C., Qiu, J., & Kucharik, C. J. (2016). Drought effects on US maize and soybean production: spatiotemporal patterns and historical changes. *Environmental Research Letters*, 11(9), 094021.

Zwiers, F. W., Alexander, L. V., Hegerl, G. C., Knutson, T. R., Kossin, J. P., Naveau, P., ... & Zhang, X. (2013). Climate extremes: challenges in estimating and understanding recent changes in the frequency and intensity of extreme climate and weather events. *Climate science for serving society: research, modeling and prediction priorities*, 339-389.

THE TERM “GREEN ECONOMY” IN SUSTAINABLE DEVELOPMENT**Ayaz Hajiyev**

Former PhD candidate at the Academy of Public Administration under the President of the Republic of Azerbaijan

ORCID ID: 0000-0003-0952-9095

ABSTRACT

New frameworks and models have been applied in the study to better understand the main factors shaping the sustainable development strategy. This approach increases the practical significance of the study and provides opportunities for the wide use of the results obtained in the areas of application.

– The study, for the first time in Azerbaijani science, has identified effective practices and successful approaches at the political level in order to implement sustainable development in the context of globalization. The results obtained can be used not only for the development strategy of Azerbaijan, but also as an exemplary approach for other countries facing similar problems.

Many countries are seriously concerned about environmental problems and have developed a complex of economic measures to combat negative anthropogenic impacts. The pioneers in this area are the countries of Western Europe, South Korea, the USA and China. In our neighbors such as Russia and Turkey, the green economy is developing steadily, although it lags behind developed countries.

Currently, interest in the concept of “green economy” is growing. Undoubtedly, the problems of this most traditional economic model stem from the economic stagnation caused by numerous crises and market failures observed since the 2000s. This is especially true of the financial and economic crisis of 2008, as well as the economic crisis that began with the COVID-19 pandemic in 2020 and has been growing year by year. Russia’s military intervention in Ukraine has also complicated the economic situation in the world.

The international Non-Governmental Organization called the “Green Economy Coalition” explains the five main principles of the green economy and its elements. In fact, these principles are explained on behalf of this NGO, but there is an international consensus on these principles. In other words, a number of leading organizations in the world, including GIZ, the Global Green Growth Institute, the International Labor Organization, PAGE, OECD, the UN, etc., are in solidarity with these principles.

The fifth principle, the principle of effective governance, put forward by the Green Economy Coalition, is based on five main propositions [103]. First, a green economy is governed by institutions that are integrated, accountable and sustainable. It supports an evidence-based approach, applies interdisciplinary approaches and integrates scientific knowledge with economics for adaptation strategies.

The main feature that distinguishes a green economy from traditional economic models is the direct valuation of natural capital and ecological services as economic value. Currently, many industries around the world are beginning to adopt green economy standards to promote “greening” practices in a globalized economy. These standards, also known as sustainability standards, are specific rules that guarantee that purchased products do not harm the environment

and the people who produce them. The number of these standards has increased recently and they can help build a new, greener economy in the modern era.

Key Words: Green economy, sustainable development, global economy.

Economic development is accompanied by a steady increase in production and consumption. Over the past 50 years, more products have been produced on Earth than during the entire period of the existence of civilization until 1950. On the one hand, the rapid development of production serves the well-being of the population, and on the other hand, has a negative impact on the environment. Every year, people destroy more than 11 million hectares of forest, emit 20 billion tons of carbon dioxide into the atmosphere, and generate more than 300 million tons of plastic waste. If this process is not intervened, humanity will face a global ecological catastrophe. In order to bridge the gap between economic growth and the protection of natural resources, a new economic concept has been developed and is based on three axioms:

1. It is impossible to expand the sphere of influence indefinitely in limited space.
2. The resources of the Earth are limited, which means that increased consumption will sooner or later exhaust them.
3. All processes on Earth are closely interconnected.

Many countries are seriously concerned about environmental problems and have developed a complex of economic measures to combat negative anthropogenic impacts. The pioneers in this field are Western European countries, South Korea, the USA and China. In our neighbors such as Russia and Turkey, the green economy is developing steadily, although it lags behind developed countries.

If we consider its history, the term “green economy” first appeared in the report “Green Economy Plan” prepared for the UK government in 1989. This report was prepared in order to form a common approach to the consequences of the concept of sustainable development in the field of measuring economic development and evaluating policies. The advisory document aimed to support the UK government in implementing the principles of sustainable development.

Currently, interest in the concept of “green economy” is growing. Undoubtedly, the problems of this most traditional economic model stem from the economic stagnation caused by numerous crises and market failures observed since the 2000s. This is particularly true of the financial and economic crisis of 2008, as well as the economic crisis that began with the COVID-19 pandemic in 2020 and has been growing year by year. Military intervention of Russia in Ukraine has also complicated the economic situation in the world.

Since 2000, that is, from the beginning of a new era to the present day, humanity has faced several crises simultaneously: the climate crisis, the biodiversity crisis, the fuel, food, water crisis, and in recent years, the crisis of the financial system and the economy as a whole. The increase in global climate change emissions indicates an increasing threat of rapid climate change, which could have dire consequences for humanity.

In October 2008, the UN Environment Program (UNEP) launched the Green Economy Initiative to provide analysis and policy support for investing in green sectors and greening environmentally unsustainable sectors. As part of this initiative, UNEP produced a report, The Global Green New Deal, published in April 2009, which proposed a set of policy actions to boost economic growth.

The report called on governments to allocate a significant portion of stimulus funding to green sectors and set out three objectives:

- (i) Economic recovery;

- (ii) (ii) Poverty eradication;
- (iii) (iii) Green stimulus programs to reduce carbon emissions.

In June 2009, on the eve of the UN Climate Change Conference in Copenhagen, the UN issued a statement endorsing the green economy as a transformational solution to multiple crises. The statement noted that the economic recovery should be based on a global green economy and hoped that the “green economy” would be a turning point in finding effective solutions to the many crises facing humanity.

The Ministers and Heads of Delegation of the UNEP Global Ministerial Environment Forum held in Nusa Dua in February 2010 acknowledged in their declaration that the concept of a green economy “can significantly address existing challenges and increase economic development opportunities for all nations and provide multiple benefits.

In this regard, it should be noted that the green economy is an economic model that aims to reduce environmental risks and environmental deficiencies and achieve sustainable development without harming the environment. In this context, we can say that it is closely related to the ecological economy.

The 2011 year’s UNEP Green Economy Report states that “to be green, an economy must be not only efficient but also fair. Fairness implies the recognition of capital dimensions at the global and country levels, in particular, ensuring a just transition to a low-carbon economy. Therefore, a green economy is resource-efficient and socially inclusive.” It is precisely because of this formulation that the green economy is being discussed as a political implementation tool and the approach to the green economy is ambiguous. In other words, it has many critics due to its potential use as a political influence mechanism.

An international non-governmental organization called the “Green Economy Coalition” explains the five main principles of the green economy and its elements. In fact, these principles are explained on behalf of this NGO, but there is an international consensus on these principles. In other words, a number of leading organizations in the world, including GIZ, the Global Green Growth Institute, the International Labor Organization, PAGE, OECD, the UN, etc., are in solidarity with these principles.

These five principles are called prosperity, equity, planetary boundaries, efficiency and sufficiency, and also efficient governance. According to their report, the first principle is the prosperity principle. This principle is built on five main directions . First, the green economy ensures that everyone can create prosperity and benefit from this prosperity equally. Second, the green economy has a people-centered approach and its main goal is to create shared and genuine prosperity. This prosperity is based not only on financial resources, but also on all forms of human, social, physical and natural capital. The third direction is to ensure access to sustainable natural systems, infrastructure opportunities, investments, knowledge and education required for the development of all people. Fourth, the green economy offers opportunities aimed at creating green and decent jobs, enterprises and livelihoods. The fifth and final direction is to rely on collective action in the provision of public goods, but also take into account individual choices.

The second principle put forward by the NGO is the justice principle. This principle is based on six propositions:

- 1) The green economy promotes equity between generations. The green economy is inclusive and non-discriminatory. It shares decision-making, benefits and costs fairly. It prevents imbalances and supports the empowerment of women in particular.
- 2) The green economy also promotes the fair distribution of outcomes, reducing disparities between people, while creating enough space for wildlife.

- 3) The green economy promises a long-term perspective for the economy, creating a long-term perspective that serves the interests of citizens, while taking urgent action to combat today's multidimensional poverty and injustice.
- 4) The green economy supports solidarity and social justice, strengthening trust and social ties, human rights, the rights of indigenous peoples and minorities, and the right to sustainable development.
- 5) The green economy promotes the empowerment of SMEs, social enterprises and institutions that create sustainable livelihoods.
- 6) The green economy strives for a swift and just transition and bears its costs. In other words, it leaves no one behind, allows vulnerable groups to be leaders of the transition, and innovates in increasing social protection spending.

The third principle put forward by the NGO is known as the planetary boundaries principle, and it is built on four main directions. First, the green economy is based on protecting, restoring and investing in nature. An inclusive green economy recognizes and develops the various values of nature - the functional values of products and services that support economic systems, the cultural values of society, and the ecological values that form the basis of all life. Second, the green economy applies the precautionary principle to protect natural capital and prevent ecological limits from being exceeded, and accepts limited substitution of natural capital with other types of capital. The third dimension involves investments aimed at protecting, enhancing and restoring biodiversity, soil, water, air and natural systems. Finally, the green economy is based on innovative approaches to managing natural systems, taking into account their properties such as circularity and promoting livelihoods that are compatible with natural systems, inspired by biodiversity.

The fourth principle presented by the NGO is the principle of efficiency and sufficiency, and it is built on four main directions. First, the green economy aims to promote sustainable consumption and production. This concept includes economic models that are low-carbon, resource-efficient, diverse and circular. It also puts forward new economic development approaches to ensure prosperity within planetary boundaries. Second, the green economy concept emphasizes that we must keep the consumption of natural resources within planetary boundaries, and that achieving this goal requires fundamental changes on a global scale. The third direction is the acceptance of a "social floor" necessary for the consumption of essential goods and services to ensure people's well-being and decent living conditions. Finally, a green economy involves aligning prices, subsidies and incentives with the real costs to society, through mechanisms where "the polluter pays" or those who achieve inclusive outcomes benefit.

The fifth principle, the principle of effective governance, put forward by the Green Economy Coalition, is based on five main premises. First, the green economy is governed by institutions that are integrated, accountable and sustainable. This economy supports an evidence-based approach, applies interdisciplinary approaches and combines scientific knowledge with economics for adaptation strategies. According to the second premise, the green economy is supported by institutions that are integrated and linked across sectors and levels of government. These institutions should have sufficient capacity to carry out their roles effectively, efficiently and in an accountable manner. The third dimension emphasizes the importance of public participation, prior informed consent, transparency and social dialogue. These principles should be implemented in public, private and civil society institutions, free from vested interests. Democratic governance, acting in accordance with the demands of an informed society, is at the heart of this process. The fourth principle promotes decision-making that enables local economies to develop and the management of natural systems, while maintaining strong standards, procedures and compliance systems. Finally, the fifth principle envisages the design of the financial system in a way that serves the interests of society in a safe and sustainable

manner. This system should ensure the achievement of the goals of prosperity and sustainability.

As we have seen, these principles and their provisions define the essence of the green economy. It should also be said that the green economy has principles according to different classifications developed by different organizations, but these principles are broader, more understandable and more applicable than others. In this regard, we were guided by these principles in our research.

The main feature that distinguishes the green economy from traditional economic models is the direct valuation of natural capital and ecological services as economic value. Currently, many industries around the world are beginning to adopt green economy standards to promote “greening” practices in a globalized economy. Also known as sustainability standards, these standards are specific rules that guarantee that purchased products do not harm the environment and the people who produce them. The number of these standards has increased recently and they can help build a new, greener economy in the modern era. They can be classified as follows:

- attention should be paid to economic sectors such as forestry, agriculture, mining or fishing;
- protection of water resources and biodiversity should be ensured;
- greenhouse gas emissions should be reduced;
- social protection and rights of workers should be supported;
- attention should be paid to ensuring that production processes do not harm nature and the environment in general.

These are not all the rules and requirements. Just the main, fundamental ones. Thus, we see that if a state tries to use the green economy as an element of its sustainable development model, it must invest heavily in various sectors. Professor Karl Burkart, in this regard, divides the green economy into six main sectors:

- Renewable energy;
- Green buildings;
- Sustainable transport;
- Water management;
- Waste management;
- Land management.

As a key element of sustainable development, renewable energy is energy that is naturally generated from renewable sources. This includes sources such as sunlight, wind, water movement and geothermal heat. Renewable energy is essential for sustainable development and provides electricity generation for independent energy systems. Renewable energy technology projects are usually large-scale, but they can also be implemented in rural and remote areas. In this regard, they are often of great importance for human development. At the same time, they are suitable for developing countries.

A green building (also known as a sustainable building) is a structure that is environmentally sound and resource efficient throughout the building's life cycle. Elements of the green economy are expected to be incorporated into the processes applied throughout the life of such buildings: planning, design, construction, operation, maintenance, repair, and demolition.

Traditional building design has problems of economy, utility, sustainability and comfort. Green Building practice solves these problems and complements the shortcomings. Green building is also functional in terms of maximizing resource saving throughout the entire life cycle of the building, including energy saving, land saving, water saving, material saving, etc. Green building technology, which is considered the future of the green economy, focuses on low consumption, high efficiency, saving, environmental protection, integration and optimization.

Sustainable transport refers to transport modes that are compatible with sustainable development in terms of social and environmental impacts. Components for assessing sustainability include specific vehicles used for road, water or air transport. The assessment also includes transport operations and logistics, as well as transit-oriented development. Transport sustainability is mainly measured by the efficiency and effectiveness of the transport system, as well as the system's environmental and climate impacts.

Indeed, today we see that transport systems have a significant impact on the environment, accounting for 20% to 25% of global energy consumption and carbon dioxide emissions. The majority of emissions, almost 97%, are caused by the direct combustion of fossil fuels. In 2019, about 95% of fuel came from dirty sources. Even in the European Union, considered the cleanest continent in the world from an ecological perspective, transport is the main source of greenhouse gas emissions. In 2019, it accounted for about 31% of global emissions and 24% of emissions in the EU.

Sustainable transport systems contribute positively to environmental, social and economic sustainability in the countries where they are implemented. It is important to remember that transport systems exist to provide social and economic connections, and people are rapidly taking advantage of the opportunities offered by increased mobility. In this regard, the benefits of increased mobility need to be weighed against the environmental, social and economic costs that transport systems generate. Short-term action often contributes to incremental improvements in fuel efficiency and vehicle emissions control, while long-term goals involve shifting transportation away from fossil-based energy to other alternatives, such as renewable energy and other renewable sources. The entire life cycle of transport systems is subject to sustainability measurement and optimization.

Regarding water management, it should be said that the issue of sustainable water management is currently relevant on a global scale. Currently, only 0.08 percent of all freshwater in the world can be used. And it should also be said that people always demand water for drinking, production, recreation and agriculture, and this demand is growing. Due to the small percentage of available freshwater, optimizing freshwater is a growing problem worldwide.

In the sustainable management of water resources, great efforts are being made to optimize water use and minimize the ecological impact of water use on the natural environment. It should be borne in mind that a sustainable ecosystem is necessary for the implementation of a green economy. Water management as an integral part of the ecosystem is based on integrated water resources management based on the 1992 Dublin Principles.

Sustainable water management requires a holistic approach based on the principles of Integrated Water Resources Management, originally articulated at the Dublin (January) and Rio (July) conferences in 1992. The four Dublin Principles, proclaimed in the Dublin Declaration, which are essential for ensuring sustainable development, are:

- 1) Freshwater is a limited and fragile resource, essential for life, development and environmental sustainability;
- 2) Water development and management should be based on an inclusive approach that involves users, planners and policymakers at all levels;
- 3) Women play a central role in the provision, management and protection of water;
- 4) Water has economic value in all its uses and should be recognized as an economic asset.

The implementation of these principles has guided the reform of national water management laws around the world since 1992. Additional challenges to the sustainable and equitable management of water resources include the international or domestic allocation of many water bodies. Otherwise, political-diplomatic and political-military conflict between countries, called water conflicts, arise.

Waste management, or waste disposal, which is one of the backbones of the green economy, encompasses the processes and measures required to manage waste from the moment it is generated to its final disposal. It includes the collection, transportation, treatment, and disposal of waste, along with the monitoring and regulation of the waste management process and waste-related laws, technologies, and economic mechanisms.

Waste can be solid, liquid or gaseous, and each type has different disposal and management methods. Waste management, which is particularly relevant for a green economy, covers all types of waste, including industrial, biological, domestic, organic, biomedical, and radioactive waste.

The green economy is guided by human health. In some cases, waste can pose a threat to human health. Health problems are associated with the working mechanism of the waste management process and the regulation of the working harmony of this process and its positive results. Health problems can arise directly or indirectly. Directly, through the processing of solid waste, and indirectly through the consumption of water, land and food. Waste is generated as a result of human activities, such as the extraction and processing of raw materials. Waste management aims to reduce the negative impacts of waste on human health, the environment, planetary resources and aesthetics. The aim of waste management is to reduce the hazardous effects of such waste on the environment and human health. A large part of waste management is accompanied by solid waste generated as a result of industrial, commercial and domestic activities.

Literature

1. Brockington, D., Ponte, S. The Green Economy in the Global South: experiences, redistributions and resistance // *Third World Quarterly*. – 2015. – 36:12. – pp. 2197-2206.
2. Buseth, J.T. Narrating Green Economies in the Global South // *Forum for Development Studies*. – 2021. – 48:1. – pp. 87-109.
3. Cabeza, G.M. The concept of weak sustainability // *Ecological Economics*. – 1996. – Vol. 17, №3. – pp. 147-156.
4. Caprotti, F., Bailey, I. Making sense of the green economy // *Geografiska Annaler: Series B, Human Geography*. – 2014. – 96:3. – pp. 195-200.
5. Chen, H. International trade, urbanization and forest resource dynamics // *Journal of Interdisciplinary Mathematics*. – 2018. – 21:4. – pp. 859-867.
6. Creutzberg, J.H., Klein, A.Z., Matos, C.A. Factors that Influence the Adoption of Mobile Government (M-gov): A Proposal of A Unified Model // *Information Systems Management*. – 2023. – Vol. 40, №1. – pp. 29-46.
7. Daly, H.E. Elements of environmental macroeconomics // *Ecological Economics: The Science and Management of Sustainability*. Columbia University Press. – 1991. – pp. 32-46.
8. Deutscher, G. The Entropy Crisis / G. Deutscher. – New Jersey: World Scientific, – 2008. – 185 pages.
9. Duina, F. Development Partners: aid and cooperation in the 1990s by Ismail Serageldin / F. Duina. – Stockholm: Swedish International Development Authority, – 1994 edn. – xviii+153 pages. URL: <https://shorturl.at/BjS8t>
10. Ekins, P., Simon, S. A framework for the practical application of the concepts of critical natural capital and strong sustainability // *Ecological Economics*. – 2003. – Vol. 44, №2-3. – pp. 165-185.

11. Escrig-Olmedo, E., Fernández-Izquierdo, M.Á., Ferrero-Ferrero, I., Rivera-Lirio, J.M., Muñoz-Torres, M.J. A framework for assessing corporate sustainability risks along global supply chains: an application in the mobile phone industry // *Journal of Environmental Planning and Management*. – 2023. DOI: 10.1080/09640568.2023.2219828.
12. Faccar, K., Nahman, A., Audouin, M. Interpreting the green economy: Emerging discourses and their considerations for the Global South // *Development Southern Africa*. – 2014. – 31:5. – pp. 642-657.
13. Filho, W.L. et al. Using the sustainable development goals towards a better understanding of sustainability challenges // *International Journal of Sustainable Development & World Ecology*. – 2019. – Vol. 26, №2. – pp. 179-190.
14. Fordham, M., Bledisloe, V. *The Land and Life: An Analysis of Problems of the Land in Relation to the Future of English Rural Life With a Policy for Agriculture After the War* / M. Fordham, V. Bledisloe. – London: Routledge, – 2019. – 92 pages.
15. Gardetti, M.A., Muthu, S.S. *The UN Sustainable Development Goals for the Textile and Fashion Industry* / M.A. Gardetti, S.S. Muthu. – Singapore: Springer Singapore, – 2020. – 81 pages.
16. Giovannoni, E., Fabietti, G. *What Is Sustainability? A Review of the Concept and Its Applications* / E. Giovannoni, G. Fabietti. – *Integrated Reporting*, – 2013. – pp. 21-40.
17. Green Economy Coalition. *Principles, priorities and pathways for inclusive green economies: Economic transformation to deliver the SDGs: [Electronic resource]* / greeneconomycoalition.org. URL: <https://www.greeneconomycoalition.org/assets/reports/GEC-Reports/Principles-priorities-pathways-inclusive-green-economies-web.pdf>

THE EFFECTS OF GREEN ROOF SYSTEMS ON ENERGY EFFICIENCY

YEŞİL ÇATI SİSTEMLERİNİN ENERJİ VERİMLİLİĞİ ÜZERİNDEKİ ETKİLERİ

Rıdvan TİK

Iğdır Üniversitesi, Ziraat Fakültesi, Bahçe Bitkileri Bölümü, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0009-0008-1102-1743>

Tuncay KAYA

Iğdır Üniversitesi, Ziraat Fakültesi, Bahçe Bitkileri Bölümü, Iğdır, Türkiye

ORCID ID: <https://orcid.org/0000-0002-9126-4567>

ÖZET

Yeşil çatılar, binaların çatılarında bitki örtüsü kullanılarak yapılan ekolojik tasarımlardır. Ayrıca çevresel sürdürülebilirliğin ve enerji verimliliğinin sağlanması bakımından önemli çözümler sunmaktadır. Bu çatılar, doğal çevre ile doğrudan etkileşime girerek enerji tüketimini azaltma ve ekosistem hizmetlerini iyileştirme potansiyeline sahiptirler. Yeşil çatılar binaların ısı yalıtımını iyileştirerek yaz aylarında aşırı ısınmayı önler ve kış aylarında iç mekânların sıcak kalmasına yardımcı olurlar. Bu şekilde, ısıtma ve soğutma ihtiyaçlarını azaltarak enerji tüketimini önemli ölçüde azaltırlar. Ayrıca, yeşil çatılar şehirlerdeki ısı adası etkisini azaltarak çevresel sıcaklıkların dengelenmesine yardımcı olurlar. Bu yapılar yağmur suyu yönetiminde de önemli rol oynarlar. Bu sayede yağmur suyunu emerek yüzey suyu birikimini ve selleri önlerler. Aynı zamanda yeşil çatıların barındırdığı bitki örtüsü kentsel biyoçeşitliliği artırarak çevresel dengeyi güçlendirir. Ayrıca yeşil çatılar çevreye görsel estetik katkılar da sağlarlar. Ancak yeşil çatılar çevresel ve ekonomik faydalar sunsa da yüksek ilk kurulum maliyetleri ve bakım gereksinimleri gibi bazı zorlukları da beraberinde getirmektedirler. Yapı taşıma kapasitesinin doğru belirlenmesi, su yalıtımı ve düzenli bitki bakımı gibi unsurlar yeşil çatının verimliliği için kritik öneme sahiptir. Başta Avrupa ve Kuzey Amerika olmak üzere dünya genelinde yaygınlaşan yeşil çatı uygulamaları, Türkiye gibi gelişmekte olan ülkelerde de giderek artan bir ivme kazanmaktadır. Yeşil çatılar sadece enerji tasarrufu sağlamakla kalmazlar, aynı zamanda çevresel etkileri en aza indirerek uzun vadeli ekonomik faydalar da sağlarlar. Uzun vadede, enerji tüketimini azaltırken, doğal çevreye olan olumlu katkılarıyla da kentsel yaşamın kalitesini arttırmaktadırlar. Bu tür uygulamaların sürdürülebilir kentleşme hedefleri doğrultusunda desteklenmesi önemlidir.

Anahtar Kelimeler: Enerji, Çevresel Sürdürülebilirlik, Yeşil Çatılar.

ABSTRACT

Green roofs are ecological designs using vegetation on the roofs of buildings. They also offer important solutions in terms of ensuring environmental sustainability and energy efficiency. These roofs have the potential to reduce energy consumption and improve ecosystem services by directly interacting with the natural environment. Green roofs improve the thermal insulation of buildings, preventing overheating in the summer and helping to keep interiors warm in

winter. In this way, they significantly reduce energy consumption by reducing heating and cooling needs. Furthermore, green roofs help stabilize environmental temperatures by reducing the heat island effect in cities. They also play an important role in stormwater management. By absorbing rainwater, they prevent surface water accumulation and flooding. At the same time, vegetation on green roofs increases urban biodiversity and strengthens the environmental balance. Green roofs also provide visual aesthetic contributions to the environment. However, although green roofs offer environmental and economic benefits, they also bring some challenges such as high initial installation costs and maintenance requirements. Factors such as the correct determination of the bearing capacity of the structure, waterproofing and regular plant maintenance are critical for the efficiency of the green roof. Green roof applications, which have become widespread around the world, especially in Europe and North America, are gaining momentum in developing countries such as Turkey. Green roofs not only save energy, but also provide long-term economic benefits by minimizing environmental impacts. In the long run, they reduce energy consumption and improve the quality of urban life through their positive contribution to the natural environment. It is important to support such practices in line with sustainable urbanization goals.

Keywords: Energy, Environmental Sustainability, Green Roofs.

GİRİŞ

Bugün, enerji kaynaklarının giderek tükenme noktasına gelmesi ve nüfus artışına bağlı olarak çok katlı yapıların sayısının hızla artması, küresel enerji talebinin arttığı bir dönemi işaret etmektedir. Özellikle binaların, toplam enerji tüketiminin yaklaşık yarısına yakın bir kısmını kullandığı göz önünde bulundurulduğunda, etkin enerji kullanımı sağlamak amacıyla ekolojik ve sürdürülebilir bina tasarımlarının ön plana çıkması kaçınılmaz bir sonuç olarak karşımıza çıkmaktadır (Külekçi, 2017). Bu bağlamda, kentlerdeki yeşil alanların, bireylerin fizyolojik ve psikolojik sağlıkları üzerindeki olumlu etkileri de önemli bir rol oynamaktadır. Kentsel rekreasyon alanları dışında, yapıların düşey yüzeyleri, avlular ve balkonlar gibi potansiyel yeşil alan kullanım alanları öne çıkmakta olup, bu alanlardan biri de özellikle son yıllarda dikkat çeken çatı yüzeyleridir (Uzun, 2007). Peyzaj mimarlığı, çevresel zorlukları ele almak ve sürdürülebilirlik ilkelerini uygulamak için giderek daha fazla enerji verimliliği ve yenilenebilir enerji çözümlerine yönelmektedir. Küresel ısınma, çevre kirliliği ve enerji tüketimi ile ilgili zorluklar, yenilenebilir enerji kaynaklarının peyzaj tasarımına entegre edilmesini daha da önemli hale getirmektedir (Tik ve Kaya, 2024). Dolayısıyla, yeşil çatılar yalnızca enerji verimliliğini artırma potansiyeline sahip olmakla kalmayıp, aynı zamanda şehirlerin ekolojik yapısını iyileştirerek bireylerin yaşam kalitesine katkı sağlamaktadır.

Bitkilendirilmiş çatılar, en ünlü örneği Mezopotamya'daki Babil bahçeleri olmak üzere, geçmişte çoğunlukla estetik amaçlarla kurulmuş eski bir teknolojidir (Ayata vd., 2011). Tarihsel olarak, özellikle olumsuz meteorolojik koşullara karşı geçerli koruma sağladıkları ve bu tür soğuk iklimlerde bina kabuğunun yalıtım etkisini ve dayanıklılığını geliştirdikleri Kuzey Avrupa ülkelerinde bina çatısı için güvenilir bir çözüm olduklarını kanıtlamışlardır. Bu teknolojinin yeniden keşfedilmesi, 1960'lı ve 1970'li yıllarda çoğunlukla Avrupa'da, binalara bitkisel yüzeylerin yerleştirilmesiyle sağlanan rekreasyonel ve estetik değer nedeniyle gerçekleşmiştir. Birbirini takip eden teknik ve mantıksal gelişmeler, daha fazla alt tabaka derinliği ve daha tutarlı bitki ve ağaçlar gerektiren yoğun tiplerin aksine, daha küçük yetiştirme ortamı ve daha düşük bitki örtüsü ile karakterize edilen ve bunun sonucunda düşük bakım operasyonu ve maliyetleri ile sonuçlanan geniş yeşil çatıların mevcut konseptlerine yol açmıştır (Korol ve Shushunova, 2020).

Mevcut kent binalarında veya yeni yapılan binalarda çatılar, çeşitli bitkilendirme yöntemleriyle yeşil çatı sistemlerine dönüştürülebilir. Bu sistemler, çatılarda kullanılacak bitki türleri ve bitkilerin optimal yaşam koşullarını sağlamak için gerekli alt sistem elemanlarının niteliklerine

göre iki ana gruba ayrılmaktadır. Literatürde, bitkilendirme temelli yeşil çatı sistemleri, bitki türlerinin özellikleri ve bitkilerin sağlıklı bir şekilde büyüebilmesi için gerekli alt sistem katmanlarının yapılarına göre yoğun (intensif) ve seyrek (ekstansif) olmak üzere iki ana tipte sınıflandırılmaktadır. Bunun yanı sıra, bu iki temel çatı sisteminin birleşimiyle oluşturulan yarı-yoğun (semi-intensif) bir ara sistem de mevcuttur (Werthmann, 2007). İntensif, ekstansif ve yarı-intensif yeşil çatı sistemlerini birbirinden ayıran başlıca faktörler arasında toprak derinlikleri, yük taşıma kapasiteleri ve işlevsel özellikleri yer almaktadır. Ekstansif yeşil çatılar, ince toprak derinliklerine ve sınırlı sayıda toprak katmanına sahip olup, bu özellikleri sayesinde daha düşük ağırlık, daha düşük maliyet ve daha az bakım gereksinimi sunmaktadır. Buna karşın, intensif yeşil çatılar, derin toprak katmanlarına sahip olup, çok daha çeşitli bitki türlerinin yetiştirilmesine olanak tanır; bu durum, çatıların ağırlığının artmasına, bakım gereksinimlerinin yükselmesine ve maliyetlerin artmasına neden olmaktadır. Bitkilendirme türündeki bu farklılıklar, yeşil çatı sistemini oluşturan yapısal katmanlarda da önemli değişikliklere yol açmaktadır (Ekşi, 2012).

Gelişen teknoloji ve yapı malzemeleri bilimi sayesinde, çatı yüzeyleri, geleneksel enerji tüketen yapı elemanları olma özelliğinden sıyrılarak, ekosisteme katkı sağlayan önemli bir yapı bileşenine dönüşmektedir. Çatılar, genellikle ısıyı emen, atıl ve kullanılmayan alanlar olarak işlev görürken, günümüzde bina içindeki yaşam alanlarına yalıtım sağlayan, aynı zamanda aktivite ve yaşam alanı olarak kullanılabilen yapılar haline gelmektedir. Bu bağlamda, yeşil çatı kavramı, yalnızca basit bir yapı unsuru olmanın ötesinde, ekolojik ve sosyal açıdan önemli bir rol oynamaktadır. Yeşil çatılar, ek ilave donanım gerektirmeksizin, binanın enerji performansını, hava kalitesini ve kent ekolojisini iyileştirirken, yağmur suyu yönetimine yönelik yenilikçi çözümler sunmaktadır. Bu özellikleriyle yeşil çatılar, sürdürülebilir binaların önemli sistemleri olarak ön plana çıkmaktadır (Tohum, 2011).

Bu çalışmada, yeşil çatı sistemlerinin enerji verimliliği üzerindeki etkilerini incelemek ve sürdürülebilir enerji tasarrufu sağlama potansiyelini değerlendirmek amaçlanmıştır. Yeşil çatılar, çevresel etkilerin azaltılması ve enerji tüketiminin optimize edilmesi açısından önemli bir rol oynamaktadır. Çalışmada, bu sistemlerin enerji verimliliği, iç mekân konforu ve çevresel sürdürülebilirlik üzerindeki faydalarını ele alacak ve yeşil çatılardan sağlanan enerji tasarruf potansiyeli incelenecektir. Ayrıca, bu teknolojilerin yaygınlaştırılması ve gelecekteki uygulama alanları üzerine bir değerlendirme sunulacaktır.

YEŞİL ÇATI SİSTEMLERİNİN TANIMI VE TÜRLERİ

Yeşil çatıların kökeni, tarihsel olarak binlerce yıl öncesine dayanmaktadır. Bu çatı sistemlerinin en bilinen örneği, Babil Asma Bahçeleri'nde (Şekil 1) görülmektedir. Çeşitli medeniyetler, buldukları coğrafi koşullara bağlı olarak yeşil çatıları estetik, yalıtım, ısıtma-soğutma, gürültü izolasyonu ve çatı yapısının korunması gibi çeşitli faydalar sağlamak amacıyla kullanmışlardır. 19. yüzyılın başlarında, Kuzey Avrupa'da çatılar, toprakla kaplanarak üzerine otsu bitkiler ve bazı bitki türlerinin eklenmesiyle ısı yalıtımı sağlanması amacıyla yeşil çatı sistemleri geliştirilmiştir. 1960'lı yılların başında İsviçre'de modern yeşil çatı teknolojileri, çatının üzerine yerleştirilen katman sistemleriyle toprak ortamını ve bitki örtüsünü destekleyecek şekilde uygulanmaya başlanmıştır. Bu sistem, özellikle Almanya'da yaygınlaşarak birçok ülkeye taşınmıştır (Ekşi ve Uzun, 2016).

Yeşil çatılar “eko-çatılar”, “yaşayan çatılar” veya “çatı bahçeleri” olarak da adlandırılır ve temel olarak son katmanında bitkiler bulunan çatılardır (Cox, 2010; Parizotto ve Lamberts, 2011). Yeşil çatılar genellikle binalarının enerji verimliliğini artırmak için inşa edilir, ancak başka birçok faydası da vardır. Aslında, bitki örtüsü katmanı fotosentez süreçlerini gerçekleştirirken, toprak katmanı yağışın emilmesini sağlar ve genellikle su akış kalitesinde bir iyileşme ile sonuçlanır (Bates vd., 2013).



Şekil 1. Babil asma bahçeleri (Anonim, 2025a)

Yeşil çatı sistemleri kök koruyucu tabakalar, su tutucu malzemeler, drenaj katmanı, filtreleme katmanları, substrat ve bitki örtülerini içeren bir yapı bileşenine sahiptir (Waldbaum, 2008).

Son yıllarda, yeşil çatılar (Şekil 2) sürdürülebilir uygulamalar olarak kendini kanıtlamış ve dünya çapında büyük bir popülerlik kazanmıştır (Berndtsson, 2010). Yeşil çatılar üzerine yapılan araştırmalar sayısız sosyal, çevresel ve ekonomik faydayı ortaya koymaktadır. Önemli kanıtlar, yeşil çatıların yağmur suyu yönetimi, kentsel ısı adasının azaltılması, kentsel bitki, yaban hayatı habitatı ve çatı yaşamının artırılması, hava ve su kalitesinin ve yaşam kalitesinin iyileştirilmesi, binanın enerji tüketim maliyetlerinin azaltılması, gürültü kirliliğinin azaltılması, rekreasyonel faaliyetlerin üretilmesi ve kentsel ortamlarda yeşil alanların ve estetik değerin artırılması gibi birçok fayda sağlayabileceğini göstermektedir (Berndtsson, 2010; Fioretti, 2010). Su kalitesinin iyileştirilmesi sonucunda yeşil çatılar bir bölgedeki su arıtma tesislerinin yükünü azaltmaktadır (Voyde vd., 2010; Beck vd., 2011; Berardi, 2014).

Bitkilerle tasarlanan bu sistemler, çatıda kullanılacak bitkilerin türlerine ve bitkilerin optimal yaşam koşullarını sağlamak için gereken alt sistem elemanlarının özelliklerine göre iki ana gruba ayrılmaktadır. Bu gruplar, bitkilendirme yöntemlerine dayalı olarak "intensif" (yoğun) çatı sistemleri ve "ekstansif" (seyrek) çatı sistemleri olarak sınıflandırılmaktadır (Liu, 2004).



Şekil 2. Binalarda 2016 mükemmellik ödülleri için seçilen yeşil çatı: konut binalarındaki yeşil çatılar (1,2); kurumsal binalar (3,4); ticari binalar (5,6) (Shafique vd., 2018)

Yoğun Bitkilendirilmiş Çatı Sistemleri (İntensif)

Yoğun bitkilendirilmiş yeşil çatılar, doğal zemin seviyesindeki peyzaj düzenlemelerine benzer şekilde tasarlanarak, 15 cm'den daha derin bir bitki büyüme katmanına sahip olup 'yoğun' bakım gerektiren sistemlerdir (Getter, 2009). Bu sistemler, bitki türleri bakımından büyük bir çeşitliliğe olanak tanıyan ve toprak derinliği ile substrat yapısının diğer sistemlere göre daha geniş olduğu intensif (yoğun) çatı sistemleri kapsamında değerlendirilir (Weiler ve Scholz-Barth, 2009).

Seyrek Bitkilendirilmiş Çatı Sistemleri (Ekstansif)

Seyrek bitkilendirilmiş yeşil çatı sistemlerinde, kullanılan bitki türlerinin bakım gereksinimlerinin minimum düzeyde olması ve doğal yaşam döngülerini sürdürebilecek, kendi kendine üreyebilen bitkiler olması tercih edilmektedir. Bu tür çatılar, genellikle çatıya sınırlı veya hiç giriş yapılmayan durumlarda tercih edilmekte olup, bitkilerin bağımsız bir şekilde varlıklarını sürdürebilmeleri için uygun ortamı sağlamaktadır. Bu tip çatılarda bitkilerin bitki taşıyıcı katman kalınlığı 2 ile 15 cm arasında değişmektedir. Çatıya vereceği toplam yük ortalama 100 kg/m²'dir. (Tohum, 2011; Külekçi, 2017). İntensif ve ekstansif çatı sistemlerinin karşılaştırılması Çizelge 1'de verilmiştir.

Çizelge 1. Yeşil çatıların sınıflandırılması (Berardi, 2014)

Ana Özellikler	Ekstansif Çatı Sistemi	İntensif Çatı Sistemi
Büyüme ortamının kalınlığı	200 mm'nin altında	200 mm'nin üstünde
Erişilebilirlik	Erişilemez (kırılğan kökler)	Erişilebilir (rekreasyon amaçlı kullanılabilir)
Ağırlık	60–150 kg/ m ²	300 kg/ m ² üzeri (güçlendirilmiş yapıya ihtiyaç duyulabilir)
Bitki çeşitliliği	Düşük (yosun, ot ve çimen)	Yüksek (çim veya çok yıllık bitkiler, çalı ve ağaç)
Yapı	Orta derecede kolay	Teknik olarak karmaşık
Sulama	Genellikle gerekli değildir	Drenaj ve sulama sistemleri gereklidir
Bakım	Basit	Karmaşık
Maliyet	Düşük	Yüksek

YEŞİL ÇATILAR VE SÜRDÜRÜLEBİLİRLİK

Yeşil çatılar binaların çatı yüzeylerinde bitki yetiştirilmesini sağlayan, su geçirmez katmanlar ve drenaj sistemleriyle donatılmış alanlardır. Bu uygulamalar, binaların enerji verimliliğini artırmanın yanı sıra doğal yalıtım sağlamak, su yönetimini iyileştirmek ve kentsel ısı adası etkisini azaltmak gibi çevresel sürdürülebilirlik açısından önemli faydalar sunar (Özbey ve Turhan, 2024).

Yeşil çatı sistemlerinin başlangıç maliyetleri klasik çatılara göre daha yüksek olsa da, uzun vadede bu maliyetler kendini amorti etmektedir. Yeşil çatılar, enerji verimliliği sağlayarak enerji tüketimini azaltır, çatı elemanlarının ömrünü uzatır ve yağmur sularını emerek altyapı üzerindeki yükleri hafifletir, bu da bakım ve onarım maliyetlerini düşürür. Ayrıca estetik açıdan binalara değer katar ve bitkilerin oluşturduğu koruyucu tabaka, malzemelerin dayanıklılığını artırarak uzun vadeli ekonomik faydalar sağlar (Tohum, 2011). Bu anlamda yeşil çatıların sürdürülebilir etkileri 7 başlık altında toplanabilir (Kıasf ve Selçuk, 2023);

Doğal Çevre ve Biyolojik Çeşitliliğin Korunmasındaki Rolü

Kişi başına düşen yeşil alan oranının düşük olduğu şehirlerde, yeşil çatılar önemli bir rol oynamaktadır. Bu çatılar, steril kentsel alanlarda biyoçeşitliliği teşvik eden doğal habitatlar oluşturarak, şehirleri daha yaşanabilir hale getirmektedir. Yeşil çatılar, kuşlar, arılar, kelebekler ve böcekler gibi canlılar için mikro ölçekte yaşam alanları sunarak, doğal çevre ile kentsel alanlar arasındaki bağlantıyı güçlendirir (Kıasf & Selçuk, 2023).

Yeşil Çatıların Enerji Verimliliği Üzerindeki Etkileri

Bina yüzeylerine bitki ve peyzaj eklenmesi, bina sıcaklığını 20 °C'ye kadar düşüren ve aynı zamanda iklimlendirme enerjisini % 25 ila % 80 arasında azaltan bir teknik olarak kabul edilmektedir. Yeşil çatılar, bina sektöründe enerji tasarrufunun en etkili araçlarından biri olarak tanıtılmıştır. Bu özel çatı türü, binaların termal performansının iyileştirilmesi yoluyla bina enerji talebini azaltmaktadır. Termal performansın iyileştirilmesi, gölgelemenin artması, daha iyi yalıtım ve çatı sisteminin daha yüksek termal kütesinden kaynaklanmaktadır. Yeşil çatıların ana işlevi, binanın iç mekânlarını ısıtan güneş radyasyonunu önlemektir. Yeşil çatılar güneş radyasyonunun % 27'sini yansıtabilir ve fotosentez yoluyla güneş radyasyonunun % 60'ını absorbe edebilir ve geri kalan % 13'lük kısmını yetiştirme ortamına iletebilir (Saadatian vd., 2013).

Yeşil çatılarda, toprak ve bitki katmanları kış aylarında ısı yalıtımı görevi görür. Kışın daha kalın olan toprak ve bitki örtüsü binanın ısı tutma özelliğini artırırken yapıyı olumsuz hava koşullarından da korur. Buna karşılık, yaz aylarında yeşil çatı katmanı güneşin ısı emme etkisini azaltarak bina içindeki soğutulmuş havanın çatıdan dışarı çıkmasını sağlar ve bina içinde gereksiz ısı birikimini önler (Kabuloğlu, 2005).

Geleneksel çatı ile yeşil çatıya dönüştürülmüş olan referans çatıdan elde edilen veriler, her iki çatı tipinin altında kalan iç ortamın ısınması ve soğutulması için gereken enerji miktarındaki farkları göstermektedir. Yapılan ölçümlere göre, geleneksel çatı için 1 günlük ortalama enerji tüketimi 6 – 7,5 kWh/gün arasında değişirken, yeşillendirilmiş çatıda bu değer 1,5 kWh/gün olduğu belirlenmiştir. Bu sonuç, yeşil çatının, geleneksel çatıya kıyasla günlük enerji tüketimini % 75 oranında azalttığını ortaya koymaktadır. Diğer bir ifadeyle, yeşil çatı, geleneksel referans çatıya göre % 75 oranında enerji tasarrufu sağlamaktadır (Aras, 2019).

Yeşil Çatılarda Isı Transferi

Yeşil çatılar güneş radyasyonunun % 20 ile % 30'unu yansıtır ve % 60'ına kadarını fotosentez yoluyla emerken, ısının % 20'sinin altındaki bir yüzdeyi de yetiştirme ortamına iletir. Kuru bir yeşil çatı için Lazzarin vd. (2005); gelen güneş ışınımının % 23'ünün güneş yansıtıcılığı, % 39'unun güneş emilimi, % 24'ünün dışarıdan iletim akısı, % 12'sinin evapotranspirasyon ve % 1,3'ünün termal birikim yoluyla dağıldığını ve % 1,8'den daha azının nihayetinde alttaki odaya girdiğini tespit etmiştir. İç sıcaklık ve bitki çeşitleri arasındaki korelasyonları arasında güçlü ilişkiler bulunmuştur (Weng vd., 2004).

Liu ve Minor (2005) yeşil çatıların ısı akışını azaltmasını yazın % 70-90, kışın ise % 10-30 aralığında değerlendirmiştir. Ayrıca, yeşil çatıların termal etkisi, yetiştirme ortamının derinliği artırıldığında yazın % 3 oranında artmıştır.

Yeşil çatılar, özellikle kentlerdeki sınırlı yeşil alan sorununa çözüm sunmanın yanı sıra, ısı adası etkisini azaltmada da önemli bir rol oynamaktadır. Geleneksel çatılar yerine yeşil çatılar kullanıldığında, hava ve yüzey sıcaklıkları arasındaki denge sağlanır. Bitkiler, evapotranspirasyon, gölgeleme ve ısı depolama gibi mekanizmalar aracılığıyla sıcaklık artışını azaltır. Yeşil çatı sisteminde, bitkinin taşıyıcı katmanında depolanan ısı, iç mekânın ısınmasını engelleyerek, bitki ve taşıyıcı katmanlardaki nemin buharlaşmasına olanak tanır. Ayrıca, sert yüzeylerin aşırı ısınmasını önleyerek ortamda nemin artmasını sağlar ve bu sayede çevre sıcaklığının düşmesine katkı sunar (Kabuloğlu, 2005).

Gürültü Etkisini Azaltma

Şehirler, yoğun nüfus nedeniyle genellikle gürültü kirliliği sorunuyla karşı karşıya kalan bölgelerdir. Yeşil çatılar, içerdikleri katmanlar ve özellikle toprağın gözenekli yapısı sayesinde ses dalgalarını emerek önemli bir ses izolasyonu sağlar ve akustik konforu artırır. Bu çatılar, sesin absorbe edilmesini ve emilmesini sağlayarak gürültü seviyelerini düşürür. Yeşil çatılarda, bitki örtüsünün kalınlığı, bitki türü, taşıyıcı katmanın kalınlığı ve yeşil çatının büyüklüğü gibi faktörler, ses kirliliği üzerinde etkili olan önemli unsurlardır. Özellikle toprak seviyesinin kalınlığı, ses yalıtımı başarısıyla doğru orantılı olarak sesin emilmesinde belirleyici bir rol oynamaktadır (Erkul, 2012).

Çatı Membranı Kullanım Ömrünü Uzatma

Bitümlü yeşil çatı malzemelerinin su geçirmezlik ömrü genellikle ortalama 25 yıl civarındadır. Ancak, yüksek sıcaklıklar, membranın ömrünü kısaltan önemli bir faktördür. Güneş ışınlarının doğrudan çatı yüzeyine gelmesi, yüzeyin kırılma yaşamasına, büzülmesine ve elastik yapının kaybolarak çatlakların oluşmasına neden olur. Öte yandan, yeşil çatılarda membranın ömrü 60 yıla kadar uzayabilmektedir. Bunun nedeni, güneş ışınlarının doğrudan membrana nüfuz edememesi, böylece çatlakların oluşmamasıdır. Bu durum, az atık üretilmesiyle ekonomiye

ve enerji tasarrufuna katkı sağlarken, aynı zamanda daha uzun bir kullanım ömrü sunar (Kabuloğlu, 2005; Koç ve Gültekin, 2010).

Yağmur Suyuna Etkisi

Yeşil çatılar, özellikle sağanak yağışlar gibi yoğun yağışlarda, suyun kanalizasyon sistemine geçişini geciktirerek önemli bir rol oynamaktadır. Yapılaşmış alanlarda, suyun yeraltı sularına karışması sınırlıdır ve bu durum, kanalizasyon tahliye sistemlerine ek yük getirmektedir. Ancak, yağmur sularının büyük bir kısmı çatılara düşmekte ve yeşil çatılar, üzerindeki toprak aracılığıyla bu suyu emer. Bitkiler, emilen suyu kullanırken, kullanılmayan su buharlaşarak çevreye geri kazandırılır. Bu mekanizma, mevcut su şebekelerinin yükünü hafifletir. İntensif yeşil çatılarda toprak kalınlığı genellikle 20-40 cm arasında değişir ve çatılar, 10-15 cm su tutma kapasitesine sahiptir. Yeşil çatılar genel olarak yağmur sularının % 10-15'ini tutabilmektedir. Toprak kalınlığı 10 cm olan çatılar suyun % 50'sini, 20 cm kalınlığındaki çatılar % 60'ını, 50 cm kalınlığındaki çatılar ise % 90'ını tutabilmektedir (Koç ve Gültekin, 2010).

Kentlerdeki yoğun yapılaşma nedeniyle yağmur sularının toprakla buluşması azalmakta, bu durum kanalizasyon ve tahliye sistemlerine ek yük bindirerek su baskınlarına yol açmaktadır. Ancak, yeşil çatılar bu sorunu çözmeye önemli bir rol oynamaktadır. Yeşil çatılarda yağışların büyük bir kısmı çatılarda bulunan toprak tarafından emilir, bu su bitkiler tarafından kullanıldıktan sonra buharlaşır. Yeşil çatı tasarımında seçilen sisteme bağlı olarak, çatıdan atılması gereken su miktarında % 90'a kadar tasarruf sağlanması mümkündür. Bu durum, yapılar ve şehir altyapılarındaki atık su borularının yükünü hafifletmektedir. Ayrıca, yeşil çatılar, filtre yatakları, yağmur bahçeleri, biyolojik depolama sistemleri, sarnıçlar ve yağmur varilleri ile birlikte kullanılabilir. Yağmur suları yeşil çatılarda toplanarak filtre edilip toprağa verilebilir ve aynı zamanda bahçe sulama veya tuvaletlerde kullanılabilir (Karıptaş, 2010).

Hava Kalitesini İyileştirme Etkisi

Şehir merkezlerinde, yüksek binalar rüzgâr hızını azaltarak ve ısıyı hava ceplerinde hapsederek havalandırmayı engelleyebilir, bu da kirleticilerin günlerce havada asılı kalmasına neden olabilir. Yeşil çatılar, toz partiküllerini filtreleyip bağlayarak ve havadaki toksinleri doğal yollarla temizleyerek bu durumu iyileştirebilir. Duman, sülfür dioksit, karbon dioksit ve diğer kirleticiler, yapraklar tarafından emilir ve filtrelenir, bu da çevredeki havanın doğal olarak temizlenmesini sağlar. Ayrıca, atmosferik toz, yağmurla birlikte yeşil çatıların toprak alt tabakasına taşınana kadar tutulur. Yeşil çatılar, ısı adası etkisini azaltarak, ozon oluşumuna katkıda bulunarak kentsel alanlardaki ozon sorununu hafifletmeye yardımcı olabilir. Örneğin, Atlanta'da ısı adası etkisi, ozon üretimini iki katına çıkarırken, yapılan araştırmalar ozon seviyelerindeki artışın astım ve diğer solunum rahatsızlıkları üzerinde olumsuz etkiler yarattığını ortaya koymuştur. Bu bağlamda, yeşil çatılar ve diğer bitki örtülü alanların artırılması, hava kalitesini büyük ölçüde iyileştirme potansiyeline sahiptir (Velazquez, 2005).

Yeşil çatılar, aynı zamanda üzerinde yer alan bitkiler sayesinde havada uçan partiküllerin filtre edilmesine yardımcı olur. Bitkiler üzerinde hapsolan havada uçan partiküller, yağmur yoluyla bitki büyüme katmanına aktarılmaktadır. Ayrıca bitkiler, gaz halindeki hava kirleticileri de fotosentez yoluyla absorbe edip yapraklarında depolayarak ileride humus olarak toprağı zenginleştirmesini sağlamaktadır (Tohum, 2011).

SÜRDÜRÜLEBİLİR PEYZAJ TASARIMINDA YEŞİL ÇATI SİSTEMLERİNİN İŞLEVİ VE ÖNEMİ

Peyzaj Mimarlığı, doğa, planlama ve tasarım kavramlarını sistematik bir çerçevede ele alarak, sanat, bilim, mühendislik ve teknolojiyi bir araya getiren bir disiplin olarak tanımlanabilir. Bu alanda, doğal ve kültürel kaynakların doğru bir şekilde değerlendirilmesi ve ekolojik,

ekonomik, fonksiyonel sürdürülebilirlik ilkeleri doğrultusunda planlama, yönetim ve alan tasarımı yapılmaktadır. Peyzaj mimarları, bu kapsamda, çevresel etmenleri göz önünde bulundurarak, alan kullanım kararlarını optimize etmekte ve sürdürülebilir yaşam alanları yaratmayı hedeflemektedir (Atıl vd., 2005). Yeşil alanların kullanımı, sürdürülebilir mimari tasarımın önemli bir ilkelerinden biridir. Bu alanlar, çevresel etkileri olumlu yönde şekillendiren bitki örtüleri, yeşil çatılar gibi tasarım stratejilerini içermektedir. Yeşil alanlar, doğal yaşamın korunmasına katkı sağlamakla birlikte, binaların enerji verimliliğini artırmak, hava kalitesini iyileştirmek ve ekosistem çeşitliliğini desteklemek gibi çeşitli faydalar sunmaktadır (Bekaroğlu ve Şentürk, 2019).

Sürdürülebilir peyzaj tasarımında yeşil çatı sistemleri çevresel, ekonomik ve sosyal sürdürülebilirliği destekleyen önemli işlevlere sahip bileşenler olarak öne çıkmaktadır. Bu sistemler binaların enerji verimliliğini artırarak ısı yalıtımını iyileştirmekte, yağmur suyu yönetimini düzenlemekte ve kentsel ısı adası etkisini azaltarak mikro iklim koşullarını iyileştirmektedir. Yeşil çatılar ayrıca biyoçeşitliliği destekleyerek ekosistem hizmetlerini güçlendirmekte, hava kalitesini iyileştirir ve karbon emilimini artırarak iklim değişikliğiyle mücadeleye katkıda bulunmaktadır. Peyzaj mimarisi bağlamında yeşil çatılar estetik değerleri artırmakta, doğal yaşam alanlarının kentlere yeniden entegre edilmesini sağlamakta ve kentsel alanlarda sürdürülebilir tasarımın gerçekleştirilmesinde kritik bir rol oynamaktadır. Bu sistemlerin uygulanması, hem tek tek binaların hem de genel kentsel yapının çevresel performansını olumlu yönde etkileyerek sürdürülebilir peyzaj tasarımına katkıda bulunmaktadır.

SONUÇ

Kentleşmenin hızla artması, kentsel sürdürülebilirlik ilkelerinin uygulanmasında önemli zorluklar yaratmakta, bu süreçte doğal çevrenin göz ardı edilmesi ve kaynakların aşırı ve sorumsuz bir şekilde kullanılması durumu yaygınlaşmaktadır. Özellikle kentlerdeki su tüketiminin artması ve betonlaşmanın hız kazanması, doğal su emme kapasitesine sahip alanların azalmasına ve bu da yağmur suyunun toprak tarafından emilmesinin önlenmesine yol açmaktadır. Bunun yanı sıra, artan nüfus yoğunluğu ve altyapı yetersizlikleri, kanalizasyon sistemlerinin taşıma kapasitesini aşmasına neden olmakta ve buna bağlı olarak, aşırı hava olayları sonucunda sel baskınları ve erozyon gibi çevresel sorunlar ortaya çıkmaktadır. Bu bağlamda, kentsel sürdürülebilirliği sağlamak için potansiyel bir çözüm olarak yeşil çatılar, ekosistem hizmetleri sunma kapasitesi ve su yönetimi konusundaki faydalarıyla önemli bir alternatif olarak öne çıkmaktadır.

Yeşil çatı sistemleri, enerji verimliliği ve sürdürülebilir enerji tasarrufu sağlama açısından önemli bir potansiyele sahiptir. Bu sistemlerin, özellikle yoğun yapılaşmanın hâkim olduğu kentsel alanlarda, geleneksel çatı sistemlerine kıyasla enerji tüketimini önemli ölçüde azalttığı yapılan araştırmalarla ortaya konmuştur. Yeşil çatılar, yağmur suyunun emilmesi, buharlaşma ve bitkiler aracılığıyla ısı transferinin düzenlenmesi gibi doğal süreçler sayesinde, binaların ısınma ve soğutma ihtiyacını en aza indirerek enerji tüketimini optimize etmektedir.

Bu çerçevede, yeşil çatılar, enerji kullanımını büyük oranda azaltma potansiyeline sahip olup, böylece binaların enerji performansını önemli ölçüde iyileştirmektedir. Ayrıca, yeşil çatı sistemleri, çevresel sürdürülebilirlik açısından da önemli katkılar sağlamaktadır. Bu sistemler, kentsel ısı adası etkisini hafifletme, suyun yönetimi ve filtrasyonu gibi çevresel faydalar sunmakta, bunun yanı sıra sera gazı emisyonlarının azaltılmasına yardımcı olmaktadır. Yeşil çatılar, binaların enerji ihtiyacını azaltarak hem çevresel hem de ekonomik açıdan olumlu etkiler yaratmakta, aynı zamanda enerji tüketimi ile ilgili küresel sorunlara çözüm olma yolunda önemli bir alternatif teşkil etmektedir.

Sonuç olarak, yeşil çatı sistemlerinin enerji verimliliği üzerindeki olumlu etkileri ve sürdürülebilir enerji tasarrufu sağlama potansiyeli, bu sistemlerin yaygınlaştırılmasının gerekliliğini vurgulamaktadır. Bu tür sistemlerin kentsel ve bireysel ölçeklerde benimsenmesi; enerji tüketiminin azaltılmasına, çevresel etkilerin minimize edilmesine ve sürdürülebilir kalkınma hedeflerine ulaşılmasına katkı sağlayacaktır.

KAYNAKLAR

- Anonim, (2025a). <https://www.gazetekadikoy.com.tr/cevre/babilin-asma-bahelerinden-bugne-yesil-atilar/> Erişim Tarihi: 01.02.2025
- Aras, B. B. (2019). Kentsel sürdürülebilirlik kapsamında yeşil çatı uygulamaları. *Manas Sosyal Araştırmalar Dergisi*, 8(1), 469-504.
- Ayata, T., Tabares-Velasco, P. C., & Srebric, J. (2011). An investigation of sensible heat fluxes at a green roof in a laboratory setup. *Building and environment*, 46(9), 1851-1861.
- Bates, A. J., Sadler, J. P., & Mackay, R. (2013). Vegetation development over four years on two green roofs in the UK. *Urban forestry & urban greening*, 12(1), 98-108.
- Beck, D. A., Johnson, G. R., & Spolek, G. A. (2011). Amending greenroof soil with biochar to affect runoff water quantity and quality. *Environmental pollution*, 159(8-9), 2111-2118.
- Bekaroğlu, E. ; Şentürk, Z. (2019). “Sürdürülebilir mimari tasarımın insan sağlığına etkisi”. *Journal of Health and Environment*, 10 (2), 98-106.
- Berardi, U., GhaffarianHoseini, A., & GhaffarianHoseini, A. (2014). State-of-the-art analysis of the environmental benefits of green roofs. *Applied energy*, 115, 411-428.
- Berndtsson, JC (2010). Yeşil çatı performansı, akış suyu miktarı ve kalitesinin yönetimine yönelik: Bir inceleme. *Ekolojik mühendislik*, 36 (4), 351-360.
- Cox, B.K. (2010). *The influence of ambient temperature on green Roof R-values*. Master Thesis. Portland State University.
- Duffy, K. (2004). NASA studies how to cool area as heat builds up. *Atlanta Journal Constitution*, 18.
- Ekşi, M. 2012.*Yeşil çatı sistemlerinin su ve enerji dengesi açısından değerlendirilmesi: İstanbul örneği*. İstanbul Üniversitesi Fen Bilimleri Enstitüsü, Doktora Tezi, İstanbul.
- Ekşi, M., & Uzun, A. (2016). Yeşil çatı sistemlerinin su ve enerji dengesi açısından değerlendirilmesi. *Journal of the Faculty of Forestry Istanbul University*, 66(1), 119-138.
- Fioretti, R., Palla, A., Lanza, L. G., & Principi, P. (2010). Green roof energy and water related performance in the Mediterranean climate. *Building and environment*, 45(8), 1890-1904.
- Getter, K.L. 2009. *Extensive green roofs: carbon sequestration potential and species evaluations*. PhD Thesis, Michigan State University, Michigan, USA.
- Kabuloğlu, K. S. (2005). Yeşil çatıların ekolojik yönden değerlendirilmesi, 2. *Ulusal Çatı Cephe Kaplamalarında Çağdaş Malzeme ve Teknolojiler Sempozyumu*. İstanbul, 1- 12.
- Karıptaş, F. S. (2010). Yeşil çatıların ekoloji bağlamında değerlendirilmesi ve Turkcell AR-GE binası örneği, 5. *Ulusal Çatı & Cephe Sempozyumu, Dokuz Eylül Üniversitesi Mimarlık Fakültesi Tınaztepe Yerleşkesi-İzmir*, 15-16.
- Kıyaslı, S., & Selçuk, E. (2023). Yeşil çatıların sürdürülebilirliğinin istanbul örneği üzerinden incelenmesi. *Mimarlık, Planlama ve Tasarım Alanında Uluslararası Araştırmalar III*, 133.
- Koç, Y. Ve Gültekin, A. B. (2010).Yeşil çatılar ve Türkiye’deki uygulamaları, 5. *Ulusal Çatı ve Cephe Sempozyumu, İzmir*, 4-9.

- Korol, E., & Shushunova, N. (2016). Benefits of a modular green roof technology. *Procedia Engineering*, 161, 1820-1826.
- Külekçi, E. A. (2017). Geçmişten günümüze yeşil çatı sistemleri ve yeşil çatılarda kalite standartlarının belirlenmesine yönelik bir araştırma. *ATA planlama ve Tasarım Dergisi*, 1(1), 35-53.
- Lazzarin, R. M., Castellotti, F., & Busato, F. (2005). Experimental measurements and numerical modelling of a green roof. *Energy and Buildings*, 37(12), 1260-1267.
- Liu, K. (2004). Sustainable building envelope – garden roof system performance, NRC – CNRC, RCI Building Symposium, Nov. 4-5, New Orleans, 1 - 14.
- Liu, K., & Minor, J. (2005). Performance evaluation of an extensive green roof. *Presentation at Green Rooftops for Sustainable Communities, Washington DC*, 1-11.
- Özbey, A. G. M. F., & Turhan, C. (2024). Yeşil binalarda yaprak alan analizi (la) ile bina enerji tüketimi arasındaki ilişkinin belirlenmesi.
- Parizotto, S., & Lamberts, R. (2011). Investigation of green roof thermal performance in temperate climate: A case study of an experimental building in Florianópolis city, Southern Brazil. *Energy and buildings*, 43(7), 1712-1722.
- Saadatian, O., Sopian, K., Salleh, E., Lim, C. H., Riffat, S., Saadatian, E., ... & Sulaiman, M. Y. (2013). A review of energy aspects of green roofs. *Renewable and sustainable energy reviews*, 23, 155-168.
- Shafique, M., Kim, R., & Rafiq, M. (2018). Green roof benefits, opportunities and challenges—A review. *Renewable and Sustainable Energy Reviews*, 90, 757-773.
- Tik, R., Kaya, T. (2024). Possibilities Of Using Renewable Energy Sources In Landscape Studies. International Congress Of High Value-Added Agricultural Products, December 1-3, Iğdır/Türkiye, 560-570.
- Tohum, N. (2011). Sürdürülebilir peyzaj tasarım aracı olarak yeşil çatılar. İstanbul Teknik Üniversitesi Fen Bilimleri Enstitüsü, Peyzaj Mimarlığı Anabilim Dalı, Yüksek Lisans Tezi, 91sy.
- Voyde, E., Fassman, E., & Simcock, R. (2010). Hydrology of an extensive living roof under sub-tropical climate conditions in Auckland, New Zealand. *Journal of hydrology*, 394(3-4), 384-395.
- Waldbaum, H. (2008). Green roofs for urban agriculture. *Dagenham: University of East London, School of Computing and Technology, MSc Architecture: Advanced Environmental and Energy Studies*.
- Weiler, S. ve Scholz-Barth, K. (2009). *Yeşil çatı sistemleri: yapı üzerindeki peyzajların planlanması, tasarımı ve inşasına yönelik bir kılavuz*. John Wiley & Sons.
- Weng, Q., Lu, D., & Schubring, J. (2004). Estimation of land surface temperature–vegetation abundance relationship for urban heat island studies. *Remote sensing of Environment*, 89(4), 467-483.
- Werthmann, C. (2007). *Green roof-a case study: michael van valkenburgh associates' design for the headquarters of the american society of landscape architects*. Princeton Architectural Press.

HOMEOPATHY IN PLANT PROTECTION

BİTKİ KORUMADA HOMEOPATI

Ezgi Günay SARGUT

Van Yüzüncü Yıl University, Institute of Science and Technology, Plant Protection
Department, Van, Turkey

ORCID ID: <https://orcid.org/0009-0003-5135-0844>

Reyyan Yergin ÖZKAN

Van Yüzüncü Yıl University, Faculty of Agriculture, Plant Protection Department,
Department of Phytopathology, Van, Turkey

ORCID ID: <https://orcid.org/0000-0003-2319-404X>

ABSTRACT

Introduction and Purpose: Homeopathy, which is used as a natural treatment method for the healing of acute and chronic diseases that apply to all living things, was introduced by Samuel Hahnemann, a German physician and pharmacist, in the early 19th century. Although the aim of homeopathy is to trigger the organism's own healing mechanisms and to put it into the healing process, its basic principle is based on the statement “A substance that produces certain findings in a healthy person provides improvement in sick people with similar findings.” Homeopathy, which is used as an alternative medicine technique in human health and animal health; Agrohmeopathy also gives successful results in plant diseases and pests in agriculture with the use of the correct remedy (homeopathic medicine). There are remedies used for the treatment of diseases caused by fungi, bacteria and viruses in plants, as well as remedies that have a repellent effect against plant pests and provide incentives such as easy germination and root development in plant development. The remedies used are natural, non-toxic and do not form residues in the soil, nor do they cause resistance gain in the plant itself. There are two sources for choosing the right remedy: the symptom guide and the remedy guide. Based on these two sources, the homeopath selects the most appropriate remedy that produces symptoms similar to the symptoms of the sick organism. In studies conducted in agrohmeopathy; remedies applied against 16 pests, 20 climatic causes, 13 environmental effects and 11 harmful pathogens have shown successful results. **Results:** With the increase in studies, remedies to be applied against different diseases and pests in agriculture will be revealed. This treatment method, which is still open-ended and will shed light on new studies, defines chemical pesticides used in good agricultural practices as poisons. Agrohmeopathy will be an economical and environmentally healthy treatment method in agriculture.

Key Words: Homeopathy, Plant Protection, Remedi, Agrohmeopathy

ÖZET

Giriş ve Amaç: Tüm canlılar için geçerli olan akut ve kronik hastalıkların iyileşmesi için doğal bir tedavi yöntemi olarak kullanılan homeopati; Alman doktor ve eczacı olan Samuel

Hahnemann tarafından 19. yy başlarında ortaya konulmuştur. Homeopatide amaç; organizmanın kendi iyileşme mekanizmalarını tetiklemek ve iyileşme sürecine sokmak olmakla beraber, temel prensibi; "Sağlıklı kişide belli bulgular ortaya çıkaran bir madde, benzer bulguya sahip hasta kişilerde iyileşme sağlar." ifadesine dayanmaktadır. İnsan sağlığında ve hayvan sağlığında alternatif bir tıp tekniği olarak kullanılan homeopati; Agrohhomeopati adı ile tarımda bitki hastalıkları ve zararlılarında da doğru remedi (homeopatik ilaç) kullanımı ile başarılı sonuçlar vermektedir. Bitkilerde fungus, bakteri ve virüslerin neden olduğu hastalıkların tedavisi amacı ile kullanılan remediler olduğu gibi, bitki zararlılarına karşı repellent etkiye sahip ve bitki gelişiminde kolay çimlenme, kök gelişimi gibi teşvikler sağlayan remediler de bulunmaktadır. Kullanılan remediler doğal, zehirsiz ve toprakta kalıntı oluşturmadığı gibi bitkinin kendisinde de dayanıklılık kazanımı gerçekleştirmemektedir. Doğru remedi seçimi için kullanılan; belirti rehberi ve ilaç rehberi olmak üzere iki kaynak vardır. Homeopat bu iki kaynaktan yola çıkarak hasta organizmanın semptomlarına benzer semptomları ortaya çıkaran en uygun remediye seçer. Agrohhomeopatide yapılan çalışmalarda; 16 zararlı, 20 iklimsel neden, 13 çevresel etki ve 11 zararlı patojene bağlı, bitki gelişiminde etkilenmeye karşı uygulanan remediler başarılı sonuçlar ortaya koymuştur. **Sonuç:** Çalışmaların artırılması ile tarımda farklı hastalık ve zararlılara karşı uygulanacak remediler ortaya konulacaktır. Henüz ucu açık, yeni çalışmalara ışık tutacak olan bu tedavi yöntemi iyi tarım uygulamalarında kullanılan kimyasal ilaçları zehir olarak tanımlamaktadır. Agrohhomeopati tarımda, ekonomik ve çevreye sağlıklı bir tedavi yöntemi olacaktır.

Anahtar Kelimeler: Homeopati, Bitki Koruma, Remedi, Agrohhomeopati

SIİRT PISTACHIO PRODUCERS' DEMOGRAPHIC CHARACTERISTICS AND PRODUCTION PROBLEMS: THE CASE OF SIİRT PROVINCE**Prof. Dr. Köksal KARADAŞ**

Iğdır Üniversitesi, Ziraat Fakültesi, Tarım Ekonomisi Bölümü, Iğdır, Türkiye

ORCID: 0000-0003-1176-3313

Dr. Öğr. Üyesi Osman Doğan BULUT

Iğdır Üniversitesi, Ziraat Fakültesi, Tarım Ekonomisi Bölümü, Iğdır, Türkiye

ORCID: 0000-0003-2682-6356

ÖZET

Antep fıstığının farklı bir çeşidi olan Siirt fıstığı tane yapısı ve tadındaki aroma nedeni ile genel olarak çerez şeklinde tüketilmektedir. Siirt'te yetiştirilen ve bölge ekonomisi için önem arz eden Siirt fıstığının ağaç başına verimi bölge illerindeki Antep fıstığı veriminin üçte birinden daha düşük seviyededir. Ürün verimini etkileyen önemli unsurlar arasında üreticilerin demografik özellikleri ve sorunları bulunmaktadır. Bu çalışmanın amacı Siirt fıstığı üreticilerinin demografik özellikleri ile üretim sorunlarını belirlemek ve bu sorunlara çözüm önerileri getirmektir. Bu amaçla Siirt İl Tarım ve Orman Müdürlüğü Çiftçi Kayıt Sistemi Verileri kullanılarak Basit Tesadüfi Örneklem Yığın Oran Tahmini Yöntemine göre belirlenen 93 adet Siirt fıstığı üreticisi ile yüz yüze yapılan anketlerden elde edilen veriler çalışmada kullanılmış olup % ve Frekans değerleri hesaplanmıştır. Çalışma sonunda ortalama değerler olarak işletme sahiplerinin yaşları 45, hane halkı sayısı 8, Erkek İş Birimi (EİB) 5,39 ve üreticilerin %76,3'ü İlk ve ortaöğretim düzeyinde eğitime sahiptir. Üreticilerin %56,6'sının tarım dışı gelirlerinin olduğu, %6,5'inin tarım sigortası yaptırdığı ve ortalama 88.8 da mülk araziye sahip oldukları tespit edilmiş olup Siirt fıstığı yanında buğday, arpa, sebze ve meyve üretimi yapmaktadırlar. Üretici sorunları arasında Siirt fıstığı üretiminde kullanılan girdilerin pahalı olması, iklimden kaynaklı verim düşüklükleri, ürün fiyatının yetersizliği gelmektedir. Devlet tarafından Siirt fıstığı üretiminin destekleme kapsamının genişletilmesi, üreticilere yetiştirme teknikleri konusunda eğitimler verilmesi ve ürün pazarlama kooperatifinin kurulması üretim ile ilgili sorunlara çözüm önerileri olarak tespit edilmiştir.

Anahtar Kelimeler: Siirt Fıstığı, Demografik Özellikler, Üretici Sorunları, Siirt.**ABSTRACT**

Siirt pistachio, a different type of pistachio, is generally consumed as a snack due to its grain structure and aroma. The yield per tree of Siirt pistachio, which is grown in Siirt and is important for the regional economy, is less than one third of the yield of pistachios in the region's provinces. The demographic characteristics of the producers and production problems are among the important factors affecting the product yield. The aim of this study is to determine the demographic characteristics and production problems of Siirt pistachio producers and to offer solutions to these problems. For this purpose, data obtained from face-to-face surveys with 93 Siirt pistachio producers determined according to the Simple Random Sampling Bulk Ratio Estimation Method using Siirt Provincial Directorate of Agriculture and Forestry Farmer

Registration System Data were used in the study and % and Frequency values were calculated. At the end of the study, it was determined that the average age of the farmers was 45, the number of household members was 8, the Male Business Unit (MBU) was 5.39 and 76.3% of the producers had primary and secondary education. It was determined that 56.6% of the producers had non-agricultural income, 93.5% did not have agricultural insurance, they owned an average of 88.8 acres of land and produced wheat, barley, vegetables and fruits in addition to Siirt pistachios. Producer problems include the high cost of inputs used in Siirt pistachio production, low yields due to climate conditions, and inadequate product prices. Expanding the scope of support for Siirt pistachio production by the state, providing training to producers on cultivation techniques and establishing a product marketing cooperative have been identified as solutions to production-related problems.

Keywords: Siirt Pistachio, Demographic Characteristics, Producer Problems, Siirt

GİRİŞ

Antepfıstığı yüksek besin değeri ve kendine özgü lezzeti ile tercih edilen ve birçok açıdan önem arz eden kabuklu bir meyvedir. Birçok tüketici Antep fıstığını çerez olarak tercih ederken, çikolata, tatlı ve yemeklerde de kullanılmakta ve benzer diğer meyvelerle karşılaştırıldığında kalori ve karbonhidrat bakımından ilk sırada yer almaktadır (Gezginç & Duman 2004). Zengin antioksidan barındırması nedeniyle insan beslenmesinde önemli bir yere sahip olan antepfıstığının içeriğinde %50 yağ, %20 protein ve %10 şeker bulunmakla birlikte, mineral ve potasyumca oldukça zengindir (Barghchi & Alderson, 1985; Bellomo & Fallico, 2007). Ayrıca Oleik asit (Omega 9), Linoleik asit (Omega 6) ve Linolenik asit (Omega 3) bakımından zengin olup kötü kolesterole, diyabete, kansızlığa ve bazı kanser türlerine iyi geldiği bilinmektedir (Demir ve Yücel, 2024).

Antep fıstığının bir çeşidi olan ve tadı, aroması ve verim özellikleri ile öne çıkan Siirt fıstığı bazı özellikleri ile Antep fıstığından ayrılmakta ve üretimde ve tüketimde daha çok tercih edilmektedir. Siirt fıstığının taneleri diğer fıstık çeşitlerine göre daha iri olup çıtlama oranı daha yüksekti ki bu durum daha çok kuru yemiş olarak tercih edilmesine sebep olmaktadır. Taze veya tuzlanmış olarak tüketilebilen ürün v veya çiğ olarak da tüketilebilmektedir. Kurağa karşı daha toleranslı olması daha az yağış alan alanlarda yetiştirilmesine olanak sağlamaktadır (TCEB, 2016). Diğer birçok çeşit %100 oranında periyodisite gösterirken Siirt fıstığı %50 oranında periyodisite göstermekte ve bu durum daha yüksek verim alınmasını sağlamaktadır (Açan ve Pakyürek, 2019; Akboğa ve Pakyürek, 2020).

2023 yılı itibarı ile dünyada 10213750 da alanda 1303461 ton fıstık elde edilmiş ve dekara verim 127 kg olarak belirlenmiştir (FAO, 2023). Dünya fıstık üretiminde ilk üç sırayı %51,9 ile ABD (6758550 ton), %23,6 ile İran (307886 ton) ve %13,5 ile Türkiye (176000 ton) almaktadır. Dünya verim sıralamasında ilk üç sırada ise ABD (361 kg/da), Irak (318 kg/da) ve Çin (311 kg) yer alırken Türkiye (42 kg/da) 16. sırada yer almaktadır.

Tablo 1. Antep Fıstığı Üretim ve Veriminde Öne Çıkan İller (2023 yılı)

	Üretim Miktarı (ton)	%		Ağaç Başına Verim (kg)
Şanlıurfa	59848	34,0	Nevşehir	16
Gaziantep	54575	31,0	Karabük	15
Siirt	32519	18,5	Antalya	14
Adıyaman	7592	4,3	Van	14
Kahramanmaraş	4325	2,5	Hatay	12
Batman	3440	2,0	Amasya	11
Mersin	3135	1,8	Mersin	9
Diyarbakır	2627	1,5	Hakkâri	7
Kilis	2531	1,4	Malatya	7
Mardin	1220	0,7	Siirt	4

Türkiye’de Antep fıstığı üretiminde %34’lük kısmı karşılayan Gaziantep (59848 ton) ilk sırada yer alırken Siirt %18,5 ile (32519 ton) 3. sırada yer almaktadır. Ağaç başına verimde ilk sıralarda Nevşehir (16 kg), Karabük (15 kg) ve Antalya (14 kg) gelirken Siirt 4 kg ile 17. sırada yer almaktadır (TÜİK, 2023).

Bölge ekonomisi için yüksek gelir getirisine sahip olan Siirt fıstığından yeterince verim alınmaması üreticilerin daha az kazanç elde etmelerine sebep olmakla birlikte bu durumun araştırılması hem bölge ekonomisi için hem de Siirt fıstığı üreticileri için önem arz etmektedir. Bu çalışmanın amacı bölge Siirt fıstığı üreticilerinin demografik özellikleri ve üretim ile ilgili sorunlarını belirlemek ve bu sorunlara çözüm önerileri getirmektir.


MATERYAL VE YÖNTEM

Araştırmanın ana materyalini 2021 yılında Siirt ilinde Siirt fıstığı yetiştiriciliği yapan 93 adet işletmeci ile yüz yüze yapılan anketlerden elde edilen veriler oluşturmaktadır.

Tablo 2. Siirt İlçelerinde Siirt Fıstığı İşletme Sayısı ve Ekim Alanı

İlçe	İşletme Sayısı	Ekilen Alan(da)	%
Baykan	513	14.612	8
Eruh	1284	34.999	19
Kurtalan	1450	65.676	35
Merkez	1238	32.962	18
Pervari	609	11.221	6
Şirvan	566	13.268	7
Tillo	401	14.274	8
Toplam	5942	187.015	100

Ayrıca araştırma ile ilgili yerli ve yabancı bilimsel çalışmalardan da faydalanılmıştır. Yukarıda Tablo 2. de Siirt İl Tarım ve Orman Müdürlüğü Çiftçi Kayıt Sistemi Verileri değerlendirilmiş ve Siirt’te fındık ekim alanının %35’inin gerçekleştirildiği Kurtalan İlçesi Gayeli Örnekleme Yöntemi ile çalışma alanı olarak seçilmiştir (Çiçek ve Erkan, 1996).



Şekil 1. Siirt İl Haritası

Siirt ili Türkiye'nin Güneydoğu Anadolu Bölgesi'nde $41^{\circ} 57'$ doğu boylamı ve $37^{\circ} -55'$ kuzey enlemi üzerinde yer almakta olup 895 m rakımda ve 2021 yılı nüfus sayımına göre 160340 kişi ilde bulunmaktadır (Anonim, 2023)

Aşağıda verilen Basit Tesadüfî Örnekleme Yığın Oran Tahmini Yöntemi kullanılarak anket sayısı hesaplanmıştır (Yamane, 2010).

$$n = \frac{Nxt^2 ppxq}{(N-1)xD^2 + t^2 ppxq}$$

n= örnek sayısı (90)

N=Küme büyüklüğü (1450)

D= Kabul edilen veya arzu edilen örnekleme hatası (0,1)

t= Tablo değeri (1,96)

p= Hesaplanması istenen oran (0,5)

q= 1-p (0,5)

$$n = \frac{1450x(1,69)^2 x(0.5)x(0.5)}{1449x(0.1)^2 + (1.69)^2 x(0.5)x(0.5)} = 69 \text{ örnek sayısı belirlenmiştir.}$$

Hesaplana anket sayısı eksik ve hatalı anketler olması ihtimaline karşı %30 artırılarak 90 adet anket yapılması planlanmış ve değerlendirme tamamlanan 93 anket üzerinden yapılmıştır. Ankete tabi tutulan işletmelerin tespitinde tesadüfî sayılar tablosu kullanılmıştır.

Anketlerde aşağıdaki konulara yönelik sorulara yer verilmiş olup anketlerden sağlanan bilgiler 2021 tarımsal üretim dönemini kapsamaktadır. Anketlerden elde edilen verilerin değerlendirilmesinde % ve frekans değerleri ile hesaplanmıştır. Arazi tasarruf şekilleri ve gelir miktarları (2021 ortalama döviz kuru 8,87 TL olarak alınmıştır (TCZM, 2022).

BULGULAR

Siirt ilinde bulunan ve Siirt fıstığı üretimi yapan 93 adet işletme ile yapılan anketlerin değerlendirilmesinde öncelikle demografik özellikler belirlenmeye çalışılmıştır. Nüfus,

tarımsal faaliyetin temel unsurlarından olan işgücünün kaynağını oluşturduğundan sosyal ve ekonomik nitelikleriyle ortaya konulmalıdır (Peker ve Ayyıldız, 1996). İlk olarak işletme sahiplerinin yaşları değerlendirilmiş en genç 29, en yaşlı 67, işletmecilerin ortalama yaşı 45 olarak belirlenmiştir. Aygün ve Gürsoy (2020) Siirt fıstığı üreticilerinin %80'inin 40 yaş üzerinde olduklarını, Akboğa & Pakyürek (2020) ise %30'unun 30-39 yaş aralığında olduğunu bildirmişlerdir.

Tablo 3. İncelenen İşletmelerde Nüfusun Yaş ve Cinsiyete Göre Dağılışı ve EİB

Yaş	Ortalama Birey sayısı	Katsayı	EİB
0-6	0,73	0	0
7-14 Kadın-Erkek	1,93	0,5	0,96
15-49 Kadın	1,39	0,75	1,04
15-49 Erkek	2,13	1	2,13
50-64 Kadın	0,45	0,5	0,23
50-64 Erkek	0,35	0,75	0,26
65 + Kadın-Erkek	0,19	0,5	0,09
Toplam	7,17	-	4,71

Üreticilerin %8,6'sı okuma yazma bilmezken %47,3'ü ilk ve orta öğretim düzeyinde, %37,6'sı ise Lise düzeyinde eğitime sahiptir (Tablo 4). Akboğa ve Pakyürek (2020) fıstık üreticilerinin %26,26'sının lise düzeyinde eğitime sahip olduklarını, Aygün ve Gürsoy (2020) fıstık üreticilerini eğitim düzeylerinin oldukça düşük olduğunu belirtmişlerdir.

Tablo 4. Üreticilerin Eğitim Seviyesi

Eğitim Seviyesi	n	%
Okuryazar değil	8	8,6
Okuryazar	8	8,6
İlkokul	19	20,4
Ortaokul	17	18,3
Lise	35	37,6
Ön lisans	5	5,4
Lisans	1	1,1
Toplam	93	100,0

Siirt fıstığı üreticilerinin %61'inin tarım dışında ek gelirleri bulunmakla birlikte bu gelir kaynaklarına sahip olanların en fazla sayıda olanlar esnaf (25,8) ve en az sayıda olanlar memur (%23,7) belirlenmiştir.

Tablo 5. Üreticilerin Tarım Dışı Gelir Miktarları Kaynakları

Eğitim Seviyesi	n	%	(\$)
Esnaf	24	25,8	687
Memur	22	23,7	564
Emekli	5	5,4	471
Özel Kurum	6	6,5	441
İşçi	4	4,3	490
Toplam	61	65,6	530,6 (Ort)

Gelir miktarı olarak ise en fazla aylık ortalama tarım dışı gelire sahip olanlar esnaflar (687 \$/ay) ve en az aylık ortalama tarım dışı gelire sahip olanlar özel kurumda çalışanlardır (441 \$/ay).

Genel olarak ise tarım dışı gelir miktarı 208-1667 \$ arasında değişmekte ve ortalama 386 \$ olarak belirlenmiştir.

Siirt fıstığı yetiştiricilerinin %6,5'i tarım sigortası yaptırmakta ve sigorta yaptıranların sigorta masrafları 580-1800 \$ arasında ve ortalama 916 \$ tarım sigortası masrafı yaptıklarını belirtmişlerdir. Üreticilerin %50,5'i pahalı olduğu için, %35,5'i gerek duymadığı için sigorta yapmadığını beyan etmiştir.

Tablo 6. Üreticilerin Sigorta Yaptırmama Nedenleri

Eğitim Seviyesi	n	%
Pahalı	47	50,5
Gerek duyulmadı	33	35,5
Bilgi sahibi değil	7	7,5
Toplam	87	93,5

Üreticiler 1250 da arasında olmak üzere ortalama 88,8 da mülk araziye sahip bulunmaktadırlar ve bu mülk arazilerin parsel sayısı 1-20 arasında değişmekte ve ortalama 5,8 adet parselde sahiptirler. Üretim deseni içerisinde alan olarak en fazla Siirt fıstığı (36,26 da), en az ise meyve-sebze (1,93 da) bulunmakla birlikte verim olarak buğday 330,5 kg/da, arpa 274,4 kg/da, Siirt fıstığı ise 88,8 kg/da ürün vermiştir. Bölge çiftçileri buğday verimi bakımından dünyadan (362,5 kg/da) daha düşük Türkiye'den (322 kg/da) daha yüksek, arpa verimi bakımından hem dünya (315,2 kg/da) hem Türkiye'den (280,6 kg/da) daha düşük verim elde ederken fıstık verimi bakımından dünyadan (127 kg/da) daha düşük Türkiye'den ise (42 kg/da) daha yüksek verim elde etmişlerdir (FAO, 2023).

Üreticiler en fazla geliri Siirt fıstığından elde ederken (24412 \$) en az geliri meyve-sebze üretiminden (203 \$) elde etmişler ve her işletmenin bitkisel üretimden ortalama 28965 \$ gelir elde ettiği belirlenmiştir. Bu durum bölge çiftçileri için Siirt fıstığının önemini gösterirken, Açı ve Pakyürek (2019) Siirt fıstığının verimlilik bakımından öne çıktığını, bölge ekonomisi için oldukça önemli olduğunu ve ayrıca üreticiler tarafından tercih edildiğini bildirmişlerdir.

Tablo 7. İşletmelerin Arazi Tasarruf Şekilleri ve Gelir Miktarları

Ürün Cinsi	Alan (da)	Elde Edilen Ürün (kg)	Verim (kg/da)	Satılanlar	
				Fiyat (\$)	Gelir (\$)
Buğday	36,3	12000	330,5	0,28	3360
Arpa	16,4	4500	274,4	0,22	990
Meyve Sebze	1,93	150,53	78	1,35	203
Siirt Fıstığı	39,26	3492,5	86,68	6,99	24412
Toplam	93,89	20143,03			28965

Üreticilere göre Siirt fıstığı üretim ve pazarlaması ile ilgili öncelikli sorunlar arasında ürünün bakım zorluğu ve masraflı olması ile girdilerin çok pahalı olması gelmektedir. Çözüm önerileri olarak bakım masrafları ve girdi fiyatlarının düşürülmesi, kooperatif kurulması, üreticilerin yetiştirme teknikleri konusunda eğitilmesi ve olumsuz hava koşullarına karşı tedbir alınması noktasında tarım sigortası yapılmasının desteklenmesi olarak belirlenmiştir. Kanber ve ark., (1993) fıstık yetiştiriciliğinde verim düşüklüğü ve periyodizitenin önemli sorunlar arasında olduğunu ve optimum verim için sulamanın gerekliliğini, Arpacı ve ark., (1995) Siirt fıstığı yetiştiriciliğinin sulu koşullarda yapılması gerektiğini, Külekçi & Aksoy (2011) Gaziantep'te antepfıstığı üreticilerine tarımsal yayım kapsamında teknik bilgi sağlanmasının verim artışı

sağlayacağını, Tahtacı (2012) Antep fıstığı ihracatında periyodisitenin önemli bir sorun olduğunu ve Siirt fıstığı çeşidi ile bu sorunun önlendiğini, Eldoğan & Şahin (2015) antepfıstığı üretimini geliştirmeye yönelik sosyal, teknik ve ekonomik çalışmalara ihtiyaç olduğunu, Ertürk ve ark., (2015) antepfıstığı üreticilerine verilen devlet desteğinin üretim miktarına göre verilmesinin üretim artışını sağlayacağını, Aydın ve Saltuk (2018) Siirt fıstığı üretiminin klasik yöntemlerle yapıldığını ve üreticilerin ilgili kurumların eğitimleri ile bilinç düzeyinin artırılması gerektiğini, Çelik (2019) Siirt fıstığında marka bilincinin ve stratejisinin oluşmadığını, Ağaç (2022) Siirt fıstığı yetiştiriciliği ile besi hayvancılığı faaliyetinin birlikte yapılmasının sürdürülebilirliğin sağlanması bakımından önemli olduğunu, Örük & Engindeniz (2022) fıstık üreticilerine girdi kullanımı ve modern üretim teknikleri konusunda eğitim verilmesi, üretici örgütlerinin oluşturulması ve su kaynaklarının artırılması gerektiğini, Pekitkan & Esgici (2022) fıstık tarımında en önemli sorunlardan birinin hasat işleminin mekanik olarak elle yapılması ve bu sebeple uzun sürmesi olduğunu bildirmişlerdir.

Tablo 8. Siirt Fıstığı Üretim-Pazarlama Sorunlar ve Çözüm Önerileri

Sorunlar	n	%	Çözüm Önerileri
Bakımı zor ve masraflı	39	41,94	Bakım masrafları düşürülmeli
Girdiler çok pahalı	34	36,56	Girdi fiyatları düşürülmeli
Fıstık fiyatları düşük	9	9,68	Kooperatif kurulmalı
İlaçların bilinçsiz kullanımı	8	8,60	Çiftçiler üretim konusunda bilgilendirilmeli
İklimden kaynaklı verim azlığı	3	3,23	Olumsuz hava koşulları için tedbir alınmalı (tarım sigortası)
Toplam	93	100,00	

SONUÇ

Siirt ilinde yaygın olarak faaliyette bulunan Siirt fıstığı işletmelerinde üreticiler ortalama 45 yaşında, hane halkı sayısı 7,17 ve üreticilerin yarısı ortaöğretim düzeyinde eğitime sahiptirler. Her işletme ortalama 530,6 \$ aylık tarım dışı gelire sahip olup üreticilerin yalnızca %6,5'i tarım sigortası yapmaktadır. Siirt fıstığı üreticisi Siirt fıstığına ilave olarak münavebeye dâhil ettikleri buğday ve arpa verimi açısından dünya verim ortalamasının altında verim elde etmişlerdir. Üreticiler Siirt fıstığı üretiminde bakımın zor ve masraflı olmasının yanında girdilerin pahalı olduğunu, ürünlerin düşük fiyattan satıldığını ve olumsuz iklim koşullarının verimi düşürdüğünü belirtmişlerdir. Üreticilerin daha düşük maliyetle Siirt fıstığı üretebilmeleri için girdi fiyatlarının düşürülmesi ve bu kapsamda ilgili resmi kurumların Siirt fıstığına verdikleri desteğin kapsamını genişletmeleri gerekmektedir. Ürün fiyatının yükseltilmesi noktasında ürün pazarlama kooperatiflerinin kurulması ve özellikle ihracat imkânlarının artırılması sağlanmalıdır. Olumsuz iklim koşullarından etkilenmenin en aza indirilebilmesi için tarım sigortasının önemi ilgili kurumlarca anlatılarak daha fazla sayıda üreticinin tarım sigortası yapması sağlanmalıdır. Ayrıca bilinçsiz girdi kullanımının önüne geçmek için üreticilere Siirt fıstığı ile ilgili yetiştirme teknikleri konusunda eğitimler verilmelidir.

KAYNAKLAR

- Açan, Y., Pakyürek, M. (2019). Siirt Fıstığı İşleme ve Değerlendirme Tesislerinin Durumu. UBAK Uluslararası Bilimler Akademisi, Uygulamalı Bilimler Tam Metin Kitabı, 543-554.
- Ağaç, M. (2022). Tarımda Yeniden Yapılanma ve Emegın Dönüşümü: Siirt Fıstığı Örneđi. *Politik Ekonomik Kuram*, 6(2), 422-453.
- Akbođa. A., Pakyürek. M. (2020). Siirt Fıstığı Yetiştiriciliğinde Üretici Davranışları. *ISPEC Tarım Bilimleri Dergisi*. 4(2)171-185.

- Arpacı, S., Akkök, F., Tekin, H. (1995). Sulu ve Kuru Koşullarda Yetiştirilen Antepfıstıklarında Gelişme ve Verim Değişimlerinin İncelenmesi. Türkiye II. Bahçe Bitkileri Kongresi. C:1, Sh.429-433, Adana.
- Aydın, Y., Saltuk, B. (2018). Siirt Yöresi Fıstık Yetiştiricilerinin Sulama Eğilimlerinin Belirlenmesi. *Süleyman Demirel Üniversitesi Ziraat Fakültesi Dergisi, 1. Uluslararası Tarımsal Yapılar ve Sulama Kongresi Özel Sayısı*:119-127.
- Aygün, M., Gürsoy, S. (2020). Antep Fıstığı (*Pistacia vera* L.) Üretimi Yapan İşletmelerin Tarımsal Mekanizasyon Düzeylerinin Belirlenmesi: Türkiye, Siirt İli Örneği. *Türkiye Tarımsal Araştırmalar Dergisi, 7(2)*: 136-142.
- Barghchi, M., Alderson, P. G. (1985). In vitro propagation of *Pistacia vera* L. and the commercial cultivars Ohadiand Kalleghochi. *Journal of Horticultural Science, 60(3)*, 423–430.
- Bellomo, M. G Fallico, B. (2007). Anthocyanins, chlorophyll and xanthophylls in pistachio nuts (*Pistacia vera*) of different geographic origin. *Journal of Food Composition and Analysis, 20(3-4)*, 352– 359.
- Çelik, M. (2019). *Uluslararası Siirt Fıstık Çalıştayı*. Siirt Üniversitesi, Ziraat Fakültesi. 29-30 Nisan 2019. Siirt. Erişim adresi <https://ziraat.siirt.edu.tr/haber-etkinlik/uluslararasi-siirt-fistigi-calistayi/796076905.html>
- Çiçek, E. Erkan, O. (1996). Tarım Ekonomisinde Araştırma ve Örnekleme Yöntemleri. Gaziosmanpaşa Üniversitesi, Ziraat Fakültesi Yayınları, No:12.
- Demir, D., Yücel, N. (2024). Kentsel Markalaşma Sürecinde Coğrafi İşaret Adayı Olarak Siirt Fıstığı. *Siirt Sosyal Araştırmalar Dergisi, 3(1)*, 42-58.
- Eldoğan, Ü., Şahin, A. (2015). Gaziantep İlinde Antepfıstığı Üretim Ekonomisi Araştırması. *KSÜ Doğa Bil. Derg.*, 18(3), 29-36.
- Ertürk, Y. E., Geçer, M. K., Gülsoy, E., ve Yalçın, S. (2015). Antepfıstığı Üretimi ve Pazarlaması. *Iğdır Üni. Fen Bilimleri Enst. Der.* 5(2): 43-62.
- FAO, 2023. Food and Agriculture Organization of the United Nations. Crops and Livestock Product. Erişim adresi <https://www.fao.org/faostat/en/#data/QCL>
- Gezginç, Y., Duman A. D, (2004). Antepfıstığı İşleme Tekniği ve Muhafazasının Kalite Üzerine Etkisi. *Gıda Dergisi, 29(5)*: 373-378.
- Kanber, R., Kırdı, C., Yazar, A., Önder, S., ve Köksal, H. (1993). Irrigation Response of old Pistachio (*Pistachio vera* L.) *Doğa-Tr.J. of Agricultural and Forestry, 17*, 659-671.
- Külelçi, M., Aksoy, A. (2011). Gaziantep İli Dağ ve Ova Köylerinde Antepfıstığı Üretim Maliyetlerinin Karşılaştırılması. *U. Ü. Ziraat Fakültesi Dergisi, 25(1)*, 41-51.
- Örük, G., Engindeniz, G. Y. (2022). Antepfıstığının Ekonomik Analizi: Siirt İli Örneği. Antepfıstığı Yetiştiriciliği. İksad Yayınevi. Ankara. Türkiye. 335-344.
- Peker, K., Ayyıldız, T. (1996). Pasinler İlçesi Tarım İşletmelerinde Atıl İşgücünün Tespiti ve Bu İşgücünü Değerlendirme İmkânları. *Tr. J. of Ariculture and Forestry 20*: 23-190.
- Pekitkan, F. G., Esgici, R. (2022). Siirt Fıstık (*Pistachio vera* L.) Çeşidinin Yük Altındaki Davranışının Belirlenmesi. *Tarım Makinaları Bilimi Dergisi, 18(3)*: 189-197.
- Tahtacı, S. A. (2012). Antep Fıstığında Geliştirilen Yeni Çeşitler. Tekin ve Bardak Yıldızı, *Antep Fıstığı Araştırma Dergisi, 1*, 6-7.
- TCEB, (2016). Antep Fıstığı- Sektör Raporları. Türkiye Cumhuriyeti Ekonomi Bakanlığı, İhracat Genel Müdürlüğü, Tarım Ürünleri Daire Başkanlığı, Ankara, 6s.

TCZM, (2022). Türkiye Cumhuriyeti Merkez Bankası. Gösterge Niteliğindeki Merkez Bankası Kurları. Erişim adresi https://www.tcmb.gov.tr/kurlar/kurlar_tr.html

TÜİK, (2023). Türkiye İstatistik Kurumu. Bitkisel Üretim İstatistikleri. Erişim adresi <https://biruni.tuik.gov.tr/medas/?kn=92&locale=tr>

Yamane, T. (2010). Temel Örnekleme Yöntemleri, Literatür Yayıncılık, İstanbul.

SOILLESS PLANT CULTURE TECHNICAL METHODS

TOPRAKSIZ BİTKİ KÜLTÜRÜ TEKNİK YÖNTEMLERİ

Dr. Öğr. Üyesi Berna BAŞ

Gaziantep Üniversitesi, Fen Edebiyat Fakültesi, Biyoloji Bölümü

ÖZET

Giriş ve Amaç: Dikey tarım (DT) kısaca topraksız bir ortamda, makro-mikro besin maddeleri içeren sulu solüsyonlarda bitki yetiştiriciliğidir. DT uygulaması teknik olarak, su-bazlı sistem ve substrat-bazlı sistem olarak iki ana sınıfa ayrılabilir. Su-bazlı sistemler temel olarak üç ana grupta incelenmektedir; hidroponik, aeroponik, akuaponik sistemler. Substrat-bazlı sistemler hidroponik uygulamalarda kullanılmakta olup katı, inert bir maddeyle besinlerin köklerden absorpsiyonu kolaylaştırmakta ve bitki köklerine mekanik destek sağlanmaktadır. Yani topraksız ortamda çeşitli substratlar kullanılarak toprağı taklit eden mini-çevre yaratılmaktadır. Bu substratların organik, inert ve kompost olmak üzere çeşitli tipleri bulunmaktadır. Hidroponik kültürde yetiştirilecek bitkilerin kökleri özel besin ihtiyaçlarına göre belli formülasyonla hazırlanan sıvı solüsyona daldırılır. Bu sistemle az miktarda su kullanarak birim alandan yüksek verim alınmaktadır. Aeroponik sistemlerde hemen hemen su bulunmayan bir ortamda bitkilere besin solüsyonu kontrollü bir şekilde püskürtülerek yetiştirilmektedir. Hidroponik sisteme göre daha az miktarda su ve gübre kullanarak bitkiler daha hızlı büyümekte ve bitkilerin besin içeriğı daha yüksek olmaktadır. Akuaponik kültür ise sürdürülebilir ve sinerjiktir, balık kültürünün hidroponikle bütünleştiğı bir sistemdir. Besin açısından zengin balık atıkları bitkiler için besin sağlarken, bitkilerde balıklar için suyu filtreleyerek kapalı devre bir ekosistem yaratılmaktadır. Bütün sistemlerin kurulum maliyetinin yüksekliğı, stabilitenin sürekliliğı, yüksek derecede düzenli bakım gerektirmesi önemli dezavantajlarıdır. Mikrobiyel kontaminasyon ve pest kontrolü diğer önemli bir sorundur. Topraktaki doğal mikrobiyel aktiviteler topraksız ortamda olmadığından dolayı mikroorganizmaların ve zararlıların hızla yayılabilme olasılığı vardır. Bu derleme de topraksız tarım tekniklerinin sürdürülebilirlik açısından avantajları ve dezavantajları karşılaştırılarak değerlendirilmiştir.

Materyal ve Metot: Google Akademik uygulama üzerinden çeşitli yayınlar derlenmiştir. **Sonuç:** Başlangıç kurulum maliyeti yüksek olan topraksız bitki yetiştiricilik tekniklerinden aeroponik sistem daha az su-besin kullanımından dolayı otsu bitkiler için uygundur. Ancak hidroponik sistem daha fazla besin ve su tedarik etmesine rağmen bitki kökleri sürekli sıvı besin solüsyonu içinde olduğu için ihtiyaç duyulan oksijene engel olmaktadır. Dolayısıyla hastalık kontaminasyonu ve hızla yayılma riski de yüksektir. Akuaponik yetiştiricilikte besin solüsyonunda balıklarda mevcut olduğu için stabilitenin korunması için daha fazla dikkat gösterilmesi gerekmektedir.

Anahtar Kelimeler: Dikey Tarım Teknikleri, Hidroponik, Aeroponik, Akuaponik, Substrat-Bazlı Sistem

ABSTRACT

Introduction and Purpose: Vertical farming (VF) is briefly the cultivation of plants in aqueous solutions containing macro and micro nutrients in a soilless environment. VF application can be technically divided into two main classes as water-based system and substrate-based system. Water-based systems are basically examined in three main groups; hydroponic, aeroponic, aquaponic systems. Substrate-based systems are used in hydroponic applications and the absorption of nutrients from the roots is facilitated with a solid, inert substance and mechanical support is provided to the plant roots. In other words, a mini-environment that mimics the soil is created by using various substrates in a soilless environment. These substrates have various types including organic, inert and compost. In hydroponic culture, plant roots are immersed in a nutrient solution prepared in certain formulated according to the specific requirements of the plants. This system obtains high yields from the unit area using a small amount of water. In aeroponics systems, the nutrient solution is sprayed to the plants in a controlled manner in an environment where there is almost no water. Compared to the hydroponic system, plants grow faster and the nutrient content of the plants is higher by using less water and fertilizer. Aquaponic culture is a sustainable and synergistic system that integrates fish farming (water culture) with hydroponics. While nutrient-rich fish waste provides nutrients for the plants, plants filter the water for the fish and create a closed-circuit ecosystem. Important advantages are the high installation cost of all systems, the need for continuous stability and high levels of regular maintenance. Microbial contamination and pest control are another important problem. Since natural microbial activities in the soil are not in a soilless environment, there is a possibility that microorganisms and pests can spread rapidly. In this review, the advantages and drawbacks of soilless farming techniques in terms of sustainability were compared and evaluated.


Material and Method: Various publications were compiled through the Google Scholar application. **Conclusion:** Aeroponic system, which is one of the soilless plant cultivation techniques with high initial installation costs, is suitable for herbaceous plants due to less water-nutrient use. However, although the hydroponic system provides more nutrients and water, it prevents the necessary oxygen as the plant roots are constantly in the liquid nutrient solution. Therefore, the risk of disease contamination and rapid spread is also high. In aquaponic farming, more care must be taken to maintain stability since the fish are present in the nutrient solution.

Key Words: Vertical Farming Techniques, Hydroponics, Aeroponics, Aquaponics, Substrate-Based System

GİRİŞ

Bitki biyoteknolojisinde inovatif tekniklerden olan dikey tarım (DT) yeni bir yöntem olmamakla beraber son yıllarda ticari olarak önem kazanmaya başlamıştır. Özellikle dünya genelinde su kaynaklarının azalması, konvansiyonel tarım girdilerinin ticari amaçlı aşırı kullanım sonucu insan eliyle ekosistem kirliliğinin artırılması, tarım alanlarının azalması, nüfus artışı, gıda güvencesi geleceğinin riske girmesi tarımsal devrimi zorunlu kılmaktadır. Topraksız bir ortamda ve kontrollü koşullarda bitki kültürü teknikleriyle bitki yetiştiriciliğinin bazı avantajları da dezavantajları da bulunmaktadır. Ürün verimliliğini artırması, su ve besin maddelerini veya gübreyi etkin bir şekilde kullanması, çevresel etkinin azalması, yeni istihdam olanakları yaratması gibi fırsatlar sunan DT, sentetik gübrelerin kullanımının yaratacağı sağlık risk ihtimalleri, yüksek enerji tüketim maliyeti, yüksek teknolojik donanım gerekliliği, uzun süreli stabilitenin sağlanması gibi bazı dezavantajlara sahiptir (Rajan, Lada & MacDonald, 2019; Klerx & Rose, 2020; Torres & ark., 2025). DT uygulaması teknik olarak, su-bazlı sistem ve substrat-bazlı sistem olarak iki ana sınıfa ayrılabilir. Su-bazlı sistemler temel olarak üç ana grupta incelenmektedir; hidroponik, aeroponik, akuaponik sistemlerdir. Substrat-bazlı sistemler hidroponik uygulamalarda kullanılmakta olup katı, inert bir maddeyle besinlerin köklerden

absorbsiyonu kolaylaşmakta ve bitki köklerine mekanik destek sağlanmaktadır. Yani topraksız ortamda çeşitli substratlar kullanılarak toprağı taklit eden mini-çevre yaratılmaktadır. Her üç yöntemin temel modüler yapısı Şekil 1 'de verilmiştir. Topraksız dikey tarımda uygulanan yöntemler dijital teknolojiler, sürdürülebilir gıda üretimi ve biyolojik güçlendirmeye bağlantılıdır (Appicciutoli & ark., 2025). Bu derleme de topraksız dikey tarım tekniklerinin sürdürülebilirlik açısından avantajları ve dezavantajları karşılaştırılarak değerlendirilmiştir.



Şekil 1. Dikey çiftçilik sistemlerinin temel modüler yapısı. Resim yazar tarafından BioRender programı kullanılarak oluşturulmuştur

SU-BAZLI KÜLTÜR SİSTEMLERİ

Hidroponik Sistem

Su-bazlı sistemlerde bitkiciklerin kökleri doğrudan besin solüsyonuna daldırılmaktadır, substrat-bazlı sistemlerde organik veya inorganik çeşitli solid materyaller içeren saksılarla kökler dolaylı olarak besin solüsyonuna daldırılmaktadır (Al-Kodmany, 2016). Hidroponik sistem de bitkiler topraksız, besin maddelerince zengin solüsyonlarda kültüre alınmaktadır (Szekely & Jijakli, 2022). İnce film tekniği, derin su kültürü gibi çeşitli sistemlere sahip olan hidroponik yöntemlerden katı-faz sistemi kökler için daha iyi aerasyon ve oksijen sağlamaktadır (Vinci & Rapa, 2019). En önemli avantajlarından biri topraklı tarımda karşılaşılan toprak kökenli hastalık ve zararlı riski elimine edilmekte veya indirgenmektedir (Fussy & Papenbrock, 2022). Ancak besin solüsyonun herhangi bir fitopatojenle kontaminasyonu kısa sürede tesisteki tüm bitkileri de etkileyecektir. Diğer sistemlere göre hidroponik yetiştiricilik daha az bakım ve emek gerektirmektedir, çevresel faktörlerin yönetimi daha kolaydır, su ve besin solüsyonunun dağılımı homojen olduğu için daha iyi verim elde edilebilir (Al-Kodmany, 2018).

Aeroponik Sistem

Aeroponik sistemde raflar üzerine yerleştirilen köpük vb. bir zemin üzerindeki deliklere bitkiler dikilerek kök boğazı bölgesinden sıkıştırılır ve aeroponik çembere yerleştirilir (Stoner & Clawson, 1997-1998). Hidroponik sistemin teknolojisinden daha iyi bir otomasyon düzenine sahip olan aeroponik sistemde su doğrudan kullanılmamaktadır, bitkilerin aşağı doğru sarkan köklerine su besin solüsyonuyla beraber spreyle veya sisleme yoluyla püskürtülmektedir (Al-

Kodmany, 2018). Böylece besin maddelerinin absorpsiyonu artmakta ve bitkiler daha hızlı gelişmektedir, kökler için daha fazla oksijenli çevre yaratılmaktadır (Stegelmeier & ark., 2022; Green.org, 2025). Bu sistemle bitki kökleri boşlukta olduğundan ve birbirleriyle temas etmediğinden dolayı hastalık ve zararlı görüldüğünde ilgili bitki hemen uzaklaştırılıp imha edilerek hastalığın tüm tesise yayılması erkenden önlenmiş olacaktır. Son yapılan çalışmalarda yeni geliştirilen plazma-destekli aeroponik sistemle su ve havanın azotu birleştirilerek nitrat, nitrit, amonyum, hidroksil iyonları gibi reaktif türler üretilmektedir (Qureshi & ark., 2025). Bu türev moleküller sistemde ki besin maddelerinin bulunabilirliğini artırarak köklerden besin emilimini ve verimliliği yükseltmektedir, böylece geleneksel azot gübrelerine bağımlılığında azaltılabileceği rapor edilmiştir.

Akuaponik Sistem

Akuaponik sistem, bitki ile balık arasındaki simbiyotik ilişkiye dayanarak balık çiftliklerindeki akuakültür ile hidroponik sistemin birleştirilmesiyle geliştirilmiştir (Al-Kodmany, 2018). Besin açısından zengin balık atıkları bitkilere besin sağlarken, bitkiler de balıklar için suyu filtreleyerek kapalı devre mini bir ekosistem yaratılmaktadır.

Ele alınan üç sistemin karmaşıklığı ve ölçeklenebilirlik performansı karşılaştırıldığında hidroponik sistem kurulumu ve tesis kapasitesinin artırılması nispeten kolaydır, aeroponik sistem daha fazla teknik deneyim ve yatırım gerektirmekle beraber büyüme potansiyeli daha yüksektir, akuaponik sistem ise karmaşıktır ancak bütüncül ve sürdürülebilir bir yaklaşıma sahiptir (Green.org, 2025).

SUBSTRAT-BAZLI KÜLTÜR SİSTEMLERİ

Substrat-bazlı sistemlerde bitki fideleri katı bir materyale dikilerek kök boğazından sabitlenmekte ve böylece köklere mekanik destek sağlanmakta olup besin maddeleri - su - O₂ için kökler tarafından yavaşça absorbe edilme olanağı yaratılmaktadır (Vinci & Rapa, 2019). Substrat olarak kullanılan materyaller fiziksel olarak topraktan daha küçük partiküllerdir ve besin solüsyonu ile oksijen için boş alan yaratarak bitkilerin gelişimine önemli katkı sağlanmaktadır (Rodríguez-Delfín, 2012). Substrat olarak tek tip materyalde tercih edilebilir ya da nem, pH, elektrik iletkenliği gibi fiziko-kimyasal özellikleri farklı materyallerin karışımı şeklinde farklı bakterilerin gelişimi için özel nişlerin oluşumunu destekleyecek şekilde kompost olarak da hazırlanabilmektedir (Grunert & ark., 2016). Substratların özellikleri havalandırma, besin ve su tutma/dağıtma kapasitesi, çürüme-ayırma, fungal-bakteriyel patojenlerin gelişimine direnç gösterme gibi niteliklere bağlı olarak inert veya inorganik, organik, kompost gibi farklı tiplere sahiptirler (Olympios, 1992; Kang, Lee & Kim, 2001; Arif & ark., 2022; Younis & ark., 2022; Zhao & ark., 2022; Subramanian & ark., 2024). Substrat tercihinde DT ile yetiştirilecek bitkiler, substratın bedeli, transport pahası, kolay ulaşılabilirliği vb. değişkenler dikkate alınabilir (Domeño, Irigoyen & Muro, 2011). Ancak her bir substratın avantajları ve dezavantajları farklı olduğu için substrat seçiminde özel veya standart bir kriter bulunmamaktadır.

Perlit, vermikülit, taş yünü gibi *inert substratlar* besin sıvısıyla reaksiyona girmez ve drenaj, aerasyon, su taşkınlıklarına engel olma gibi özellikleri nedeniyle bitki köklerinin sağlıklı gelişimini desteklemekte, hafif olmalarından dolayı kullanımları kolay olup besin maddesi olmadıkları için de çözültideki besin maddeleri ve pH 'nın kontrol edilmesi gerekmektedir (Olympios, 1992; Sonneveld & ark., 2009). Taş yününde yetiştirilen salatalık kökleri genellikle *Pseudomonas* spp. türleri tarafından kolonize edilmekte ve bu kolonizasyon salatalık köklerindeki eksudatların miktarını sınırlayarak bir toprak patojeni olan *Pythium aphanidermatum* 'a karşı antagonistik etki göstermektedir (Postma & ark., 2000). Hindistan cevizi lifi, turba yosunu, hayvansal atıklar gibi doğal substrat kaynakları inert substratlara göre

daha sürdürülebilir özelliktedir, su tutma kapasiteleri daha yüksektir, hindistan cevizi lifinin fungal-bakteriyel patojenlerin gelişimine direnç gösterme özellikleri substrat olarak üstünlük sağlamakta ancak organik materyaller zamanla çürüdüğü için substratın da yapısal bütünlüğü bozularak su-besin tutma kapasitesi zayıflamaktadır (Ferreira & ark., 2020; Younis & ark., 2022).

Diğer substrat örneklerine göre hindistan cevizi lifi, turba yosunu gibi **organik substratlar** 'ın en önemli özelliklerinden biri de kation değişim kapasitesine sahip olmalarıdır (Abad & ark., 2002). Organik substratlar inert olanlara göre daha yoğun yapılı olduğu için uygun bir ayarlama yapıldığında bitki kökleri için yeterli aerasyon sağlayabilirler (Chan & ark., 2009). Ancak organik substratlar perlit, kaba kum gibi malzemelerle karıştırılarak aerasyon kapasitesi iyileştirilmekte dolayısıyla köklerin besin alım özelliği artırılarak daha güçlü kök gelişimi desteklenmektedir (Chan & ark., 2009; De Castro & ark., 2022). Bu şekilde farklı substratların karışımıyla elde edilen **kompost substratlar** besinlerin verimli kullanımı ve kök sağlığı üzerinde hassas bir kontrole sahip olduğu için ticari hidroponik tesislerde özellikle tercih edilmektedir (Rajendran & ark., 2024). Komposit substratların amacı da her bir substratın dezavantajını azaltmak, bitkilerin ihtiyaç anında besin maddelerinin mevcudiyetini artırmak ve kök bölgesi nemini muhafaza etmektir (Zhao & ark., 2022). Ağaç kabuklarından hazırlanan kompost materyaller yeşil atık kompostuna kıyasla daha fazla CO₂ üreterek mikrobiyel aktiviteyi de etkileyebilmektedir (Van Gerrewey & ark., 2020). Kısacası hidroponik substrat-bazlı sistemlerde dengeli bir çevre yaratmak için substrat seçiminde ince ayarlamalarla bitki köklerinin su-besin-oksijen ihtiyacı optimize edilerek bitkisel üretim verime olumlu katkı sağlanabilir.

SONUÇ

Ele alınan derlemede DT yöntemlerinin uygulama metotları karşılaştırılarak incelenmiştir. Akuaponik sistemle iki farklı besin kaynağı, hem balık hem bitki yetiştiriciliği yapılacağı için daha ideal bir model sunulmakla beraber konuyla ilgili deneyimli eleman yetersizliğinden dolayı ilk başlangıç adımı olarak uygulama kolaylığı açısından hidroponik sistem tercih edilebilir. Yapay zeka ve IoT (Internet of Things) sistemleriyle proaktif çalışmalar yapılarak sistem kurulumu, işletimi, yönetimi ve sürdürülebilirliğiyle ilgili karar süreci kolaylaştırılabilir. Toplumların beslenme alışkanlıkları dikey yetiştiriciliğin yönünü belirlemede kritik rol oynayacaktır. Her bir sistemin avantajları ve dezavantajları bulunmakla beraber çiftçiler, araştırmacılar, politikacılar ve özellikle ticari girişimcilerin işbirliğiyle DT toplumlarda daha kolay kabul görebilir. DT özellikle tarımda dışa bağımlı ülkelerde, su kaynakları yetersiz olan ve tarıma elverişli olmayan coğrafyalarda dış bağımlılığı azaltıcı ve kendi kendine yeterliliği destekleyici potansiyele sahiptir. Halen maliyeti pahalı olan bir teknik yöntem olan DT doğal kaynakları koruyarak ve çevre kontrolünü sağlayarak, gıda güvencesinin geleceğini garantileyerek sürdürülebilir bir tarım sistemine katkı sunabilir. Sistemin sorunları ve dezavantajları gelişen teknolojilerin entegrasyonu ile çözülecek ve DT 'ı daha avantajlı yapacaktır. Enerji maliyeti yüksek olan dikey tarımla ülkemizde çeşitli ticari işletmelerde üretim yapılmaktadır. Ancak ülkemizde elektrik üretiminin önemli kısmı kömür ve hidroelektrik santralleriyle gerçekleştirildiği için çevre ve azalan su kaynaklarını korumak için alternatif doğal kaynaklarla enerji üretimine yönelmediğimiz sürece dikey üretimin yaygınlaşması zor görünmektedir.

KAYNAKLAR

Abad, M., Noguera, P., Puchades, R., Maquieira, A., & Noguera, V. (2002). Physico-chemical and chemical properties of some coconut coir dusts for use as a peat substitute for containerised ornamental plants. *Bioresource technology*, 82(3), 241-245.

- Al-Kodmany, K. (2016). Sustainable tall buildings: cases from the global south. *Archnet-IJAR: International Journal of Architectural Research*, 10(2), 52.
- Al-Kodmany, K. (2018). The vertical farm: A review of developments and implications for the vertical city. *Buildings*, 8(2), 24.
- Appicciutoli, D., Amici, A. S., Bentivoglio, D., Chiaraluce, G., Staffolani, G. & Finco, A. (2025). Cultivating the Future: A Bibliometric Review of Emerging Trends in Soilless Farming. *Horticulturae*, 11(2), 140.
- Arif, Z. U., Khalid, M. Y., Sheikh, M. F., Zolfagharian, A. & Bodaghi, M. (2022). Biopolymeric sustainable materials and their emerging applications. *Journal of Environmental Chemical Engineering*, 10(4), 108159.
- Chan, Y. J., Chong, M. F., Law, C. L. & Hassell, D. G. (2009). A review on anaerobic–aerobic treatment of industrial and municipal wastewater. *Chemical engineering journal*, 155(1-2), 1-18.
- De Castro, J., Hill, R. D., Stasolla, C. & Badea, A. (2022). Waterlogging stress physiology in barley. *Agronomy*, 12(4), 780.
- Domeño, I., Irigoyen, I. & Muro, J. (2011). Comparison of traditional and improved methods for estimating the stability of organic growing media. *Scientia horticulturae*, 130(1), 335-340.
- Ferreira, V., Elozegi, A., D. Tiegs, S., von Schiller, D. & Young, R. (2020). Organic matter decomposition and ecosystem metabolism as tools to assess the functional integrity of streams and rivers—a systematic review. *Water*, 12(12), 3523.
- Fussy, A. & Papenbrock, J. (2022). An overview of soil and soilless cultivation techniques—chances, challenges and the neglected question of sustainability. *Plants*, 11(9), 1153.
- GREEN.ORG, (2025). <https://green.org/2024/01/30/hydroponics-vs-aeroponics-vs-aquaponics/> (Son erişim tarihi: 25.02.2025)
- Grunert, O., Hernandez-Sanabria, E., Vilchez-Vargas, R., Jauregui, R., Pieper, D. H., Perneel, M., Van Labeke, M. C., Reheul, D. & Boon, N. (2016). Mineral and organic growing media have distinct community structure, stability and functionality in soilless culture systems. *Scientific reports*, 6(1), 18837.
- Kang, J. Y., Lee, H. H. & Kim, K. H. (2001). Physical and chemical properties of inorganic horticultural substrates used in Korea. In *International Symposium on Growing Media and Hydroponics 644*, (237-241).
- Klerkx, L. & Rose, D. (2020). Dealing with the game-changing technologies of Agriculture 4.0: How do we manage diversity and responsibility in food system transition pathways?. *Global Food Security*, 24, 100347.
- Olympios, C. M. (1992). Soilless media under protected cultivation rockwool, peat, perlite and other substrates. In *Symposium on Soil and Soilless Media under Protected Cultivation in Mild Winter Climates 323*, (215-234).
- Postma, J., Willemsen-de Klein, M. J. & van Elsas, J. D. (2000). Effect of the indigenous microflora on the development of root and crown rot caused by *Pythium aphanidermatum* in cucumber grown on rockwool. *Phytopathology*, 90(2), 125-133.
- Qureshi, W. A., Gao, J., Elsherbiny, O., Mosha, A. H., Tunio, M. H. & Qureshi, J. A. (2025). Boosting Aeroponic System Development with Plasma and High-Efficiency Tools: AI and IoT—A Review. *Agronomy*, 15(3), 546.
- Rajan, P., Lada, R. R. & MacDonald, M. T. (2019). Advancement in indoor vertical farming for microgreen production. *American Journal of Plant Sciences*, 10(08), 1397.

- Rajendran, S., Domalachenpa, T., Arora, H., Li, P., Sharma, A. & Rajauria, G. (2024). Hydroponics: Exploring innovative sustainable technologies and applications across crop production, with Emphasis on potato mini-tuber cultivation. *Heliyon*, *10*(5), e26823.
- Rodríguez-Delfín, A. (2012). Advances of hydroponics in Latin America. In *II International Symposium on Soilless Culture and Hydroponics 947* (pp. 23-32).
- Sonneveld, C., Voogt, W., Sonneveld, C. & Voogt, W. (2009). Substrates: Chemical characteristics and preparation. In *Plant nutrition of greenhouse crops*, 227-256.
- Stegelmeier, A. A., Rose, D. M., Joris, B. R. & Glick, B. R. (2022). The use of PGPB to promote plant hydroponic growth. *Plants*, *11*(20), 2783.
- Stoner, R.J. & J.M. Clawson (1997-1998). *A High Performance, Gravity Insensitive, Enclosed Aeroponic System for Food Production in Space*. Principal Investigator, NASA SBIR NAS10-98030.
- Subramanian, P., Gupta, A., Gopal, M., Selvamani, V., Mathew, J., Surekha & Indhuja, S. (2024). Coconut (*Cocos nucifera* L.). In *Soil Health Management for Plantation Crops: Recent Advances and New Paradigms*, (37-109). Singapore: Springer Nature Singapore.
- Szekely, I. & Jijakli, M. H. (2022). Bioponics as a promising approach to sustainable agriculture: A review of the main methods for producing organic nutrient solution for hydroponics. *Water*, *14*(23), 3975.
- Torres, J. D., Jaeger, S. R., Puerta, P. & Tárrega, A. (2025). How do Spanish consumers perceive different lettuce cultivation systems? Insights from explicit and implicit methods. *Applied Food Research*, *5*(1), 100709.
- Van Gerrewey, T., Ameloot, N., Navarrete, O., Vandecruys, M., Perneel, M., Boon, N. & Geelen, D. (2020). Microbial activity in peat-reduced plant growing media: Identifying influential growing medium constituents and physicochemical properties using fractional factorial design of experiments. *Journal of Cleaner Production*, *256*, 120323.
- Vinci, G., & Rapa, M. (2019). Hydroponic cultivation: life cycle assessment of substrate choice. *British Food Journal*, *121*(8), 1801-1812.
- Younis, A., Ahsan, M., Akram, A., Lim, K. B., Zulfiqar, F. & Tariq, U. (2022). Use of organic substrates in sustainable horticulture. *Hasanuzzaman, M., Hawrylak Nowak, B., Islam, T. and Fujita M. Biostimulants for Crop Production and Sustainable Agriculture*, 122-138.
- Zhao, Z., Xu, T., Pan, X., Susanti, White, J. C., Hu, X., Miao, Y., Demokritou, P. & Ng, K. W. (2022). Sustainable nutrient substrates for enhanced seedling development in hydroponics. *ACS Sustainable Chemistry & Engineering*, *10*(26), 8506-8516.

PESTS OF GREENHOUSE VEGETABLE GROWING IN IĞDIR PROVINCE

IĞDIR İLİ ÖRTÜ ALTI SEBZE YETİŞTİRİCİLİĞİ ZARARLILARI

Öğr. Gör. Nagihan KILIÇ

Iğdır Üniversitesi, Teknik Bilimler Meslek Yüksekokulu Bitkisel ve Hayvansal Üretim
Bölümü Seracılık Programı

Prof. Dr. Celalettin GÖZÜAÇIK

Iğdır Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü,

ABSTRACT

This study, conducted in the greenhouses of Iğdır Province's central villages (Obaköy, Küllük, Evcı) during the years 2023-2024, aimed to identify and determine the definition and detection of vegetable pests. Sampling was carried out at regular 15-day intervals throughout the growing season, from May to July. A total of 50-100 samples were taken from cucumber, tomato, pepper, squash, and bean plants grown in plastic greenhouses. In all the greenhouses where sampling was conducted, it was determined that the vegetables were infected with pests at some point during the season. Among the pests commonly found in the greenhouses in the examined villages, whitefly (*Bemisia tabaci* Gennadius), aphids (Aphididae), thrips (Thripidae), red spider mite (*Tetranychus urticae* Koch), and leaf miners (*Liriomyza* spp.) were identified in cucumber, pepper, squash, and beans. Additionally, the tomato leaf miner (*Tuta absoluta* Meyrick) was observed in tomatoes along with these pests. The pest population density peaked towards the end of the growing season. However, the tomato leaf miner can cause damage throughout all growth stages, especially from the seedling stage to harvest. The most critical damage occurs during flowering and fruit formation because the tomato leaf miner enters the fruit and causes it to rot, which deteriorates the product's quality and reduces its market value. Besides, aphids mostly damage young shoots, leaves, and during the flowering stage. These pests cause leaves to yellow, negatively affect growth, and, through their feeding and virus vectoring, lead to the rapid spread of these diseases. To successfully manage these pests, it is essential to have a thorough understanding of their identification and life cycle.

Keywords: Greenhouse, vegetables, pests

ÖZET

Iğdır ili merkez köylerde (Obaköy, Küllük, Evcı) sebze üretimi yapılan seralarda bulunan sebze zararlılarının tanımı ve tespitinin belirlenmesi üzerine yapılan bu çalışma 2023-2024 yıllarında yürütülmüştür. Örneklemeler Mayıs-Temmuz ayları arasında yetiştirme mevsimi boyunca 15'er günlük düzenli periyotlarla yapılmıştır. Çalışmada plastik seralarda yetiştirilen hıyar, domates, biber, kabak ve fasulye bitkilerinden toplam 50-100 arası örnekleme yapılmıştır. Örnekleme yapılan seraların tümünde sezonun herhangi bir döneminde sebzelerin zararlılarla bulaşık olduğu tespit edilmiştir. İncelenen köylerdeki seralarda yaygın olarak görülen zararlılar arasında hıyar, biber, kabak ve fasulyede beyaz sinek (*Bemisia tabaci* Gennadius) (Hemiptera: Aleyrodidae), yaprak bitleri (Hemiptera: Aphididae), thripsler (Tyhsanoptera: Thripidae),

kırmızı örümcek (*Tetranychus urticae* Koch) (Acarina: Tetranychidae), galerisineklere (*Liriomyza* spp.) (Diptera: Agromyzidae), Yeşilkurt (*Helicoverpa armigera* Hubn.) (Lepidoptera: Noctuidae) tespit edilmiş. Domateste de ise bu zararlılara ek olarak domates güvesi (*Tuta absoluta* Meyrick) (Lepidoptera: Gelechiidae) yoğun olarak görülmüştür. Zararlı böceklerin yoğunlukları vejetasyon sonuna doğru maksimuma ulaşmıştır. Ancak Domates güvesi, özellikle fide döneminden hasat sonuna kadar tüm gelişim evrelerinde zarar verebilir. Ancak en kritik zarar çiçeklenme ve meyve oluşum döneminde görülür. Çünkü domates güvesi meyvenin içine girerek iç kısımlarını çürütür ve ürün kalitesini bozarak pazar değerini düşürür. Bunun dışında yaprak biti ise en çok genç sürgünler, yapraklar ve çiçeklenme döneminde zarar yapar. Bu zararlılar bitki yapraklarını sarartarak, gelişimini olumsuz etkileyerek ve virüs vektörlüğü taşımaları ve beslenmeleri sonucunda bu hastalıkların hızlı bir şekilde yayılmasına neden olmaktadır. Söz konusu zararlılar ile mücadeleyi başarılı bir şekilde yapılabilmesi için zararlıların tanımının ve yaşayışlarının çok iyi bilinmesi gerekmektedir.

Anahtar kelimeler: Örtü altı, sebze, zararlı

GİRİŞ

Tarım sektörü, küresel ölçekte artan nüfus, hızla değişen iklim koşulları ve doğal kaynakların sınırlılığı nedeniyle giderek daha fazla sürdürülebilir üretim modellerine yönelmektedir. Bu kapsamda, tarımsal üretimde yeni teknolojilerin kullanımı ve kontrollü üretim ortamlarının yaygınlaştırılması büyük bir önem kazanmıştır. Geleneksel tarım yöntemleri, açık alanda değişken iklim koşullarına doğrudan maruz kaldığından, üretim süresinin kısıtlı olması, hastalık ve zararlı etmenlerin kontrolünün zor olması ve su kullanım verimliliğinin düşük olması gibi çeşitli sınırlamalarla karşı karşıya kalmaktadır (Tüzel ve Gül, 2008).

Örtü altı tarımı, kontrollü çevre koşulları sayesinde bu olumsuzlukları minimize eden bir üretim modeli olarak öne çıkmaktadır. Sera tarımı olarak da adlandırılan bu sistem, bitkilerin optimum sıcaklık, nem, ışık ve karbon dioksit seviyelerine erişmesini sağlayarak yıl boyunca kesintisiz üretime olanak tanımaktadır (Kaşka ve ark., 2011). Ayrıca, örtü altı tarımı, modern sulama teknikleri (damla sulama, hidroponik sistemler vb.) ile su kullanım verimliliğini artırmakta, böylece doğal su kaynaklarının korunmasına da katkı sağlamaktadır.

Özellikle iklim koşullarının dış tarımsal üretimi sınırladığı bölgelerde, örtü altı yetiştiriciliği büyük bir avantaj sağlamaktadır. Soğuk iklime sahip bölgelerde seralar sayesinde üretim sezonu uzatılırken, sıcak ve kurak bölgelerde ise aşırı sıcaklık ve su kıtlığı gibi sorunlar kontrollü ortamlarla dengelenebilmektedir. Aynı zamanda, örtü altı tarımı hastalık ve zararlı yönetiminde de avantaj sunarak daha düşük kimyasal kullanımı ile daha sağlıklı ürünlerin yetiştirilmesine olanak tanımaktadır (Tüzel ve Gül, 2008).

Türkiye, iklim çeşitliliği ve coğrafi yapısı sayesinde örtü altı tarımı açısından önemli bir potansiyele sahiptir. Türkiye’de 2022 yılı itibarıyla toplam 800 bin dekarın üzerinde bir alanda örtü altı sebze üretimi yapılmakta olup, bu alanlarda başlıca domates, biber, salatalık ve patlıcan gibi sebzeler üretilmektedir (TÜİK, 2022). Ülke genelinde örtü altı tarımın büyük bir kısmı Akdeniz ve Ege Bölgeleri’nde yoğunlaşmış olsa da, mikroklima özellikleri gösteren bazı bölgelerde de yaygınlaşmaya başlamıştır (Demirtaş ve ark., 2020). Bu bağlamda, Doğu Anadolu Bölgesi’nde yer alan Iğdır ili, sahip olduğu özel iklim koşulları sayesinde bölgedeki diğer illerden farklı bir tarımsal üretim yapısına sahiptir. Iğdır, Karasal Doğu Anadolu İklimi ile Subtropikal Akdeniz İklimi arasında bir geçiş özelliği taşıyan mikroklima yapısı sayesinde geniş bir ürün yelpazesine ev sahipliği yapmaktadır (Çelik ve ark., 2015). Özellikle son yıllarda bölgedeki çiftçiler, açık tarımın yanı sıra örtü altı sebze yetiştiriciliğine yönelmiş ve bu kapsamda sera alanları artış göstermiştir.

Resmi verilere göre, Iğdır ilinde örtü altı tarımı yapılan alan yaklaşık 350-400 dekar olup, bu alanlarda yıllık ortalama 5.000-6.000 ton sebze üretimi gerçekleştirilmektedir (TÜİK, 2022;

Tarım ve Orman Bakanlığı, 2023). Üretimin büyük bir kısmını domates, biber ve salatalık oluşturmakta, ancak son yıllarda marul, ıspanak ve kabak gibi ürünlerin de örtü altı tarımına dahil edildiği görülmektedir. Iğdır'daki örtü altı sebze üretimi, hem iç piyasa ihtiyacını karşılamakta hem de bölge ekonomisine katkı sağlamaktadır.

Ancak, Iğdır'da örtü altı sebze yetiştiriciliği bazı sorunlarla da karşı karşıyadır. Yüksek sera kurulumu maliyetleri, sulama suyu kaynaklarının yetersizliği ve pazar erişiminde yaşanan zorluklar ve genel olarak bunun başında da entomolojik sorunlar olarak yer alan örtü altı sebze zararlıları yer almaktadır. (Demirtaş ve ark., 2020).

Yapılan bu çalışma ile, Iğdır ili merkez köylerde (Obaköy, Küllük, Evcı) bulunan zararlı türlerin belirlenmesi hedeflenmiştir. Elde edilecek sonuçların Entegre mücadele uygulamasında temel veriler oluşturacağı umulmaktadır.

ARAŞTIRMA BULGULARI

İncelenen köylerdeki seralarda yaygın olarak görülen zararlılar arasında hıyar, biber, kabak ve fasulyede beyaz sinek (*Bemisia tabaci* Gennadius) (Hemiptera: Aleyrodidae), yaprak bitleri (Hemiptera: Aphididae), thripsler (Tyhsanoptera: Thripidae), kırmızı örümcek (*Tetranychus urticae* Koch) (Acarina: Tetranychidae), galerisinekleri (*Liriomyza* spp.) (Diptera: Agromyzidae), Yeşilkurt (*Helicoverpa armigera* Hubn.) (Lepidoptera: Noctuidae) tespit edilmiş. Domateste de ise bu zararlılara ek olarak domates güvesi (*Tuta absoluta* Meyrick) (Lepidoptera: Gelechiidae) yoğun olarak görülmüştür. Zararlı böceklerin yoğunlukları vejetasyon sonuna doğru maksimuma ulaşmıştır. Ancak Domates güvesi, özellikle fide döneminden hasat sonuna kadar tüm gelişim evrelerinde zarar verebilir. Ancak en kritik zarar çiçeklenme ve meyve oluşum döneminde görülür. Çünkü domates güvesi meyvenin içine girerek iç kısımlarını çürütür ve ürün kalitesini bozarak pazar değerini düşürür. Bunun dışında yaprak biti ise en çok genç sürgünler, yapraklar ve çiçeklenme döneminde zarar yapar. Bu zararlılar bitki yapraklarını sarartarak, gelişimini olumsuz etkileyerek ve virüs vektörlüğü taşımaları ve beslenmeleri sonucunda bu hastalıkların hızlı bir şekilde yayılmasına neden olmaktadır.

SAPTANAN ZARARLILAR

Yaprak bitleri

(Hemiptera: Aphididae)

Tanımı:

Yaprak bitleri (Aphidoidea), bitkiler üzerinde zararlı olan, genellikle yumuşak vücutlu, küçük ve sokucu emici ağız yapısına sahip böceklerdir. Hemiptera (Yarımkanatlılar) takımına ait olan bu böcekler, genellikle bitkilerin yaprak, sürgün ve tomurcuklarında yoğunlaşır orda beslenerek öz suyunu emer ve tarımsal ürünlerde önemli verim kayıplarına neden olurlar. (Blackman & Eastop, 2000). Yaprak bitleri biyolojileri gereği oldukça hızlı çoğalabilen zararlılardır. Özellikle partenogenez (eşsiz üreme) yoluyla dişiler kısa sürede çok sayıda yavru üretebilirler. Ayrıca, çevresel koşullara bağlı olarak kanatlı ve kanatsız formlar oluşturabilmeleri, yeni konukçulara hızla yayılmalarını sağlar (Dixon, 1998). Tarımsal üretimde büyük kayıplara yol açabilen virüslerin taşınmasında kritik rol oynarlar (Van Emden & Harrington, 2007). Ayrıca, beslenme sırasında salgıladıkları tatlımsı madde, fumajin adı verilen mantarların gelişmesine neden olarak bitki yüzeyinde bir örtü oluşturur ve fotosentezi engelleyerek dolaylı zararları artırır (Dedryver et al., 2010).

Biyolojisi:

Yaprak bitleri, üreme stratejileri açısından son derece esnek ve çevre koşullarına bağlı olarak farklı üreme şekilleri sergileyebilirler.

Partenogenez (Eşeysiz Üreme)

Yaprak bitleri çoğunlukla partenogenez ile ürer. Bu yöntemde dişi bireyler, döllenmeye ihtiyaç duymadan doğrudan yeni bireyler meydana getirir. Özellikle ilkbahar ve yaz aylarında partenogenetik üreme baskındır. Bu üreme stratejisi sayesinde popülasyonları çok hızlı bir şekilde artabilir ve kısa sürede geniş alanlara yayılabilir (Van Emden & Harrington, 2007).

Canlı Doğum (Viviparite)

Yaprak bitleri yumurta yerine doğrudan yavru doğurur (viviparite). Bu özellik, nesillerin çok hızlı bir şekilde çoğalmasına ve koloniler oluşturmasına olanak tanır. Yeni doğan bireyler genellikle birkaç gün içinde olgunlaşarak üremeye başlar.

Cinsel Üreme ve Yumurta Üretimi


Sonbaharda çevresel koşulların kötüleşmesiyle birlikte yaprak bitleri cinsel üremeye yönelir. Dişi ve erkek bireyler çiftleşerek yumurta bırakır. Yumurtalar kış boyunca dayanıklı bir şekilde bekleyerek ilkbaharda tekrar aktif hale gelir (Blackman & Eastop, 2000).

Kanatlı ve Kanatsız Formlar

Yaprak bitleri, popülasyon yoğunluğu ve çevre koşullarına bağlı olarak kanatlı ve kanatsız bireyler oluşturabilir.

Kanatsız bireyler: Beslenme ve üreme için daha uygun koşullar sağlandığında çoğunluktadır.

Kanatlı bireyler: Popülasyon yoğunluğu arttığında veya besin kaynakları azaldığında ortaya çıkar ve yeni konukçulara göç ederler (Dedryver et al., 2010).



Şekil 1. Yaprak bitlerinin yaşam döngüsü A: fundatrix, B: apter, C: Alate, D: sexupare, E: ovipare, F: male G: egg (After Jones, 1961)

Zararı

Yaprak bitleri, bitkinin floem dokularına ulaşarak öz suyunu emer. Bu beslenme şekli bitkinin gelişiminde stres oluşturur, su kaybına ve besin dengesizliğine neden olur (Dixon, 1998). Beslenme sırasında bitkiye zarar veren enzimler ve salgılar üreterek bitkinin büyümesini engelleyebilir (Van Emden & Harrington, 2007). Özellikle genç yapraklar ve sürgünler üzerinde yoğunlaşarak yaprak kıvrılması, sararma ve bodur büyüme gibi belirtilere yol açarlar (Dedryver et al., 2010). Özellikle meyve ve sebze üretiminde kaliteyi düşürerek ekonomik zarara neden olur (Kennedy et al., 1962).

Yaprak bitleri, bitki hastalıklarına neden olan **200'den fazla virüsün** en önemli taşıyıcılarıdır (*vector organisms*) (Nault, 1997).

Non-persistent (Geçici) Virüsler: Yaprak bitleri kısa süreli beslenme sırasında virüsü alıp hızla yayabilir (örneğin, Potyvirus grubu).

Persistent (Kalıcı) Virüsler: Virüs, yaprak bitinin sindirim sisteminde çoğalarak uzun süre taşınabilir (örneğin, Luteovirüsler) (Feres & Moreno, 2009).

Yaprak bitleri, beslenme sırasında şekerli salgılar üretir. Bu salgılar bitki yüzeyine yapışarak fumajin (karaballık mantarı) enfeksiyonlarına neden olur (Michaud, 1998). Fumajin, bitkilerin fotosentez yapmasını engelleyerek büyümeyi yavaşlatır ve kaliteyi düşürür.



Şekil 2. Yaprak Biti a) Nimfi, b) Çeşitli biyolojik dönem ve formları, c) Yaprakbiti kolonisi, d) Yaprak biti nimf gömlekleri

Mücadele yöntemleri

Kültürel Mücadele Yöntemi

Yaprak bitleri belirli bitki türlerini tercih ettiğinden, dayanıklı bitki türleri ve çeşitleri yetiştirilerek zarar en aza indirilebilir. Alternatif konukçu bitkilerin yok edilmesi veya farklı ekim sistemleri kullanılması yaprak biti popülasyonlarının gelişimini sınırlandırır (Kennedy et al., 1962). Ekim nöbeti, konukçu bitkilerin değişmesini sağlayarak popülasyonları düşürebilir (Van Emden, 2007). Aşırı azotlu gübreleme yaprak biti popülasyonlarını artırabilir, bu nedenle gübrelemenin dengeli yapılması gereklidir (Dixon, 1998).

Biyolojik Mücadele Yöntemi

Biyolojik mücadele yöntemi ile yaprak bitlerine karşı doğada predatörler (avcı) böcekler, parazitoitler ve entomopatojen funguslar kullanılarak kontrol altına alınabilir.

Uğur böcekleri (Coccinellidae): Hem larva hem de ergin aşamada yaprak bitleriyle beslenir.
Syrphidae (Çiçek sinekleri): Larvaları yaprak bitlerini tüketerek popülasyonları baskılar.
Chrysopidae (Yeşil kanatlı böcekler): Larvaları aktif olarak yaprak bitlerini avlar (Obrycki & Kring, 1998).
Aphidius spp. (Hymenoptera: Braconidae), yaprak bitlerinin içine yumurta bırakarak larvaların konakçıyı tüketmesini sağlar. **Aphelinus spp.**, yaprak biti larvalarını enfekte eden diğer önemli parazitoitlerdir (Starý, 1993). **Beauveria bassiana**, **Metarhizium anisopliae** ve **Verticillium lecanii** gibi entomopatojen funguslar yaprak bitlerini enfekte ederek popülasyonlarını düşürebilir (Shah & Pell, 2003).

Kimyasal Mücadele

Kimyasal mücadele, yaprak bitleri için hızlı ve etkili bir kontrol yöntemi olmakla birlikte, aşırı kullanımı direnç gelişimine ve doğal düşmanların yok olmasına neden olabilir.

Beyz sinek (Bemisia tabaci Genn., Trialeurodes vaporariorum West.)

(Hemiptera: Aleyrodidae)**Tanımı:**

Bemisia tabaci (Gennadius) (Hemiptera: Aleyrodidae), dünya çapında tarımsal üretimde ciddi zararlara neden olan bir **beyaz sinek (whitefly) türüdür**. Hem doğrudan bitki öz suyunu emerek hem de virüs vektörü olarak büyük ekonomik kayıplara yol açmaktadır (Byrne & Bellows, 1991). Bu zararlı tür **yüksek üreme kapasitesi, geniş konukçu yelpazesi ve insektisitlere karşı hızla direnç geliştirme yeteneği** ile tanınır (Brown et al., 1995). *Bemisia tabaci*, özellikle tropikal ve subtropikal bölgelerde seracılık ve açık tarla yetiştiriciliğinde önemli sorunlara neden olmaktadır (Oliveira et al., 2001). Ergini 2-3 mm uzunluğunda, küçük ve ince yapılıdır. Kanatları beyaz renklidir ve dinlenme halinde vücuda paralel konumlanır. Gövde sarı renkte, gözler ise kırmızıdır.

Biyolojisi:

Yumurtadan yeni çıkan larvalar oval yapılı olup hareketlidir. Soluk sarı renkte olan larva, yaklaşık 8 saat içinde kendini yaprağa sabitleyerek hareketsiz hale gelir. Beslendikçe gelişen larva, dört dönem geçirir ve dördüncü dönemde pupa evresine girer.

Beyaz sinek (*Bemisia tabaci*), öncelikle sebzelerde görülür ve Mayıs ayından itibaren pamuk bitkisine geçiş yapar. Yumurtalarını bırakmak için taze yaprakları ve yaprakların alt yüzeyini tercih eder. Yoğun popülasyon durumlarında, yaprakların üst yüzeyine ve saplara da yumurta bırakabilir. Yumurtalarını tek tek veya küçük gruplar halinde bırakır ve her yumurta, bir sapçık (*pedicel*) aracılığıyla yaprağa tutturulur. Bu sapçık, yumurtanın bitki hücrelerinden su almasını sağlayarak gelişimini hava nemine bağlı olmaksızın sürdürebilmesine yardımcı olur. Bir dişi, yaklaşık 300 yumurta bırakabilir (Gerling & Mayer, 1996). Beyaz sineğin yumurta bırakması için en uygun sıcaklık 26-27°C olup, %60'ın üzerindeki orantılı nem koşulları üreme için idealdir. 14°C'nin altında yumurta bırakma süreci yavaşlar, 10°C'nin altında ise ergin bireylerin hareketleri büyük ölçüde azalır veya tamamen durur. 30°C sıcaklıkta yumurtalar yaklaşık 4 günde açılır ve çıkan larvalar yaz aylarında 8 saat içinde kendilerini sabitleştirir. Pupa evresinde beslenme gerçekleşmez. Larva ve pupa dönemi yaz aylarında yaklaşık 10-11 gün sürer. Beyaz sinek, yaz aylarında yaklaşık iki haftada bir döl verir ve yılda 9-10 nesil oluşturur. Üreme ve gelişim için en uygun koşullar; rüzgarsız ortam, 30°C'nin üzerindeki sıcaklık ve %60'tan fazla orantılı nemdir. Yağışlar ergin bireyleri olumsuz etkileyebilse de, ortam nemini artırarak üremeyi ve larvaların gelişimini teşvik eder (Jones, 2003).



Şekil 3. Life cycle of *B. tabaci* (Ghosh et al., 2019)

Zararı

Bitki öz suyunu emerek yapraklarda sararma, solma ve gelişme geriliğine neden olur. Aşırı beslenme sonucunda bitkinin büyümesini yavaşlatır ve ciddi popülasyonlarda ölüme yol açabilir. Tatlımsı madde (honeydew) salgılayarak fumajin (karaballık mantarı) gelişimini teşvik eder. Tatlımsı maddenin stomaları kapatması sonucu bitkide fotosentez yapma düzeni bozulur ve bu halin devamı bitkinin erken ölmesine neden olur (Brown & Czosnek, 2002). *Bemisia tabaci*, 100'den fazla bitki virüsünün vektörüdür.

Mücadele yöntemleri

Kültürel Mücadele Yöntemi

Bitkideki aşırı azot, zararlıların üreme gücünü artırdığı için, önerilen dozların üzerinde azotlu gübre kullanılmamalıdır. Sulama miktarı gerektiği kadar olmalı ve aşırı sulamadan kaçınılmalıdır. Bitki sıklığı, önerilen aralıklara uygun olmalı ve bitkiler fazla sık dikilmemelidir. Sebze, bostan ve pamuk tarlalarının iç içe olmasına özen gösterilmeli, eğer böyle bir ekim düzeni varsa, hasat sonrası bitki artıkları toplanarak yok edilmelidir. Ayrıca, zararlıların kışlamasına olanak sağlayan narenciye bahçeleri gibi alanlardaki yabancı otlar, toprak işleme yöntemleriyle ortadan kaldırılmalıdır (Naranjo & Ellsworth, 2009). Pamuk, biber ve domates tarlalarında fasulye veya ayçiçeği gibi çekici bitkilerin ekilmesi, zararlıların ana üründen uzaklaştırılmasına yardımcı olabilir (Mason et al., 2000). Beyaz sineklerin seralara girişini önlemek için 400-600 mikronluk gözenek çapına sahip sineklik tüllerinin kullanılması önerilir (Bethke et al., 1991).

Biyolojik Mücadele Yöntemi

Beyazsineğe karşı *Encarsia formosa* (Hym.: Aphelinidae) biyolojik mücadelede kullanılmaktadır. Serada erginlerin uçuş zamanı belirlendikten sonra, seranın büyüklüğüne bağlı olarak 20 ila 50 bitki seçilir. Bu bitkilerin alt, orta ve üst yapraklarından birer örnek alınarak beyazsinek larvaları sayılır ve yaprak başına düşen ortalama larva sayısı hesaplanır. Eğer yaprak başına 5 veya daha fazla larva tespit edilirse, *E. formosa* pupaları, yaprak başına bir adet düşecek şekilde seraya eşit olarak dağıtılır (Gerling & Mayer, 1996).

Kimyasal Mücadele Yöntemi

Kültürel önlemler ve doğal düşmanların baskısına rağmen beyazsinek popülasyonu **Ekonomik Zarar Eşiğini** aşarsa, ilaçlı mücadeleye başvurulmalıdır. Bu mücadelede, yüzey ilaçlamaları tercih edilmelidir. Zararlı yoğunluğunun eşiği aştığı durumlarda, ilk sulamadan önce veya hemen sonra, yer aletleriyle etkili bir ilaçlama yapılması hedeflenmelidir.

Beyazsineğe karşı ilaçlı mücadelede en önemli kriter olan **Ekonomik Zarar Eşiği**, çeşitli faktörlere bağlı olarak değişse de genel olarak yaprak başına **5 ergin veya 10 larva** olarak kabul edilmektedir. İlaç uygulamalarında, tarla pülverizatörlerine yaprak altı meme setleri eklenmelidir. İlaç seçiminde, yararlı türlerin korunması için **böcek gelişim engelleyicileri ve böcek gelişim düzenleyicileri** öncelikli olarak tercih edilmelidir. Ayrıca, zararlının kimyasallara karşı direnç geliştirmesini önlemek veya geciktirmek için ilaç grupları dönüşümlü olarak kullanılmalıdır (Mart ve ark., 1995).

Yaprak galerisinekleri (*Liriomyza* spp.)


(Diptera: Agromyzidae)

Tanımı

Erginleri yaklaşık 1,3–2,5 mm uzunluğunda, genellikle siyah ve sarı renkte sineklerdir (Murphy & LaSalle, 1999). Yumurtayı yaprak dokusu içine bırakılır ve gözle görülemeyecek kadar küçüktür (Waterhouse & Norris, 1987). Larvası sarımsak renkte, bacaksız ve silindirik yapıdadır. Larvaların beslenme sırasında oluşturduğu galeriler yapraklarda karakteristik zarar oluşturur (Parrella & Jones, 1987). Pupalari yaprak üzerinde veya toprakta kahverengimsi oval bir yapı oluştururlar (Weintraub & Horowitz, 1995).

Biyolojisi

Yetişkin dişi bireyler, yumurta koyma iğneleri (ovipozitörleri) ile yaprak yüzeyini delerek yaprak dokularıyla beslenir ve sıvı hücre içeriklerini emerler. Dişiler ayrıca yaprak dokusunu delerek yumurtalarını bırakır ve 7 ila 10 gün boyunca yaşayabilirler. Hem beslenme hem de yumurtlama sonucu oluşan delikler beyaza döner ve zarar gören yapraklar benekli veya lekeli bir görünüm alır. Yumurtalar, bırakıldıktan 2 ila 5 gün sonra çatlar. Larvalar, yaprak dokusu içinde beslenerek kıvrımlı galeriler veya geniş lekeler oluşturur. Larvalar büyüdükçe yaprak galerileri genişler ve larvalar üç farklı gelişim evresinden (instar) geçer. Beslenme tamamlandığında, olgun larvalar yaprak yüzeyinde bir yarık açarak yapraktan çıkar ve yere (veya saksı toprağına) düşerek pupa haline gelir. Sıcaklık 10°C'nin üzerinde olduğu sürece yılda birçok nesil meydana gelebilir (Infonet Biovision, 2016).



Şekil 4. Life cycle for *Liriomyza* spp. (Kephis)

Zararı

Yaprak galerisineikleri, hem beslenme hem de yumurta bırakma aşamalarında yaprak dokusunda zarar oluşturur. Dişi bireyler, ovipozitörleri ile yaprak epidermisini delerek beslenir ve hücre öz suyunu emer. Bu beslenme aktivitesi, yaprak üzerinde beyazımsı veya sarımsı küçük noktalar şeklinde lekeler oluşturur (Parrella, 1987). Aynı zamanda, yumurta bırakma sırasında meydana gelen delikler de benekli bir görünüm oluşturur (Spencer, 1990). Larvalar, yaprak epidermisinin altına girdikten sonra mezofil dokusunda tüneller açarak beslenirler. Bu tüneller, ilk başta ince ve dar, ancak larvalar büyüdükçe genişleyen kıvrımlı galeriler şeklindedir (Weintraub & Horowitz, 1995). Galeriler, yaprakta su kaybına ve fotosentez kapasitesinin azalmasına yol açar (Leibee, 1984). Yaprak dokusunda meydana gelen zarar nedeniyle bitkinin fotosentez kapasitesi ciddi oranda azalır. Özellikle yoğun bulaşma durumlarında, bitkiler büyüme geriliği gösterir ve verim kaybı yaşanır (Minkenberg, 1988). Yaprak kaybı nedeniyle bitki stres altında kalır ve ikinci derecede enfeksiyonlara açık hale gelir (Murphy & LaSalle, 1999).



Şekil 5. *Liriomyza* spp. (Kephıs) hıyar da zararı

Mücadele Yöntemleri

Bazı bitki türleri yaprak galeri sineklerine karşı doğal direnç gösterebilir. Özellikle sert yaprak dokusuna sahip bitkiler, dişi sineklerin yumurta bırakmasını zorlaştırır (Parrella, 1987). Aynı alanda sürekli olarak konukçu bitkilerin ekilmesi, zararlı populasyonunun artmasına neden olur. Bu yüzden hassas bitkilerle dayanıklı türlerin dönüşümlü olarak yetiştirilmesi, sineklerin çoğalmasını sınırlandırabilir (Leibee, 1984). Yaprak galerisineklerinin pupası toprakta olduğu için düzenli olarak işlenmesi, pupaların yok edilmesine yardımcı olur (Spencer, 1990).

Kimyasal mücadele, diğer kontrol yöntemleri yetersiz kaldığında ve zararlı populasyonu ekonomik zarar eşiğini aştığında uygulanmalıdır. Ancak, yaprak galerisinekleri birçok insektisite karşı direnç geliştirebilir, bu yüzden dikkatli bir şekilde kullanılmalıdır (Leibee, 1984).

Domates Güvesi (*Tuta absoluta* Meyrick)


(Lepidoptera: Gelechiidae)

Tanımı

Tuta absoluta (Meyrick), domates güvesi olarak da bilinen, özellikle domates (*Solanum lycopersicum*) başta olmak üzere Solanaceae (patlıcangiller) familyasındaki birçok bitkiye zarar veren önemli bir tarımsal zararlıdır (Desneux et al., 2010). Güney Amerika kökenli olup, 2006 yılında Avrupa'ya, ardından Asya ve Afrika kıtalarına yayılmıştır ve günümüzde dünya genelinde büyük ekonomik kayıplara yol açan en tehlikeli tarım zararlılarından biri olarak kabul edilmektedir (Biondi et al., 2018). *Tuta absoluta* erginleri küçük boyutlu güveler olup gri-kahverengi renkte ve yaklaşık 6-7 mm uzunluğundadır. Kanat açıklığı 8-10 mm arasında değişir ve kanatlarında siyah noktalar bulunur (Urbaneja et al., 2012). Yumurtalar yaklaşık 0.35 mm uzunluğunda, silindirik ve kremi beyaz renkte olup, zamanla sarımsı renge dönüşür (Guedes & Picanço, 2012). Larvalar başlangıçta krem renğinde olup daha sonra yeşilimsi bir renk alır ve son evrede pembemsi bir görünüme sahip olur. Pupa evresi genellikle toprakta veya bitki dokusu içinde gerçekleşir (Caparros Megido et al., 2013).

Biyolojisi

Tuta absoluta, sıcak iklimlerde çok hızlı üreyebilen ve yılda 10-12 nesil verebilen bir türdür (Desneux et al., 2010). Bir dişi ömrü boyunca yaklaşık 250-300 yumurta bırakabilir ve uygun koşullarda yumurtalar 4-5 gün içinde açılır. Larvalar, yaprak, meyve ve gövde dokularında tüneller açarak beslenir, bu da ciddi ürün kayıplarına neden olur (Biondi et al., 2018). Larva evresi 12-15 gün sürerken, pupa süresi sıcaklığa bağlı olarak 7-10 gün arasında değişebilir (Urbaneja et al., 2012).



Şekil 6. Life cycle of *T. absoluta*, photographs compiled from Arno et al. (2010) and developmental periods from Erdogan and Babaroglu (2014)

Zararı

Tuta absoluta (Meyrick) (Lepidoptera: Gelechiidae), özellikle domates (*Solanum lycopersicum*) ve diğer Solanaceae familyasına ait bitkilerde ciddi ekonomik kayıplara yol açan, dünya çapında yayılmış bir zararlıdır (Desneux et al., 2010). Zararı, larvaların beslenme şekli ve bitki üzerindeki etkileri ile ilişkilidir. *Tuta absoluta* larvaları, yaprak dokularında tüneller (galeriler) açarak beslenir. Bu beslenme sonucu yapraklar solgunlaşır, fotosentez kapasitesi azalır ve bitkinin genel büyüme hızı düşer (Urbaneja et al., 2012). *Tuta absoluta* larvaları, domates meyvelerine doğrudan zarar verir. Yumurtadan çıkan larvalar meyveye girerek iç kısımda beslenir ve çürümeye neden olur (Caparros Megido et al., 2013). Meyve zararları, pazar değeri kaybına ve üreticiler için ekonomik kayıplara neden olur. Özellikle fide aşamasındaki bitkilerde, larvaların gövde içinde beslenmesi sonucu bitkinin büyümesi durur veya ölür (Urbaneja et al., 2012).



Şekil 7. *T. absoluta*'nın meyve deki ve yapraktaki zararı (İreç, Altuntekin, 2024)

Mücadele Yöntemleri

Bulaşık bitki artıkları temizlenmeli ve yok edilmelidir. Seralar iyi izole edilmeli, giriş-çıkışlar kontrol altına alınmalıdır. Erken ekim yapılmalı ve bitkiler sağlıklı büyütülmelidir. *Tuta absoluta* feromon tuzakları, ergin güveleri yakalayarak popülasyonu izleme ve kitle yakalama yöntemiyle azaltma amacıyla kullanılır (Desneux et al., 2010). Özellikle seralarda, uçan ergin bireylerin yakalanması için sarı yapışkan tuzaklar kullanılır. Biyolojik mücadelede predatör böceklerden, *Chrysoperla carnea*, *Nabis punctatus punctatus*, *Orius* spp., ve *Geocoris megacephalus* gibi türler Domates güvesinin predatörü olduğu bildirilen avcı böceklerdir. Öztemiz (2012) tarafından yapılan bir araştırmada *Chrysoperla* sp., *Nabis pseudoferus*, *Orius* sp., ve *Geocoris* sp., türlerinin *T. absoluta*'nın doğal düşmanları olduğu tespit edilmiştir. Bunun dışında parazitoit olarak da iki larva parazitoidi *Bracon hebetor* (Say, 1836) (Hymenoptera Braconidae, Braconinae) ve *Apanteles* sp. (Hymenoptera Braconidae, Microgasterinae) ve bir larva pupa parazitoidi *Campoplex* sp. (Hymenoptera: Ichneumonidae) tespit edilmiştir (Öztemiz, 2012)

Thrips spp.


(Thysanoptera: Thripidae)

Tanımı

Erginler, uzun, dar ve hafif yassı bir vücuda sahiptir. **Saman sarısı** veya **kirli sarı** renkte olup vücutlarının bazı bölgelerinde **grimsi lekeler** bulunur. Kanatları **sarı** veya **sarımsı gri** renkte olup uzun ve kenarları saçaklıdır. Antenleri **7 segmentli** olup uç segmentler daha koyu renktedir. Abdomen uzun, açık veya kirli sarı renkte olup dişilerde belirgin ve kıvrık bir **ovipozitör** bulunmaktadır. Vücut uzunluğu **0.8-1.0 mm** arasındadır (Lewis, 1997). Erkek bireyler doğada oldukça nadir görülür ve kanatsızdır. Yumurtalar **saydam ve beyazımsı** olup yaklaşık **0.25 mm** uzunluğunda ve **fasulye şeklindedir**. **I. ve II. dönem nimfler** sarı renkte olup **0.8-0.9 mm** boyutlarındadır. **Polifag** bir zararlı olup ülkemizin hemen her bölgesinde farklı yoğunluklarda bulunur (Mound, 2005).

Biyolojisi

Ülkemizin **soğuk bölgelerinde**, bu zararlı kışı **ergin** olarak **bitki sapları, kuru otlar, bitki kökleri** etrafı, **bitki artıkları, ağaç kabukları** altı veya **toprak içinde pupa** ya da **ergin** formda geçirir. Buna karşılık, **sıcak bölgelerde** ve **seralarda** yıl boyunca aktif olarak yaşamını sürdürebilir. Dişiler, **eşesiz üreme** yoluyla çoğalır ve doğada erkek bireylere oldukça ender rastlanır. Yumurtlama sırasında, dişiler **ovipozitörleri** yardımıyla bitkinin **epidermisini** delerek doku içinde **torba şeklinde bir oyuk** oluşturur ve yumurtalarını buraya bırakır. Bir dişi, yaşamı boyunca yaklaşık **80 yumurta** bırakabilir. **Yaz aylarında**, yumurtalar **3-5 gün** içinde açılır ve çıkan nimfler kısa sürede beslenmeye başlar. Bitki üzerinde **iki nimf dönemi** geçirdikten sonra toprağa inerek **prepupa** ve **pupa** evrelerine girerler. Yaklaşık **bir hafta** sonra ergin bireyler çıkar ve tekrar bitkilere geçerek beslenmeye devam eder. Bir döl, **26°C'de yaklaşık 18 günde** tamamlanırken, uygun koşullarda yılda **10 döl** verebilir (Ravinder, 2022).



Şekil 8. Thrips spp. Lyfe cycle (Alec Blume)

Zararı

Ergin ve nimfleri bitkilerin yaprak, sap ve meyvelerinin epidermis tabakasını ağız parçaları ile yırtarak ya da zedeleyerek çıkan öz suyu emerler. Thripslerin yapraklardaki karakteristik zararı gümüşümsü renk açılmalarıdır. Yoğunluğun fazla olduğu yerlerde yapraklarda başlangıçta koyu yeşil renk, daha sonra yanıklar oluşacak şekilde renk açılmaları meydana gelir. Zamanla yaprak solar, bükülür ve beyazımsı bir hal alır. Buna halk arasında akdamar hastalığı denir.

Bitkilerde böceğin beslendiği yerler bir süre sonra beyazımsı veya gümüşü bir renk alır. Yoğun saldırılarda yapraklar görevini yapamaz, vaktinden önce dökülür. Bitkinin verimi azaldığı gibi ürün kalitesi de bozulur (Mound, Teulon, 1995).

Mücadele yöntemi

Tütün tripsinin aşırı çoğalmasını önlemek için sık sulanan fidelik ve tarlalar kuru bırakılmamalıdır. Zararlıların barınabileceği bitki artıkları ve yabancı otlar temizlenerek yok edilmelidir. Daha önce bulaşık olduğu bilinen alanlara fidelik kurulmamalı ve dikim yapılmamalıdır. Ayrıca, kontrol edilmeden bulaşık fideler tarlaya dikilmemelidir. Tripsin biyoteknik mücadelesinde, sarı yapışkan tuzaklar yaygın olarak kullanılmaktadır. Doğada, tripslerin popülasyonunu kontrol altında tutan çeşitli doğal düşmanlar bulunmaktadır. Bunların çoğunu avcı anthocorid, mirid ve akar türleri oluşturmaktadır. Özellikle, *Macrolophus costalis* (=rubi), tripslerin en önemli predatörlerinden biridir ve hem nimf hem de erginleriyle beslenmektedir. Bunun yanı sıra, *Orius* spp. biyolojik mücadelede önemli bir rol oynamaktadır. Tripslerin kimyasal mücadelesine karar vermek ve doğru ilaçlama zamanını belirlemek için seradaki zararlı yoğunluğu ve doğal düşman popülasyonu düzenli olarak izlenmelidir (Mound & Marullo, 1996). Seranın büyüklüğüne bağlı olarak 20-50 bitki seçilir ve her bitkinin alt, orta ve üst yapraklarından birer adet alınarak zararlı yoğunluğu belirlenir. Kimyasal mücadele, küçük yapraklı bitkilerde (örneğin, fasulye ve biber) yaprak başına 10 trips, büyük yapraklı bitkilerde (domates, patlıcan, hıyar, kabak) yaprak başına 20 trips veya çiçek başına 3 nimf + ergin olduğunda uygulanmalıdır (Weintraub & Berlinger, 2004)

Yeşilkurt (*Helicoverpa armigera* Hubn.)


(Lepidoptera: Noctuidae)

Tanımı

Erginin Ön kanatları kahverengimsi, sarımsı veya yeşilimsi renkte olup dalgalı desenlere sahiptir. Arka kanatlar açık renkte olup kenar kısımlarında koyu renkli bant bulunur (Zalucki et al., 1994). Yeni çıkan larvalar açık yeşil veya sarı renkte olup büyüdükçe kahverengimsi-yeşil renge dönüşebilir. Larvaların vücudunda belirgin uzunlamasına çizgiler bulunur (Sharma, 2005). Küçük, yuvarlak ve krem rengindedir. Yumurtalar yaprak, meyve veya tomurcuklara bırakılır (Fitt, 1989). Toprakta, bitki kalıntıları arasında kahverengimsi renkte bulunur (Zalucki et al., 1994).

Biyolojisi

Helicoverpa armigera, sıcak iklimlere adapte olmuş, yüksek üreme potansiyeline sahip bir zararlıdır. Yaşam döngüsü çevresel koşullara, özellikle sıcaklığa bağlı olarak değişiklik gösterir. Optimum sıcaklık aralığında (25-30°C) gelişim süresi daha kısa olurken, daha düşük sıcaklıklarda gelişim süreci uzar (Fitt, 1989). Dişiler yumurtalarını tek tek veya gruplar halinde yaprak, çiçek tomurcuğu, meyve ve gövdeye bırakır. Yumurtalar genellikle yuvarlak, parlak beyaz renkte olup, gelişim ilerledikçe sarımsı veya turuncumsu renge dönüşebilir. Dişi bireyler günde 100-200, yaşamları boyunca toplamda 1.500-3.000 yumurta bırakabilir (Reed & Pawar, 1982). Erkek bireyler ise 2-10 gün yaşar ve dişilerle çiftleşerek popülasyon artışına katkı sağlar. Larvalar yumurtadan çıktıktan sonra 6 evrede gelişim gösterir. Son evrede larva yaklaşık 30-40 mm uzunluğa ulaşabilir ve toprağa inerek pupaya dönüşmek için kendine bir çukur hazırlar. Büyük larvalar, bitkinin generatif organlarını (çiçek ve meyve) hedef alarak büyük ekonomik kayıplara neden olur. Larvalar olgunlaştığında toprağın 2-10 cm derinliğine girerek pupa olur. Pupa rengi başlangıçta açık kahverengi olup, zamanla koyu kahverengiye dönüşür (Zalucki et al., 1994). Yoğun popülasyonlarda, pupaların büyük kısmı toprakta diyapoz (dinlenme evresi) geçirerek sonraki yıllarda çıkış yapabilir.



Şekil 9. Life cycle of *Helicoverpa armigera* (Genç ve Yücel, 2017)

Zararı

Yeni çıkmış larvalar bitkinin genç yapraklarını tahrip eder. Yaprak yüzeyinde düzensiz delikler ve parçalanmalar görülür (Fitt, 1989). Bitkinin fotosentez kapasitesi azalır, büyüme ve gelişme olumsuz etkilenir. Larvalar çiçek tomurcuklarını delerek içeride beslenir, böylece çiçek açmadan dökülür (Zalucki et al., 1994). Çiçekler zarar gördüğünde döllenme azalır, bu da verimde önemli düşümlere neden olur. En büyük ekonomik kayıp meyvelerde oluşur. Larvalar doğrudan meyveleri delerek içeri girer ve beslenmeye devam eder. Dışarıdan bakıldığında küçük giriş delikleri görülse de, meyve içinde büyük boşluklar oluşturur (Sharma, 2005).



Şekil 10. Yeşilkurt' un domates meyvedesindeki zararı (İreç, 2024)

Mücadele yöntemi

Bulaşık bitki artıkları temizlenmeli ve yok edilmelidir. Seralar iyi izole edilmeli, giriş-çıkışlar kontrol altına alınmalıdır. Erken ekim yapılmalı ve bitkiler sağlıklı büyütülmelidir. Feromon tuzakları, kelekleri yakalayarak popülasyonu izleme ve kitle yakalama yöntemiyle azaltma amacıyla kullanılır. Biyolojik mücadelesinde ise *Deraeocoris pallens* (Reuter, 1904) *Macrolophus pygmaeus* (Rambur, 1839), *Chrysoperla carnea*, *Coccinella septempunctata*

(Linnaeus, 1758) predatörleri kullanılır (Khan, 2016). Yeşilkurt'un 3 dölü bitkide zararlı olmakta ve her dölü yaklaşık 30 günde tamamlanmaktadır. Bırakılan yumurta sayısı arttığı sürece ilaçlama geciktirilmeli, ancak bu süre Yeşilkurt larvalarının çok fazla büyümesine imkan verecek kadar uzun olmamalıdır. İlaçlı mücadelede yüksek etki elde etmek için her dölde yumurtadan yeni çıkmış larvaların çoğunlukta olduğu dönem iyi belirlenmelidir. Larvalar gündüzleri genellikle çiçek ve meyve içinde beslendiklerinden sabahın erken saatlerinde veya akşamın geç saatlerinde ilaçlama yapılmalıdır (Jaber, 2017).

SONUÇ

Iğdır ili, sahip olduğu iklim avantajları ve verimli toprakları sayesinde örtü altı sebze yetiştiriciliğinde önemli bir üretim merkezi konumundadır. Ancak, bu üretim süreci çeşitli zararlılar nedeniyle ciddi tehdit altındadır. Beyaz sinek (*Bemisia tabaci*), yaprak bitleri (*Aphididae*), thripsler (*Thripidae*), kırmızı örümcek (*Tetranychus urticae*) ve galerisineği (*Liriomyza spp.*) gibi ana zararlılar, bitkilerin gelişimini engelleyerek hem doğrudan zarar vermekte hem de virüs hastalıkların taşınmasına aracılık etmektedir. Bu durum, ürün kayıplarına yol açarak üreticilerin ekonomik kayıplar yaşamasına neden olmaktadır.

Zararlılarla mücadelede yalnızca kimyasal yöntemlere başvurmak, kısa vadeli bir çözüm sunarken, uzun vadede direnç gelişimi ve çevresel zararlar gibi olumsuz etkiler ortaya çıkarmaktadır. Bu nedenle, biyolojik mücadele, kültürel önlemler ve fiziksel yöntemlerin entegre edilmesi büyük önem taşımaktadır. Entegre Zararlı Yönetimi (IPM) çerçevesinde, doğal düşmanların kullanımı, zararlılar için uygun olmayan çevre koşullarının oluşturulması ve düzenli izleme sistemlerinin kurulması, mücadele sürecinin daha etkili olmasını sağlayacaktır.

Ayrıca, üreticilerin bilinçlendirilmesi ve eğitim programlarının yaygınlaştırılması, zararlılarla mücadelede başarıyı artıracak önemli faktörler arasındadır. Çiftçilerin, zararlıların biyolojisini ve yaşam döngülerini iyi tanınması, mücadele yöntemlerini doğru zaman ve dozda uygulaması hem ürün kayıplarını önleyecek hem de çevreye duyarlı üretim yapılmasını sağlayacaktır.

Sonuç olarak, Iğdır'da örtü altı sebze yetiştiriciliğinde karşılaşılan zararlılarla etkili bir şekilde mücadele edebilmek için entegre yönetim stratejilerinin benimsenmesi gerekmektedir. Zararlılara karşı tek yönlü mücadele yerine, biyolojik, kültürel ve kimyasal yöntemleri içeren çok yönlü yaklaşımlar tercih edilmelidir. Bu sayede, üretimde sürdürülebilirlik sağlanarak hem çevreye duyarlı hem de ekonomik açıdan verimli bir sebze yetiştiriciliği mümkün olacaktır. Gelecekte yapılacak bilimsel çalışmalar ve teknolojik gelişmelerle, daha etkili ve çevre dostu mücadele yöntemlerinin geliştirilmesi, bölgedeki tarımsal üretimi daha ileri seviyeye taşıyacaktır.

KAYNAKÇA

Bass, C., Field, L. M., & Williamson, M. S. (2014). The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochemistry and Molecular Biology*, 51, 41-51.

Biondi, A., Guedes, R. N. C., Wan, F. H., & Desneux, N. (2018). Ecology, worldwide spread, and management of the invasive South American tomato pinworm, *Tuta absoluta*: past, present, and future. *Annual Review of Entomology*, 63(1), 239-258.

Blackman, R. L., & Eastop, V. F. (2000). *Aphids on the world's crops: An identification and information guide*. John Wiley & Sons.

Brown, J. K., & Czosnek, H. (2002). Whitefly transmission of plant viruses. *Advances in Botanical Research*, 36, 65-100.

- Brown, J. K., Frohlich, D. R., & Rosell, R. C. (1995). The sweetpotato or silverleaf whiteflies: Biotypes of *Bemisia tabaci* or a species complex? *Annual Review of Entomology*, 40, 511-534.
- Butler, G. D., & Henneberry, T. J. (1990). Plant-derived oils and detergents as control agents for whiteflies, *Bemisia tabaci* and *Trialeurodes vaporariorum*. *Southwestern Entomologist*, 15(3), 257-263.
- Byrne, D. N., & Bellows, T. S. (1991). Whitefly biology. *Annual Review of Entomology*, 36, 431-457.
- Cahill, M., Gorman, K., Day, S., & Denholm, I. (1996). Resistance to the pyrethroid insecticide bifenthrin in the B biotype of *Bemisia tabaci*. *Pesticide Science*, 48(3), 257-263.
- Caparros Megido, R., Haubruge, É., & Verheggen, F. J. (2013). Tomato-infesting *Tuta absoluta*: Pest management and control strategies. *Insect Science*, 20(3), 177-193.
- Çelik, A., et al. (2015). Iğdır ili tarımsal yapısı ve örtü altı üretim potansiyeli. *Tarım Ekonomisi Dergisi*, 21(3), 189-202.
- De Barro, P. J., Liu, S. S., Boykin, L. M., & Dinsdale, A. B. (2011). *Bemisia tabaci*: A statement of species status. *Annual Review of Entomology*, 56, 1-19.
- Dedryver, C. A., Le Ralec, A., & Fabre, F. (2010). The biology of aphids and their role in crop protection. *Comptes Rendus Biologies*, 333(6-7), 461-470.
- Demirtaş, M., et al. (2020). Doğu Anadolu Bölgesi'nde seracılık: Iğdır ili örneği. *Türkiye Tarım ve Ormancılık Dergisi*, 26(1), 45-58.
- Desneux, N., Luna, M. G., Guillemaud, T., & Urbaneja, A. (2010). The invasive South American tomato pinworm, *Tuta absoluta*, continues to spread in Afro-Eurasia and beyond: the new threat to tomato world production. *Journal of Pest Science*, 83(1), 1-13.
- Dixon, A. F. G. (1998). *Aphid ecology: An optimization approach*. Springer Science & Business Media.
- Fereres, A., & Moreno, A. (2009). Behavioural aspects influencing plant virus transmission by homopteran insects. *Virus Research*, 141(2), 158-168.
- Fitt, G. P. (1989). The ecology of *Helicoverpa armigera* (Hübner) in Australia and implications for management. *Annual Review of Entomology*, 34(1), 17-52.
- Gerling, D., & Mayer, R. T. (1996). *Bemisia: 1995 taxonomy, biology, damage, control and management*. Intercept Limited.
- Gerling, D., Alomar, O., & Arnó, J. (2001). Biological control of *Bemisia tabaci* using predators and parasitoids. *Crop Protection*, 20(9), 779-799.
- Guedes, R. N. C., & Picanço, M. C. (2012). The tomato borer *Tuta absoluta* in South America: pest status, management, and insecticide resistance. *EPPO Bulletin*, 42(2), 211-216.
- Gurr, G. M., Wratten, S. D., & Luna, J. M. (2004). Multi-function agricultural biodiversity: Pest management and other benefits. *Basic and Applied Ecology*, 5(2), 107-116.
- Horowitz, A. R., Kontsedalov, S., Khasdan, V., & Ishaaya, I. (2020). Biotypes B and Q of *Bemisia tabaci* and their relevance to neonicotinoid and pyriproxyfen resistance. *Journal of Pest Science*, 93(3), 897-907.
- Jaber, L.R., et al. (2017). "Use of entomopathogenic fungi to control *Helicoverpa armigera* (Hübner)." *Biocontrol Science and Technology*.
- Jones, D. R. (2003). Plant viruses transmitted by whiteflies. *European Journal of Plant Pathology*, 109, 195-219.

- Kaşka, N., et al. (2011). Seracılıkta modern üretim teknikleri. Akdeniz Üniversitesi Ziraat Fakültesi Yayınları.
- Kennedy, J. S., Day, M. F., & Eastop, V. F. (1962). Aphis migration and virus diseases. Methuen & Co Ltd.
- Khan, A., et al. (2016). "Biological control of *Helicoverpa armigera* using *Trichogramma* species." *Biocontrol Science and Technology*.
- Leibee, G. L. (1984). Influence of temperature on development and fecundity of *Liriomyza trifolii* (Diptera: Agromyzidae). *Annals of the Entomological Society of America*, 77(3), 280-283.
- Lewis, T. (1997). Thrips as Crop Pests. *CAB International*, 740 pp.
- Mart, C., Kismir, A., Belli, A., Tunc, A., Turhan, N., Pala, Y. Aktura, T. (1995). Researches on the chemical control of whitefly (*Bemisia tabaci* Genn.) on cotton fields in Mediterranean Region.
- Michelakis, S.E., 1986. Problems in the application of biological control against *Trialeurodes vaporariorum* in unheated plastic glasshouses in Crete. *EPPO Bulletin*, 16 (2): 423-427 (Abst. in: *Rev. Appl. Ento* (1988): 76: 3448).
- Minkenbergh, O. P. J. M. (1988). Dispersal of *Liriomyza bryoniae* and its parasitoid *Dacnusa sibirica* in tomato greenhouses. *Entomologia Experimentalis et Applicata*, 46(3), 291-300.
- Mound, L. A. (2005). Thysanoptera: Diversity and Interactions. *Annual Review of Entomology*, 50, 247-269.
- Mound, L. A., & Teulon, D. A. (1995). Thysanoptera as phytophagous opportunists. In *Thrips biology and management* (pp. 3-19). Boston, MA: Springer US.
- Murphy, S. T., & LaSalle, J. (1999). Balancing biological control strategies in greenhouse systems. *Biocontrol News and Information*, 20(3), 91N-104N.
- Nauen, R., & Denholm, I. (2005). Resistance of insect pests to neonicotinoid insecticides. *Annual Review of Entomology*, 50, 249-278.
- Nault, L. R. (1997). Arthropod transmission of plant viruses: a new synthesis. *Annals of the Entomological Society of America*, 90(5), 521-541.
- Navas-Castillo, J., Fiallo-Olivé, E., & Sánchez-Campos, S. (2011). Emerging virus diseases transmitted by whiteflies. *Annual Review of Phytopathology*, 49, 219-248.
- Obrycki, J. J., & Kring, T. J. (1998). Predaceous coccinellidae in biological control. *Annual Review of Entomology*, 43, 295-321.
- Oliveira, M. R. V., Henneberry, T. J., & Anderson, P. (2001). History, current status, and collaborative research projects for management of *Bemisia tabaci*. *Crop Protection*, 20(9), 709-723.
- Oliveira, M. R. V., Henneberry, T. J., & Anderson, P. (2001). History, current status, and collaborative research projects for management of *Bemisia tabaci*. *Crop Protection*, 20(9), 709-723.
- Öztemiz S. (2012). Domates güvesi [(*Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae)] ve Biyolojik Mücadelesi. *KSÜ Doğa Bil. Derg.*, 15(4).
- Parrella, M. P. (1987). Biology of *Liriomyza*. *Annual Review of Entomology*, 32(1), 201-224.
- Parrella, M. P., & Jones, V. P. (1987). Development of *Liriomyza trifolii* on greenhouse ornamentals. *Journal of Economic Entomology*, 80(1), 58-62.

- Parrella, M.P., W.W. Allen and P. Morishita, 1981. Leaf miner species causes California Mum growers new problems. Calif. Agric., 35 (9-10): 28-30 (Abst. in: Rev. Appl. Ent. (1982), 70 (4): 2274).
- Ravinder, S. (2022). *Biology and population dynamics of Thrips tabaci Lindeman (Thysanoptera: Thripidae) on onion* (Doctoral dissertation, Punjab Agricultural University, Ludhiana).
- Reed, W., & Pawar, C. S. (1982). Heliothis: a global problem. Horticultural Reviews, 4, 221-258.
- Sharma, H. C. (2005). *Heliothis/Helicoverpa* management: Emerging trends and strategies for future research. Oxford & IBH Publishing Co. Pvt. Ltd., India.
- Spencer, K. A. (1973). Agromyzidae (Diptera) of economic importance. Series Entomologica 9. Dr. W. Junk B.V., The Hague.
- Spencer, K. A. (1990). Host specialization in the world Agromyzidae (Diptera). Kluwer Academic Publishers, Dordrecht.
- Tarım ve Orman Bakanlığı. (2023). Türkiye'de Örtü Altı Üretim Alanları Raporu.
- Türkiye İstatistik Kurumu (TÜİK). (2022). Tarımsal Üretim Verileri.
- Urbaneja, A., González-Cabrera, J., Arnó, J., & Gabarra, R. (2012). Prospects for the biological control of *Tuta absoluta* in tomatoes of the Mediterranean basin. Pest Management Science, 68(9), 1215-1222.
- Güzel, Y., & Gül, A. (2008). Örtü altı sebze yetiştiriciliği ve çevre. Tarım Bilimleri Dergisi, 14(2), 67-75.
- Van Emden, H. F., & Harrington, R. (2007). Aphids as crop pests. CABI Publishing.
- Waterhouse, D. F., & Norris, K. R. (1987). Liriomyza species as crop pests in South-East Asia and the Pacific. ACIAR Monograph No. 3.
- Weintraub, P. G., & Berlinger, M. J. (2004). Physical control of insect pests in agriculture. *Pest Management Science*, 60(6), 609-614.
- Weintraub, P. G., & Horowitz, A. R. (1995). The newest leafminer pest in Israel, *Liriomyza huidobrensis*. *Phytoparasitica*, 23(2), 177-184.
- Yano, E., 1983. Constraints on the use of *Encarsia formosa* in tomatoes in Japan with special reference to the effect of temperature on its efficiency. Bull. SROP, 6 (3): 49-54.
- Zalucki, M. P., Darglish, G., Firepong, S., & Twine, P. H. (1994). The biology and ecology of *Helicoverpa armigera* (Hübner) in relation to its status as a pest in cotton. Crop Protection, 13(8), 539-560.

SUPERFOOD OF THE FUTURE: SEAWEED

GELECEĞİN SÜPER BESİNİ: DENİZ YOSUNLARI

Emine NAKİLCİOĞLU

Ege University, Faculty of Engineering, Department of Food Engineering, Izmir, Turkey

ORCID ID: <https://orcid.org/0000-0003-4334-2900>

Gizem TİRYAKİ

Ege University, Institute of Science and Technology, Food Engineering, Izmir, Turkey

ORCID ID: <https://orcid.org/0000-0001-5608-9815>

ABSTRACT

Introduction and Purpose: Global food production has been seeking new resources due to increasing population and the need for sustainability. Seaweeds have emerged as a sustainable food source due to their high nutritional value, bioremediation properties, and ability to be cultivated without the need for agricultural land. Additionally, they provide environmental benefits by offering ecosystem services such as carbon sequestration and improving water quality. This study aims to examine the nutritional value, health benefits, and industrial potential of seaweeds, and to explore how they can be more widely used in the food and other industries in the future. **Materials and Methods:** This study was conducted based on a literature review. The composition, health effects, and industrial uses of brown (*Phaeophyceae*), red (*Rhodophyceae*), and green (*Chlorophyceae*) algae species were examined, and the commercial production of seaweeds and their place in the global market were analyzed. Current scientific studies and market data on the nutritional composition, functional properties, and sectoral applications of seaweeds were evaluated. **Results:** It was determined that seaweed production reached 36 million tons in 2020, with a global market size of 9.9 billion USD. The sector is expected to continue growing at a rate of 2.3% until 2030. Seaweeds are a nutritious food source due to their high protein, mineral, and dietary fiber content. They also contain bioactive components with antiviral, antimicrobial, antioxidant, and anti-inflammatory properties. Brown algae are rich in omega-3 and omega-6 fatty acids, supporting gut health and helping to prevent obesity and hypertension. Red algae contain gelling agents and are widely used in the food and biotechnology industries. Green algae, with their high carbohydrate content and functional bioactive components, are seen as a potential resource in the food and energy sectors. **Discussion and Conclusion:** The evaluation of seaweeds as a sustainable food source is becoming increasingly important due to their environmental and nutritional advantages. Seaweeds offer benefits compared to traditional food production, as they can be cultivated without the need for agricultural land and support ecosystem health. Additionally, their positive health effects due to the functional components they contain increase their potential use in the pharmaceutical and cosmetic industries. With their growing use in the food, pharmaceutical, cosmetic, and animal feed sectors, seaweeds are expected to gain a broader place in the industrial field with further research and development in the future.

Key Words: Sustainable food, seaweed, brown algae, red algae, green algae

ÖZET

Giriş ve Amaç: Küresel gıda üretimi, artan nüfus ve sürdürülebilirlik ihtiyacı nedeniyle yeni kaynaklar arayışına girmiştir. Deniz yosunları, yüksek besin değerleri, biyoremediasyon özellikleri ve tarım alanlarına ihtiyaç duymadan yetiştirilebilmeleri nedeniyle sürdürülebilir bir gıda kaynağı olarak öne çıkmaktadır. Ayrıca, karbon tutma ve su kalitesini iyileştirme gibi ekosistem hizmetleri sunmaları nedeniyle çevresel faydalar sağlamaktadır. Bu çalışma, deniz yosunlarının besin değeri, sağlık yararları ve endüstriyel potansiyelini inceleyerek, gelecekte gıda ve diğer sektörlerde nasıl daha yaygın kullanılabilceğini ortaya koymayı amaçlamaktadır.

Gereç ve Yöntem: Bu çalışma, literatür taramasına dayalı olarak gerçekleştirilmiştir. Kahverengi (*Phaeophyceae*), kırmızı (*Rhodophyceae*) ve yeşil (*Chlorophyceae*) alg türlerinin bileşimi, sağlık etkileri ve endüstriyel kullanımları incelenerek deniz yosunlarının ticari üretimi ve küresel pazardaki yeri analiz edilmiştir. Deniz yosunlarının besinsel bileşimi, fonksiyonel özellikleri ve sektörel uygulamaları ile ilgili güncel bilimsel çalışmalar ve pazar verileri değerlendirilmiştir. **Bulgular:** Deniz yosunlarının 2020 yılında 36 milyon ton üretildiği ve küresel pazar büyüklüğünün 9,9 milyar USD'ye ulaştığı belirlenmiştir. 2030 yılına kadar bu sektörün %2,3'lük bir büyüme oranı ile gelişmeye devam etmesi öngörülmektedir. Deniz yosunları, içerdiği yüksek protein, mineral ve diyet lifi sayesinde besleyici bir gıda kaynağıdır. Ayrıca, antiviral, antimikrobiyal, antioksidan ve anti-inflamatuar özellikler taşıyan biyoaktif bileşenler içermektedir. Kahverengi algler, omega-3 ve omega-6 yağ asitleri bakımından zengindir ve bağırsak sağlığını destekleyerek obezite ve hipertansiyonu önlemeye yardımcı olmaktadır. Kırmızı algler jel oluşturucu bileşenler içermekte olup, gıda ve biyoteknoloji endüstrisinde yaygın olarak kullanılmaktadır. Yeşil algler ise yüksek karbonhidrat içeriği ve fonksiyonel biyoaktif bileşenleri ile dikkat çekmekte olup, gıda ve enerji sektörlerinde potansiyel bir kaynak olarak görülmektedir. **Tartışma ve Sonuç:** Deniz yosunlarının sürdürülebilir bir gıda kaynağı olarak değerlendirilmesi, hem çevresel hem de besinsel avantajları nedeniyle giderek önem kazanmaktadır. Yosunlar, tarımsal alan ihtiyacı duymadan yetiştirilebilmeleri ve ekosistem sağlığını destekleyici etkileri nedeniyle geleneksel gıda üretimine kıyasla avantajlar sunmaktadır. Ayrıca, içerdiği fonksiyonel bileşenler sayesinde sağlık üzerindeki olumlu etkileri, bu bileşenlerin farmasötik ve kozmetik endüstrisinde kullanım potansiyelini arttırmaktadır. Gıda, ilaç, kozmetik ve hayvan yemi sektörlerinde kullanımı artan deniz yosunlarının, gelecekte daha fazla araştırma ve geliştirme ile endüstriyel alanda daha geniş bir yer edinmesi beklenmektedir.

Anahtar Kelimeler: Sürdürülebilir gıda, deniz yosunu, kahverengi algler, kırmızı algler, yeşil algler

WEED CONTROL IN CANNABIS**Dr. Öğr. Üyesi Murat KARACA**

Selçuk Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Konya, Türkiye

ORCID ID: <https://orcid.org/0000-0002-8561-5199>**Zir. Müh. Hanife ÇEÇEN**

Sector Tarım, Konya, Türkiye

ORCID ID: <https://orcid.org/0000-0001-8615-2520>**ÖZET**

Dünyadaki nüfusun hızlı bir artış göstermesi sonucunda meydana gelen gıda açığının karşılanabilmesi için yapılan bitkisel üretimde birim alandan alınan verimin yükseltilmesi gerekmektedir. Tarımın, iktisadi kalkınmanın başlangıç aşamasında sermaye birikim süreci ve ekonomiye çok önemli katkılar sağlayan bir sektör olduğu bilinmektedir. Bu katkılar başlıca şu şekilde sıralamak mümkündür; piyasa, üretim faktörleri, üretim katkısı olarak ifade edilir. Kenevir (*Cannabis sativa* L.), tıbbi ve endüstriyel kullanım alanlarında giderek artan bir önem taşırken, başarılı bir yetiştiricilik için yabancı ot kontrolü önemli bir rol oynamaktadır. Kenevir bitkisi yetiştiriciliğinde çeşitli etmenler (yabancı ot, hastalık, zararlı) verimde kayba neden olurlar. Bunların en önemlilerinden birisi ise yabancı otlardır. Özellikle çok yıllık yabancı otlar, toprak işleme işlemlerini zorlaştırabilir ve tohumluk üretiminde hasat sürecini güçleştirebilir. Ayrıca, bu otlar kontrol maliyetlerini artırabilir ve hem insanlar hem de hayvanlar için zehirli olabilir. Bu nedenlerle, kenevir tarımında yabancı otların etkin bir şekilde kontrol edilmesi önemlidir. Kenevir bitkisi, özellikle başlangıç aşamasında, yabancı otlarla rekabet etme konusunda zayıf olduğundan dolayı yabancı otlar kenevirin büyümesini ve verimini önemli oranlarda olumsuz etkileyebilir. Bu nedenle yabancı ot mücadelesi, kenevirin sağlıklı gelişimi ve yüksek verimliliği için gereklidir. Kenevir yetiştiriciliğinde fiziksel yöntemler içerisinde mekanik veya elle yabancı otların temizlenmesi yer alırken, kimyasal yöntemler ise herbisitlerin uygulanmasını içermektedir. Kimyasal mücadele hızlı sonuç ve maliyet etkinliği nedeniyle sıklıkla tercih edilir. Kimyasal mücadelede herbisitlerin doğru zaman ve dozda uygulanması büyük önem arz etmektedir. Toprak hazırlığı ve ekim stratejileri de yabancı ot mücadelesinde önemli rol oynamaktadır. Ekim öncesinde toprak işleme ve yabancı ot kontrolü, kenevirin gelişim sürecini olumlu yönde etkiler. Ayrıca, kenevirde yapılan münavebeler, yabancı ot baskısını azaltabilir ve toprak verimliliğini artırabilir. Bu derlemede kenevir yetiştiriciliğinde sorun olan önemli yabancı ot türleri ile kenevirde yabancı ot mücadele uygulamaları hakkında bilgi verilecektir.

Anahtar Kelimeler: Kenevir, *Cannabis sativa* L., Yabancı Ot, Fiziksel, Kimyasal, Mücadele.

ABSTRACT

In order to meet the food deficit resulting from the rapid increase in the world population, the yield per unit area in plant production must be increased. It is known that agriculture is a sector that makes significant contributions to the economy and capital accumulation process in the initial stages of economic development. These contributions are mainly; It is expressed as the contribution of production, market and production factors. While hemp (*Cannabis sativa* L.) is

increasingly important in medical and industrial uses, weed control plays an important role for successful cultivation. Various factors (diseases, pests, weeds) cause yield loss in cannabis. One of the most important of these is weeds. Particularly, perennial weeds can complicate soil cultivation and the harvesting process in seed production. Additionally, these weeds can increase control costs and be toxic to both humans and animals. For these reasons, it is important to effectively control weeds in hemp farming. Since the cannabis plant is weak in competing with weeds, especially in the initial stage, weeds can significantly negatively affect the growth and yield of cannabis. Therefore, weed control is necessary for the healthy development and high productivity of hemp. While physical methods in hemp cultivation include mechanical or manual weed removal, chemical methods include the application of herbicides. Chemical control is often preferred due to rapid results and cost effectiveness. In chemical control, it is of great importance to apply herbicides at the right time and dose. Soil preparation and planting strategies also play an important role in weed control. Soil cultivation and weed control before planting positively affects the development process of hemp. Additionally, rotations in hemp can reduce weed pressure and increase soil yield. In this review, information will be given about the important weed species that are problematic in hemp cultivation and weed control practices in hemp.

Keywords: Hemp, *Cannabis sativa* L., Weed, Physical, Chemical, Management.

GİRİŞ

Dünyada nüfusun hızlı bir artış göstermesi sonucunda meydana gelen gıda açığını karşılayabilmek adına bitkisel üretimde birim alandan alınan verimin yükseltilmesi gerekmektedir. Tarımın, iktisadi kalkınmanın başlangıç aşamasında sermaye birikim sürecine ve ekonomiye önemli derecede katkı sağlayan bir sektör olduğu bilinmektedir. Bu katkılar başlıca şu şekildedir; piyasa, üretim faktörleri ve üretim katkısı olarak ifade etmek mümkündür. Ülke ekonomisine katkısından dolayı kenevir bitkisinin ekonomide yan ürünleriyle birlikte birçok alanda katkı sağladığı görülmektedir. İlaç, kâğıt, kozmetik, kumaş gibi dokuma, biyoyakıt, lif, otomotiv sektörü, iplik olmak üzere oldukça geniş bir kullanım potansiyeline sahip olan kenevir bitkisi, petrol ve petrokimyanın yanı sıra tahıl yapımında kullanıldığı her alanda da değerlendirilen ve kullanılabilen endüstriyel bir kültür bitkisidir. Kenevir tohumlarından elde edilen yağlar, sabun ve losyon gibi cilt bakım ürünlerinde yaygın olarak kullanılır. Yağ çıkarıldıktan sonra kalan küspe ise hayvan yemi olarak değerlendirilebilir.

Kenevir (*Cannabis sativa* L.) bitkisinin kökeni Orta Asya olup, Cannabaceae familyası içerisinde yer almaktadır. Lif ve tahıl yapımında kullanılabilen ve yetiştiriciliğinin ilk kez Çin'de 8500 yıl öncesine dayandığı düşünülen bir kültür bitkisidir Daha sonra kenevir bitkisi kıtalara yayılarak özellikle lif kısmı Avrupa'da denizle ilişkili olan ülkelerde ip, halat, çadır ve yelken yapımında kullanılmıştır (Schultes ve ark., 1970; Fike, 2016).

Bazı kaynaklarda Kenevir bitkisinin kökeninin Güney Asya dahil olmak üzere Hindistan alt bölgesi ile Orta Asya olduğu bilinmektedir. Bununla beraber, kenevirin ekim alanıysa Ekvator'dan Kutuplara kadar oldukça geniş bir coğrafik alana yayılım sağlamıştır. Endüstriyel bir kültür bitkisi olarak ıslah edilmeye ve kullanılmaya başlanan ilk bitkilerden biri olan kenevir bitkisi, tarih boyunca birçok ürünün bitkisel hammadde kaynağı olarak ve keyif verici bir madde olarak kullanılmıştır. Bununla beraber kenevir bitkisinden psikolojik bir ilaç olarak Afrika ile Avrasya'da tarih öncesindeki toplumlarda kullanılmasına ilişkin bulgular, kanıtlar arkeolojik kalıntılardan elde edilmiştir. Bunlarla beraber 16. yy.'dan 18. yy.'a kadar keten ve kenevir bitkisi Kuzey Amerika, Avrupa ve Rusya, Avrupa'nın başlıca lif bitkilerinden olmuştur. Yeni Dünya bitkilerinin keşfinden sonra, jüt, pamuk ve diğer tropikal lif bitkilerinin ekim alanlarının artış göstermesiyle dünyadaki kenevir ekim alanları azalmalar görülmüştür. Sentetik liflerin ortaya çıkmasıyla ve birçok ülkede kenevir ekiminin yasal olarak sınırlandırılması sonucu kenevir ekim alanlarındaki azalmalar devamlılık göstermiştir. İkinci

dünya savaşından sonra lif amaçlı kenevir üretimi Doğu Avrupa, Sovyetler Birliği ve Çin ülkelerinde önem kazanmaya başlamıştır. Bindokuzyüztümüş yıllarında kâğıt sanayiinde hammadde olarak kenevir kullanımı artmaya başlamasına rağmen tekstil endüstrisinde kenevir lifi kullanımı azalmıştır. Bunun yanı sıra ürün esrarın yasadışı hele gelmesinden dolayı 1960 yılında yasaklanmıştır. Son yıllardaysa sentetik liflerin yerine doğal liflere olan ilginin artmasının bir sonucu olarak ve kullanım çeşitliliğinin fazla olması sayesinde kenevire karşı ilgi yeniden artış göstermiştir. Son yıllarda kenevir, yüksek biyokütle üretimi nedeniyle bir enerji ürünü ve bir yapı ve inşaat malzemesi olarak da yetiştirilmektedir. Küresel olarak kenevir yetiştiriciliği, karbon tutma özelliği ve fitoremediasyon kapasitesi, kâğıt üretimi için sürdürülebilir bir selüloz kaynağı olması ve bitkinin tüm kısımlarının daha fazla işlenebilmesi nedeniyle sıfır atık üretmesi nedeniyle başarılı bir ticari ürün olarak ortaya çıkmıştır.

Dünyada Kenevir birçok ülkede lif ve tohum amacıyla üretilmektedir. FAO' nun 2020 verilerine göre dünyada 6 ülke tohum amacıyla kenevir üretimi yapılmaktadır. Dünyada 2019 yılında 23.339 ha alanda 76.730 ton kenevir tohumu üretimi yapılmıştır. Son yıllarda, lif için kenevir yetiştiriciliğine ayrılan alan, AB'de 2015'te 20.540 hektardan (ha) 2022'de 33.020 hektara (yüzde 60 artış) önemli ölçüde artmıştır. Aynı dönemde, kenevir üretimi 97.130 tondan 179.020 tona (yüzde 84,3 artış) çıkmıştır. Fransa, AB üretiminin yüzde 60'ından fazlasını oluşturan en büyük üreticidir, bunu Almanya (%17) ve Hollanda (%5) takip etmektedir.

Çin 12.603 ha ile dünya kenevir tohumu üretiminin %93,08'ini oluşturduğu bilinmektedir. Çin'i sırasıyla %3,77 ile Rusya (5.992 ha), %2'yle Şili (3.323 ha) takip etmektedir. Türkiye toplam üretimin sadece %0.03 (48 ha)'ünü oluşturmaktadır. Hektara kenevir tohumu verimi Çin'de 5667,1 kg ile dünyada ilk sırada yer alırken, Rusya'da 482,8 kg., Şili'de 463,1 kg ve Türkiye'de 416,7 kg'dır (Anonim, 2021).

Kenevirin kullanıldığı önemli, karlı sektörlerden biri ilaç sektördür. Kenevir yaprak ve çiçeklerinden elde edilen CBD, THC, CBG ve CBDA gibi kannabinoidler kullanılarak üretilen ilaçlar tedavi amaçlı olarak kullanılmaktadır. İlaç sanayisinde katma değeri en yüksek CBD içerikli ilaçlar en fazladır (Grinspoon, 2007). Dünyada kenevir bitkisinin bir ürün haline gelmesi (Adamovics ve Zeverte-rivza, 2015) ayrıca üretiminin çok büyük ölçüde izine bağlı olması, yasa dışı üretimlerinin de dahil olduğu kenevir verilerine erişimini oldukça zorlaştırmaktadır. Bu sebeple Kanada gibi büyük önem taşıyan kenevir üreticilerine ait veriler, FAO veri tabanlarına ne yazık ki yansımamıştır. Kanada'nın 2018 yılında 31484 ha'lık alanda kenevir bitkisi üretimi gerçekleştirdiği tahmin edilmektedir (Johnson, 2018).

Dünyada 2019 yılında 69.342 ha alanda 174.027 ton kenevir bitkisi lifi üretimi yapılmıştır. Fransa 14.550 ha alan ile dünyada lif amaçlı üretim alanının %20,98'ini oluşturmuştur. Fransa'yı sırasıyla %6,32 ile Şili, %5,79 ile Çin ve %4,47 ile Rusya takip etmektedir. Türkiye toplam kenevir lifi üretim alanının %0.04'ünü oluşturmaktadır. Hektara kenevir lif verimi İtalya'da 8.472,5 kg, Hollanda'da 7.884 kg, Polonya'da 7.628,4 kg, Türkiye'de 964,3 kg'dır. Dünya kenevir lifi üretiminde Fransa %44,85, Çin %8,35, Hollanda %8,08, Polonya %8,02, Avusturya %5,13, İtalya %4,43'lük bir paya sahiptir. Türkiye'nin kenevir lifi üretimindeki payı sadece %0.02'dir.

	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	Değişim (%) ²
Alan ¹ (ha)	19.488	19.621	25.000	24.640	25.768	24.983	36.291	49.907	27.694	23.339	-15,73
Verim ¹ (hg/ha)	35.970	34.875	44.988	37.400	46.151	36.458	43.976	51.919	39.969	32.876	-17,75
Üretim ¹ (ton)	70.099	68.429	112.470	92.153	118.923	91.084	159.595	259.111	110.691	76.730	-30,68

Kaynak: ¹/FAO 2021, ²/Verisi bulunan son iki pazarlama yılının değişimini göstermektedir.

Tablo 1. Dünya kenevir tohumu verileri


	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	Değişim (%) ²
Alan ¹ (ha)	40.638	40.689	40.901	42.261	43.911	44.943	42.204	40.557	59.743	69.342	16,07
Verim ¹ (hg/ha)	11.703	12.758	13.525	14.064	17.504	17.672	16.627	14.410	32.620	25.097	-23,06
Üretim ¹ (ton)	47.557	51.909	55.320	59.435	76.863	79.422	70.173	58.441	194.880	174.027	-7,01

Kaynak: ¹/FAO 2021, ²/Verisi bulunan son iki pazarlama yılının değişimini göstermektedir.

Tablo 2. Dünya kenevir lifi verileri

Küresel Kenevir Pazarı büyüklüğünün 2023 yılında 24,55 milyar ABD doları değerinde olduğu ve 2024 yılında 25,16 milyar ABD dolarından 2031 yılına kadar 33,73 milyar ABD dolarına çıkacağı ve tahmin dönemi boyunca %4,28'lik bir Bileşik Büyüme Oranı sergileyeceği tahmin edilmektedir. İnşaatçılar ve geliştiriciler geleneksel inşaat malzemelerine sürdürülebilir alternatifler aradıkça kenevir pazarı genişlemektedir.

Şekil 1. AB' de kenevir lifi yetiştiriciliğinde kullanılan arazi alanı



Küresel kenevir pazarı:

Kuzey Amerika	Avrupa	Pasifik-Asya	Afrika ve Orta Doğu	Latin
Meksika	Fransa	Çin	GCC	Brezilya
Kanada	İngiltere	Rusya	Kuzey Afrika	Arjantin
ABD	İspanya	İtalya	Güney Afrika	
	Almanya		Güney Kore	
	Hindistan			
	Japonya			

Türkiye’de kenevir bitkisinin üretimi 2004 yılına kadar sadece Samsun, Çorum, Kütahya ve Kastamonu il ve ilçelerinde yapılmıştır (Aydoğan ve ark., 2019). Kenevirin yeniden bir tarım ürünü olarak gündeme gelmesi, 9 Ocak 2019 tarihinde gerçekleştirilen “Cumhurbaşkanlığı Hükümet Sisteminde Yerel Yönetimler Sempozyumunda” gerçekleşmiştir. Bu kapsamda 11 Ocak 2019 tarihinde Tarım ve Orman Bakanlığı, Sanayi ve Teknoloji Bakanlığı, Sağlık Bakanlığı, TÜBİTAK ve Samsun Ondokuz Mayıs Üniversitesi’nin katılımıyla “Türkiye’de Endüstriyel Kenevir Yetiştiriciliği Raporu ve Eylem Planı” hazırlanmıştır. Eylem planında öne çıkan konular; kenevir üretiminin kısa vadede yerli popülasyonlarla artırılması, yerli Narlısaray popülasyonundan tohum çoğaltma işlemlerinin gerçekleştirilmesi, kenevire dayalı sanayinin ve mekanizasyon sistemlerinin geliştirilmesi, yerli çeşitlerin olması ve geliştirilmesi için tohum tescil çalışmalarının en kısa zamanda tamamlanarak sertifikalı tohum kullanımına geçiş yapılması ve öncelikli yatırım alanlarının belirlenmesi şeklindedir (TİGEM, 2019).

Diğer birçok ülkede olduğu gibi Türkiye’de de kenevir bitkisi tohumunun üretilmesi çok uzun bir süre arka planda kalmıştır. Bunun başlıca nedenlerinden birisi de yapraklarının esrar yapımında kullanılması nedeniyle çiftçilerin bu bitkiye karşı önyargı besleyerek yaklaşımının etkisi fazla olduğu görülmüştür. Ekstra olarak mevzuatlar, kanun ve yönetmelik de üretimin sınırlı kalmasında başlıca rol oynamışlardır. Böylece kenevir bitkisi ekimi-üretimi yapacak üreticilerin il veya ilçe müdürlüklerinden yasal olarak izin alması zorunlu olmasının yanı sıra üretim sürecinde sürekli olarak düzenli bir şekilde kontrollerin yapılması zorunluluğu da bulunmaktadır. İzinsiz bir şekilde üretimi yapılan kenevir bitkisi hangi sebebe dayalı olursa olsun kanun hükümlerine göre imha edilmekte olup üretici hakkında cezai işlemler uygulanmakta ve yasal kovuşturma yapılmaktadır. İzinli kenevir ekimi yapılan alanlar da il ve ilçe müdürlükleri teknik elemanları tarafından ekimden hasat zamanına kadar düzenli olacak şekilde kontroller gerçekleştirilmektedir. Tarım Orman Bakanlığı tarafından yapılan değerlendirmeler sonucunda Türkiye’de 2019 yılı itibariyle endüstriyel amaçlı kenevir ekiminin 19 ilde serbest bırakılmasına karar verilmiştir. Bu kararın ardından Türkiye’de lif amacıyla kenevir üretiminde artış beklenmektedir. 29.09.2016 tarihli Resmî Gazetede yayınlanan yönetmelik kapsamında Türkiye’deki 19 il; Antalya, Amasya, Burdur, Bartın, Çorum, İzmir, Kayseri, Kastamonu, Kayseri, Karabük, Malatya, Ordu, Rize, Sinop, Samsun, Tokat, Uşak, Yozgat ve Zonguldak. Bu illerin bütün ilçelerinde de üretim yapılabilmektedir. İzin verilen il ve ilçelerin dışında kenevir bitkisi yetiştiriciliği kesinlikle yasaktır. Fakat yönetmelik çerçevesinde belirlenen hükümlere uymak şartıyla bilimsel araştırma amacıyla ana veya tali bitki olarak kenevir bitkisi yetiştiriciliğine birinci fıkrada belirlenen bölgeler dışında da Bakanlıkça izin verilebilir. Ayrıca kenevir bitkisi tarımının geliştirilmesine yönelik olarak Yozgat Bozok Üniversitesi ve Samsun ilinde Ondokuz Mayıs Üniversitesi bünyesinde olmak üzere iki adet "Kenevir Araştırmaları Enstitüsü" kurulmuştur. Türkiye’de kenevir üretim alanı ve üretim miktarları iller bazında değerlendirildiğinde; tohum amaçlı ekim ve üretimin Kastamonu, Kayseri, Kütahya, Samsun ve Yozgat illerinde olmak üzere 5 ilde yapıldığı, bu

iller arasında Samsun ilinin diğer 4 ilin ekim ve üretim miktarından daha fazla paya sahip olduğu görülmektedir.

Şekil 2. Türkiye’de izin verilen kenevir ekim alanları



Türkiye’de tohum amacı ile kenevir bitkisi üretimi 2005 yılında 650 da iken, hızlı bir azalış göstererek 2013 yılında 7 da’a kadar düşmüştür. Üretimse yalnızca Samsun ili ve ilçelerinde yapılmıştır. 2018 yılında ekim alanı 59 da’a, 2019 yılında 536 da’a 2020 yılında ise, 4.252 da’a yükselmiştir. Kenevir tohumu verimi dalgalı bir seyir izleyerek 2005 yılında 20 kg/da, 2011 yılında 57 kg/da, 2013 yılında ise 143 kg/da’a kadar yükselmiştir. Kenevir tohumu verimi 2018 yılında 51 kg/da, 2019 yılında 42 kg/da’a inmiştir. 2020 yılındaysa, 64 kg/da’a yükselmiştir. Türkiye’de kenevir tohumu üretim miktarı 2005 yılında 13 ton iken, üretim alanında azalmaya bağlı olarak 2013 yılında 1 tona kadar düşmüştür. Üretim miktarı, ekim alanlarında artışa bağlı olarak 2018 yılında 3 tona, 2019 yılında 20 tona, 2020 yılında ise, 273 tona yükselmiştir.

Türkiye’de lif amaçlı kenevir bitkisi ekim alanı, verimi ve üretim miktarı yıllara göre farklılık gösterdiği görülmüştür. Lif amaçlı kenevir bitkisi üretimi 2005 yılında 650 dekar iken, 2011 yılında 157 dekara, 2014 yılında ise 10 dekara kadar gerilediği görülmüştür. Kenevirin tekrar gündeme gelmesi ile beraber kenevir lifi ekim alanı 2018 yılında 55 dekara, 2019 yılında ise 151 dekara, 2020 yılında ise 96 dekara yükselmiştir. Kenevir lifi verimi ise 2005 yılında 85 kg/da iken, 2014 yılında 100 kg/da’a, 2019 yılında ise 126 kg/da’a, yükselmiştir. 2020 yılında ise 94 kg/da’a düşmüştür. Yapılan ıslah ve yeni çeşit çalışmalarının etkisiyle kenevir lifi veriminde artış sağlanmış fakat beklenildiği kadar etkili olmamıştır. Kenevir lifi üretim miktarı ise 2005 yılında 55 ton iken, 2012 yılında 6 tona, 2015 yılında ise 1 tona kadar gerilemiştir. Kenevir lifi üretimi 2016-2018 dönemi ortalaması olarak 7 ton iken, üretim 2020 yılında 9 ton olarak kaydedilmiştir.

	2005	2010	2015	2018	2019	2020	Değişim (%) ²
Kenevir Tohumu Ekilen Alanı' (da)	650	221	10	59	536	4.252	396,64
Kenevir Tohumu Üretim Miktarı' (ton)	13	7	1	3	20	273	682,5
Kenevir Tohumu Verimi (kg/da)	20	32	100	51	42	64	47,62
Kenevir Lifi Ekilen Alanı' (da)	650	221	10	55	160	101	-36,87
Kenevir Lifi Üretim Miktarı' (ton)	55	10	1	7	19	9	-52,63
Kenevir Lifi Verimi (kg/da)	85	45	100	127	126	94	-25,40

Kaynak: ¹TÜİK 2021, ²Verisi bulunan son iki pazarlama yılının değişimini göstermektedir

Tablo 3. Türkiye kenevir tohumu ve kenevir lifi verileri

TÜİK verilerine göre 2022 yılında 365 dekar alana ekim yapılmışken 2023 yılında 2111 dekar alan kenevir ekimi yapılmıştır. Yine 2022 yılında üretim 31 ton iken 2023 yılında bu üretim 359 ton olarak ciddi bir artış sağlamıştır.

Yıllar	Ekim alanı (da)	Üretim (Ton)	Verim (kg/da)
2019	160	19	119
2020	101	9	89
2021	324	21	65
2022	365	31	85
2023	2117	359	170

Kaynak: Tarım ve Orman Bakanlığı

Tablo 4. Türkiye 2019-2023 yılları kenevir ekim alanı, üretimi ve verimi (TÜİK, 2023)

*FAO (2021) verileri göre 2018 ve 2019 yılları Türkiye'nin kenevir lifi ekim alanı, üretim miktarı ve veriminin TÜİK (2020) verilerinde farklılık gösterdiği tespit edilmiştir.

YABANCI OT MÜCADELESİ

Yabancı Ot Nedir?

“Tarım alanları içinde veya dışında bulunan ve yetişmesi arzu edilmeyen bitkiler” şeklinde tanımlandığı gibi “İnsanoğlunun istemediği yerde yetişen ve zararı yararından fazla olan bitkiler” şeklinde de tanımlanır (Uygur ve ark., 1984; Günçan, 2019). Yabancı otlar kültür bitkileri ve diğer yabancı otlarla yüksek bir rekabete girmektedirler. Bu rekabet daha çok mineral maddeler, ışık ve su yönünden olmaktadır ve bunun sonucu olarak da kültür bitkilerinin veriminde azalmalar meydana gelmektedir (Günçan ve Karaca, 2023). Diğer taraftan yabancı otların neden olduğu verim kaybı kültür bitkisinin türüne bağlı olarak da değişmektedir. Yabancı otlar sahip oldukları rekabet güçleri, vejetatif ve generatif yolla üremeleri, allelopatik etkileri ve genetik çeşitlilikleri sayesinde çok farklı ekosistemlere uyum sağlayabilmekte, parazitik etkileri ile kültür bitkilerini etkilemekte, tarım alanlarında büyük verim ve kalite kayıplarına neden olmakta, hastalık ve zararlılara konukçuluk yapmaktadırlar. Diğer taraftan yabancı otların neden olduğu verim kaybı kültür bitkisinin türüne bağlı olarak da değişmektedir.

Ayrıca mücadeleleri için yapılan faaliyetler nedeniyle zaman ve kaynak israfına neden olabilmekte, insan sağlığı ve çevre için risk oluşturabilmektedirler (Uygur ve ark., 1984; Özer ve ark., 2001; Günçan, 2009).

Kenevir Bitkisinde Yabancı Ot Zararı

Tarlalarda yabancı otların gelişmesinin önlenmesi durumunda kenevir gelişiminin hızlı olduğu, kalın yaprak oluşturduğu, gölgeleme yaparak yabancı otlara karşı rekabet gücünü yükselttiği ve verimde artış sağladığı bildirilmiştir (Poisa ve Adamovic, 2010; Rehman ve ark., 2013). Kenevir son derece rekabetçi olmasına rağmen uygulanan ekim sistemine bağlı olarak yabancı otların etkinliğinde farklılık olduğu görülmektedir. Dolayısıyla kenevirin rekabet gücü çok farklı unsurlardan etkilenmektedir. Bu çerçeveden üretimde kullanılan kenevir tohumunun çeşit özellikleri büyük öneme sahiptir. Nitekim genel olarak fenotipik olarak lif üretimi için kullanılan çeşitlerden farklı olan bazı tohumluk çeşitlerin kanopi (bitki örtüsü) gelişimi daha düşük olduğundan, yabancı ot rekabetine hassas olabilmektedirler (McPartland ve ark., 2004). Özellikle kısa boylu tohum çeşitleri (FIN-314 gibi) hasat döneminde dahi toprak yüzeyini tamamen örterek gölgeleyememektedirler. Bunun bir sonucu olarak da bu çeşitler yabancı ot rekabetine karşı son derece duyarlı hale gelmektedirler (McPartland ve ark., 2000). Tohum veya lif üretimi amacıyla kullanılan çeşitlerin rekabet gücü bölgesel/ülkesel olarak da farklılık göstermektedir. Örneğin Avrupa ve Kanada'nın nispeten yüksek bölgelerinde geliştirilen çeşitler; nispeten ılıman veya daha düşük yükseltilere sahip bölgelerde/ülkelerde yetiştirildiklerinde yabancı otlara karşı rekabet güçlerinde farklılık olduğu saptanmıştır (Sandler ve ark., 2019). Zira iklim ve toprak koşulları bölgedeki yabancı ot tür ve yoğunluklarını etkilediği gibi kültür bitkisinin gelişimini de etkileyebilmektedir. Yabancı otların rekabet gücü de doğal olarak tarladaki/bölgedeki yabancı ot florasına göre farklılık göstermektedir (Önen vd., 2016). Nitekim tek yıllık yabancı otlardan farklı olarak, çok yıllık *Ambrosia trifida* ile bazı bambu türlerinin (*Bambusa* spp., *Dendrocalamus* spp. ve *Phyllostachys* spp.) kenevirden daha hızlı büyüebildiği ve keneviri gölgeleyebildiği belirtilmektedir (McPartland ve ark., 2000).

Diğer yandan bazı yabancı ot türleri kenevire son derece benzer morfolojik özellikler gösterirler. Keneviri taklit eden bu yabancı otları ise üreticiler genel olarak gözden kaçırabilmektedir. Bu nedenle de bu bitkiler tarlada sorun haline gelebilmektedir. Örneğin (*Galeopsis* spp.) fidelerini kenevir fidelerinde ayırmak oldukça güçtür. Small (1975) tarafından da bildirildiği üzere renk otunun (*Datisca cannabina* L.) erkek bitkileri de profesyonel taksonomistleri dahi yanıltacak kadar keneviri andırır (McPartland ve ark., 2000).

Geç ekilen ve özellikle tohumluk üretimi amacıyla sıra araları geniş tutulan, bu nedenle de birim alandaki bitki yoğunluğu düşük olan ekim sistemlerinde de yabancı otlar önemli bir sorun olarak karşımıza çıkabilmektedir (Darby ve ark., 2016). Zira tarla yüzeyi kenevir tarafından tamamen örtülmediğinden yabancı otlar açık kalan alanlarda hızla gelişerek sorun haline gelebilmektedirler. Kenevir geç ekildiğinde ise özellikle dar yapraklı yabancı ot türleri genç kenevir fidelerini rahatlıkla bastırmakta ve fideleri adeta boğabilmektedirler (McPartland ve ark., 2000). Nitekim Bocsa ve Karus (1998) lif üretimi için de olsa düşük tohum yoğunluğunda ekilen veya zayıf çıkış görülen tarlalarda kenevirin toprak yüzeyini ancak 100-150 cm boyu ulaştığında örttüğünü, toprak yüzeyinin daha geç kaplanması ise yabancı otların gelişimine izin verdiğini belirtmektedir. Sık ekimde kenevirde ise bitkiler 50 cm boya varmadan toprağı bitki örtüsü ile tamamen kapladığından yabancı otlar sorun olmaktan çıkmaktadır.

Daha önce ülkemizde kenevir alanlarında sorun olan yabancı otların belirlenmesine yönelik olarak herhangi bir çalışmaya rastlanmamış olup nispeten az sayıda tarım alanından elde edilmiştir. Tunçdemir (1987) tarafından Kastamonu (Taşköprü)'da kenevirde sorun olan soğan sak nematodu ile mücadele olanaklarının araştırılması amacıyla yürütülen bir çalışmada deneme parsellerinde yabancı tere (*Lepidim draba*), köygöçüren (*Cirsium arvense*), akkazayağı (*Chenopodium album*), tarla sarmaşığı (*Convolvulus arvensis.*), Ballıbaba (*Lamium sp.*), hindiba (*Cichorium intybus*) ve sarmaşık çobandegneği (*Polygonum convolvulus*) türlerinin yaygın olarak bulunduğu belirtilmektedir. Kastamonu, Sakarya, Tokat ve Samsun'da küçük üretim ve deneme alanlarında yapılan gözlemler sonucunda özellikle Poaceae, Chenopodiaceae, Polygonaceae, Asteraceae, Amaranthaceae ve Brassicaceae familyalarına dahil bazı türlerin erken dönemde önemli yoğunluk oluşturabildiği saptanmıştır.

Sonuç olarak; genel olarak lif üretiminde kullanılan çeşitlerin rekabet gücünün yüksek olması ile erken ve sık ekim nedeniyle yabancı otlar önemli ölçüde baskı altına alınmasına rağmen özellikle kenevir tohumu elde etmek için yapılan üretimde yabancı otlar sorun olabilmektedir. Zira tohum üretimi için yapılan kenevir tarımında, genel olarak daha geç ekim ve daha düşük kültür bitkisi yoğunluğu veya düzensiz ekimin bir sonucu olarak toprak yüzeyi kısa zamanda örtülerek güneşin geçişi engellenmemektedir. Bu durum yabancı ot tohumlarının çimlenmesine ve fide gelişimine yardımcı olmaktadır. Dolayısıyla kenevir tohumlarının ekim zamanı ve ekim oranı ile bu iki faktörün etkileşimleri, toprak ve iklim koşulları, alanda bulunan yabancı ot türleri vb unsurlar (kültür bitkisinin verim ve kalitesi yanında) yabancı ot yoğunlukları dolayısıyla da yabancı ot rekabetinden kaynaklanan verim kayıpları üzerinde büyük etkiye sahiptir (Van der Werf ve ark., 1995; McPartland ve ark., 2000). Diğer bitki koruma etmenleri için de vurgulandığı üzere; ülkemizde kenevir ekim alanlarının genişlemesiyle beraber bölgesel yabancı ot tür ve yoğunluklarında önemli farklılıkların olabileceği düşünülmektedir. Muhtemel farklılıklara bağlı olarak da bölgesel düzeyde çeşitli yabancı ot kültür bitkisi etkileşimlerinin ortaya çıkabileceği (parazitik etki, rekabet vb) ve yabancı otların verim ve kalite açısından beklenenden daha büyük sorunlara yol açabileceği öngörülmektedir.

Yabancı otların parazitik etkileri

Çiçekli tam parazitik yabancı ot türleri Kenevirde sorun oluşturan en önemli yabancı otlardır. Genel olarak klorofil içermeyen bu bitkiler gelişimleri için ihtiyaç duydukları besin maddeleri ve suyu farklılaşan kökleri (haustorium) vasıtasıyla konukçularının kök veya gövdelerinden alırlar (Özer ve ark., 1999; Özer ve ark., 2001). Dünyada 2500'den fazla parazit bitki türü bulunmasına rağmen şu ana kadar on türden daha az parazitik bitkinin kenevirde sorun oluşturabildiği rapor edilmiştir (McPartland ve ark., 2000). Dünya genelinde *Phelipanche* (*Orobanch*) cinsine (canavar otları) dâhil 3 ve *Cuscuta* cinsine (kükütler) dahil 5 60 çiçekli parazitik bitki türünün kenevirde sorun oluşturabildiği görülmektedir.

Cuscuta L. (Küskütgiller - Cuscutaceae)

- *Cuscuta campestris* (L.) Yunck. Küsküt
- *C. europea* L. Avrupa küskütü
- *C. pentagona* Engelm. Beş köşeli küsküt
- *C. epilinum* Weihe Küsküt
- *C. suaveolens* Ser. Püsküllü küsküt

Orobanche L. (*Phelipanche* - Canavar otugiller - Orobanchaceae)

- *Orobanche aegyptiaca* Pers. Mısırlı canavar otu
- *O. ramosa* L. Mavi çiçekli canavar otu
- *O. cernua* Loeffl. Boğumlu canavar otu



Amaranthus retroflexus (Kırmızı köklü tilki kuyruğu)
(Darıcan)



Echinochloa crus-galli



Solanum nigrum (Köpek üzümü)
sarmaşığı)



Convolvulus arvensis (Tarla



Chenopodium album (Sirken)
pıtrağı)



Xanthium macrocarpom (Domuz)



Cynodon dactylon (Köpek dişi ayrığı)
arpası)



Hordeum murinum (Duvar)



Potulaca oleracea (Semiz otu)



Setaria spp. (Kirpi darı)



Sorghum halepense (Kanyaş)



Cyperus rotundus (Topalak)



Sinapsis arvensis (Yabani hardal)
(Köygöçüren)



Cirsium arvense L.



Capsella bursa-pastoris L. (Çoban çantası)
değneği)



Polygonum aviculare L. (Çoban
değneği)

Şekil 2. Kenevir bitkisinde sorun olan önemli bazı yabancı otlar

Kenevirde Yabancı Ot Mücadele Yöntemleri

Bütün kültür bitkilerinde olduğu gibi kenevirde de yabancı otlarla mücadelede tek bir yabancı ot mücadelesi şekli yerine, entegre mücadelenin uygulanması daha başarılıdır.

Kültürel önlemler

- **Sertifikalı ve temiz tohumluk kullanımı:** Tarlalarda ekim için kullanılacak kenevir tohumluğunun temiz ve yabancı ot tohumlarıyla bulaşık olmaması gerekmektedir. Bu nedenle tarlalarda üretimi geciktiren yabancı ot türleri kenevir liflerine yapışarak lif kalitesini düşürebilir ya da kenevir tohum miktarının azalmasına sebep olabilmektedir. Yabancı ot tohumlarıyla karışık olarak ithal edilen kenevir tohumluklarına dikkat edilmelidir. Gerekirse uygun selektörlerden geçirildikten sonra tarlalara ekim yapılmalıdır. Eğer sertifikalı ve yabancı ottan ari tohumluk var ise tercih edilmelidir.

- **Doğru ve uygun kenevir çeşidi seçimi:** Bölgeye, lokasyona hatta tarlaya göre uygun ve geliştirilmiş olan kenevir çeşitleri kullanılmalıdır. Eğer belirlenen kenevir çeşidi yabancı ot türleriyle rekabet (su, ışık, besin vb.) edemezse kenevire ait verim ve verim unsurları azalabilmektedir. Kenevir bitkisinin tercih ettiği topraklarda (pH, tekstür, su tutma kapasitesi vs.), uygun iklim koşullarına sahip bölgelerde (sıcaklık, yağmur, nem vb.) ve yabancı otlarla rekabet gücü yüksek çeşitler kullanılarak yetiştiricilik yapılmalıdır (Özer ve ark., 2001; Jankauskiene ve ark., 2014).

- İnokulum kaynaklarının uzaklaştırılması: Hasattan sonra tarlalarda inokulum olabilecek bitki artıkları bırakılmayıp, tarlalardan uzaklaştırılmalıdır. Çünkü çevrede bulunan yabancı otlar kenevirde hastalığa neden olan patojenlerin ya da zararlı böceklerin hayat döngülerini devam ettirerek (konukçuluk özelliği) bir sonraki ekilen kenevir bitkilerine patojenleri ya da zararlı böcekleri taşırlar.

- Alet ve ekipman temizliği: Yetiştiricilik yapılan tarlalarda bir tarladan diğerine geçerken yabancı ot tohumları veya vejetatif çoğalma organları taşınabilmektedir. Bu yüzden kullanılacak olan çapalama ve mibzer gibi aletlerin temizliği önemlidir.

- Erken veya geç ekim ile ekim sıklığı: Tohum üretimi amacıyla yapılan ekimlerde, bitkinin fazla uzamasını engellemek için geç ekim tercih edilerek vejetasyon süresi kısaltılmalıdır (Basbag ve ark., 2020; Kale, 2023). Erken veya geç ekim zamanında kenevir kültür bitkisi yabancı ot türlerine göre daha hızlı ve gür gelişebilmekte, dolayısıyla yabancı ot tohumlarının çimlenme kabiliyetini azaltabilmektedir. Dolayısıyla erken ve sık ekimde kenevir bitkisi yabancı ot çıkışlarını baskılamaktadır. Yabancı otlarla mücadelede en önemli yöntemin ise sık ekim olduğu kenevirde, bitki sayısının artmasıyla yabancı ot popülasyonlarının baskılanabildiği ve farklı kenevir çeşitlerinin kullanılmasının önemli olduğu kaydedilmiştir (Vera ve ark., 2006; Hall ve ark., 2014). Geç ekimde ise tarlalarda sorun olan yabancı ot popülasyonları ortamdaki uzaklaştırıldıktan sonra ekim rahatlıkla yapılabilir. Sık ekimde birim alanda kenevir bitki sayısının artışı sayesinde toprak yüzeyi kaplanır ve yabancı ot popülasyonlarının gelişmesi önlenir (Ontario, 2024a). Bu durumda, kenevir bitkisi hem vejetatif hem de generatif olarak yeterli gelişim gösteremediğinden, lif ve tohum verimi azalır (Van der Werf, 2002). Daha dar sıra aralıkları lif verimi ve kalitesini, daha geniş sıra aralıkları ise tohum verimi ve kalitesini artırmaktadır (Göre ve Kurt, 2021; Kale, 2023). Tabii ki kenevir kültür bitkisinin istediği iklimsel ve tarımsal istekleri baz alınarak erken veya geç ekim yöntemi uygulanmalıdır.

- Gübreleme ve sulama: Kenevir gübrelemesi toprak analizlerine göre, tavsiye edilen gübreleme dozunda ve zamanında yapılmalıdır. Kenevir bitkisi toprakta yer alan bütün mineralleri toplamaktadır, buna ek olarak zengin gübreleme işlemleri gerektirmektedir. Gübreleme yaparken de yabancı ot popülasyonlarının tarlalara uygulanan gübrelere ortak olacağı da unutulmamalıdır. Ehrensing (1998), dünya çapındaki verilere dayanarak kenevir tarlalarında tavsiye edilen gübre miktarlarını 4-20 kg da-1 (NH₄ NH₃), 3-12 kg da-1 (P₂O₅) ve 0-20 kg da-1 (K₂O) olarak ifade etmektedir. Fosfor ve potasyum içeren gübreler genellikle tohum yatağı hazırlığı esnasında veya ekim sırasında toprağa verilir. Azotlu gübreler ise hızlı yıkanma özelliği nedeniyle ekimle birlikte ve sonraki büyüme dönemlerinde uygulanır (Basbag ve ark., 2020). Bu nedenle yabancı otun gelişme dönemi ve gübreleme sayısı önemlidir. Kullanılacak olan doğal veya kompost çiftlik gübrelerinin iyi yanmış olması gerekmektedir. Hayvanların dışkıсында sindirilemeyen yabancı ot tohumlarının tarlalara çiftlik gübresi şeklinde uygulanması yabancı ot popülasyonlarının tarlalarda çoğalmasını sağlar. Kurak olmayan bölgelerde bir veya iki kez sulama yapmak yeterliyken, kurak bölgelerde sulama sayısı artabilmektedir (Ontario, 2024b). Sulama yapılacak tarlalarda suyla bulaşabilecek tohumların yayılmasını önlemek için su kanal ve kanaletlerin kenarında bulunan yabancı otlar tohum bağlamadan önce biçilmelidir. Sulama borularının önüne tül benzeri vs. materyaller konularak suyla bulaşan yabancı ot tohumlarının taşınması önlenmelidir. Kök gelişimini büyük ölçüde tamamlamış bitkiler kuraklığa dayanıklı olsa da, kuraklık aşırı seviyelere ulaştığında bitkiler zayıf kalır ve normal gelişim sürecinden daha erken çiçeklenme ve tohum üretimi gösterirler (Basbag ve ark., 2020).

Mekanik mücadele

- Tohum yatağı ve ekim hazırlığı: Yetersiz toprak hazırlığı yapılmış, kötü drenaj özelliklerine sahip, tınlı, asidik, kumlu veya ağır bünyeli topraklar, kenevir yetiştiriciliği için uygun değildir (Struik ve ark., 2000). Kenevir tohumu hızlı çimlenebilmek için uygun toprak tavında olmalıdır. Aksi halde, kenevir tohumunun çimlenmesi gecikerek, diğer rekabet gücü yüksek olan yabancı

ot tohumlarının çimlenmesine imkan verilir. Kenevir ekimi genellikle bahar döneminde (Nisan) mibzer kullanılarak gerçekleştirilmektedir. Ekim işleminin sıralı ve çok düzgün bir şekilde gerçekleştirilmesi, kenevir fide çıkışlarında ve gelişmesinde etkilidir. Kenevir tohumunun 2-3 cm derinlikte toprağa yüzeysel ekimi yeterlidir (Desanlis ve ark., 2013). Ancak kenevirde lif ya da tohum üretimi için gerekli olan ekim şekli değişebilmektedir.

- **Çapalama:** Uygun çapalama aralığına sahip aletlerle, kenevir bitkisi düzenli olarak çapalanmalıdır. Toprak özelliğine bağlı olarak, toprak hazırlığı için kullanılan aletler ve toprak işleme sayısı yabancı ot popülasyonlarının mücadelesinde önemlidir. Kenevirde ilk çapalama kenevir bitkileri 5- 10 cm boya ulaşınca gerçekleşir (Anonim, 2025). Kenevirde uygun ekim zamanının saptanması yanında, yeterli bitki yoğunluğu bir araya geldiğinde herhangi bir çapalama işlemine dahi gereksinim duyulmayabilir (Jankauskiene ve ark., 2015). Tarlalarda kenevirin erken gelişme döneminde küçük lokal alanlarda yabancı otlar el çapası yardımıyla da alınabilmektedir. Ancak sık ekimde bu tarz çapalama işlemleri yapılamamaktadır. Fiziksel mücadele

- **Solarizasyon:** Bu işlem boş ve küçük tarım alanlarında şeffaf plastik örtü benzeri malzemelerle güneş enerjisi kullanılarak gerçekleştirilmektedir. Burada amaç toprak ısısını yükselterek yabancı ot tohumlarını öldürmektir, hatta sorun olan toprak kökenli patojenlerin ve nematodların azalmasını sağlar.

- **Malçlama:** Daha geniş ekim aralığına sahip tarlalarda toprak yüzeyi farklı malç materyalleri kullanılarak (saman malçı, siyah malç tekstili, talaş vs.) örtülebilmektedir. Burada da amaç solarizasyona benzer şekilde toprağın ısısını yükseltmek ve nem kaybını azaltmaktır. Bu sayede yabancı ot çıkışları önlenerek kontrolü sağlanmış olur.

Ekim nöbeti sistemlerinin uygulanması

Günümüzde yabancı otlarla direk mücadelede ekim nöbeti en önemli yöntem haline gelmiştir. Sadece tarlalara kenevir ekimini yapmak yeterli olmamaktadır. Kenevir ekilen tarlalarda farklı ürünlerin değişimi, kültür bitkilerinin ekim ve hasat tarihlerini değiştirmekte, gübreleme isteklerini azaltmakta ve ana zararlı yabancı ot türlerinin biyolojilerinin tamamlanmasını önlemektedir (O'Donovan ve ark., 2007; Vencill ve ark., 2012). Çünkü ana zararlı yabancı otlar kültür bitkilerine göre özelleşmiştir (Labrada, 2006). Bu oluşan döngünün kırılmasında ise ekim nöbeti önemli bir mücadele yöntemi olmuştur. Ekim nöbetinde kültür bitkisinin önemi, ekosisteme olan katkısı ve tarıma sağladığı faydalar unutulmamalıdır. Münavebe uygulamaları, yağ verimini artırabilir ve hastalıkları azaltabilir. Kenevir, tahıllar ve sebzelerle uyumlu olarak münavebe edilebilir (Gorchs ve ark., 2017). Ayrıca, azot açısından zengin baklagil bitkileri, özellikle yonca veya üçgül, kenevirle yapılan münavebede oldukça faydalı olabilir (Basbag ve ark., 2020).

Kimyasal Mücadele

Kimyasal mücadele, özellikle herbisit kullanımı hem dünya genelinde hem de ülkemizde yabancı ot kontrolünde yaygın olarak tercih edilen bir yöntemdir. Diğer yöntemlere göre uygulanmasının kolay olması, hızlı sonuç vermesi ve maliyetinin düşük olması nedeniyle bu yöntem sıklıkla kullanılmaktadır (Kiely ve ark., 2004). Üreticiler tarafından kısa sürede sonuç alınması, ekonomik olması ve uygulanabilirliğinin kolay olmasından ötürü çok fazla kimyasal (herbisit) kullanımı tercih edilmektedir. Ancak ülkemizde kenevir yetiştiriciliğinin devlet kontrolünde ve dar alanlarda yapılması, bunun yanında yaygın ekilen ürün olmaması herbisit tüketimini önlemektedir. Tarım ve Orman Bakanlığı-Gıda ve Kontrol Genel Müdürlüğü (GKGM) tarafından şuan ruhsatlandırılmış bir herbisit bulunmamaktadır (GKGM, 2025). Fakat istilacı ve potansiyel durumda bulunan kenevirde sorun oluşturabilecek ana zararlı yabancı ot türleri bilinmeli, ekonomik zarar eşiği çalışmaları yapılmalıdır.

Türkiye'de de Kale (2023) tarafından yapılan çalışmada pendimethalin, pyroxasulfone herbisitlerinin kenevirde yabancı ot mücadelesinde kullanılabileceği gibi lif veriminin

artmasına katkı sağladığı kaydedilmiştir. Kenevir yetiştiriciliğinde benfluralin, dimethachlor + clomazone, pendimetalin + clomazone, S-metalochlor, linuron, herbisitleri yabancı otları kontrol altına sağlamakta başarılı olup fitotoksisite oluşturmadığı ve kenevir yetiştiriciliğinde kullanılabileceği belirlenmiştir.

Amerika ve Kanada'da kenevir yetiştiriciliğinde ruhsatlandırılmış herbisitler bulunabilmektedir.

Amerika'da *Amaranthus* spp., *Chenopodium* spp., *Echinochloa* spp., *Polygonum* spp., *Sinapis* spp. ve *Solanum* spp. gibi bazı türler üzerinde denenilen pendimethalin, S-metalochlor ve fomesafen gibi aktif maddelerinin çıkış öncesi (Byrd, 2019), MSMA (monosodium methanearsonate), bromoxynil, sethoxydim, halosulfuron, clopyralid ve quizalofop gibi aktif maddelerin çıkış sonrası (Maxwell, 2016) kenevir yetiştiriciliğinde kullanılabileceği bildirilmiştir.

Diğer pek çok ülkeden farklı olarak İngiltere'de kenevir üretiminde kullanılmak üzere ruhsatlandırılmış çok sayıda pestisit bulunmaktadır. Endüstriyel kenevirde kimyasal yabancı ot kontrolüne ilişkin sınırlı araştırma olması nedeniyle, Yunanistan'da halihazırda ruhsatlı olan ancak endüstriyel kenevir bitkisi için ruhsatlandırılmayan pendimetalin, s-metolachlor ve aclonifen herbisitleri de kullanılmaktadır.

Dünyanın pek çok ülkesinde olduğu gibi ülkemizde de endüstriyel kenevirde ruhsatlı herhangi bir herbisit bulunmamaktadır.

Dünyada ve Türkiye'de Kenevir Bitkisinde Yabancı Otlara Karşı Kimyasal Mücadele Çalışmalarından Bazı Örnekler

ABD'de çıkış öncesi (pre-emergence) ve çıkış sonrası (postemergence) kullanılan, farklı etki mekanizmalarına sahip bazı herbisitler kenevirde fitotoksite yönüyle test edilmiştir. Çıkış öncesi herbisitler ekim zamanında toprağa uygulanırken çıkış sonrası herbisitler bitki çıkışından 22- 24 gün sonra uygulanmıştır. Çalışmada mesotrione ve trifloxysulfuron içeren herbisitlerin kenevir fidelerine % 78 oranından daha fazla zarar verdiği ve bu nedenle kenevir için uygun olmadığı saptanmıştır. Ancak denemede yer verilen bromoxynil, pendimethalin ve Monosodium methanearsonate (MSMA)'ın kenevir fidelerinde son derece düşük oranda fitotoksiteye neden olduğu saptanmıştır. Çalışma sonucunda bu 3 etken maddeye sahip herbisitlerin kenevirde yabancı ot kontrolü için mükemmel birer aday oldukları belirlenmiştir (Maxwell, 2016).

Kanada'da sera ve tarla çalışmaları şeklinde yürütülen bir çalışma sonucunda; çıkış öncesi uygulanan pendimethalin, S-metalochlor ve fomesafen içeren herbisitlerin düşük fitotoksisite göstermeleri ve kenevir gelişimini kabul edilebilir düzeyde etkilemeleri nedeniyle kenevir üretiminde kullanımlarının uygun olabileceği ortaya konmuştur. Ayrıca çıkış sonrası kullanılan; sethoxydim, bromoxynil, clopyralid ve quizalofop içeren herbisitlerin de endüstriyel kenevir üretimi için uygun olabileceği ancak bu herbisitlerin bazı durumlarda fitotoksite gösterebildikleri ortaya konulmuştur (Byrd, 2019)

Kenevir bitkisinde iki vasküler bitki paraziti oluşur: gövde, dal ve tomurcuklarda küsküt (*Cuscuta*) ve köklerde canavar otu (*Orobancha*). Dallanmış canavar otu (*O. ramosa* L.) kenevir bitkisinin en önemli vasküler bitki parazitiyken, *O. cernua* Loefling ve *O. aegyptiaca* Persoon de kenevir üzerinde görüldüğü kayıtlarda bulunmuştur (Mcpartland, 1996; Tanase vd., 2012).

Canavar otu parazitik bitkisi Avrupa'da önemli derecede bir yabancı ottur. Yaşam döngüsünün büyük bir kısmını yeraltında geçiren ve her bir bitki kenevir tohumlarına iyi yapışan ve yapışkan olan 500.000 toza kadar tohum üretebilmektedir. Kenevir bitkisinin üzerinde, ayrıca Amerika'da ekili ve yabani kenevirde en sık kaydedilmiş olan *Cuscuta campestris* ile küskütün çeşitli türleri de bulunmuştur (McPartland ve ark., 2000).

SONUÇ VE ÖNERİLER

Kenevir bitkisi, yabancı ot rekabeti bakımından üstün performansı, adaptasyon kabiliyeti ve bitki besin elementi bakımından aza kanaat etmesi sebebiyle çevre dostu bir endüstriyel bitkidir. Besin kompozisyonu bakımından hayvan ve insan beslenmesi bakımından oldukça yüksek oranda bir potansiyeli bulunmaktadır. Biyoyakıt, gıda, tekstil, kâğıt, kozmetik, ilaç, inşaat ve otomotiv sektörüne kadar olmak üzere oldukça geniş bir kullanım aralığına sahip olan kenevir bitkisi, petrol ve petrokimyanın kullanıldığı her alanda da kullanılabilen alternatif bir endüstriyel kültür bitkisidir. Narkotik- eğlence maddesi olarak da kullanılabilmesi özelliğinden dolayı 19. asrın ortalarında birçok ülkede üretimi yasaklanan kenevir bitkisi, son yıllarda özellikle endüstriyel kenevir olarak üretimi yapılabilmesi için yeniden önem kazanmaya başlamıştır. Türkiye'de 2019 yılında endüstriyel kenevir üretim amacı ile 19 ilde kenevir tarımına izin verilmiştir. Bu illerde kenevir üretiminin artırılmasına yönelik olarak teknik ve yasal altyapının tamamlanması yönünde önemli çalışmalar yapılmaya devam etmektedir. Birçok kaynakta geleceğin bitkileri arasında gösterilen kenevir bitkisi bilhassa endüstriyel kenevir olarak tarımsal üretimde mutlaka hak ettiği yeri alacak olup geniş kullanım alanı sayesinde ulusların ekonomilerine katkı sağlayacak önemli bir kaynak haline gelecektir.

Ülkemizin çoğu bölgesi kenevir bitkisi tarımı için uygun toprak ve iklim özelliklerine sahiptir. Fakat ülkemizde ilaç ve lif tipi kenevirin üretimi için tescil edilmiş çeşitler mevcut değildir (Şu an sadece iki çeşit tescil aşamasındadır). Bu kapsamda kenevir üretiminin yapılabilmesi için, yetiştiriciliğin yapılacağı bölgenin ekolojik koşullarına çeşitlerin geliştirilmesi ya da dünyanın muhtelif ülkelerindeki endüstriyel kenevir çeşitlerinin tohumlarının ve ülkemizde bulunan yerel varyetelerin tohumlarının toplanarak, öncelikle muhtelif bölgelerde çok yıllık kenevir bitkisinin adaptasyon çalışmalarının başlatılması gereklidir. Bununla birlikte ülkemiz koşullarına uygun çeşitlerin geliştirilmesine yönelik olarak melezleme ve seleksiyon çalışmalarının bir an önce başlatılması, büyük önem taşımaktadır. Farklı amaçlara yönelik olarak kullanılacak kenevir bitkisi çeşitlerinin geliştirilmesi ve bunların sektörün talepleri doğrultusunda iyileştirilme programlarına tabi tutulması büyük önem arz etmektedir. Programdaki başlıca araştırma disiplinleri; bitki fizyolojisi, , ekonomi ve pazar araştırmaları , bitki ıslahı ve yetiştirme tekniği, zararlılar ve hastalık, ülkemizde yeni geliştirilmekte olan otomotiv sektörü için kalıplanmış ve preslenmiş plastik kompozit teknolojisi, hasat ve depolama teknolojisi, kâğıt hamuru teknolojisi, kannabinoid ekstraksiyon teknolojisi gibi saymak mümkündür. Özetle kenevir; çok farklı amaçlı kullanım alanlarına sahip bir bitki türü olarak tohumlarından yemeklik yağ ve farklı besin maddeleri üretimi ve lifinden tekstil ve kâğıt üretim potansiyeline sahip olması ve özellikle kontrollü koşullarda yetiştirildiğinde yılda 5-6 defa hasat yapılarak tedavi amaçlı çok sayıda farmasötik bileşik/kannabinoidler üretilerek sürdürülebilir tarım sistemleri için doğru profile sahip ve potansiyel olarak kârlı bir endüstriyel bitkidir.

KAYNAKLAR

Anonim, (2021). Ordu Ticaret Borsası Kenevir (Kendir) Raporu, 24s. <https://www.ordutb.org.tr/wp-content/uploads/2021/03/Kenevir-Raporu.pdf> (erişim tarihi: 17.01.2025).

Anonim, (2025). Tarım, Lif Bitkileri. T.C. Millî Eğitim Bakanlığı, Ankara. https://megep.meb.gov.tr/mte_program_modul/moduller/Lif%20Bitkileri.pdf (Erişim tarihi: 12.01.2025).

Adamovics, A., Zeverte-Rivza, S., (2015). Industrial hemp (*Cannabis sativa* L.) Productivity and risk assessment in hemp production. The 25th NJF Congress, 16-18 june 2015, Riga, Latvia, 243-248.

Aydoğan, M, Terzi, Y.E., Gizlenci, Ş, Acar, M., Esen, A., Meral, B., (2019). Türkiye’de Kenevir Yetiştiriciliğinin Ekonomik Olarak Yapılabilirliği: Samsun ili Vezirköprü ilçesi örneği. *Anadolu Tarım Bilimleri Dergisi*, 35.

Aytaç, S. (2018). Endüstriyel kenevir gerçeği ve ülkemizdeki durum. Karadeniz Tarımsal Araştırma Enstitüsü Müdürlüğü Endüstriyel Kenevir Gerçeği Paneli, 2-3 Mayıs, Samsun.

Akpınar D, Nizamoğlu A, (2019), International Balkan University - Social Sciences - Osmanlı’dan Cumhuriyet’e Kenevir Üretimi Araştırma Makalesi, <https://doi.org/10.29228/TurkishStudies.23448>, Macedonia.

Basbag, S., Ekinci, R., Yaşar, M. (2020). Kenevirin ekolojik istekleri. *Kenevir (Cannabis sativa L.)*, Palme Yayıncılık, s. 45-60.

Bocsa, I. & M. Karus. (1998). The cultivation of hemp: Botany, varieties, cultivation and harvesting. (translated by Chris Filben) Hemptech, Sebastopol.

Byrd, J. (2019). Industrial hemp (*Cannabis sativa* L.) germination temperatures and herbicide tolerance screening. MSc. Thesis, Crop and Soil Environmental Sciences, Faculty of the Virginia Polytechnic Institute and State University, USA.

Darby, H., Gupta, A., Cummings, E., Cubins, J., Emick, H., Post, J., Ruhl, L, & Ziegler, S. (2016). Industrial Hemp Weed Control Trial. *Northwest Crops & Soils Program*. 92.

Desanlis, F., Cerruti, N., Warner, F. (2013). Hemp Agronomics and Cultivation, Chapter 6. Hemp Industrial Production and Uses (Boulloc, P., Allegret, S., Arnaud, L., eds.), CAB International, London, UK.

Ehrensing, D.T. (1998). Feasibility of industrial hemp production in the United States Pacific Northwest.

https://ir.library.oregonstate.edu/concern/administrative_report_or_publications/j3860729t (Erişim tarihi: 13.01.2025)

Fike, J. (2016). Industrial hemp: Renewed opportunities for an ancient crop. *Critical Reviews in Plant Sciences*, 35 (5-6), 406-424. <https://doi.org/10.1080/07352689.2016.1257842>

GKGM (2025). T.C. Tarım ve Orman Bakanlığı, Gıda ve Kontrol Genel Müdürlüğü, Bitki Koruma Ürünleri Daire Başkanlığı Veri Tabanı. <https://bku.tarimorman.gov.tr/> (Erişim tarihi: 08.01.2025)

Gorchs, G., Lloveras, J., Serrano, L., Cela, S. (2017). Hemp yields and its rotation effects on wheat under rainfed Mediterranean conditions. *Agronomy Journal*, 109 (4), 1551-1560. <https://doi.org/10.2134/agronj2016.11.0676>

Göre, M., Kurt, O. (2021). Bitkisel üretimde yeni bir trend: Kenevir. *International Journal of Life Sciences and Biotechnology*, 4 (1), 138- 157.

Güncan, A., (2009). Yabancı Otlar ve Mücadele Prensipleri. Selçuk Üniversitesi Ziraat Fakültesi yayınları, Konya.

Güncan, A. (2019). Yabancı Otlar ve Mücadele Prensipleri, Selçuk Üniversitesi, Ziraat Fakültesi, KONYA.

Güncan, A. ve Karaca, M. (2023). Yabancı Ot Mücadelesi. Güncellenmiş ve ilaveli beşinci baskı. Akıncı Ofset Matbaa. 368 s. Konya.

Grinspoon, I. (2007). On the future of cannabis as medicine. *Cannabinoids* 2007;2(2):13-15

Hall, J., Bhattarai, S.P., Midmore, D.J. (2014). Effect of industrial hemp (*Cannabis sativa* L.) planting density on weed suppression, crop growth, physiological responses, and fibre yield in the subtropics. *Renewable Bioresources*, 2 (1),1. <https://doi.org/10.7243/2052-6237-2-1>

- Jankauskiene, Z., Gruzdevienė, E., Lazauskas, S. (2014). Potential of industrial hemp (*Cannabis sativa* L.) genotypes to suppress weeds. *Zemdirbyste-Agriculture*, 101 (3), 265-270.
- Jankauskiene, Z., Gruzdevienė, E., Burbulis, N., Maumevičius, E., Layko, I.M. (2015). Investigation of hemp (*Cannabis sativa* L.) crop weediness. *Proceedings of the International Scientific and Practical Conference*, Vol. 2, 120-123. Environment. Technology. Resources. Rezekne, Latvia. <http://journals.rta.lv/index.php/ETR/article/view/272/667>
- Johnson, R. (2018). *Hemp as an Agricultural Commodity*; Congressional Research Service: 7, 5700. Washington, DC, USA.
- Kale, K. (2023). Endüstriyel kenevir (*Cannabis sativa* L.)’de yabancı ot mücadelesi üzerine araştırmalar. Doktora tezi, Erciyes Üniversitesi Fen Bilimleri Enstitüsü, Kayseri.
- Kiely, T., Donaldson, D., Grube, A. (2004). *Pesticide Industry Sales and Usage 2000 and 2001 Market Estimates*. U.S.A Environmental Protection Agency, Washington.
- Labrada, R. (2006). Weed management: a basic component of modern crop production. Chapter 2. *Handbook of Sustainable Weed*.
- Maxwell, B.A. (2016). Effects of Herbicides on Industrial Hemp (*Cannabis sativa* L.) Phytotoxicity, Biomass, and Seed Yield. Masters Theses & Specialist Projects. Paper 1742.
- McPartland, J.M. (1996). A review of Cannabis diseases. *Journal of the International Hemp Association* 3: 19-23.
- McPartland, J.M., Clarke, R.C., & Watson, D.P. (2000). *Hemp diseases and pests: management and biological control: an advanced treatise*. CABI Publishing.
- McPartland, J.M., Cutler, S. & McIntosh, D.J. (2004). Hemp Production in Aotearoa. *Journal of Industrial Hemp* 9(1): 105-115.
- O’Donovan, J.T., Blackshaw, R.E., Harker, K.N., Clayton, G.W, Moyer, J.R., Dossall L.M., Maurice, D.C., Turkington, T.K. (2007). Integrated approaches to managing weeds in spring-sow
- ONTARIO (2024a). *Agronomy guide for field crops, Chapter 7. Other Crops, Hemp*. p.179-180, <https://www.ontario.ca/files/2022-10/omafraagronomy-guide-for-field-crops-chapter-7-en-2022-10-13.pdf> (Erişim tarihi: 10.01.2025).
- ONTARIO (2024b). *Growing Industrial Hemp in Ontario*. ISSN 1198-712X <https://files.ontario.ca/omafra-growing-industrial-hemp-in-ontario-22-020-en2023-07-21.pdf> (Erişim tarihi: 11.01.2025).
- Özer, Z., Önen, H., Tursun, N., Uygur, F.N. (1999). Türkiye’nin Bazı Önemli Yabancı Otları. Gaziosmanpaşa Üniversitesi, Ziraat Fakültesi Yayınları, No:38 Kitap seri No:16, Tokat.
- Özer, Z., Kadioğlu, İ., Önen, H., Tursun, N., (2001). Herboloji (Yabancı Ot Bilimi) Gaziosmanpaşa Üniversitesi Ziraat Fakültesi Yayınları, No:20 Kitap seri No:10 Tokat.
- Poisa, L., & Adamovics, A. (2010). Hemp (*Cannabis sativa* L.) as an environmentally friendly energy plant. *Scientific Journal of Riga Technical University. Environmental and Climate Technologies*, 5 (1), 80-85. <https://doi.org/10.2478/v10145-010-0038-z>
- Rehman, M.S.U., Rashid, N., Saif, A., Mahmood, T., Han, J.I. (2013). Potential of bioenergy production from industrial hemp (*Cannabis sativa*): Pakistan perspective. *Renewable and Sustainable Energy Reviews*, 18, 154-164. <https://doi.org/10.1016/j.rser.2012.10.019>
- Sandler, L. N., Beckerman, J. L., Whitford, F., and Gibson, K. A. (2019). Cannabis as conundrum. *Crop Prot.* 117, 37–44. doi: 10.1016/j.cropro.2018.11.003

- Schultes, R., Joyce, C., Curry, S.H. (1970). Random thoughts and queries on the botany of cannabis. *Botany and Chemistry of Cannabis* 1-38.
- Small, E. (1975). The case of the curious “*Cannabis*”. *Economic Botany* 29, 254.
- Struik, P.C., Amaducci, S., Bullard, M.J., Stutterheim, N.C., Venturi, G., Cromack H.T.H. (2000). Agronomy of fibre hemp (*Cannabis sativa* L.) in Europe. *Industrial Crops and Products*, 11 (2-3), 107-118. [https://doi.org/10.1016/S0926-6690\(99\)00048-5](https://doi.org/10.1016/S0926-6690(99)00048-5)
- TİGEM, 2019. Kenevir üretim çalışmalarına ilişkin kamuoyuna bilgilendirme. <https://www.tigem.gov.tr/haber/haberdetay/9079d190-1793-44ec-958b-77d150b82094> Erişim: 16.01.2025.
- TÜİK, (2020). Türkiye istatistik enstitüsü. <http://www.tuik.gov.tr/Start.do> Erişim: 15.09.2020.
- TÜİK, (2023). Türkiye istatistik enstitüsü. <http://www.tuik.gov.tr/Start.do> Erişim: 11.10.2023.
- Tunçdemir, Ü. (1987). Investigations on the Control Possibilities of Stem Nematode *Ditylenchus dipsaci* Kühn.) Causing Damage to Hemps in the Black Sea Region. *Plant Protection Bulletin*, 27(3).
- Uygur, F.N., Koch, W., Walte, H., (1984). Plits, Yabancı ot bilimine giriş (kurs notu) 1984 / 2 (1) Josef Margraf, Stuttgart.
- Van der Werf, H. M. G., van Geel, W. C. A., van Gils, L. J. C., & Haverkort, A. J. (1995). Nitrogen fertilization and row width affect self-thinning and productivity of fibre hemp (*Cannabis sativa* L.). *Field Crops Research*, 42, 27-37. [https://doi.org/10.1016/0378-4290\(95\)00017-k](https://doi.org/10.1016/0378-4290(95)00017-k)
- Van der Werf, H.M.G. (2002). Hemp production in France. *Journal of Industrial Hemp*, 7 (2), 105-109. https://doi.org/10.1300/J237v07n02_12
- Vencill, W.K., Nichols, R.L., Webster, T.M., Soteris, J.K., Mallory-Smith, C., Burgos, N.R., Johnson, W.G., McClelland, M.R. (2012). Herbicide resistance: Toward an understanding of resistance development and the impact of herbicide-resistant crops. *Weed Science* 60 (SP1), 2-30. <https://doi.org/10.1614/WS-D-11-00206.1>
- Vera, C.L., Woods, S.M., Raney, J.P. (2006). Seeding rate and row spacing effect on weed competition, yield and quality of hemp in the Parkland region Saskatchewan. *Canadian Journal of Plant Science*, 86 (5), 911-915. <https://doi.org/911-915.10.4141/P05-177>

**VALORISATION OF BIOPESTICIDE FROM DITTRICHIA VISCOSA (L.)
GREUTER 1973 FROM ALGERIAN ARID REGION**

Deghiche-Diab Nacima

Scientific and Technical Research Center on Arid Regions (CRSTRA), 07000 Biskra, Algeria

<https://orcid.org/0000-0003-4544-2083>

Saad Somia

Scientific and Technical Research Center on Arid Regions (CRSTRA), 07000 Biskra, Algeria

Belhamra Youcef Islem

Plant Production Biotechnology Laboratory, Department of Biotechnology and Agro-Ecology, Faculty of Natural and Life Sciences, Blida 1 University, PoBox. 270, Soumaa road, Blida, Algeria.

Ouamene Tarek Abdelmoneim

Scientific and Technical Research Center on Arid Regions (CRSTRA), 07000 Biskra, Algeria

Karoune Samira

Scientific and Technical Research Center on Arid Regions (CRSTRA), 07000 Biskra, Algeria

Djoudi Madjed

Scientific and Technical Research Center on Arid Regions (CRSTRA), 07000 Biskra, Algeria

Kechebar Mohamed Seif Allah

Scientific and Technical Research Center on Arid Regions (CRSTRA), 07000 Biskra, Algeria

Djazouli Zahr Eddine

Plant Production Biotechnology Laboratory, Department of Biotechnology and Agro-Ecology, Faculty of Natural and Life Sciences, Blida 1 University, PoBox. 270, Soumaa road, Blida, Algeria.

Abstract

To investigate the effects of *Dittrichia viscosa* L. Greuter 1973 extract as a biopesticide on beneficial entomofauna in a greenhouse setting in the Biskra region, our study was conducted at the experimental site (CRSTRA). The aerial parts of *D. viscosa* were collected, dried, and analyzed in laboratory. The antioxidant potential of the plant extract was assessed by 2, 2-diphenyl-1-picrylhydrazyl (DPPH), alkaline DMSO superoxide, O-phenanthroline chelating (Phen) and iron reducing power (RP) methods. Enzyme inhibitory was studied using the

iodine/potassium iodide method. A 200 m² greenhouse was established in November 2021 and planted with tomato (Cecilia), divided into four blocks, each equipped with sticky traps for monitoring entomofauna from May 1st to June 1st. The pulverized *D. viscosa* extract was applied at different concentrations (D1=5 ml/l; D2=10 ml/l; D3=15 ml/l) every 10 days. Analysis of the hydro-methanolic extract of *D. viscosa* revealed significant antioxidant activities and effective α -amylase inhibition, indicating its potential as a free radical scavenger and a useful therapeutic agent for addressing radical-related pathological damage. Additionally, the extract's biopesticide effects on beneficial insects were evaluated. Results showed a total richness of 75 species across 7 orders and 44 families, with the most abundant species belonging to Hymenoptera (37 species), Coleoptera (14 species), and Diptera (11 species). The predominant categories included parasitoids (29%), predators (24%), pests (15%), and pollinators (11%). As a biopesticide, the extract proved effective in reducing pest populations by 50% at the highest concentration (15 ml/L). The Pielou evenness index values (H5 ml/L = 0.8787, H10 ml/L = 0.8506, H15 ml/L = 0.836, compared to the control = 0.4179) indicated a nearly homogeneous distribution of individuals among species. The Shannon diversity index suggested that increasing concentrations of *D. viscosa* extract (E5 ml/L = 0, E10 ml/L = 1.33, and E15 ml/L = 0.5623) did not significantly impact the presence of pollinator species in the tomato greenhouse in the Biskra region.

Keywords: naturel ennemis, Biopesticides, *Dittrichia viscosa* (L.) Greuter 1973, Antioxidant, arid regions.

Species Index: *Dittrichia viscosa* L. Greuter 1973, *Solanum lycopersicum* L.



2nd International Congress on

SUSTAINABLE AGRICULTURE

March 04-05, 2025 / Iğdır University, Türkiye



ISBN: 979-8-89695-046-2